

Wald Chapter 4: Einstein's Equation

Source: R. M. Wald, General Relativity, Chapter 4; selected worked problems and related solutions.

CONTENTS

- 1 Chapter 4, Problem 1: Charge Conservation
- 2 Chapter 4, Problem 2: Hodge Duality and Maxwell Theory
- 3 Chapter 4, Problem 3: Gravitomagnetism and Frame Dragging
- 4 Chapter 4, Problem 4: Quadratic Ricci Tensor
- 5 Chapter 4, Problem 5: A Superpotential for a Conserved Tensor
- 6 Chapter 4, Problem 7: Energy in Linearised Gravity
- 7 Chapter 4, Problem 8: Radiation from an Oscillating Spring
- 8 Chapter 4, Problem 9: Orbital Decay of a Binary System

Problem prompts are paraphrased from Robert M. Wald, *General Relativity* (University of Chicago Press, 1984); the equations, notation, numbering, and guidance follow the source. See the official publisher page.

1 Chapter 4, Problem 1: Charge Conservation

Show that Maxwell's equation (4.3.12) implies strict charge conservation,

$$\nabla_a j^a = 0. \quad (1.1)$$

Solution. Although it is a lot easier to work in form language, but we shall stick with covariant derivatives here. First, note that $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = \nabla_\mu A_\nu - \nabla_\nu A_\mu$. The equation of motion becomes

$$\nabla^\mu \nabla_\mu A_\nu - \nabla^\mu \nabla_\nu A_\mu = -4\pi j_\nu. \quad (1.2)$$

Taking covariant derivative on both side, the LHS is

$$\nabla^\nu \nabla^\mu \nabla_\mu A_\nu - \nabla^\nu \nabla^\mu \nabla_\nu A_\mu = [\nabla^\nu, \nabla^\mu] \nabla_\mu A_\nu. \quad (1.3)$$

Using (3.2.21), we have

$$\nabla^\nu \nabla^\mu \nabla_\mu A_\nu - \nabla^\nu \nabla^\mu \nabla_\nu A_\mu = R_{\nu\mu}{}^{\mu\delta} \nabla_\delta A^\nu + R_{\nu\mu}{}^{\nu\delta} \nabla_\delta A^\mu. \quad (1.4)$$

Relabelling the second term, and using the antisymmetric property of Riemann tensor we find that the LHS vanishes identically. Therefore, we find

$$\nabla_\mu j^\mu = 0. \quad (1.5)$$

2 Chapter 4, Problem 2: Hodge Duality and Maxwell Theory

1. Let α be a p -form on an oriented n -dimensional manifold with metric g_{ab} . Thus $\alpha_{a_1 \dots a_p}$ is totally antisymmetric. Define its dual by

$$\star \alpha_{b_1 \dots b_{n-p}} = \frac{1}{p!} \alpha^{a_1 \dots a_p} \epsilon_{a_1 \dots a_p b_1 \dots b_{n-p}}, \quad (2.1)$$

where $\epsilon_{a_1 \dots a_n}$ is the natural volume element, fixed up to sign by equation (B.2.9). Prove

$$\star \star \alpha = (-1)^{s+p(n-p)} \alpha, \quad (2.2)$$

with s the number of minus signs in the signature of g_{ab} .

Solution.

$$\begin{aligned}
 (\star \star \alpha)_{c_1 \dots c_p} &= \frac{1}{(n-p)!} (\star \alpha)^{b_1 \dots b_{n-p}} \epsilon_{b_1 \dots b_{n-p} c_1 \dots c_p} \\
 &= \frac{1}{p!} \frac{1}{(n-p)!} \alpha_{a_1 \dots a_p} \epsilon^{a_1 \dots a_p b_1 \dots b_{n-p}} \epsilon_{b_1 \dots b_{n-p} c_1 \dots c_p} \\
 &= \frac{1}{p!} \frac{1}{(n-p)!} \alpha_{a_1 \dots a_p} (-1)^{s+p(n-p)} p! (n-p)! \delta_{c_1}^{[a_1} \dots \delta_{c_p}^{a_p]} \\
 &= (-1)^{s+p(n-p)} \alpha_{c_1 \dots c_p}.
 \end{aligned} \tag{2.3}$$

2. Rewrite Maxwell's equations (4.3.12) and (4.3.13) in differential-form notation as

$$d \star F = 4\pi \star j, \quad dF = 0. \tag{2.4}$$

Explain how Stokes's theorem applied to the first equation gives Gauss's law on a three-dimensional hypersurface Σ with boundary S :

$$\int_{\Sigma} \star j = \frac{1}{4\pi} \int_S \star F, \tag{2.5}$$

together with

$$-\int_{\Sigma} \star j = -\int_{\Sigma} j^a t_a d\Sigma = e, \quad -\int_S \star F = \int_S E^a n_a dA, \tag{2.6}$$

where t^a is the unit normal to Σ and $E_a = F_{ab} t^b$.

Solution. First, we start with the covariant derivative form,

$$\nabla_{\mu} F^{\mu\nu} = \partial_{\mu} F^{\mu\nu} + \Gamma^{\mu}_{\mu\sigma} F^{\sigma\nu} + \Gamma^{\nu}_{\mu\sigma} F^{\mu\sigma}, \tag{2.7}$$

where the last term vanishes. Then, note that

$$\Gamma^{\mu}_{\mu\sigma} = \frac{1}{2} g^{\mu\rho} \partial_{\sigma} g_{\mu\rho} = \frac{1}{2} \partial_{\sigma} \text{tr} \log g_{\mu\rho} = \frac{1}{2} \partial_{\sigma} \log g = \partial_{\sigma} \log \sqrt{|g|} = \frac{1}{\sqrt{|g|}} \partial_{\sigma} \sqrt{|g|}, \tag{2.8}$$

which follows

$$\nabla_{\mu} F^{\mu\nu} = \frac{1}{\sqrt{|g|}} \partial_{\mu} \left(\sqrt{|g|} F^{\mu\nu} \right) = -4\pi j^{\nu}. \tag{2.9}$$

To see that this is equivalent to the alternative formulation, we compute

$$d \star F = \frac{1}{4} \partial_{\lambda} (F_{\mu\nu} \epsilon^{\mu\nu\rho\sigma}) dx^{\lambda} \wedge dx^{\rho} \wedge dx^{\sigma}, \tag{2.10}$$

while

$$\star j = \frac{1}{3!} j_{\mu} \epsilon^{\mu}_{\nu\rho\sigma} dx^{\nu} \wedge dx^{\rho} \wedge dx^{\sigma}. \tag{2.11}$$

Using $\epsilon_{\mu\nu\rho\sigma} = \sqrt{|g|} \varepsilon_{\mu\nu\rho\sigma}$, where ε is the totally antisymmetric tensor, we find

$$d \star F = 4\pi \star j \iff \frac{1}{4} \partial_\lambda \left(\sqrt{|g|} F^{\mu\nu} \right) \varepsilon_{\mu\nu\rho\sigma} = 4\pi \frac{1}{3!} \sqrt{|g|} j^\mu \varepsilon_{\mu\lambda\rho\sigma}. \quad (2.12)$$

Contracting both side with $\varepsilon^{\delta\lambda\rho\sigma}$, we find

$$\frac{1}{4} \partial_\lambda \left(\sqrt{|g|} F^{\mu\nu} \right) 2! \left(\delta_\mu^\delta \delta_\nu^\lambda - \delta_\mu^\lambda \delta_\nu^\delta \right) = 4\pi \frac{1}{3!} \sqrt{|g|} j^\mu 3! \delta_\mu^\delta \iff \frac{1}{\sqrt{|g|}} \partial_\mu \left(\sqrt{|g|} F^{\mu\nu} \right) = -4\pi j^\nu. \quad (2.13)$$

Therefore, we have shown

$$\nabla_\mu F^{\mu\nu} = -4\pi j^\nu \iff d \star F = 4\pi \star j. \quad (2.14)$$

Now, using stokes theorem, we have

$$\int_\Sigma \star j = \frac{1}{4\pi} \int_\Sigma d \star F = \frac{1}{4\pi} \int_{\partial\Sigma} \star F. \quad (2.15)$$

If we take the Σ to be spacelike, then n_a is a timelike unit vector, and hence the above reduces to Gauss's law.

3 Chapter 4, Problem 3: Gravitomagnetism and Frame Dragging

1. Derive equation (4.4.24).

Solution. We start with

$$\partial^a \partial_a \bar{\gamma}_{0\mu} = 16\pi J_\mu. \quad (3.1)$$

Identifying $A_\mu \equiv -T_{ab} t^a$, where $t^a = (\partial/\partial x^0)^a$, we find that the equation reduces to Maxwell's equation. Now, we shall evaluate the geodesic equation while neglecting the time derivative of the metric. In this case, the perturbed metric is given by

$$\bar{\gamma}_{0\mu} = \bar{\gamma}_{\mu 0} = -4A_\mu, \quad \bar{\gamma} = \bar{\gamma}^0_0 = 4A_0, \quad (3.2)$$

and hence

$$\gamma_{00} = -2A_0, \quad \gamma_{0i} = 4A_i. \quad (3.3)$$

To first order in γ , the Christoffel connection is given by

$$\Gamma^\mu_{\rho\sigma} = \frac{1}{2} (\partial_\rho \gamma^\mu_\sigma + \partial_\sigma \gamma^\mu_\rho - \partial^\mu \gamma_{\rho\sigma}). \quad (3.4)$$

In our case, the relevant connections are

$$\Gamma^i_{00} = \partial^i A_0 = E^i, \quad \Gamma^i_{0j} = \Gamma^i_{j0} = 2(\partial_i A_j - \partial_j A_i), \quad (3.5)$$

which gives

$$\ddot{x}_i = -E_i - 4(\partial_i A_j - \partial_j A_i) \dot{x}^j. \quad (3.6)$$

This is precisely

$$\ddot{\mathbf{x}} = -\mathbf{E} - 4\mathbf{v} \times \mathbf{B}. \quad (3.7)$$

2. Consider a spherical shell of mass M and radius R , with $M \ll R$, rotating slowly with angular velocity $\boldsymbol{\omega}$. Show that the gravitational electric and magnetic fields inside it are

$$\mathbf{E} = 0, \quad \mathbf{B} = \frac{2M}{3R}\boldsymbol{\omega}. \quad (3.8)$$

Solution. The current is given by

$$J^0 = \rho = \frac{M}{4\pi R^2}\delta(r - R), \quad \mathbf{J} = \frac{M}{4\pi R^2}\delta(r - R)\boldsymbol{\omega} \times \mathbf{r}. \quad (3.9)$$

By Gauss's law, it is easy to see that $\mathbf{E} = 0$ inside the shell. For the gravitational magnetic field, we use the Green's function method, which gives

$$\mathbf{A}(\mathbf{x}) = \int \frac{J(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} d^3\mathbf{x}' = \frac{M}{4\pi R^2}\boldsymbol{\omega} \times \int \frac{\delta(r' - R)\mathbf{x}'}{|\mathbf{x} - \mathbf{x}'|} d^3\mathbf{x}' = \frac{M}{4\pi R^2}\boldsymbol{\omega} \times \int d\Omega' \frac{R^3 \hat{\mathbf{r}}'}{|\mathbf{x} - R\hat{\mathbf{r}}'|}. \quad (3.10)$$

Since $M \ll R$, we can carry out dipole expansion, which gives

$$\mathbf{A}(\mathbf{x}) = \frac{MR}{4\pi}\boldsymbol{\omega} \times \int d\Omega' \left[\frac{\hat{\mathbf{r}}'}{R} + \frac{\hat{\mathbf{r}}'(\mathbf{x} \cdot \hat{\mathbf{r}}')}{R^2} \right] = \frac{M}{3R}\boldsymbol{\omega} \times \mathbf{x}, \quad (3.11)$$

where we have used the identities

$$\int \hat{r}_i \hat{r}_j d\Omega = \frac{4\pi}{3} \delta_{ij}.$$

and

$$\int \hat{\mathbf{r}} d\Omega = 0.$$

Therefore, we find

$$\mathbf{B} = \frac{2M}{3R}\boldsymbol{\omega}. \quad (3.12)$$

3. An observer at rest at the shell's centre parallelly propagates a vector S^a along the observer's geodesic world line, with $S^a u_a = 0$. Show that its inertial components $\mathbf{S}$ precess according to

$$\frac{d\mathbf{S}}{dt} = \boldsymbol{\Omega} \times \mathbf{S}, \quad \boldsymbol{\Omega} = 2\mathbf{B} = \frac{4}{3} \frac{M}{R}\boldsymbol{\omega}. \quad (3.13)$$

Interpret this as the rotating shell dragging the local inertial frame, in the sense discussed in connection with Mach's principle.

Solution.

$$u^b \nabla_b S^a = 0 \implies \frac{dS^a}{dt} + u^\nu \Gamma^\mu_{\nu\rho} S^\rho = 0. \quad (3.14)$$

Note that we have already computed the Christoffel symbol, and hence we have

$$\frac{dS^i}{dt} = -\Gamma^i_{0j} S^j = -2(\partial_i A_j - \partial_j A_i) S^j \implies \frac{d\mathbf{S}}{dt} = \frac{4M}{3R} \boldsymbol{\omega} \times \mathbf{S}. \quad (3.15)$$

Therefore, $\mathbf{S}$ precesses around the rotating axis of the spherical shell with an angular velocity Ω .

4 Chapter 4, Problem 4: Quadratic Ricci Tensor

Starting from equation (3.4.5) for R_{ab} , substitute

$$g_{ab} = \eta_{ab} + \gamma_{ab} \quad (4.1)$$

and retain exactly the terms quadratic in γ_{ab} . In this way, derive the formula (4.4.51) for $R_{ab}^{(2)}$.

Solution. See my worked solution to David Tong GR Problem Sheet 4, Question 4.

5 Chapter 4, Problem 5: A Superpotential for a Conserved Tensor

Let T_{ab} be symmetric and conserved in Minkowski spacetime:

$$T_{ab} = T_{ba}, \quad \partial^a T_{ab} = 0. \quad (5.1)$$

Show that there is a tensor U_{acbd} with symmetries

$$U_{acbd} = U_{[ac]bd} = U_{ac[bd]} = U_{bdac} \quad (5.2)$$

such that

$$T_{ab} = \partial^c \partial^d U_{acbd}. \quad (5.3)$$

Hint: If a Minkowski-space vector field obeys $\partial_a v^a = 0$, the converse of the Poincaré lemma applied to the three-form $\epsilon_{abcd} v^d$ gives an antisymmetric field $s^{ab} = -s^{ba}$ such that

$$v^a = \partial_b s^{ab}. \quad (5.4)$$

First use this fact to obtain

$$T_{ab} = \partial^c W_{cab}, \quad W_{cab} = W_{[ca]b}, \quad (5.5)$$

and then use $\partial^c W_{c[ab]} = 0$ to reach the required result.

Solution. Let

$$J = \frac{1}{3!} v^d \epsilon_{abcd} dx^a \wedge dx^b \wedge dx^c, \quad (5.6)$$

then

$$dJ = 0 \iff \partial^a v_a = 0. \quad (5.7)$$

By Poincaré lemma, we have

$$dJ \iff J = d\omega, \quad \omega \in \Omega^2(\mathbb{R}^{1,n-1}), \quad (5.8)$$

which is equivalent to

$$v^d \epsilon_{abcd} = \partial_a \omega_{bc} \implies v^a = \partial_b \omega_{cd} \epsilon^{abcd} = \partial_b s^{ab}, \quad (5.9)$$

where we defined $s^{ab} \equiv \omega_{cd} \epsilon^{abcd}$. Now take $\partial^a T_{ab} = 0$, where we can treat b as a dummy label. This gives

$$T_{ab} = \partial^c W_{cab}, \quad W_{cab} = W_{[ca]b}. \quad (5.10)$$

Now using the symmetric property of T^{ab} , we have $\partial^c W_{c[ab]}$. Now, we can treat $[ab]$ as dummy labels and apply the result from Poincaré lemma once more, which gives

$$W_{c[ab]} = \partial^d V_{dc[ab]}, \quad V_{dc[ab]} = V_{[dc][ab]}. \quad (5.11)$$

Defining $U_{acbd} \equiv -V_{acbd} - V_{bdac}$, we find the required tensor.

6 Chapter 4, Problem 7: Energy in Linearised Gravity

1. Show that the total energy E of equation (4.4.55) is independent of time: the same value is obtained by integrating over a time translate Σ' of Σ .

Solution. Since, $\partial^a t_{ab} = 0$, we have

$$\frac{dE}{dt} = \int_{\Sigma} \frac{dt_{00}}{dt} d^3x = \int_{\Sigma} \partial^i t_{i0} d^3x = 0. \quad (6.1)$$

2. Let ξ_a generate a gauge transformation which is zero outside a bounded spatial region. Prove

$$E[\gamma_{ab}] = E[\gamma_{ab} + 2\partial_{(a}\xi_{b)}]. \quad (6.2)$$

Compare both sides with

$$E[\gamma_{ab} + 2\partial_{(a}\xi'_{b)}], \quad (6.3)$$

where ξ'_a agrees with ξ_a near Σ but vanishes near another hyperplane Σ' .

Solution. Since ξ'_a vanishes at Σ' , we have

$$E[\gamma_{ab}] = \int_{\Sigma'} t_{00}[\gamma_{ab}] d^3x = \int_{\Sigma'} t_{00}[\gamma_{ab} + 2\partial_{(a}\xi'_{b)}] d^3x. \quad (6.4)$$

By conservation of total energy, this is equivalent to

$$E[\gamma_{ab}] = \int_{\Sigma} t_{00}[\gamma_{ab} + 2\partial_{(a}\xi'_{b)}] d^3x. \quad (6.5)$$

Finally, since ξ'_a agrees with ξ_a near Σ , we have

$$E[\gamma_{ab}] = E[\gamma_{ab} + 2\partial_{(a}\xi_{b)}]. \quad (6.6)$$

7 Chapter 4, Problem 8: Radiation from an Oscillating Spring

Two point particles, each of mass M , are fixed to opposite ends of a spring with spring constant K and set oscillating. Using the quadrupole approximation (4.4.58), determine the fraction of their oscillation energy radiated in one cycle.

Solution. The masses are oscillating with frequency ω and amplitude A , and separated by a distance $2L$ at equilibrium. Then, the mass density is given by

$$T_{00}(\mathbf{x}, t) = \rho(\mathbf{x}, t) = M\delta(y)\delta(z)[\delta(x - L - A\cos\omega t) + \delta(x + L + A\cos\omega t)]. \quad (7.1)$$

The only nonzero quadrupole moment of energy is given by

$$I_{11} = 2M(L + A\cos\omega t)^2 \equiv 2Mx^2(t), \quad (7.2)$$

and hence its traceless part:

$$Q_{ij} = \begin{pmatrix} \frac{4}{3}Mx^2 & 0 & 0 \\ 0 & -\frac{2}{3}Mx^2 & 0 \\ 0 & 0 & -\frac{2}{3}Mx^2 \end{pmatrix}. \quad (7.3)$$

Taking the third order time derivative

$$\ddot{x}^2(t) = 2A\omega^3(2A\sin 2\omega t + L\sin\omega t) \quad (7.4)$$

and we find the power of emission to be

$$\mathcal{P} = \frac{32}{15}GM^2A^2\omega^6(2A\sin 2\omega t + L\sin\omega t)^2. \quad (7.5)$$

Averaging over one period, we find

$$\langle \mathcal{P} \rangle = \frac{16}{15}GM^2A^2\omega^6(4A^2 + L^2) \quad (7.6)$$

The total energy we started with is

$$E = M\omega^2A^2, \quad (7.7)$$

and hence

$$\frac{\Delta E}{E} = \frac{32\pi}{15} GM\omega^3 (4A^2 + L^2). \quad (7.8)$$

8 Chapter 4, Problem 9: Orbital Decay of a Binary System

Two negligibly small stars, each of mass M , move in a nearly Newtonian circular orbit of radius R about one another. Assuming equation (4.4.58) applies, compute the rate at which gravitational radiation increases their orbital frequency.

Solution. See my related worked solution to David Tong GR Problem Sheet 4, Question 6.