

Renormalization Group for ϕ - χ Theory — Srednicki Q28.3

Source: M. Srednicki, *Quantum Field Theory*, Chapter 28 (Renormalization Group), Problem 28.3.

1 Renormalization Group for ϕ - χ Theory, Srednicki Q28.3

Consider the Lagrangian density

$$\begin{aligned}\mathcal{L} = & -\frac{1}{2}Z_\phi\partial^\mu\phi\partial_\mu\phi - \frac{1}{2}Z_m m^2\phi^2 + Y\phi \\ & -\frac{1}{2}Z_\chi\partial^\mu\chi\partial_\mu\chi - \frac{1}{2}Z_M M^2\chi^2 \\ & + \frac{1}{6}Z_g g\mu^{\varepsilon/2}\phi^3 + \frac{1}{2}Z_h h\mu^{\varepsilon/2}\phi\chi^2\end{aligned}\quad (1.1)$$

in $d = 6 - \varepsilon$ dimensions, where ϕ and χ are real scalar fields, and Y is adjusted to make $\langle 0|\phi(x)|0\rangle = 0$. (Why is no such term needed for χ ?)

1. Compute the one-loop contributions to each of the Z 's in the $\overline{\text{MS}}$ renormalization scheme.

Solution. The self-interaction of ϕ^3 theory is already computed in the on-shell renormalisation scheme. We shall switch to $\overline{\text{MS}}$ renormalisation scheme. Let us first go back to the two-point correlation function at one-loop level,

$$\Pi_\phi(k^2) \sim \frac{\alpha_g}{2} \left[-(2/\varepsilon + 1)(m^2 + k^2/6) + \int_0^1 dx D \ln(D/\mu^2) \right] - (Ak^2 + Bm^2) + \mathcal{O}(\alpha_g^2). \quad (1.2)$$

However, we also have a contribution by replacing the internal propagators by ϕ with χ , which is equivalent to replacing the vertices. Therefore, the full propagator in this case is

$$\Pi_\phi(k^2) \sim \frac{g^2 + h^2}{2(4\pi)^3} \left[-(2/\varepsilon + 1)(m^2 + k^2/6) + \int_0^1 dx D \ln(D/\mu^2) \right] - (Ak^2 + Bm^2) + \mathcal{O}(\alpha_g^2)$$

Now, we let A and B absorb only the UV divergence part of the theory. That is

$$A = -\frac{g^2 + h^2}{6(4\pi)^3} \frac{1}{\varepsilon} + \mathcal{O}(g^2), \quad B = -\frac{g^2 + h^2}{(4\pi)^3} \frac{1}{\varepsilon} + \mathcal{O}(g^2). \quad (1.4)$$

The story for vertices is more complicated, and we shall postpone that to later.

Let us compute the exact propagator for χ fields. Since the $\Pi_\phi(k^2)$ is of order g^2 , to one loop we simply have one ϕ and one χ propagator connecting two vertices. That is, we simply replace one of the g by h , and Z_ϕ, Z_m by Z_χ, Z_M respectively in (1.3), which gives

$$D = -\frac{gh}{3(4\pi)^3} \frac{1}{\varepsilon} + \mathcal{O}(h^2), \quad E = -\frac{gh}{(4\pi)^3} \frac{1}{\varepsilon} + \mathcal{O}(h^2), \quad (1.5)$$

where $D = Z_\chi - 1$, and $E = Z_M - 1$. Note that the factor of $1/6$ becomes $1/3$ by breaking the internal line exchange symmetry.

For the three-point ϕ^3 vertex, we had

$$V_3(k_1, k_2, k_3)/g \sim 1 + \alpha_g \left[\frac{1}{\varepsilon} + \frac{1}{2} \int dF_3 \ln(\mu^2/D) + \dots \right] + C + \mathcal{O}(\alpha_g^2), \quad (1.6)$$

where $C = Z_g - 1$. However, we could replace the ϕ^3 vertices with $\phi\chi\chi$. Therefore, we should have

$$\begin{aligned} V_3(k_1, k_2, k_3)/g \sim 1 + \frac{h^3/g}{(4\pi)^3} \left[\frac{1}{\varepsilon} + \frac{1}{2} \int dF_3 \ln(\mu^2/D) + \dots \right] \\ + \frac{g^2}{(4\pi)^3} \left[\frac{1}{\varepsilon} + \frac{1}{2} \int dF_3 \ln(\mu^2/D) + \dots \right] + C + \mathcal{O}(g^4), \end{aligned} \quad (1.7)$$

which gives

$$C = -\frac{1}{\varepsilon} \left(\frac{g^2}{(4\pi)^3} + \frac{h^3/g}{(4\pi)^3} \right) + \mathcal{O}(g^2), \quad (1.8)$$

where we have assumed $g \sim h$.

For the three-point vertex, we could also have two types of vertices. One with three $\phi\chi\chi$ vertices and one with two $\phi\chi\chi$ vertices and a ϕ^3 vertex. Therefore, we have

$$\begin{aligned} V_3(k_1, k_2, k_3)/h \sim 1 + \frac{gh}{(4\pi)^3} \left[\frac{1}{\varepsilon} + \frac{1}{2} \int dF_3 \ln(\mu^2/D) + \dots \right] \\ + \frac{h^2}{(4\pi)^3} \left[\frac{1}{\varepsilon} + \frac{1}{2} \int dF_3 \ln(\mu^2/D) + \dots \right] + F + \mathcal{O}(g^4), \end{aligned} \quad (1.9)$$

where $F = Z_h - 1$. Hence, we find

$$F = -\frac{1}{\varepsilon} \left(\frac{h^2}{(4\pi)^3} + \frac{gh}{(4\pi)^3} \right) + \mathcal{O}(h^2). \quad (1.10)$$

2. The bare couplings are related to the renormalised ones via

$$\begin{aligned} g_0 &= Z_\phi^{-3/2} Z_g g \mu^{\varepsilon/2}, \\ h_0 &= Z_\phi^{-1/2} Z_\chi^{-1} Z_h h \mu^{\varepsilon/2}. \end{aligned} \quad (1.11)$$

Define

$$\begin{aligned} G(g, h, \varepsilon) &= \sum_{n=1}^{\infty} G_n(g, h) \varepsilon^{-n} \equiv \ln \left(Z_\phi^{-3/2} Z_g \right), \\ H(g, h, \varepsilon) &= \sum_{n=1}^{\infty} H_n(g, h) \varepsilon^{-n} \equiv \ln \left(Z_\phi^{-1/2} Z_\chi^{-1} Z_h \right). \end{aligned} \quad (1.12)$$

By requiring g_0 and h_0 to be independent of μ , and by assuming that $dg/d\mu$ and $dh/d\mu$ are finite as $\varepsilon \rightarrow 0$, show that

$$\begin{aligned}\mu \frac{dg}{d\mu} &= -\frac{1}{2}\varepsilon g + \frac{1}{2}g \left(g \frac{\partial G_1}{\partial g} + h \frac{\partial G_1}{\partial h} \right), \\ \mu \frac{dh}{d\mu} &= -\frac{1}{2}\varepsilon h + \frac{1}{2}h \left(g \frac{\partial H_1}{\partial g} + h \frac{\partial H_1}{\partial h} \right).\end{aligned}\tag{1.13}$$

Solution. From (1.11), we have

$$\ln g_0 = \ln g + \frac{\varepsilon}{2} \ln \mu + G(g, h, \varepsilon),\tag{1.14}$$

$$\ln h_0 = \ln h + \frac{\varepsilon}{2} \ln \mu + H(g, h, \varepsilon).\tag{1.15}$$

Since the bare coupling constant is fixed for any values of μ , we have

$$0 = \frac{1}{g} \frac{dg}{d \ln \mu} + \frac{\varepsilon}{2} + \frac{\partial G}{\partial h} \frac{dh}{d \ln \mu} + \frac{\partial G}{\partial g} \frac{dg}{d \ln \mu},\tag{1.16}$$

$$0 = \frac{1}{h} \frac{dh}{d \ln \mu} + \frac{\varepsilon}{2} + \frac{\partial H}{\partial h} \frac{dh}{d \ln \mu} + \frac{\partial H}{\partial g} \frac{dg}{d \ln \mu}.\tag{1.17}$$

Since the rate at which the coupling constants change should be finite for $\varepsilon \rightarrow 0$, we should have

$$\frac{dg}{d \ln \mu} = -\frac{\varepsilon}{2}g + \beta_g(g, h),\tag{1.18}$$

$$\frac{dh}{d \ln \mu} = -\frac{\varepsilon}{2}h + \beta_h(g, h),\tag{1.19}$$

where β functions can be fixed by matching higher order terms in $1/\varepsilon$. Rearranging and expanding G and H , we find

$$0 = \left(1 + \frac{g}{\varepsilon} \frac{\partial G_1}{\partial g} + \dots \right) \frac{dg}{d \ln \mu} + \left(\frac{g}{\varepsilon} \frac{\partial G_1}{\partial g} + \dots \right) \frac{dh}{d \ln \mu} + \frac{\varepsilon}{2}g.\tag{1.20}$$

Matching the $\mathcal{O}(\varepsilon^0)$ terms, we find

$$-\frac{g^2}{2} \frac{\partial G_1}{\partial g} - \frac{gh}{2} \frac{\partial G_1}{\partial h} + \beta_g(g, h) = 0.\tag{1.21}$$

Then, G_n for $n \geq 2$ are obtained by matching higher order terms, which are fixed by G_1 . In other words, we have implicitly required β functions to have no ε dependence. Therefore, the above equation is exact. Hence, we find

$$\mu \frac{dg}{d\mu} = -\frac{1}{2}\varepsilon g + \frac{1}{2}g \left(g \frac{\partial G_1}{\partial g} + h \frac{\partial G_1}{\partial h} \right),\tag{1.22}$$

and similarly

$$\mu \frac{dh}{d\mu} = -\frac{1}{2}\varepsilon h + \frac{1}{2}h \left(g \frac{\partial H_1}{\partial g} + h \frac{\partial H_1}{\partial h} \right). \quad (1.23)$$

3. Use your results from part (a) to compute the beta functions $\beta_g(g, h) \equiv \lim_{\varepsilon \rightarrow 0} \mu dg/d\mu$ and $\beta_h(g, h) \equiv \lim_{\varepsilon \rightarrow 0} \mu dh/d\mu$. You should find terms of order g^3 , gh^2 , and h^3 in β_g , and terms of order g^2h , gh^2 , and h^3 in β_h .

Solution. By definition of G and H , we have

$$\frac{G_1}{\varepsilon} + \dots = \ln \left[\left(1 - \frac{g^2 + h^2}{6(4\pi)^3} \frac{1}{\varepsilon} + \dots \right)^{-3/2} \left(1 - \left(\frac{g^2}{(4\pi)^3} + \frac{h^3/g}{(4\pi)^3} \right) \frac{1}{\varepsilon} + \dots \right) \right] \quad (1.24)$$

and

$$\frac{H_1}{\varepsilon} + \dots = \ln \left[\left(1 - \frac{g^2 + h^2}{6(4\pi)^3} \frac{1}{\varepsilon} + \dots \right)^{-1/2} \left(1 - \frac{gh}{3(4\pi)^3} \frac{1}{\varepsilon} + \dots \right)^{-1} \left(1 - \left(\frac{h^2}{(4\pi)^3} + \frac{g}{4(4\pi)^3} \right) \frac{1}{\varepsilon} + \dots \right) \right]$$

Therefore, we have

$$G_1(g, h, \varepsilon) = \frac{1}{(4\pi)^3} (-3g^2/4 + h^2/4 - h^3/g), \quad (1.26)$$

and

$$H_1(g, h, \varepsilon) = \frac{1}{(4\pi)^3} (g^2/12 - 11h^2/12 + gh/6). \quad (1.27)$$

Therefore, we find

$$\beta_g(g, h) = \frac{1}{2(4\pi)^3} (-3g^3/2 - 2h^3 + h^2g/2) = \frac{1}{4(4\pi)^3} (-3g^3 - 4h^3 + h^2g), \quad (1.28)$$

and

$$\beta_h(g, h) = \frac{1}{12(4\pi)^3} (g^2h - 12gh^2 - 7h^3). \quad (1.29)$$

4. Without loss of generality, we can choose g to be positive; h can then be positive or negative, and the difference is physically significant. (You should understand why this is true.) For what numerical range(s) of h/g are β_g and β_h/h both negative? Why is this an interesting question?

Solution. Since the bare coupling constant is constant and determined by g and h . Therefore, by fixing the value of g , we have constrained the value of h .

Solving the equality using Mathematica, we find

$$h/g > \frac{1}{11}(-6 + \sqrt{47}) \simeq 0.079634. \quad (1.30)$$

If the above is satisfied, then the theory is asymptotically free. We can use the above to infer the range of the bare coupling constant for which the theory is asymptotically free.