

Functional Methods & Grassmann Integrals — Osborn AQFT PS2

Source: *H. Osborn, Advanced Quantum Field Theory, Problem Sheet 2 (functional methods, effective action, Grassmann integrals, SUSY QM).*

ON THIS PAGE

Legendre Transform and Perturbative Expansion, Q1

Effective Action for Free Scalar Field, Q2

Relations Between Green's Functions and 1PI Vertices, Q3

Grassmann Gaussian Integral with Bosonic Background, Q4

Fermionic Gaussian Integral, Q5

Pfaffian and Real Grassmann Integrals, Q6

Fermionic Path Integral and Partition Function, Q7

Supersymmetric Quantum Mechanics, Q8

Legendre Transform and Perturbative Expansion, Q1

Let, with $x_i, b_i, a_i \in \mathbb{R}^n$,

$$e^{W(b)} = \int d^n x e^{-S+b \cdot x}, \quad S = \frac{1}{2} x \cdot \mathbf{A} x + V(x), \quad \frac{\partial}{\partial b_i} W(b) = a_i, \quad \Gamma(a) + W(b) = b \cdot a. \quad (1.1)$$

Show that

$$\frac{\partial}{\partial a_i} \Gamma(a) = b_i, \quad W''_{ij} \Gamma''_{jk} = \delta_{ik}, \quad \frac{\partial}{\partial b_i} = W''_{ij} \frac{\partial}{\partial a_j}, \quad (1.2)$$

where W'' , Γ'' are matrices formed by $\frac{\partial^2}{\partial b_i \partial b_j} W$ and $\frac{\partial^2}{\partial a_i \partial a_j} \Gamma$. Find the first few terms in the perturbative expansion of $W(b)$ and correspondingly for $\Gamma(a)$.

Solution.

$$\frac{\partial}{\partial a_i} (\Gamma(a) + W(b)) = \frac{\partial}{\partial a_i} \Gamma(a) = b_i. \quad (1.3)$$

$$W''_{ij} \Gamma''_{jk} = \frac{\partial a_i}{\partial b_j} \frac{\partial b_j}{\partial a_k} = \delta_{ik}. \quad (1.4)$$

$$\frac{\partial}{\partial b_i} = \frac{\partial a_j}{\partial b_i} \frac{\partial}{\partial a_j} = W''_{ij} \frac{\partial}{\partial a_j}. \quad (1.5)$$

Now, Taylor expands the zero-dimensional Euclidean path integral, we find

$$\begin{aligned}
e^{W(b)} &= e^{-V(\partial/\partial b)} \int_{\mathbb{R}^n} d^n x e^{-\frac{1}{2}(x_i - A_{il}^{-1}b_l)A_{ij}(x_j - A_{jk}^{-1}b_k) + \frac{1}{2}b_i A_{ij}^{-1}b_j} \\
&= e^{-V(\partial/\partial b)} \int_{\mathbb{R}^n} d^n \xi e^{-\frac{1}{2}\xi_i A_{ij}\xi_j + \frac{1}{2}b_i A_{ij}^{-1}b_j} \\
&= \left(\int_{\mathbb{R}^n} d^n \xi e^{-\frac{1}{2}\xi_i A_{ij}\xi_j} \right) e^{-V(\partial/\partial b)} e^{\frac{1}{2}b_i A_{ij}^{-1}b_j} \\
&= \frac{(2\pi)^{n/2}}{\sqrt{\det A}} \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} \left[V \left(\frac{\partial}{\partial b} \right) \right]^j e^{\frac{1}{2}b_i A_{ij}^{-1}b_j}.
\end{aligned} \tag{1.6}$$

where we have made the substitution $\xi_i = x_i - A_{ij}^{-1}b_j$. Now, we shall take natural log on both side, which gives

$$W(b) = \ln \left(\frac{(2\pi)^{n/2}}{\sqrt{\det A}} \right) + \ln \left\{ \left(1 + \sum_{j=1}^{\infty} \frac{(-1)^j}{j!} \left[V \left(\frac{\partial}{\partial b} \right) \right]^j \right) e^{\frac{1}{2}b_i A_{ij}^{-1}b_j} \right\}. \tag{1.7}$$

Now, given

$$e^{-\frac{1}{2}b_i A_{ij}^{-1}b_j} \frac{\partial}{\partial b_i} e^{\frac{1}{2}b_i A_{ij}^{-1}b_j} = \frac{\partial}{\partial b_i} + A_{ij}^{-1}b_j, \tag{1.8}$$

we have

$$e^{-\frac{1}{2}b_i A_{ij}^{-1}b_j} V \left(\frac{\partial}{\partial b} \right) e^{\frac{1}{2}b_i A_{ij}^{-1}b_j} = V \left(\frac{\partial}{\partial b} + A^{-1}b \right). \tag{1.9}$$

Therefore,

$$W(b) = W_0(0) + \frac{1}{2}b_i A_{ij}^{-1}b_j + \sum_{l=1}^{\infty} \frac{1}{l} \sum_{j=1}^{\infty} \frac{(-1)^{j+2l+1}}{(j!)^l} \left[V \left(\frac{\partial}{\partial b} + A^{-1}b \right) \right]^{j+l}, \tag{1.10}$$

where the derivative is understood to act on 1.

Let us take the potential to be

$$V(x) = \frac{\lambda}{4!} x^4, \tag{1.11}$$

the connected generating function becomes

$$W(b) = W_0(0) + \frac{1}{2}b_i A_{ij}^{-1}b_j + \sum_{n=1}^{\infty} \frac{1}{n} \left\{ \sum_{m=1}^{\infty} \frac{(-1)^{m+1} \lambda^m}{m!4!} \left[\left(\frac{\partial}{\partial b_i} + A_{ij}^{-1}b_j \right)^4 \right]^m \right\}^n. \tag{1.12}$$

Now, let us compute the first few terms of the above,

$$\begin{aligned}
\left(\frac{\partial}{\partial b_i} + A_{ij}^{-1}b_j \right)^4 &= \left(\frac{\partial}{\partial b_i} + A_{ij}^{-1}b_j \right)^3 A_{ik}^{-1}b_k \\
&= \left(\frac{\partial}{\partial b_i} + A_{ij}^{-1}b_j \right)^2 (A_{ii}^{-1} + A_{ik}^{-1}A_{il}^{-1}b_l b_k) \\
&= \left(\frac{\partial}{\partial b_i} + A_{ij}^{-1}b_j \right) (3A_{ii}^{-1}A_{ip}^{-1}b_p + A_{ik}^{-1}A_{il}^{-1}A_{ip}^{-1}b_l b_k b_p) \\
&= (3A_{ii}^{-1}A_{ii}^{-1} + 6A_{ii}^{-1}A_{ij}^{-1}A_{il}^{-1}b_j b_l + A_{ik}^{-1}A_{il}^{-1}A_{ip}^{-1}A_{iq}^{-1}b_l b_k b_p b_q),
\end{aligned} \tag{1.13}$$

which corresponds to the diagrams

$$\frac{\lambda}{4!} \left(\frac{\partial}{\partial b_i} + A_{ij}^{-1} b_j \right)^4 = \frac{1}{8} \text{ (two circles joined at a point)} + \frac{1}{4} \text{ (circle on a line)} + \frac{1}{24} \text{ (X diagram)} \quad (1.14)$$

Therefore, to one-loop order, we have

$$W(b) = W_0 + \frac{1}{2} \text{ (line)} - \frac{1}{8} \text{ (two circles joined at a point)} - \frac{1}{4} \text{ (circle on a line)} - \frac{1}{24} \text{ (X diagram)} + O(\lambda^2). \quad (1.15)$$

The quantum action can be obtained by performing a Legendre transformation, which is

$$\Gamma(a) = -W(b) + b \cdot a, \quad (1.16)$$

where

$$a_i = \frac{\partial W(b)}{\partial b_i} = A_{ij}^{-1} b_j - \frac{1}{2} \lambda A_{il}^{-1} A_{li}^{-1} A_{lj}^{-1} b_j - \frac{1}{6} \lambda A_{li}^{-1} A_{lj}^{-1} A_{lk}^{-1} A_{lp}^{-1} b_j b_k b_p + O(\lambda^2). \quad (1.17)$$

Therefore, to first order in λ , we can replace b_i with $A_{ij} a_j$

$$\Gamma(a) = -W_0(0) + \frac{1}{2} A_{ij} a_i a_j + \frac{\lambda}{8} (A_{ii}^{-1})^2 + \frac{\lambda}{4} A_{ii}^{-1} a_i^2 + \frac{\lambda}{24} a_i^4 + O(\lambda^2). \quad (1.18)$$

Effective Action for Free Scalar Field, Q2

Starting from $Z[J]$, determine $W[J]$ and $\Gamma[\phi]$ for the free field theory for a scalar field $\phi(x)$ of mass m . What are

$$\frac{\delta^2}{\delta J(x) \delta J(y)} W[J] \Big|_{J=0}, \quad \frac{\delta^2}{\delta \phi(x) \delta \phi(y)} \Gamma[\phi] \Big|_{\phi=0} \quad (2.1)$$

in this case? Verify that they satisfy the expected inverse relations.

Solution. For a free scalar field theory,

$$Z[J] = \int \mathcal{D}\phi e^{iS[\phi] + i \int d^d x J\phi}, \quad (2.2)$$

where

$$S[\phi] = \int d^d x \left[\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} (m^2 - i\epsilon) \phi^2 \right]. \quad (2.3)$$

We define

$$\phi_J(x) = \phi(x) - i \int d^d y D_F(x-y) J(y), \quad (2.4)$$

where

$$D_F(x-y) = \int \frac{d^d k}{(2\pi)^d} \frac{i e^{-ik \cdot (x-y)}}{k^2 - m^2 + i\epsilon}. \quad (2.5)$$

The exponent of the generating functional becomes

$$\begin{aligned} & -\frac{i}{2} \int d^d x \phi(x) (\square + m^2 - i\epsilon) \phi(x) + i \int d^d x J(x) \phi(x) \\ &= -\frac{i}{2} \int d^d x \phi_J(x) (\square + m^2 - i\epsilon) \phi_J(x) - \frac{1}{2} \int d^d x \int d^d y J(x) D_F(x-y) J(y). \end{aligned} \quad (2.6)$$

Therefore, we have

$$Z[J] = \exp \left[-\frac{1}{2} \int d^d x \int d^d y J(x) D_F(x-y) J(y) \right] \frac{(2\pi)^{\infty/2}}{\sqrt{\text{Det}(m^2 - \bar{\square})}}, \quad (2.7)$$

where $\bar{\square} = \partial_\tau^2 + \nabla^2$ is the Euclidean Laplace operator. Then, the connected diagram generating functional is given by

$$iW[J] = \ln Z[J] - \frac{1}{2} \int d^d x \int d^d y J(x) D_F(x-y) J(y). \quad (2.8)$$

The quantum action is defined by the Legendre transformation

$$\Gamma[\phi_J] = -W[J] + \int d^d x J(x) \phi_J(x), \quad (2.9)$$

where

$$\frac{\delta \Gamma[\phi_J]}{\delta \phi_J(x)} = J(x), \quad \frac{\delta W[J]}{\delta J(x)} = \phi_J(x). \quad (2.10)$$

Therefore,

$$\phi_J(x) = i \int d^d y D_F(x-y) J(y), \quad (2.11)$$

and hence

$$\Gamma[\phi_J] = i \ln Z[0] + \frac{1}{2} \int d^d x \phi_J(x) (\square + m^2 - i\epsilon) \phi_J(x). \quad (2.12)$$

Now, we compute

$$\frac{\delta^2}{\delta J(x)\delta J(y)}W[J]\Big|_{J=0} = iD_F(x-y), \quad \frac{\delta^2}{\delta\phi(x)\delta\phi(y)}\Gamma[\phi]\Big|_{\phi=0} = (\square + m^2 - i\epsilon), \quad (2.13)$$

which verifies

$$\frac{\delta^2}{\delta J(x)\delta J(y)}W[J]\Big|_{J=0} \frac{\delta^2}{\delta\phi(y)\delta\phi(z)}\Gamma[\phi]\Big|_{\phi=0} = \delta^d(x-z). \quad (2.14)$$

Relations Between Green's Functions and 1PI Vertices, Q3

Starting from the definition of the effective action $\Gamma[\phi]$ in terms of the generating functional for connected graphs $W[J]$,

$$W[J] + \Gamma[\phi] = \int d^d x J\phi, \quad \frac{\delta}{\delta J(x)}W[J] = \phi(x), \quad (3.1)$$

obtain

$$-\int d^d z G_2(x, z) \Gamma_2(z, y) = \delta^d(x - y), \quad -i \frac{\delta}{\delta J(x)} = \int d^d y G_2(x, y) \frac{\delta}{\delta\phi(y)}, \quad (3.2)$$

where

$$G_n(x_1, \dots, x_n) = (-i)^{n-1} \frac{\delta}{\delta J(x_1)} \cdots \frac{\delta}{\delta J(x_n)} W[J], \quad \Gamma_n(x_1, \dots, x_n) = -i \frac{\delta}{\delta\phi(x_1)} \cdots \frac{\delta}{\delta\phi(x_n)} \Gamma[\phi]. \quad (3.3)$$

Show that

$$G_3(x_1, x_2, x_3) = \int d^d y_1 d^d y_2 d^d y_3 G_2(x_1, y_1) G_2(x_2, y_2) G_2(x_3, y_3) \Gamma_3(y_1, y_2, y_3). \quad (3.4)$$

Also obtain relations in diagrammatic form for G_4 , G_5 in terms of G_2 , Γ_3 , Γ_4 and Γ_5 .

Solution.

$$\frac{\delta\phi(x)}{\delta\phi(y)} = \delta^d(x - y) = \int d^d z \frac{\delta\phi(x)}{\delta J(z)} \frac{\delta J(z)}{\delta\phi(y)}. \quad (3.5)$$

Recall

$$\frac{\delta\Gamma[\phi]}{\delta\phi(x)} = J(x), \quad \frac{\delta W[J]}{\delta J(x)} = \phi(x), \quad (3.6)$$

which gives

$$\delta^d(x - y) = \int d^d z \frac{\delta^2 W[J]}{\delta J(x)\delta J(z)} \frac{\delta^2 \Gamma[\phi]}{\delta\phi(z)\delta\phi(y)} = - \int d^d z G_2(x, z) \Gamma_2(z, y). \quad (3.7)$$

Similarly,

$$-i \frac{\delta}{\delta J(x)} = -i \int d^d y \frac{\delta\phi(y)}{\delta J(x)} \frac{\delta}{\delta\phi(y)} = \int d^d y G_2(x, y) \frac{\delta}{\delta\phi(y)}. \quad (3.8)$$

Now, we turn to the exact connected three point function

$$\begin{aligned}
 G_3(x_1, x_2, x_3) &= - \int d^d y_1 d^d y_2 d^d y_3 \left(iG_2(x_1, y_1) \frac{\delta}{\delta \phi(y_1)} \right) \left(iG_2(x_2, y_2) \frac{\delta}{\delta \phi(y_2)} \right) \\
 &\quad \left(iG_2(x_3, y_3) \frac{\delta}{\delta \phi(y_3)} \right) \left(-\Gamma[\phi] + \int d^d x J(x) \phi(x) \right) \\
 &= \int d^d y_1 d^d y_2 d^d y_3 G_2(x_1, y_1) G_2(x_2, y_2) G_2(x_3, y_3) \Gamma_3(y_1, y_2, y_3),
 \end{aligned} \tag{3.9}$$

where we have jumped a few lines of algebra.

For higher-order connected correlation functions, we have the diagrammatic expressions:

$$\begin{aligned}
 & \text{Diagrammatic equation (3.10):} \\
 & G_4(x_1, x_2, x_3, x_4) = \text{1PI diagram} + \sum_{(ij)(kl)} \dots
 \end{aligned} \tag{3.10}$$

$$\begin{aligned}
 & \text{Diagrammatic equation (3.11):} \\
 & G_5(x_1, x_2, x_3, x_4, x_5) = \text{1PI diagram} + \sum_{10 \text{ ch.}} \dots + \sum_{15 \text{ ch.}} \dots
 \end{aligned} \tag{3.11}$$

Grassmann Gaussian Integral with Bosonic Background, Q4

Evaluate, for $\theta_i, \bar{\theta}_i$ Grassmann variables,

$$Z = \frac{1}{(2\pi)^{n/2}} \int d^n x \prod_{i=1}^n d\bar{\theta}_i d\theta_i e^{-S(x, \theta, \bar{\theta})}, \quad S(x, \theta, \bar{\theta}) = \frac{1}{2} w_i(x) w_i(x) + \bar{\theta}_i w_{i,j}(x) \theta_j, \tag{4.1}$$

where $w_{i,j}(x) = \frac{\partial}{\partial x_j} w_i(x)$ and assuming $\det[w_{i,j}(x)] > 0$.

If $w_i(x) = x_i + \frac{1}{2}g_{ijk}x_jx_k$, $g_{ijk} = g_{ikj}$, what are the Feynman rules for a perturbative expansion? Work out the two-loop contributions to $Z(g)$ (this is much simpler if g_{ijk} is completely symmetric).

If $w_i(x) \rightarrow \lambda w_i(x)$ in S , show that

$$\frac{d}{d\lambda} e^{-S(x,\theta,\bar{\theta})} = - \left(\theta_j \frac{\partial}{\partial x_j} + \lambda w_j(x) \frac{\partial}{\partial \bar{\theta}_j} \right) \bar{\theta}_i w_i(x) e^{-S(x,\theta,\bar{\theta})}. \quad (4.2)$$

By letting $x_i \rightarrow \lambda x_i$, $g_{ijk} \rightarrow g_{ijk}/\lambda$, what does this imply for $Z(g)$?

Solution. Let us first evaluate the Grassmann integral by making the substitution $\theta'_i = w_{i,j}\theta_j$

$$\int \prod_{k=1}^n d\bar{\theta}_k d\theta_k e^{-\bar{\theta}_i w_{i,j}\theta_j} = \prod_{k=1}^n \frac{\partial}{\partial \bar{\theta}_k} \frac{\partial}{\partial \theta_k} (1 - \bar{\theta}_i w_{i,j}\theta_j) = \det w_{i,j} \left(\frac{\partial}{\partial \bar{\theta}} \frac{\partial}{\partial \theta'} (1 - \bar{\theta}\theta') \right)^n = \det w_{i,j}. \quad (4.3)$$

Then, we make the substitution $d^n x = d^n w / \det w_{i,j}$, which gives

$$Z = \frac{1}{(2\pi)^{n/2}} \int d^n w e^{-w_i w_i/2} = 1. \quad (4.4)$$

For $w_i(x) = x_i + \frac{1}{2}g_{ijk}x_jx_k$, $w_{i,j}(x) = \delta_{ij} + g_{ijk}x_k$ the action becomes

$$S = \frac{1}{2}x_i x_i + \frac{1}{2}g_{ijk}x_i x_j x_k + \frac{1}{8}g_{ijk}g_{ilm}x_j x_k x_l x_m + \bar{\theta}_i \theta_i + g_{ijk}\bar{\theta}_i x_j \theta_k. \quad (4.5)$$

We can read off the Feynman rules:

$$i \text{ --- } j = \delta_{ij}, \quad (4.6)$$

$$i \longrightarrow j = \delta_{ij}, \quad (4.7)$$

$$\begin{array}{c} j \\ \diagup \\ i \text{ --- } \bullet \\ \diagdown \\ k \end{array} = -g_{ijk}, \quad (4.8)$$

$$\begin{array}{c} j \quad k \\ \diagdown \quad \diagup \\ \times \\ \diagup \quad \diagdown \\ l \quad m \end{array} = -g_{ijk}g_{ilm}, \quad (4.9)$$

$$\begin{array}{c} j \\ | \\ i \longrightarrow \bullet \longrightarrow k \end{array} = -g_{ijk}. \quad (4.10)$$

Since we have already shown that $Z = 1$, the loop expansions should be cancelled order by order in g . There are no diagrams of order g . For order g^2 , the diagram should cancel, but we shall not show it explicitly here.

Now, we take $w_i \rightarrow \lambda w_i$, which the action becomes

$$S = \frac{1}{2} \lambda^2 w_i(x) w_i(x) + \lambda \bar{\theta}_i w_{i,j}(x) \theta_j. \quad (4.11)$$

Then,

$$\frac{d}{d\lambda} e^{-S(x,\theta,\bar{\theta})} = -(\lambda w_i(x) w_i(x) + \bar{\theta}_i w_{i,j}(x) \theta_j) e^{-S(x,\theta,\bar{\theta})}, \quad (4.12)$$

while

$$\begin{aligned} & - \left(\theta_j \frac{\partial}{\partial x_j} + \lambda w_j(x) \frac{\partial}{\partial \theta_j} \right) \bar{\theta}_i w_i(x) e^{-S(x,\theta,\bar{\theta})} \\ &= - \left(\theta_j w_{i,j}(x) \bar{\theta}_i - \theta_j \bar{\theta}_i w_i(x) [\lambda^2 w_k(x) w_{k,j}(x) + \lambda \bar{\theta}_k w_{i,lj}(x) \theta_l] \right. \\ & \quad \left. + \lambda w_i(x) w_i(x) - \lambda^2 \bar{\theta}_i w_j(x) w_i(x) w_{l,j}(x) \theta_l \right) e^{-S(x,\theta,\bar{\theta})} \\ &= - (\lambda w_i(x) w_i(x) + \bar{\theta}_i w_{i,j}(x) \theta_j) e^{-S(x,\theta,\bar{\theta})}, \end{aligned} \quad (4.13)$$

where the term containing $w_{i,jl}$ vanishes by anti-commutation of Grassmann numbers.

Letting $x_i \rightarrow \lambda x_i$ and $g_{ijk} \rightarrow g_{ijk}/\lambda$ is equivalent to letting $w_i \rightarrow \lambda w_i$. In other words,

$$w_i(\lambda x, g/\lambda) = \lambda w_i(x). \quad (4.14)$$

Fermionic Gaussian Integral, Q5

Show that, for $\theta_i, \bar{\theta}_i, \eta_i, \bar{\eta}_i, i = 1, 2, \dots, n$ independent Grassmann variables and B an invertible $n \times n$ matrix,

$$\int \prod_{i=1}^n d\bar{\theta}_i d\theta_i \exp(-\bar{\theta}_i B_{ij} \theta_j + \bar{\eta}_i \theta_i + \bar{\theta}_i \eta_i) = \det B \exp(\bar{\eta}_i (B^{-1})_{ij} \eta_j), \quad (5.1)$$

and

$$\langle \theta_i \bar{\theta}_j \rangle = - \frac{\partial}{\partial \bar{\eta}_i} \frac{\partial}{\partial \eta_j} \exp(\bar{\eta}_i (B^{-1})_{ij} \eta_j) \Big|_{\eta=\bar{\eta}=0} = (B^{-1})_{ij}. \quad (5.2)$$

Solution. Making the substitution $\chi_i = \theta_i - B_{ij}^{-1} \eta_j$, and $\bar{\chi}_i = \bar{\theta}_i - \bar{\eta}_j B_{ji}^{-1}$, the integral becomes

$$\int \prod_{i=1}^n d\bar{\chi}_i d\chi_i \exp(-\bar{\chi}_i B_{ij} \chi_j + \bar{\eta}_j B_{ji}^{-1} \eta_i) = \exp(\bar{\eta}_i (B^{-1})_{ij} \eta_j) \int \prod_{i=1}^n d\bar{\chi}_i d\chi_i \exp(-\bar{\chi}_i B_{ij} \chi_j), \quad (5.3)$$

where the latter integral is equal to $\det B$. Then,

$$\langle \theta_i \bar{\theta}_j \rangle = - \frac{\partial}{\partial \bar{\eta}_i} \frac{\partial}{\partial \eta_j} \exp(\bar{\eta}_i (B^{-1})_{ij} \eta_j) \Big|_{\eta=\bar{\eta}=0} = - \frac{\partial}{\partial \bar{\eta}_i} \frac{\partial}{\partial \eta_j} (1 + \bar{\eta}_i B_{ij}^{-1} \eta_j) \Big|_{\eta=\bar{\eta}=0} = B_{ij}^{-1}. \quad (5.4)$$

Pfaffian and Real Grassmann Integrals, Q6

Show that

$$\int d^{2n} \theta \exp\left(\frac{1}{2} \theta_i A_{ij} \theta_j + \eta_i \theta_i\right) = \text{Pf}(A) \exp\left(\frac{1}{2} \eta_i (A^{-1})_{ij} \eta_j\right), \quad d^{2n} \theta = d\theta_{2n} \cdots d\theta_1, \quad (6.1)$$

where θ_i and η_i are Grassmann variables, A is an invertible antisymmetric matrix and the Pfaffian is

$$\text{Pf}(A) = \frac{1}{2^n n!} \varepsilon_{i_1 \cdots i_{2n}} A_{i_1 i_2} \cdots A_{i_{2n-1} i_{2n}}. \quad (6.2)$$

Why does $\text{Pf}(A)^2 = \det A$? Hence obtain

$$\langle \theta_i \theta_j \rangle = \frac{\partial}{\partial \eta_i} \frac{\partial}{\partial \eta_j} \exp\left(\frac{1}{2} \eta_i (A^{-1})_{ij} \eta_j\right) \Big|_{\eta=0} = -(A^{-1})_{ij}. \quad (6.3)$$

Show that $\delta(\theta) = \theta_1 \cdots \theta_{2n}$ plays the role of a δ -function for Grassmann integrals over $\theta_1, \dots, \theta_{2n}$.

Solution. Similarly, we perform the substitution $\chi_i = \theta_i - A_{ij}^{-1} \eta_j$, which gives

$$\int d^{2n} \chi \exp\left(\frac{1}{2} (\chi_i + A_{il}^{-1} \eta_l) A_{ij} (\chi_j + A_{jk}^{-1} \eta_k) + \eta_i (\chi_i + A_{ij}^{-1} \eta_j)\right). \quad (6.4)$$

The exponent is

$$\frac{1}{2} \chi_i A_{ij} \chi_j + \frac{1}{2} \chi_i A_{ij} A_{jk}^{-1} \eta_k + \frac{1}{2} A_{il}^{-1} \eta_l A_{ij} \chi_j + \frac{1}{2} A_{il}^{-1} \eta_l A_{ij} A_{jk}^{-1} \eta_k + \eta_i \chi_i + \eta_i A_{ij}^{-1} \eta_j. \quad (6.5)$$

For the $\theta A \theta$ term to be non-vanishing, we need A to be antisymmetric. Therefore,

$$\frac{1}{2} A_{il}^{-1} \eta_l A_{ij} \chi_j = -\frac{1}{2} \eta_i \chi_i. \quad (6.6)$$

The integral becomes

$$\int d^{2n} \theta \exp\left(\frac{1}{2} \theta_i A_{ij} \theta_j + \eta_i \theta_i\right) = \exp\left(\frac{1}{2} \eta_i (A^{-1})_{ij} \eta_j\right) \int d^{2n} \chi \exp\left(\frac{1}{2} \chi_i A_{ij} \chi_j\right). \quad (6.7)$$

The integral on the right-hand side is

$$\begin{aligned} \int d^{2n} \chi \exp\left(\frac{1}{2} \chi_i A_{ij} \chi_j\right) &= \frac{\partial}{\partial \chi_{2n}} \cdots \frac{\partial}{\partial \chi_1} \sum_{m=0}^n \frac{1}{m!} \left(\frac{1}{2} \chi_i A_{ij} \chi_j\right)^m \\ &= \frac{1}{2^n n!} \frac{\partial}{\partial \chi_{2n}} \cdots \frac{\partial}{\partial \chi_1} (\chi_i A_{ij} \chi_j)^n \\ &= \frac{1}{2^n n!} \varepsilon_{i_1 \cdots i_{2n}} A_{i_1 i_2} \cdots A_{i_{2n-1} i_{2n}} \\ &= \text{Pf}(A). \end{aligned} \quad (6.8)$$

Now, we consider

$$\begin{aligned} \text{Pf}^2(A) &= \left(\int d^{2n} \chi \exp \left(\frac{1}{2} \chi_i A_{ij} \chi_j \right) \right)^2 \\ &= \int d^{2n} \chi d^{2n} \chi' \exp \left(\frac{1}{2} \chi_i A_{ij} \chi_j + \frac{1}{2} \chi'_i A_{ij} \chi'_j \right). \end{aligned} \quad (6.9)$$

Let $\eta_i = \frac{1}{\sqrt{2}}(\chi_i + \chi'_i)$ and $\eta'_i = \frac{i}{\sqrt{2}}(\chi_i - \chi'_i)$, so that the above becomes

$$\text{Pf}^2(A) = \int \prod_{i=1}^{2n} d\eta_i d\eta'_i \exp(\eta'_i A_{ij} \eta_j) = \det(-A) = (-1)^{2n} \det(A) = \det(A). \quad (6.10)$$

It is easy to see that

$$\langle \theta_i \theta_j \rangle = \frac{\partial}{\partial \eta_i} \frac{\partial}{\partial \eta_j} \exp \left(\frac{1}{2} \eta_i (A^{-1})_{ij} \eta_j \right) \Big|_{\eta=0} = -(A^{-1})_{ij} \quad (6.11)$$

is true.

Now, consider a Grassmann function, which can be expanded as $f(\theta) = f(0) + (\text{terms with } \theta_i)$. Therefore,

$$\int d^{2n} \theta \theta_1 \cdots \theta_{2n} f(\theta) = \int d^{2n} \theta \theta_1 \cdots \theta_{2n} f(0) = f(0). \quad (6.12)$$

Fermionic Path Integral and Partition Function, Q7

For the simple fermionic oscillator with energy eigenstates $|0\rangle, |1\rangle$ and $H = b^\dagger b \omega$, let $|\theta\rangle = |0\rangle + \theta|1\rangle$, $\langle \bar{\theta}| = \langle 0| + \bar{\theta}\langle 1|$ where $\theta, \bar{\theta}$ are Grassmann variables. Show that

$$\text{tr } e^{-\beta H} = \int d\theta d\bar{\theta} e^{\theta \bar{\theta}} \langle \bar{\theta} | e^{-\beta H} | \theta \rangle, \quad \int d\bar{\theta} d\theta e^{\theta \bar{\theta}} | \theta \rangle \langle \bar{\theta} | = 1. \quad (7.1)$$

Let $\beta = (N+1)\epsilon$ and show that this can be expressed for large N as a path integral over products of $\langle \bar{\theta}_{i+1} | e^{-H} | \theta_i \rangle \approx \exp((1 - \epsilon\omega)\bar{\theta}_{i+1}\theta_i)$ so that

$$\text{tr } e^{-\beta H} \approx \int \prod_{i=1}^{N+1} d\bar{\theta}_i d\theta_i \exp(-\bar{\theta}_i B_{ij} \theta_j) = \det(B), \quad (7.2)$$

where we take $\bar{\theta}_{N+1} = \bar{\theta}$, $\theta_{N+1} = -\theta$ and B is an $(N+1) \times (N+1)$ matrix of the form

$$B = \begin{pmatrix} 1 & 0 & \cdots & 0 & 1 - \epsilon\omega \\ -1 + \epsilon\omega & 1 & \cdots & 0 & 0 \\ 0 & -1 + \epsilon\omega & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & \cdots & -1 + \epsilon\omega & 1 \end{pmatrix}. \quad (7.3)$$

Show that $\det B = 1 + (1 - \epsilon\omega)^{N+1} \rightarrow 1 + e^{-\beta\omega}$ as $N \rightarrow \infty$.

Solution. The completeness relation is given by

$$|0\rangle\langle 0| + |1\rangle\langle 1| = 1 = \int d\bar{\theta} d\theta e^{\theta\bar{\theta}} |\theta\rangle\langle\bar{\theta}|. \quad (7.4)$$

Then

$$\begin{aligned} \text{tr } e^{-\beta H} &= \langle 0|e^{-\beta H}|0\rangle + \langle 1|e^{-\beta H}|1\rangle \\ &= \int d\bar{\theta} d\theta e^{\theta\bar{\theta}} (\langle\bar{\theta}|e^{-\beta H}|0\rangle + \theta\langle\bar{\theta}|e^{-\beta H}|1\rangle) \\ &= \int d\bar{\theta} d\theta e^{\theta\bar{\theta}} \langle\bar{\theta}|e^{-\beta H}|\theta\rangle. \end{aligned} \quad (7.5)$$

Taking $\beta = N + 1$, so that the path integral becomes

$$\text{tr } e^{-\beta H} = \int d\bar{\theta} d\theta e^{\theta\bar{\theta}} \int \prod_{i=1}^N (d\bar{\theta}_i d\theta_i e^{\theta_i\bar{\theta}_i}) \langle\bar{\theta}|e^{-H}|\theta_1\rangle \langle\bar{\theta}_1|e^{-H}|\theta_2\rangle \cdots \langle\bar{\theta}_N|e^{-H}|\theta\rangle. \quad (7.6)$$

For large N ,

$$\langle\bar{\theta}_{i+1}|e^{-\epsilon H}|\theta_i\rangle = \langle\bar{\theta}_{i+1}|e^{-\epsilon\omega b^\dagger b}|\theta_i\rangle \approx e^{-\epsilon\omega\bar{\theta}_{i+1}\theta_i} \langle\bar{\theta}_{i+1}|\theta_i\rangle = e^{-\epsilon\omega\bar{\theta}_{i+1}\theta_i} (1 + \bar{\theta}_{i+1}\theta_i) = e^{(1-\epsilon\omega)\bar{\theta}_{i+1}\theta_i}, \quad (7.7)$$

where we have neglected terms of order ϵ^2 or higher. Plugging this back into the path integral, we find

$$\text{tr } e^{-\beta H} \approx \int_{\theta_1=\theta_{N+2}} \prod_{i=1}^{N+1} d\bar{\theta}_i d\theta_i e^{\theta_i\bar{\theta}_i} e^{(1-\epsilon\omega)\bar{\theta}_{i+1}\theta_i} = \int \prod_{i=1}^{N+1} d\bar{\theta}_i d\theta_i \exp(-\bar{\theta}_i B_{ij} \theta_j) = \det(B). \quad (7.8)$$

The determinant is given by

$$\begin{aligned} \det(B) &= \det \begin{pmatrix} 1 & \cdots & 0 & 0 \\ -1 + \epsilon\omega & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 \\ 0 & \cdots & -1 + \epsilon\omega & 1 \end{pmatrix} + (-1)^N (1 - \epsilon\omega) \det \begin{pmatrix} -1 + \epsilon\omega & 1 & \cdots & 0 \\ 0 & -1 + \epsilon\omega & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \\ 0 & 0 & \cdots & -1 + \epsilon\omega \end{pmatrix} \\ &= 1 + (-1)^N (1 - \epsilon\omega)(-1 + \epsilon\omega)^N = 1 + (1 - \epsilon\omega)^{N+1} \rightarrow 1 + e^{-\beta\omega}. \end{aligned}$$

Supersymmetric Quantum Mechanics, Q8

Let

$$S = \int_0^\beta d\tau \left(\frac{1}{2} \dot{x}^2 + \bar{\theta} \dot{\theta} + \frac{1}{2} \lambda^2 w'(x)^2 + \lambda w''(x) \bar{\theta} \theta \right). \quad (8.1)$$

Assume $x(\tau)$ and the Grassmannian $\theta(\tau)$, $\bar{\theta}(\tau)$ are all periodic with period β . Show that

$$Q(\bar{\theta} \cdot w' e^{-S}) = -\frac{\partial}{\partial \lambda} e^{-S}, \quad Q = \int_0^\beta d\tau \left(\theta \frac{\delta}{\delta x} + (\lambda w'(x) + \dot{x}) \frac{\delta}{\delta \bar{\theta}} \right), \quad \bar{\theta} \cdot w' = \int_0^\beta d\tau \bar{\theta} w'(x). \quad (8.2)$$

Hence show that

$$Z = \int \mathcal{D}x \mathcal{D}\bar{\theta} \mathcal{D}\theta e^{-S} \quad (8.3)$$

is independent of λ if $\lambda \neq 0$. Calculate Z for $\lambda w'(x) = \omega x$ if $Z = 1$ for $\omega = 0$.

Solution.

$$\begin{aligned} Q(\bar{\theta} \cdot w' e^{-S}) &= \int_0^\beta d\tau \left(\theta \frac{\delta}{\delta x} + (\lambda w'(x) + \dot{x}) \frac{\delta}{\delta \bar{\theta}} \right) \left(\int_0^\beta d\tau' \bar{\theta} w'(x) e^{-S} \right) \\ &= \int_0^\beta d\tau d\tau' \left(\theta(\tau) \bar{\theta}(\tau') w''(x(\tau')) \delta(\tau - \tau') + \theta(\tau) \bar{\theta}(\tau') w'(x(\tau')) \right. \\ &\quad \times \int_0^\beta d\tau'' [\ddot{x}(\tau) \delta(\tau - \tau'') - \lambda^2 w'(x(\tau'')) w''(x(\tau'')) \delta(\tau - \tau'') \\ &\quad \left. - \lambda w'''(x(\tau'')) \bar{\theta}(\tau'') \theta(\tau'') \delta(\tau - \tau'')] \right) e^{-S} \\ &\quad + \int_0^\beta d\tau d\tau' [\lambda w'(x(\tau)) + \dot{x}(\tau)] \left(\delta(\tau - \tau') w'(x(\tau')) \right. \\ &\quad \left. + \bar{\theta}(\tau') w'(x(\tau')) \int_0^\beta d\tau'' [\dot{\theta}(\tau'') \delta(\tau - \tau'') + \lambda w''(x(\tau'')) \theta(\tau'') \delta(\tau - \tau'')] \right) e^{-S} \\ &= \int_0^\beta d\tau \theta \bar{\theta} w''(x) e^{-S} \\ &\quad + \int_0^\beta d\tau d\tau' \theta(\tau) \bar{\theta}(\tau') w'(x(\tau')) [\ddot{x}(\tau) - \lambda^2 w'(x(\tau)) w''(x(\tau))] e^{-S} \\ &\quad + \int_0^\beta d\tau [\lambda w'(x) + \dot{x}] w'(x) e^{-S} \\ &\quad + \int_0^\beta d\tau d\tau' [\lambda w'(x(\tau)) + \dot{x}(\tau)] \bar{\theta}(\tau') w'(x(\tau')) [\dot{\theta}(\tau) + \lambda w''(x(\tau)) \theta(\tau)] e^{-S} \\ &= \int_0^\beta d\tau [w''(x) \bar{\theta} \theta + \lambda w'(x) w'(x) + \dot{x} w'(x)] e^{-S} \\ &= -\frac{d}{d\lambda} e^{-S}, \end{aligned}$$

where we have used the fact that $\dot{x} w'(x) = \frac{d}{d\tau} w(x)$ is a total derivative, and thus vanishes.

Now consider

$$\frac{dZ}{d\lambda} = - \int \mathcal{D}x \mathcal{D}\bar{\theta} \mathcal{D}\theta Q(\bar{\theta} \cdot w' e^{-S}), \quad (8.5)$$

which only contains boundary terms of x and $\bar{\theta}$. Therefore, we have $dZ/d\lambda = 0$, which means Z is independent of λ for SUSY QM. Now, take $\lambda w'(x) = \omega x$, and then

$$S = \int_0^\beta d\tau \left(\frac{1}{2} \dot{x}^2 + \bar{\theta} \dot{\theta} + \frac{1}{2} \omega^2 x^2 + \omega \bar{\theta} \theta \right). \quad (8.6)$$

The partition function is then given by

$$\begin{aligned}
Z &= \int \mathcal{D}x e^{-\int d\tau \frac{1}{2}(\dot{x}^2 + \omega^2 x^2)} \int \mathcal{D}\bar{\theta} \mathcal{D}\theta e^{-\int d\tau (\bar{\theta} \dot{\theta} + \omega \bar{\theta} \theta)} \\
&= \int \mathcal{D}x e^{-\int d\tau \frac{1}{2}x(-\frac{d^2}{d\tau^2} + \omega^2)x} \int \mathcal{D}\bar{\theta} \mathcal{D}\theta e^{-\int d\tau \bar{\theta}(\frac{d}{d\tau} + \omega)\theta} \\
&= \frac{\text{Det}_{\text{AP}}(\frac{d}{d\tau} + \omega)}{\sqrt{\text{Det}_{\text{P}}(-\frac{d^2}{d\tau^2} + \omega^2)}}.
\end{aligned} \tag{8.7}$$

The determinants can be evaluated as

$$\text{Det}_{\text{AP}}\left(\frac{d}{d\tau} + \omega\right) = \prod_n \lambda_n = \prod_n \left[\frac{(2n-1)\pi i}{\beta} + \omega\right] = \text{Det}_{\text{AP}}\left(\frac{d}{d\tau}\right) \prod_n \left[1 - \frac{i\omega\beta}{(2n-1)\pi}\right], \tag{8.8}$$

and

$$\text{Det}_{\text{P}}\left(-\frac{d^2}{d\tau^2} + \omega^2\right) = \prod_n \left[\left(\frac{2\pi n}{\beta}\right)^2 + \omega^2\right] = \text{Det}_{\text{P}}\left(-\frac{d^2}{d\tau^2}\right) \prod_n \left[1 + \left(\frac{\omega\beta}{2\pi n}\right)^2\right]. \tag{8.9}$$

Using the normalisation, $Z = 1$ for $\omega = 0$, we find

$$Z = \prod_n \frac{1 - \frac{i\omega\beta}{(2n-1)\pi}}{\sqrt{1 + \left(\frac{\omega\beta}{2\pi n}\right)^2}}. \tag{8.10}$$