

Differential Geometry

Source: D. Tong, *General Relativity, Problem Sheet 1* (manifolds, tensors, Lie and exterior derivatives).

ON THIS PAGE

Integral Curves of Vector Fields, Q1

Tensors from Linear Maps, Q2

The Hessian, Q3

Transformation of the Metric Determinant, Q4

Lie Derivatives and Cartan's Magic Formula, Q5

Properties of the Exterior Derivative, Q6

Coordinate-Free Definition of the Exterior Derivative, Q7

Maurer-Cartan Structure Equations on S^3 , Q8

Coordinate-Induced Bases and the Poincaré Lemma, Q9

Integral Curves of Vector Fields, Q1

Draw the integral curves corresponding to a vector field in $\mathbb{R}^2$ with components, in Cartesian coordinates, given by

- $X^\mu = (y, -x)$
- $X^\mu = (x - y, x + y)$

[Hint: you might find life easier in polar coordinates.]

Solution. Recall that a vector field can be defined as the tangent to the streamlines at each point, which means

$$X^\mu = \frac{dx^\mu(t)}{dt} \implies \begin{cases} \dot{x} = y, \\ \dot{y} = -x, \end{cases} \quad (1.1)$$

which are solved by the circles on $\mathbb{R}^2$.

For the second vector field, we have

$$\begin{cases} \dot{x} = x - y, \\ \dot{y} = x + y. \end{cases} \quad (1.2)$$

This is solved by spirals of the form

$$x^2 + y^2 = r^2 e^{2 \tan^{-1}(y/x)}. \quad (1.3)$$

Tensors from Linear Maps, Q2

Let $\hat{H} : T_p(M) \rightarrow T_p^*(M)$ be a linear map. Define

$$H(X, Y) = [\hat{H}(Y)](X). \quad (2.1)$$

Show that this map is linear in both arguments (e.g. $H(fX + gY, Z) = fH(X, Z) + gH(Y, Z)$ for $f, g \in C^\infty$ and $X, Y, Z \in T_p(M)$) and hence defines a rank (0, 2) tensor. Similarly, show that a linear map $T_p(M) \rightarrow T_p(M)$ defines a tensor of rank (1, 1). What tensor δ arises from the identity map?

Solution. Proof of linearity:

$$H(fX + gY, Z) = [\hat{H}(Z)](fX + gY). \quad (2.2)$$

Since $\hat{H}(Z)$ is a covariant vector, we have

$$H(fX + gY, Z) = f[\hat{H}(Z)](X) + g[\hat{H}(Z)](Y) = fH(X, Z) + gH(Y, Z). \quad (2.3)$$

$$H(X, fY + gZ) = [\hat{H}(fY + gZ)](X). \quad (2.4)$$

By linearity of the linear map $\hat{H}$, we have

$$H(X, fY + gZ) = f[\hat{H}(Y)](X) + g[\hat{H}(Z)](X) = fH(X, Y) + gH(X, Z). \quad (2.5)$$

Therefore, H defines a multi-linear map, ■

$$H : T_p(M) \times T_p(M) \rightarrow \mathbb{R}, \quad (2.6)$$

which is a rank (0, 2) tensor.

Use the linear map,

$$\hat{G} : T_p(M) \rightarrow T_p(M), \quad (2.7)$$

to define

$$G(\omega, X) = \omega(\hat{G}(X)), \quad (2.8)$$

with $\omega \in T_p^*(M)$. Now, we shall check the linearity of the G constructed in both entries:

$$G(f\omega + g\eta, X) = fG(\omega, X) + gG(\eta, X) \quad (2.9)$$

by linearity of covariant vectors, and

$$G(\omega, fX + gY) = fG(\omega, X) + gG(\omega, Y) \quad (2.10)$$

by linearity of $\hat{G}$. Therefore, G is a rank (1, 1) tensor

$$G : T_p^*(M) \times T_p(M) \rightarrow \mathbb{R}. \quad (2.11)$$

The identity map is

$$\hat{\delta}(X) = X. \quad (2.12)$$

Therefore, it defines the tensor

$$\delta(\omega, X) = \omega_\mu X^\mu \quad (2.13)$$

with

$$\delta(f^\mu, e_\nu) = \delta^\mu_\nu. \quad (2.14)$$

The Hessian, Q3

Let M be a manifold and $f : M \rightarrow \mathbb{R}$ be a smooth function such that $df = 0$ at some point $p \in M$. Let x^μ be a coordinate chart defined in a neighbourhood of p . Define

$$F_{\mu\nu} = \frac{\partial^2 f}{\partial x^\mu \partial x^\nu}. \quad (3.1)$$

By considering the transformation law for components show that $F_{\mu\nu}$ defines a rank $(0, 2)$ tensor. (This is called the Hessian of f at p .) Construct also a coordinate-free definition and demonstrate its tensorial properties.

Solution. Under the transformation $\tilde{x}^\mu \rightarrow x^\mu$, $F_{\mu\nu}$ transforms as

$$\tilde{F}_{\mu\nu} \Big|_p \rightarrow F_{\mu\nu} \Big|_p = \frac{\partial x^\rho}{\partial \tilde{x}^\mu} \frac{\partial}{\partial x^\rho} \left(\frac{\partial x^\sigma}{\partial \tilde{x}^\nu} \frac{\partial f}{\partial x^\sigma} \right) \Big|_p = \frac{\partial x^\rho}{\partial \tilde{x}^\mu} \frac{\partial x^\sigma}{\partial \tilde{x}^\nu} \frac{\partial^2 f}{\partial x^\rho \partial x^\sigma} \Big|_p + \frac{\partial^2 x^\sigma}{\partial \tilde{x}^\mu \partial \tilde{x}^\nu} \frac{\partial f}{\partial x^\sigma} \Big|_p. \quad (3.2)$$

Given $df|_p = 0$, the second term vanishes, so

$$\tilde{F}_{\mu\nu} \Big|_p \rightarrow \frac{\partial x^\rho}{\partial \tilde{x}^\mu} \frac{\partial x^\sigma}{\partial \tilde{x}^\nu} F_{\rho\sigma} \Big|_p. \quad (3.3)$$

This is exactly the transformation law of a rank $(0, 2)$ tensor. Therefore, f defines a rank $(0, 2)$ tensor at p .

We can also write the tensor in a coordinate-free way as

$$F(X_p, Y_p) = X(Y(f))|_p = X^\mu Y^\nu \frac{\partial^2 f}{\partial x^\mu \partial x^\nu} \Big|_p + X^\mu \frac{\partial Y^\nu}{\partial x^\mu} \frac{\partial f}{\partial x^\nu} \Big|_p. \quad (3.4)$$

Since $df|_p = 0$, this reduces to

$$F(X_p, Y_p) = X^\mu Y^\nu \frac{\partial^2 f}{\partial x^\mu \partial x^\nu} \Big|_p, \quad (3.5)$$

which is bilinear and symmetric.

Transformation of the Metric Determinant, Q4

Let $g_{\mu\nu}$ be a rank $(0, 2)$ tensor. In a basis, one can regard the components $g_{\mu\nu}$ as elements of an $n \times n$ matrix, so that one may define the determinant $g = \det(g_{\mu\nu})$. How does g transform under a change of basis?

Solution. Under a coordinate transformation $x^\mu \rightarrow \tilde{x}^\mu$, we have

$$g \rightarrow \tilde{g} = \det \left(\frac{\partial x^\rho}{\partial \tilde{x}^\mu} \frac{\partial x^\sigma}{\partial \tilde{x}^\nu} g_{\rho\sigma} \right) = \left[\det \left(\frac{\partial x}{\partial \tilde{x}} \right) \right]^2 g. \quad (4.1)$$

Lie Derivatives and Cartan's Magic Formula, Q5

Use the Leibniz rule to derive the formula for the Lie derivative of a 1-form ω , valid in any coordinate basis:

$$(\mathcal{L}_X \omega)_\mu = X^\nu \partial_\nu \omega_\mu + \omega_\nu \partial_\mu X^\nu. \quad (5.1)$$

[Hint: consider $(\mathcal{L}_X \omega)(Y)$ for a vector field Y .] Show that the Lie derivative of a $(0, 2)$ tensor g is

$$(\mathcal{L}_X g)_{\mu\nu} = X^\rho \partial_\rho g_{\mu\nu} + g_{\mu\rho} \partial_\nu X^\rho + g_{\rho\nu} \partial_\mu X^\rho. \quad (5.2)$$

For a p -form η , define $\iota_X \eta$ to be the $(p-1)$ -form that results from contracting a vector field X with the first index of η . Show that for a 1-form ω ,

$$\mathcal{L}_X \omega = \iota_X(d\omega) + d(\iota_X \omega). \quad (5.3)$$

Solution. Consider

$$(\mathcal{L}_X \omega)(Y) = \mathcal{L}_X(\omega_\mu dx^\mu)(Y^\nu \partial_\nu) = X^\sigma (\partial_\sigma \omega_\mu) Y^\mu + \omega_\mu \mathcal{L}_X(dx^\mu)(Y^\nu \partial_\nu), \quad (5.4)$$

and use the fact that d and $\mathcal{L}_X$ commute,

$$\mathcal{L}_X(dx^\mu) = d\mathcal{L}_X(x^\mu) = dX^\mu = \partial_\nu X^\mu dx^\nu. \quad (5.5)$$

Therefore, we find

$$(\mathcal{L}_X \omega)(Y) = X^\sigma (\partial_\sigma \omega_\mu) Y^\mu + \omega_\mu (\partial_\sigma X^\mu) Y^\sigma \implies (\mathcal{L}_X \omega)_\mu = X^\nu \partial_\nu \omega_\mu + \omega_\nu \partial_\mu X^\nu. \quad (5.6)$$

For a rank $(0, 2)$ tensor, the Lie derivative is defined as

$$\mathcal{L}_X g = \lim_{t \rightarrow 0} \frac{((\sigma_{-t})^* g)_p - g_p}{t}, \quad (5.7)$$

where the push-forward is given by

$$(\varphi_*g)(X, Y)\Big|_p = g(\varphi_*^{-1}X, \varphi_*^{-1}Y)\Big|_{\varphi^{-1}(p)}, \quad (5.8)$$

and σ_{-t} is the flow generated by $X \in \mathfrak{X}(M)$. The flow equation, in a local coordinate, is given by

$$X^\mu(x) = \frac{dx^\mu}{dt}, \quad (5.9)$$

which is satisfied by

$$x^\mu(t) = x^\mu(0) + tX^\mu(x) + \mathcal{O}(t^2) \quad (5.10)$$

to first order in t (note the slight abuse of notation $x^\mu(\sigma_t) = x^\mu(t)$). In a chart, the push-forward is given by

$$(\varphi_*g)_{\mu\nu}\Big|_p = \frac{\partial x^\sigma}{\partial x^\mu(\varphi)} \frac{\partial x^\rho}{\partial x^\nu(\varphi)} g_{\sigma\rho}\Big|_{\varphi^{-1}(p)}, \quad (5.11)$$

which gives

$$\begin{aligned} ((\sigma_{-t})_*g)_{\mu\nu}\Big|_p &= (\delta_\mu^\sigma - t\partial_\mu X^\sigma)^{-1}(\delta_\nu^\rho - t\partial_\nu X^\rho)^{-1}g_{\sigma\rho}\Big|_{\sigma_t(p)} + \mathcal{O}(t^2) \\ &= (\delta_\mu^\sigma + t\partial_\mu X^\sigma)(\delta_\nu^\rho + t\partial_\nu X^\rho)(g_{\mu\nu} + tX^\lambda\partial_\lambda g_{\mu\nu})\Big|_p + \dots \\ &= g_{\mu\nu}\Big|_p + t(X^\rho\partial_\rho g_{\mu\nu} + g_{\mu\rho}\partial_\nu X^\rho + g_{\rho\nu}\partial_\mu X^\rho)\Big|_p + \mathcal{O}(t^2), \end{aligned} \quad (5.12)$$

where we have used (5.10) to Taylor expand g around p . Therefore, we find

$$(\mathcal{L}_X g)_{\mu\nu} = X^\rho\partial_\rho g_{\mu\nu} + g_{\mu\rho}\partial_\nu X^\rho + g_{\rho\nu}\partial_\mu X^\rho. \quad (5.13)$$

First, consider

$$\begin{aligned} \iota_X(d\omega) &= \iota_X \frac{1}{2}(\partial_\mu\omega_\nu - \partial_\nu\omega_\mu)dx^\mu \wedge dx^\nu \\ &= \frac{1}{2}(\partial_\mu\omega_\nu - \partial_\nu\omega_\mu)X^\mu dx^\nu - \frac{1}{2}(\partial_\mu\omega_\nu - \partial_\nu\omega_\mu)X^\nu dx^\mu \\ &= X^\mu\partial_\mu\omega_\nu dx^\nu - X^\mu\partial_\nu\omega_\mu dx^\nu. \end{aligned} \quad (5.14)$$

Then,

$$d(\iota_X\omega) = d(\omega_\mu X^\mu) = X^\mu\partial_\nu\omega_\mu dx^\nu + \omega_\mu\partial_\nu X^\mu dx^\nu. \quad (5.15)$$

Therefore, we find

$$\mathcal{L}_X\omega = \iota_X(d\omega) + d(\iota_X\omega). \quad (5.16)$$

Properties of the Exterior Derivative, Q6

Let ω be a p -form and η a q -form. Show that the exterior derivative satisfies the properties

- $d(d\omega) = 0$
- $d(\omega \wedge \eta) = (d\omega) \wedge \eta + (-1)^p \omega \wedge d\eta$
- $d(\varphi^* \omega) = \varphi^*(d\omega)$ where $\varphi : M \rightarrow N$ for some manifolds M and N

Solution.

$$\begin{aligned} d(d\omega) &= d \left(\frac{1}{p!} \frac{\partial \omega_{\mu_1 \dots \mu_p}}{\partial x^\nu} dx^\nu \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \right) \\ &= \frac{1}{p!} \frac{\partial^2 \omega_{\mu_1 \dots \mu_p}}{\partial x^\rho \partial x^\nu} dx^\rho \wedge dx^\nu \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \\ &= 0 \end{aligned} \tag{6.1}$$

by antisymmetry of the wedge product.

$$\begin{aligned} d(\omega \wedge \eta) &= d \left(\frac{1}{p!q!} \omega_{[\mu_1 \dots \mu_p} \eta_{\nu_1 \dots \nu_q]} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \wedge dx^{\nu_1} \wedge \dots \wedge dx^{\nu_q} \right) \\ &= \frac{1}{p!q!} ((\partial_{[\mu} \omega_{\mu_1 \dots \mu_p}]) \eta_{\nu_1 \dots \nu_q]} + \omega_{[\mu_1 \dots \mu_p} \partial_{[\mu} \eta_{\nu_1 \dots \nu_q]}) dx^\mu \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \wedge dx^{\nu_1} \wedge \dots \wedge dx^{\nu_q}. \end{aligned} \tag{6.2}$$

Now consider the wedge product

$$\begin{aligned} (d\omega \wedge \eta)_{\mu\mu_1 \dots \mu_p \nu_1 \dots \nu_q} &= \frac{(p+1)}{(p+1)!q!} (\partial_{[\mu} \omega_{\mu_1 \dots \mu_p}]) \eta_{\nu_1 \dots \nu_q]} \\ &= \frac{1}{p!q!} (\partial_{[\mu} \omega_{\mu_1 \dots \mu_p}]) \eta_{\nu_1 \dots \nu_q]}, \end{aligned} \tag{6.3}$$

which is exactly the first term in (6.2). Similarly, the second term of (6.2) can also be written as $\omega \wedge d\eta$ by permuting μ to the left, which gives a factor of $(-1)^p$. Therefore, we find

$$d(\omega \wedge \eta) = (d\omega) \wedge \eta + (-1)^p \omega \wedge d\eta. \tag{6.4}$$

The pull-back can be written as $\partial \tilde{x}^\mu / \partial x^\nu$. By antisymmetry of the exterior derivative, any term involving $d(\partial \tilde{x}^\mu / \partial x^\nu)$ must vanish. Therefore, the exterior derivative must pass through the pull-back and act on the p -form, ω . That is

$$d(\varphi^* \omega) = \varphi^* d\omega. \tag{6.5}$$

Coordinate-Free Definition of the Exterior Derivative, Q7

The exterior derivative of $\omega \in \Lambda^p(M)$ can be defined as

$$\begin{aligned}
d\omega(X_1, \dots, X_{p+1}) &= \sum_{j=1}^{p+1} (-1)^{j-1} X_j \left(\omega(X_1, \dots, \hat{X}_j, \dots, X_{p+1}) \right) \\
&\quad + \sum_{j < k} (-1)^{j+k} \omega([X_j, X_k], X_1, \dots, \hat{X}_j, \dots, \hat{X}_k, \dots, X_{p+1})
\end{aligned} \tag{7.1}$$

where a hat denotes omission. In a coordinate basis $\omega = \frac{1}{p!} \omega_{\mu_1 \dots \mu_p} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}$, use this definition to determine the components of $d\omega$ in the cases $p = 1$ and $p = 2$.

Solution. For $p = 1$, we have

$$\begin{aligned}
d\omega(X, Y) &= X(\omega(Y)) - Y(\omega(X)) - \omega([X, Y]) \\
&= X^\mu \partial_\mu (\omega_\nu Y^\nu) - Y^\nu \partial_\nu (\omega_\mu X^\mu) - \omega_\rho (X^\mu \partial_\mu Y^\rho - Y^\mu \partial_\mu X^\rho) \\
&= (\partial_\mu \omega_\nu - \partial_\nu \omega_\mu) X^\mu Y^\nu,
\end{aligned} \tag{7.2}$$

so the components read $(d\omega)_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu = 2 \partial_{[\mu} \omega_{\nu]}$, and

$$d\omega = \partial_\mu \omega_\nu dx^\mu \wedge dx^\nu. \tag{7.3}$$

For $p = 2$, with $\omega = \frac{1}{2} \omega_{\mu\nu} dx^\mu \wedge dx^\nu$, we have $\omega(Y, Z) = \omega_{\mu\nu} Y^\mu Z^\nu$, and

$$\begin{aligned}
d\omega(X, Y, Z) &= X(\omega(Y, Z)) - Y(\omega(X, Z)) + Z(\omega(X, Y)) \\
&\quad - \omega([X, Y], Z) + \omega([X, Z], Y) - \omega([Y, Z], X).
\end{aligned} \tag{7.4}$$

Expanding the first piece,

$$X(\omega(Y, Z)) = X^\rho (\partial_\rho \omega_{\mu\nu}) Y^\mu Z^\nu + \omega_{\mu\nu} X^\rho (\partial_\rho Y^\mu) Z^\nu + \omega_{\mu\nu} X^\rho Y^\mu (\partial_\rho Z^\nu), \tag{7.5}$$

and similarly for the other two, while each commutator contributes terms such as

$$\omega([X, Y], Z) = \omega_{\rho\nu} (X^\sigma \partial_\sigma Y^\rho - Y^\sigma \partial_\sigma X^\rho) Z^\nu. \tag{7.6}$$

All terms involving derivatives of the vector fields cancel pairwise after relabeling dummy indices and using antisymmetry of $\omega_{\mu\nu}$. What remains is

$$d\omega(X, Y, Z) = (\partial_\rho \omega_{\mu\nu} + \partial_\mu \omega_{\nu\rho} + \partial_\nu \omega_{\rho\mu}) X^\rho Y^\mu Z^\nu, \tag{7.7}$$

so the components read

$$(d\omega)_{\rho\mu\nu} = \partial_\rho \omega_{\mu\nu} + \partial_\mu \omega_{\nu\rho} + \partial_\nu \omega_{\rho\mu} = 3 \partial_{[\rho} \omega_{\mu\nu]}, \tag{7.8}$$

which gives

$$d\omega = \frac{1}{2} \partial_\rho \omega_{\mu\nu} dx^\rho \wedge dx^\mu \wedge dx^\nu. \tag{7.9}$$

Both cases fit the general pattern $(d\omega)_{\mu_0 \mu_1 \dots \mu_p} = (p+1) \partial_{[\mu_0} \omega_{\mu_1 \dots \mu_p]}$.

Maurer-Cartan Structure Equations on S^3 , Q8

A three-sphere can be parameterized by Euler angles (θ, ϕ, ψ) where $0 < \theta < \pi$, $0 < \phi < 2\pi$, $0 < \psi < 4\pi$. Define the following 1-forms

$$\begin{aligned}\sigma^1 &= -\sin \psi d\theta + \cos \psi \sin \theta d\phi, \\ \sigma^2 &= \cos \psi d\theta + \sin \psi \sin \theta d\phi, \\ \sigma^3 &= d\psi + \cos \theta d\phi.\end{aligned}$$

Show that $d\sigma^1 = \sigma^2 \wedge \sigma^3$ with analogous results for $d\sigma^2$ and $d\sigma^3$.

Solution.

$$d\sigma^1 = -\cos \theta d\psi \wedge d\theta - \sin \psi \sin \theta d\psi \wedge d\phi + \cos \psi \cos \theta d\theta \wedge d\phi, \quad (8.1)$$

and

$$\begin{aligned}\sigma^2 \wedge \sigma^3 &= (\cos \psi d\theta + \sin \psi \sin \theta d\phi) \wedge (d\psi + \cos \theta d\phi) \\ &= \cos \psi d\theta \wedge d\psi + \sin \psi \sin \theta d\phi \wedge d\psi + \cos \psi \cos \theta d\theta \wedge d\phi,\end{aligned} \quad (8.2)$$

which clearly shows

$$d\sigma^1 = \sigma^2 \wedge \sigma^3. \quad (8.3)$$

Similarly, one can obtain analogous results for $d\sigma^2$ and $d\sigma^3$.

Coordinate-Induced Bases and the Poincaré Lemma, Q9

Let $\{e_\mu\}$ be a basis of vector fields, with

$$[e_\mu, e_\nu] = \gamma^\rho_{\mu\nu} e_\rho. \quad (9.1)$$

The functions $\gamma^\rho_{\mu\nu}$ are known as commutator components. For a choice of coordinates $\{x^\mu\}$, we often work with the coordinate induced basis $e_\mu = \{\partial_\mu\}$. Show that, in this case, $[e_\mu, e_\nu] = 0$.

The purpose of this question is to show the converse: that $[e_\mu, e_\nu] = 0$ only for a coordinate induced basis. Consider a general basis $\{e_\mu\}$ and the dual basis $\{f^\mu\}$ of one-forms. In general, these can be expanded as

$$e_\mu = e_\mu^\rho \frac{\partial}{\partial x^\rho} \quad \text{and} \quad f^\mu = f^\mu_\rho dx^\rho \quad (9.2)$$

where $e_\mu^\rho f^\nu_\rho = \delta^\nu_\mu$. Show that

$$e_\mu^\sigma \frac{\partial e_\nu^\lambda}{\partial x^\sigma} - e_\nu^\sigma \frac{\partial e_\mu^\lambda}{\partial x^\sigma} = \gamma^\rho_{\mu\nu} e_\rho^\lambda. \quad (9.3)$$

Hence deduce that

$$e_{\mu}^{\sigma} e_{\nu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} - e_{\nu}^{\sigma} e_{\mu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} = -\gamma^{\rho}_{\mu\nu}, \quad (9.4)$$

and finally that

$$\frac{\partial f^{\rho}_{\sigma}}{\partial x^{\lambda}} - \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} = -\gamma^{\rho}_{\mu\nu} f^{\mu}_{\lambda} f^{\nu}_{\sigma}. \quad (9.5)$$

Use this result, together with the Poincaré lemma, to show that if $[e_{\mu}, e_{\nu}] = 0 \forall \mu, \nu$ then the basis is coordinate induced.

Solution. For coordinate bases, we clearly have

$$[\partial_{\mu}, \partial_{\nu}] = 0. \quad (9.6)$$

Now, consider the commutation relation

$$[e_{\mu}^{\rho} \partial_{\rho}, e_{\nu}^{\sigma} \partial_{\sigma}] = \gamma^{\lambda}_{\mu\nu} e_{\lambda}^{\delta} \partial_{\delta}. \quad (9.7)$$

The left-hand side is

$$[e_{\mu}^{\rho} \partial_{\rho}, e_{\nu}^{\sigma} \partial_{\sigma}] = e_{\mu}^{\rho} (\partial_{\rho} e_{\nu}^{\sigma}) \partial_{\sigma} - e_{\nu}^{\sigma} (\partial_{\sigma} e_{\mu}^{\rho}) \partial_{\rho}. \quad (9.8)$$

After relabeling, we find

$$e_{\mu}^{\sigma} \frac{\partial e_{\nu}^{\lambda}}{\partial x^{\sigma}} - e_{\nu}^{\sigma} \frac{\partial e_{\mu}^{\lambda}}{\partial x^{\sigma}} = \gamma^{\rho}_{\mu\nu} e_{\rho}^{\lambda}. \quad (9.9)$$

Then, consider the expression

$$e_{\mu}^{\sigma} \frac{\partial (e_{\nu}^{\lambda} f^{\rho}_{\lambda})}{\partial x^{\sigma}} - e_{\nu}^{\sigma} \frac{\partial (e_{\mu}^{\lambda} f^{\rho}_{\lambda})}{\partial x^{\sigma}}. \quad (9.10)$$

Since $e_{\mu}^{\rho} f^{\nu}_{\rho} = \delta^{\mu}_{\nu}$, the above expression must vanish. Therefore, we must have

$$\begin{aligned} 0 &= e_{\mu}^{\sigma} \frac{\partial (e_{\nu}^{\lambda} f^{\rho}_{\lambda})}{\partial x^{\sigma}} - e_{\nu}^{\sigma} \frac{\partial (e_{\mu}^{\lambda} f^{\rho}_{\lambda})}{\partial x^{\sigma}} \\ &= e_{\mu}^{\sigma} f^{\rho}_{\lambda} \frac{\partial e_{\nu}^{\lambda}}{\partial x^{\sigma}} - e_{\nu}^{\sigma} f^{\rho}_{\lambda} \frac{\partial e_{\mu}^{\lambda}}{\partial x^{\sigma}} + e_{\mu}^{\sigma} e_{\nu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} - e_{\nu}^{\sigma} e_{\mu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} \\ &= \gamma^{\rho}_{\mu\nu} + e_{\mu}^{\sigma} e_{\nu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} - e_{\nu}^{\sigma} e_{\mu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}}. \end{aligned} \quad (9.11)$$

Therefore, we find

$$e_{\mu}^{\sigma} e_{\nu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} - e_{\nu}^{\sigma} e_{\mu}^{\lambda} \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} = -\gamma^{\rho}_{\mu\nu}. \quad (9.12)$$

Then, contracting with f , we find

$$\frac{\partial f^{\rho}_{\sigma}}{\partial x^{\lambda}} - \frac{\partial f^{\rho}_{\lambda}}{\partial x^{\sigma}} = -\gamma^{\rho}_{\mu\nu} f^{\mu}_{\lambda} f^{\nu}_{\sigma}. \quad (9.13)$$

The Poincaré lemma states that, on $M = \mathbb{R}^n$, closed implies exact. Multiplying the previous expression by $dx^\lambda \wedge dx^\sigma$ and using antisymmetry of the wedge,

$$d\theta^\rho = \frac{1}{2} \left(\frac{\partial f^\rho_\sigma}{\partial x^\lambda} - \frac{\partial f^\rho_\lambda}{\partial x^\sigma} \right) dx^\lambda \wedge dx^\sigma = -\frac{1}{2} \gamma^\rho_{\mu\nu} \theta^\mu \wedge \theta^\nu, \quad (9.14)$$

where $\theta^\rho = f^\rho_\sigma dx^\sigma$ is the dual basis to $e_\mu = e_\mu^\rho \partial_\rho$. Now suppose $\gamma^\rho_{\mu\nu}$ vanish. Then $d\theta^\rho = 0$, so θ^ρ is a closed 1-form. By the Poincaré lemma, on $M = \mathbb{R}^n$ (or any contractible neighbourhood) closed forms are exact, so there exist functions y^ρ with

$$\theta^\rho = dy^\rho. \quad (9.15)$$

The y^ρ are functionally independent because θ^ρ are linearly independent at every point, so they form a coordinate system. In these coordinates, $\theta^\rho(e_\mu) = \delta^\rho_\mu$ becomes $\partial y^\rho / \partial y^\mu_{(e)} = \delta^\rho_\mu$, i.e. $e_\mu = \partial / \partial y^\mu$, a coordinate basis.

Conversely, if $e_\mu = \partial / \partial y^\mu$ for some coordinates y^μ , then $[e_\mu, e_\nu] = [\partial_{y^\mu}, \partial_{y^\nu}] = 0$, so $\gamma^\rho_{\mu\nu} = 0$. Therefore $\{e_\mu\}$ is (locally) a coordinate basis if and only if $\gamma^\rho_{\mu\nu} = 0$.