

Contour Integration via Residue Theorem and Cauchy's Integral Formula

J. E. Santos, Cambridge Part IB Complex Methods, Example Sheet 2, Question 9.

Yucheng (Luca) Jin

Mathematical Methods

1. By integrating the function

$$z^n(z-a)^{-1}(z-a^{-1})^{-1}$$

around the unit circle and applying the residue theorem, evaluate

$$\int_0^{2\pi} \frac{\cos n\theta}{1-2a\cos\theta+a^2} d\theta, \quad (1)$$

where a is real, $a > 1$, and n is a non-negative integer.

2. Obtain the same result using Cauchy's integral formula instead of the residue theorem.

Solution Parametrise the integration by $z = e^{i\theta}$ with $\theta \in [0, 2\pi)$. The integral becomes

$$I = \int_0^{2\pi} d\theta \frac{aie^{in\theta}}{-1+2a\cos\theta-a^2} = \int_0^{2\pi} d\theta \frac{aie^{-in\theta}}{-1+2a\cos\theta-a^2} = -2\pi i \frac{a^{-n+1}}{a^2-1} \quad (2)$$

Therefore,

$$\int_0^{2\pi} \frac{\cos n\theta}{1-2a\cos\theta+a^2} d\theta = -\frac{I}{ai} = 2\pi \frac{a^{-n}}{a^2-1}. \quad (3)$$