

**EEIGM**  
**École Européenne d'Ingénieurs en Génie des Matériaux**

*2<sup>ème</sup>* Année, 1<sup>er</sup> Semestre

**MÉCANIQUE DU SOLIDE DÉFORMABLE**

**NOTES DE COURS**

**Luca Di Stasio**



Cette oeuvre est mise à disposition selon les termes de la  
Licence Creative Commons Attribution - Pas d'Utilisation Commerciale  
4.0 International.

**4 septembre 2020**

## Table des matières

<b>1</b>	<b>Systèmes de coordonnées curvilignes</b>	<b>1</b>
1.1	Formulation analytique . . . . .	1
1.2	Le déplacement infinitésimal d'un point et le jacobien de la transformation	1
1.3	Les vecteurs du repère local naturel (base covariante) . . . . .	2
1.4	L'élément infinitésimal de ligne et le tenseur métrique . . . . .	2
1.5	L'élément infinitésimal de volume . . . . .	3
1.6	Les vecteurs de la base contravariante et le gradient d'une fonction scalaire $\nabla f$ . . . . .	3
1.7	L'opérateur laplacien $\nabla^2$ . . . . .	5



# 1 Systèmes de coordonnées curvilignes

## 1.1 Formulation analytique

$$\begin{cases} x = x(\xi, \eta, \zeta) \\ y = y(\xi, \eta, \zeta) \\ z = z(\xi, \eta, \zeta) \end{cases} \longleftrightarrow \begin{cases} \xi = \xi(x, y, z) \\ \eta = \eta(x, y, z) \\ \zeta = \zeta(x, y, z) \end{cases} \quad (1)$$

## 1.2 Le déplacement infinitésimal d'un point et le jacobien de la transformation

$$\mathbf{r} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \xi \\ \eta \\ \zeta \end{bmatrix} \quad (2)$$

$$d\mathbf{r} = \begin{bmatrix} dx \\ dy \\ dz \end{bmatrix} \quad (3)$$

$$\begin{cases} dx = \frac{\partial x}{\partial \xi} d\xi + \frac{\partial x}{\partial \eta} d\eta + \frac{\partial x}{\partial \zeta} d\zeta \\ dy = \frac{\partial y}{\partial \xi} d\xi + \frac{\partial y}{\partial \eta} d\eta + \frac{\partial y}{\partial \zeta} d\zeta \\ dz = \frac{\partial z}{\partial \xi} d\xi + \frac{\partial z}{\partial \eta} d\eta + \frac{\partial z}{\partial \zeta} d\zeta \end{cases} \quad (4)$$

$$d\mathbf{r} = \begin{bmatrix} dx \\ dy \\ dz \end{bmatrix} = \underbrace{\begin{bmatrix} \frac{\partial x}{\partial \xi} & \frac{\partial x}{\partial \eta} & \frac{\partial x}{\partial \zeta} \\ \frac{\partial y}{\partial \xi} & \frac{\partial y}{\partial \eta} & \frac{\partial y}{\partial \zeta} \\ \frac{\partial z}{\partial \xi} & \frac{\partial z}{\partial \eta} & \frac{\partial z}{\partial \zeta} \end{bmatrix}}_{\mathbf{J}} \begin{bmatrix} d\xi \\ d\eta \\ d\zeta \end{bmatrix} = \mathbf{J} \begin{bmatrix} d\xi \\ d\eta \\ d\zeta \end{bmatrix} \quad (5)$$

### 1.3 Les vecteurs du repère local naturel (base covariante)

$$\begin{aligned} & \left[ \begin{array}{ccc} \frac{\partial x}{\partial \xi} & \frac{\partial x}{\partial \eta} & \frac{\partial x}{\partial \zeta} \\ \frac{\partial y}{\partial \xi} & \frac{\partial y}{\partial \eta} & \frac{\partial y}{\partial \zeta} \\ \frac{\partial z}{\partial \xi} & \frac{\partial z}{\partial \eta} & \frac{\partial z}{\partial \zeta} \end{array} \right] & \left[ \begin{array}{ccc} \frac{\partial x}{\partial \xi} & \frac{\partial x}{\partial \eta} & \frac{\partial x}{\partial \zeta} \\ \frac{\partial y}{\partial \xi} & \frac{\partial y}{\partial \eta} & \frac{\partial y}{\partial \zeta} \\ \frac{\partial z}{\partial \xi} & \frac{\partial z}{\partial \eta} & \frac{\partial z}{\partial \zeta} \end{array} \right] & \left[ \begin{array}{ccc} \frac{\partial x}{\partial \xi} & \frac{\partial x}{\partial \eta} & \frac{\partial x}{\partial \zeta} \\ \frac{\partial y}{\partial \xi} & \frac{\partial y}{\partial \eta} & \frac{\partial y}{\partial \zeta} \\ \frac{\partial z}{\partial \xi} & \frac{\partial z}{\partial \eta} & \frac{\partial z}{\partial \zeta} \end{array} \right] \\ & \frac{\partial \mathbf{r}}{\partial \xi} & \frac{\partial \mathbf{r}}{\partial \eta} & \frac{\partial \mathbf{r}}{\partial \zeta} \end{aligned}$$

$$\begin{aligned} \mathbf{i}_\xi &= \frac{1}{\|\frac{\partial \mathbf{r}}{\partial \xi}\|} \frac{\partial \mathbf{r}}{\partial \xi} = \frac{1}{\sqrt{\left(\frac{\partial x}{\partial \xi}\right)^2 + \left(\frac{\partial y}{\partial \xi}\right)^2 + \left(\frac{\partial z}{\partial \xi}\right)^2}} \begin{bmatrix} \frac{\partial x}{\partial \xi} \\ \frac{\partial y}{\partial \xi} \\ \frac{\partial z}{\partial \xi} \end{bmatrix} \\ \mathbf{j}_\eta &= \frac{1}{\|\frac{\partial \mathbf{r}}{\partial \eta}\|} \frac{\partial \mathbf{r}}{\partial \eta} = \frac{1}{\sqrt{\left(\frac{\partial x}{\partial \eta}\right)^2 + \left(\frac{\partial y}{\partial \eta}\right)^2 + \left(\frac{\partial z}{\partial \eta}\right)^2}} \begin{bmatrix} \frac{\partial x}{\partial \eta} \\ \frac{\partial y}{\partial \eta} \\ \frac{\partial z}{\partial \eta} \end{bmatrix} \\ \mathbf{k}_\zeta &= \frac{1}{\|\frac{\partial \mathbf{r}}{\partial \zeta}\|} \frac{\partial \mathbf{r}}{\partial \zeta} = \frac{1}{\sqrt{\left(\frac{\partial x}{\partial \zeta}\right)^2 + \left(\frac{\partial y}{\partial \zeta}\right)^2 + \left(\frac{\partial z}{\partial \zeta}\right)^2}} \begin{bmatrix} \frac{\partial x}{\partial \zeta} \\ \frac{\partial y}{\partial \zeta} \\ \frac{\partial z}{\partial \zeta} \end{bmatrix} \end{aligned} \quad (6)$$

### 1.4 L'élément infinitésimal de ligne et le tenseur métrique

$$dl^2 = d\mathbf{r}^T d\mathbf{r} = \begin{bmatrix} dx & dy & dz \end{bmatrix} \begin{bmatrix} dx \\ dy \\ dz \end{bmatrix} = dx^2 + dy^2 + dz^2 \quad (7)$$

$$\begin{aligned}
dl^2 &= d\mathbf{r}^T d\mathbf{r} = [d\xi \quad d\eta \quad d\zeta] \mathbf{J}^T \mathbf{J} \begin{bmatrix} d\xi \\ d\eta \\ d\zeta \end{bmatrix} = \\
&= [d\xi \quad d\eta \quad d\zeta] \begin{bmatrix} \frac{\partial x}{\partial \xi} & \frac{\partial y}{\partial \xi} & \frac{\partial z}{\partial \xi} \\ \frac{\partial x}{\partial \eta} & \frac{\partial y}{\partial \eta} & \frac{\partial z}{\partial \eta} \\ \frac{\partial x}{\partial \zeta} & \frac{\partial y}{\partial \zeta} & \frac{\partial z}{\partial \zeta} \end{bmatrix} \begin{bmatrix} \frac{\partial x}{\partial \xi} & \frac{\partial x}{\partial \eta} & \frac{\partial x}{\partial \zeta} \\ \frac{\partial y}{\partial \xi} & \frac{\partial y}{\partial \eta} & \frac{\partial y}{\partial \zeta} \\ \frac{\partial z}{\partial \xi} & \frac{\partial z}{\partial \eta} & \frac{\partial z}{\partial \zeta} \end{bmatrix} \begin{bmatrix} d\xi \\ d\eta \\ d\zeta \end{bmatrix} = \\
&= [d\xi \quad d\eta \quad d\zeta] \underbrace{\begin{bmatrix} \left(\frac{\partial x}{\partial \xi}\right)^2 + \left(\frac{\partial y}{\partial \xi}\right)^2 + \left(\frac{\partial z}{\partial \xi}\right)^2 & \frac{\partial x}{\partial \xi} \frac{\partial x}{\partial \eta} + \frac{\partial y}{\partial \xi} \frac{\partial y}{\partial \eta} + \frac{\partial z}{\partial \xi} \frac{\partial z}{\partial \eta} & \frac{\partial x}{\partial \xi} \frac{\partial x}{\partial \zeta} + \frac{\partial y}{\partial \xi} \frac{\partial y}{\partial \zeta} + \frac{\partial z}{\partial \xi} \frac{\partial z}{\partial \zeta} \\ \frac{\partial x}{\partial \eta} \frac{\partial x}{\partial \xi} + \frac{\partial y}{\partial \eta} \frac{\partial y}{\partial \xi} + \frac{\partial z}{\partial \eta} \frac{\partial z}{\partial \xi} & \left(\frac{\partial x}{\partial \eta}\right)^2 + \left(\frac{\partial y}{\partial \eta}\right)^2 + \left(\frac{\partial z}{\partial \eta}\right)^2 & \frac{\partial x}{\partial \eta} \frac{\partial x}{\partial \zeta} + \frac{\partial y}{\partial \eta} \frac{\partial y}{\partial \zeta} + \frac{\partial z}{\partial \eta} \frac{\partial z}{\partial \zeta} \\ \frac{\partial x}{\partial \zeta} \frac{\partial x}{\partial \xi} + \frac{\partial y}{\partial \zeta} \frac{\partial y}{\partial \xi} + \frac{\partial z}{\partial \zeta} \frac{\partial z}{\partial \xi} & \frac{\partial x}{\partial \zeta} \frac{\partial x}{\partial \eta} + \frac{\partial y}{\partial \zeta} \frac{\partial y}{\partial \eta} + \frac{\partial z}{\partial \zeta} \frac{\partial z}{\partial \eta} & \left(\frac{\partial x}{\partial \zeta}\right)^2 + \left(\frac{\partial y}{\partial \zeta}\right)^2 + \left(\frac{\partial z}{\partial \zeta}\right)^2 \end{bmatrix}}_{\mathbf{g}} \begin{bmatrix} d\xi \\ d\eta \\ d\zeta \end{bmatrix} \quad (8)
\end{aligned}$$

## 1.5 L'élément infinitésimal de volume

$$dV = \frac{\partial \mathbf{r}}{\partial \xi} d\xi \cdot \left( \frac{\partial \mathbf{r}}{\partial \eta} d\eta \wedge \frac{\partial \mathbf{r}}{\partial \zeta} d\zeta \right) = \det \begin{bmatrix} \left(\frac{\partial \mathbf{r}}{\partial \xi}\right)^T \\ \left(\frac{\partial \mathbf{r}}{\partial \eta}\right)^T \\ \left(\frac{\partial \mathbf{r}}{\partial \zeta}\right)^T \end{bmatrix} d\xi d\eta d\zeta = \det(\mathbf{J}^T) d\xi d\eta d\zeta \quad (9)$$

## 1.6 Les vecteurs de la base contravariante et le gradient d'un fonction scalaire $\nabla f$

$$f(x, y, z) = f(x(\xi, \eta, \zeta), y(\xi, \eta, \zeta), z(\xi, \eta, \zeta)) \quad (10)$$

$$f(\xi, \eta, \zeta) = f(\xi(x, y, z), \eta(x, y, z), \zeta(x, y, z)) \quad (11)$$

$$\nabla f_{xyz} = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} = \frac{\partial f}{\partial x} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \frac{\partial f}{\partial y} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + \frac{\partial f}{\partial z} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \frac{\partial f}{\partial x} \mathbf{i}^x + \frac{\partial f}{\partial y} \mathbf{j}^y + \frac{\partial f}{\partial z} \mathbf{k}^z \quad (12)$$

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} = \begin{bmatrix} \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial x} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial x} \\ \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial y} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial y} \\ \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial z} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial z} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial z} \end{bmatrix} = \frac{\partial f}{\partial \xi} \begin{bmatrix} \frac{\partial \xi}{\partial x} \\ \frac{\partial \xi}{\partial y} \\ \frac{\partial \xi}{\partial z} \end{bmatrix} + \frac{\partial f}{\partial \eta} \begin{bmatrix} \frac{\partial \eta}{\partial x} \\ \frac{\partial \eta}{\partial y} \\ \frac{\partial \eta}{\partial z} \end{bmatrix} + \frac{\partial f}{\partial \zeta} \begin{bmatrix} \frac{\partial \zeta}{\partial x} \\ \frac{\partial \zeta}{\partial y} \\ \frac{\partial \zeta}{\partial z} \end{bmatrix} \quad (13)$$

$$\begin{aligned} \mathbf{i}^\xi &= \frac{1}{\sqrt{\left(\frac{\partial \xi}{\partial x}\right)^2 + \left(\frac{\partial \xi}{\partial y}\right)^2 + \left(\frac{\partial \xi}{\partial z}\right)^2}} \begin{bmatrix} \frac{\partial \xi}{\partial x} \\ \frac{\partial \xi}{\partial y} \\ \frac{\partial \xi}{\partial z} \end{bmatrix} \\ \mathbf{j}^\eta &= \frac{1}{\sqrt{\left(\frac{\partial \eta}{\partial x}\right)^2 + \left(\frac{\partial \eta}{\partial y}\right)^2 + \left(\frac{\partial \eta}{\partial z}\right)^2}} \begin{bmatrix} \frac{\partial \eta}{\partial x} \\ \frac{\partial \eta}{\partial y} \\ \frac{\partial \eta}{\partial z} \end{bmatrix} \\ \mathbf{k}^\zeta &= \frac{1}{\sqrt{\left(\frac{\partial \zeta}{\partial x}\right)^2 + \left(\frac{\partial \zeta}{\partial y}\right)^2 + \left(\frac{\partial \zeta}{\partial z}\right)^2}} \begin{bmatrix} \frac{\partial \zeta}{\partial x} \\ \frac{\partial \zeta}{\partial y} \\ \frac{\partial \zeta}{\partial z} \end{bmatrix} \end{aligned} \quad (14)$$

$$\begin{aligned} \nabla_{\xi\eta\zeta} f &= \frac{\partial f}{\partial \xi} \sqrt{\left(\frac{\partial \xi}{\partial x}\right)^2 + \left(\frac{\partial \xi}{\partial y}\right)^2 + \left(\frac{\partial \xi}{\partial z}\right)^2} \cdot \mathbf{i}^\xi + \\ &\quad + \frac{\partial f}{\partial \eta} \sqrt{\left(\frac{\partial \eta}{\partial x}\right)^2 + \left(\frac{\partial \eta}{\partial y}\right)^2 + \left(\frac{\partial \eta}{\partial z}\right)^2} \cdot \mathbf{j}^\eta + \\ &\quad + \frac{\partial f}{\partial \zeta} \sqrt{\left(\frac{\partial \zeta}{\partial x}\right)^2 + \left(\frac{\partial \zeta}{\partial y}\right)^2 + \left(\frac{\partial \zeta}{\partial z}\right)^2} \cdot \mathbf{k}^\zeta = \\ &= \begin{bmatrix} \frac{\partial f}{\partial \xi} \sqrt{\left(\frac{\partial \xi}{\partial x}\right)^2 + \left(\frac{\partial \xi}{\partial y}\right)^2 + \left(\frac{\partial \xi}{\partial z}\right)^2} \\ \frac{\partial f}{\partial \eta} \sqrt{\left(\frac{\partial \eta}{\partial x}\right)^2 + \left(\frac{\partial \eta}{\partial y}\right)^2 + \left(\frac{\partial \eta}{\partial z}\right)^2} \\ \frac{\partial f}{\partial \zeta} \sqrt{\left(\frac{\partial \zeta}{\partial x}\right)^2 + \left(\frac{\partial \zeta}{\partial y}\right)^2 + \left(\frac{\partial \zeta}{\partial z}\right)^2} \end{bmatrix} \end{aligned} \quad (15)$$

## 1.7 L'opérateur laplacien $\nabla^2$

$$f(x, y, z) = f(x(\xi, \eta, \zeta), y(\xi, \eta, \zeta), z(\xi, \eta, \zeta)) \quad (16)$$

$$f(\xi, \eta, \zeta) = f(\xi(x, y, z), \eta(x, y, z), \zeta(x, y, z)) \quad (17)$$

$$\nabla_{xyz}^2 f(x, y, z) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \quad (18)$$

$$\begin{cases} \frac{\partial f}{\partial x} = \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial x} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial x} \\ \frac{\partial f}{\partial y} = \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial y} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial y} \\ \frac{\partial f}{\partial z} = \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial z} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial z} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial z} \end{cases} \quad (19)$$

$$\begin{cases} \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial x} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial x} \right) = \\ \quad = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial \xi} \right) \frac{\partial \xi}{\partial x} + \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial \eta} \right) \frac{\partial \eta}{\partial x} + \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial \zeta} \right) \frac{\partial \zeta}{\partial x} + \frac{\partial f}{\partial \xi} \frac{\partial^2 \xi}{\partial x^2} + \frac{\partial f}{\partial \eta} \frac{\partial^2 \eta}{\partial x^2} + \frac{\partial f}{\partial \zeta} \frac{\partial^2 \zeta}{\partial x^2} \\ \frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial y} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial y} \right) = \\ \quad = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial \xi} \right) \frac{\partial \xi}{\partial y} + \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial \eta} \right) \frac{\partial \eta}{\partial y} + \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial \zeta} \right) \frac{\partial \zeta}{\partial y} + \frac{\partial f}{\partial \xi} \frac{\partial^2 \xi}{\partial y^2} + \frac{\partial f}{\partial \eta} \frac{\partial^2 \eta}{\partial y^2} + \frac{\partial f}{\partial \zeta} \frac{\partial^2 \zeta}{\partial y^2} \\ \frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} \left( \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial z} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial z} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial z} \right) = \\ \quad = \frac{\partial}{\partial z} \left( \frac{\partial f}{\partial \xi} \right) \frac{\partial \xi}{\partial z} + \frac{\partial}{\partial z} \left( \frac{\partial f}{\partial \eta} \right) \frac{\partial \eta}{\partial z} + \frac{\partial}{\partial z} \left( \frac{\partial f}{\partial \zeta} \right) \frac{\partial \zeta}{\partial z} + \frac{\partial f}{\partial \xi} \frac{\partial^2 \xi}{\partial z^2} + \frac{\partial f}{\partial \eta} \frac{\partial^2 \eta}{\partial z^2} + \frac{\partial f}{\partial \zeta} \frac{\partial^2 \zeta}{\partial z^2} \end{cases} \quad (20)$$

$$\begin{cases} \frac{\partial}{\partial x} = \left( \frac{\partial \xi}{\partial x} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial x} \frac{\partial}{\partial \eta} + \frac{\partial \zeta}{\partial x} \frac{\partial}{\partial \zeta} \right) \\ \frac{\partial}{\partial y} = \left( \frac{\partial \xi}{\partial y} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial y} \frac{\partial}{\partial \eta} + \frac{\partial \zeta}{\partial y} \frac{\partial}{\partial \zeta} \right) \\ \frac{\partial}{\partial z} = \left( \frac{\partial \xi}{\partial z} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial z} \frac{\partial}{\partial \eta} + \frac{\partial \zeta}{\partial z} \frac{\partial}{\partial \zeta} \right) \end{cases} \quad (21)$$



(22)

(23)

(24)

(25)

$$\begin{aligned}
\frac{\partial^2 f}{\partial x^2} = & \left( \frac{\partial \xi}{\partial x} \right)^2 \frac{\partial^2 f}{\partial \xi^2} + \left( \frac{\partial \eta}{\partial x} \right)^2 \frac{\partial^2 f}{\partial \eta^2} + \left( \frac{\partial \zeta}{\partial x} \right)^2 \frac{\partial^2 f}{\partial \zeta^2} + \\
& + 2 \left( \frac{\partial \xi}{\partial x} \right) \left( \frac{\partial \eta}{\partial x} \right) \frac{\partial^2 f}{\partial \xi \partial \eta} + 2 \left( \frac{\partial \xi}{\partial x} \right) \left( \frac{\partial \zeta}{\partial x} \right) \frac{\partial^2 f}{\partial \xi \partial \zeta} + 2 \left( \frac{\partial \eta}{\partial x} \right) \left( \frac{\partial \zeta}{\partial x} \right) \frac{\partial^2 f}{\partial \eta \partial \zeta} + \\
& + \frac{\partial f}{\partial \xi} \frac{\partial^2 \xi}{\partial x^2} + \frac{\partial f}{\partial \eta} \frac{\partial^2 \eta}{\partial x^2} + \frac{\partial f}{\partial \zeta} \frac{\partial^2 \zeta}{\partial x^2}
\end{aligned} \quad (26)$$

$$\begin{aligned}
\frac{\partial^2 f}{\partial y^2} = & \left( \frac{\partial \xi}{\partial y} \right)^2 \frac{\partial^2 f}{\partial \xi^2} + \left( \frac{\partial \eta}{\partial y} \right)^2 \frac{\partial^2 f}{\partial \eta^2} + \left( \frac{\partial \zeta}{\partial y} \right)^2 \frac{\partial^2 f}{\partial \zeta^2} + \\
& + 2 \left( \frac{\partial \xi}{\partial y} \right) \left( \frac{\partial \eta}{\partial y} \right) \frac{\partial^2 f}{\partial \xi \partial \eta} + 2 \left( \frac{\partial \xi}{\partial y} \right) \left( \frac{\partial \zeta}{\partial y} \right) \frac{\partial^2 f}{\partial \xi \partial \zeta} + 2 \left( \frac{\partial \eta}{\partial y} \right) \left( \frac{\partial \zeta}{\partial y} \right) \frac{\partial^2 f}{\partial \eta \partial \zeta} + \\
& + \frac{\partial f}{\partial \xi} \frac{\partial^2 \xi}{\partial y^2} + \frac{\partial f}{\partial \eta} \frac{\partial^2 \eta}{\partial y^2} + \frac{\partial f}{\partial \zeta} \frac{\partial^2 \zeta}{\partial y^2}
\end{aligned} \quad (27)$$

$$\begin{aligned}
\frac{\partial^2 f}{\partial z^2} = & \left( \frac{\partial \xi}{\partial z} \right)^2 \frac{\partial^2 f}{\partial \xi^2} + \left( \frac{\partial \eta}{\partial z} \right)^2 \frac{\partial^2 f}{\partial \eta^2} + \left( \frac{\partial \zeta}{\partial z} \right)^2 \frac{\partial^2 f}{\partial \zeta^2} + \\
& + 2 \left( \frac{\partial \xi}{\partial z} \right) \left( \frac{\partial \eta}{\partial z} \right) \frac{\partial^2 f}{\partial \xi \partial \eta} + 2 \left( \frac{\partial \xi}{\partial z} \right) \left( \frac{\partial \zeta}{\partial z} \right) \frac{\partial^2 f}{\partial \xi \partial \zeta} + 2 \left( \frac{\partial \eta}{\partial z} \right) \left( \frac{\partial \zeta}{\partial z} \right) \frac{\partial^2 f}{\partial \eta \partial \zeta} + \\
& + \frac{\partial f}{\partial \xi} \frac{\partial^2 \xi}{\partial z^2} + \frac{\partial f}{\partial \eta} \frac{\partial^2 \eta}{\partial z^2} + \frac{\partial f}{\partial \zeta} \frac{\partial^2 \zeta}{\partial z^2}
\end{aligned} \quad (28)$$

$$\begin{aligned}
\nabla_{\xi\eta\zeta}^2 f(\xi, \eta, \zeta) = & \left[ \left( \frac{\partial \xi}{\partial x} \right)^2 + \left( \frac{\partial \xi}{\partial y} \right)^2 + \left( \frac{\partial \xi}{\partial z} \right)^2 \right] \frac{\partial^2 f}{\partial \xi^2} + \\
& + \left[ \left( \frac{\partial \eta}{\partial x} \right)^2 + \left( \frac{\partial \eta}{\partial y} \right)^2 + \left( \frac{\partial \eta}{\partial z} \right)^2 \right] \frac{\partial^2 f}{\partial \eta^2} + \\
& + \left[ \left( \frac{\partial \zeta}{\partial x} \right)^2 + \left( \frac{\partial \zeta}{\partial y} \right)^2 + \left( \frac{\partial \zeta}{\partial z} \right)^2 \right] \frac{\partial^2 f}{\partial \zeta^2} + \\
& + 2 \left[ \left( \frac{\partial \xi}{\partial x} \right) \left( \frac{\partial \eta}{\partial x} \right) + \left( \frac{\partial \xi}{\partial y} \right) \left( \frac{\partial \eta}{\partial y} \right) + \left( \frac{\partial \xi}{\partial z} \right) \left( \frac{\partial \eta}{\partial z} \right) \right] \frac{\partial^2 f}{\partial \xi \partial \eta} + \quad (29) \\
& + 2 \left[ \left( \frac{\partial \eta}{\partial x} \right) \left( \frac{\partial \zeta}{\partial x} \right) + \left( \frac{\partial \eta}{\partial y} \right) \left( \frac{\partial \zeta}{\partial y} \right) + \left( \frac{\partial \eta}{\partial z} \right) \left( \frac{\partial \zeta}{\partial z} \right) \right] \frac{\partial^2 f}{\partial \eta \partial \zeta} + \\
& + 2 \left[ \left( \frac{\partial \xi}{\partial x} \right) \left( \frac{\partial \zeta}{\partial x} \right) + \left( \frac{\partial \xi}{\partial y} \right) \left( \frac{\partial \zeta}{\partial y} \right) + \left( \frac{\partial \xi}{\partial z} \right) \left( \frac{\partial \zeta}{\partial z} \right) \right] \frac{\partial^2 f}{\partial \xi \partial \zeta} + \\
& + \left( \frac{\partial^2 \xi}{\partial x^2} + \frac{\partial^2 \xi}{\partial y^2} + \frac{\partial^2 \xi}{\partial z^2} \right) \frac{\partial f}{\partial \xi} + \left( \frac{\partial^2 \eta}{\partial x^2} + \frac{\partial^2 \eta}{\partial y^2} + \frac{\partial^2 \eta}{\partial z^2} \right) \frac{\partial f}{\partial \eta} + \\
& + \left( \frac{\partial^2 \zeta}{\partial x^2} + \frac{\partial^2 \zeta}{\partial y^2} + \frac{\partial^2 \zeta}{\partial z^2} \right) \frac{\partial f}{\partial \zeta}
\end{aligned}$$