

**EEIGM**  
**École Européenne d'Ingénieurs en Génie des Matériaux**

*2<sup>ème</sup>* Année, 1<sup>er</sup> Semestre

**MÉCANIQUE DU SOLIDE DÉFORMABLE**

**TRAVAUX DIRIGÉS**

**Luca Di Stasio**



Cette oeuvre est mise à disposition selon les termes de la  
Licence Creative Commons Attribution - Pas d'Utilisation Commerciale  
4.0 International.

**6 août 2020**

# Contents

<b>List of Acronyms</b>	<b>iii</b>
<b>List of Symbols</b>	<b>v</b>
<b>Abstract</b>	<b>vii</b>
<b>1. Systèmes de coordonnées curvilignes</b>	<b>1</b>
1.1. Énoncé . . . . .	1
1.1.1. Problème A . . . . .	1
1.1.2. Problème B . . . . .	1
1.2. Corrigé . . . . .	1
1.2.1. Problème A . . . . .	1
1.2.2. Problème B . . . . .	4
<b>2. Second section</b>	<b>7</b>
<b>A. First appendix</b>	<b>9</b>



## **List of Acronyms**

---

## *List of Acronyms*

---

## List of Symbols



## **Abstract**





# 1. Systèmes de coordonnées curvilignes

## 1.1. Énoncé

### 1.1.1. Problème A

### 1.1.2. Problème B

## 1.2. Corrigé

### 1.2.1. Problème A

$$\begin{cases} x = x(\xi, \eta, \zeta) \\ y = y(\xi, \eta, \zeta) \\ z = z(\xi, \eta, \zeta) \end{cases} \longleftrightarrow \begin{cases} \xi = \xi(x, y, z) \\ \eta = \eta(x, y, z) \\ \zeta = \zeta(x, y, z) \end{cases} \quad (1)$$

$$f(x, y, z) = f(x(\xi, \eta, \zeta), y(\xi, \eta, \zeta), z(\xi, \eta, \zeta)) \quad (2)$$

$$f(\xi, \eta, \zeta) = f(\xi(x, y, z), \eta(x, y, z), \zeta(x, y, z)) \quad (3)$$

$$\nabla_{xyz}^2 f(x, y, z) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \quad (4)$$

$$\begin{cases} \frac{\partial f}{\partial x} = \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial x} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial x} \\ \frac{\partial f}{\partial y} = \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial y} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial y} \\ \frac{\partial f}{\partial z} = \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial z} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial z} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial z} \end{cases} \quad (5)$$





$$\begin{aligned}
\nabla_{\xi\eta\zeta}^2 f(\xi, \eta, \zeta) = & \left[ \left( \frac{\partial \xi}{\partial x} \right)^2 + \left( \frac{\partial \xi}{\partial y} \right)^2 + \left( \frac{\partial \xi}{\partial z} \right)^2 \right] \frac{\partial^2 f}{\partial \xi^2} + \\
& + \left[ \left( \frac{\partial \eta}{\partial x} \right)^2 + \left( \frac{\partial \eta}{\partial y} \right)^2 + \left( \frac{\partial \eta}{\partial z} \right)^2 \right] \frac{\partial^2 f}{\partial \eta^2} + \\
& + \left[ \left( \frac{\partial \zeta}{\partial x} \right)^2 + \left( \frac{\partial \zeta}{\partial y} \right)^2 + \left( \frac{\partial \zeta}{\partial z} \right)^2 \right] \frac{\partial^2 f}{\partial \zeta^2} + \\
& + 2 \left[ \left( \frac{\partial \xi}{\partial x} \right) \left( \frac{\partial \eta}{\partial x} \right) + \left( \frac{\partial \xi}{\partial y} \right) \left( \frac{\partial \eta}{\partial y} \right) + \left( \frac{\partial \xi}{\partial z} \right) \left( \frac{\partial \eta}{\partial z} \right) \right] \frac{\partial^2 f}{\partial \xi \partial \eta} + \\
& + 2 \left[ \left( \frac{\partial \eta}{\partial x} \right) \left( \frac{\partial \zeta}{\partial x} \right) + \left( \frac{\partial \eta}{\partial y} \right) \left( \frac{\partial \zeta}{\partial y} \right) + \left( \frac{\partial \eta}{\partial z} \right) \left( \frac{\partial \zeta}{\partial z} \right) \right] \frac{\partial^2 f}{\partial \eta \partial \zeta} + \\
& + 2 \left[ \left( \frac{\partial \xi}{\partial x} \right) \left( \frac{\partial \zeta}{\partial x} \right) + \left( \frac{\partial \xi}{\partial y} \right) \left( \frac{\partial \zeta}{\partial y} \right) + \left( \frac{\partial \xi}{\partial z} \right) \left( \frac{\partial \zeta}{\partial z} \right) \right] \frac{\partial^2 f}{\partial \xi \partial \zeta}
\end{aligned} \tag{13}$$

### Coordonnées cylindriques

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases} \longleftrightarrow \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \tan^{-1} \left( \frac{y}{x} \right) \\ z = z \end{cases} \tag{14}$$

$$\begin{cases} \frac{\partial r}{\partial x} = \frac{1}{2} \frac{2x}{\sqrt{x^2 + y^2}} = \frac{r' \cos \theta}{r'} \\ \frac{\partial r}{\partial y} = \frac{1}{2} \frac{2y}{\sqrt{x^2 + y^2}} = \frac{r' \sin \theta}{r'} \\ \frac{\partial r}{\partial z} = 0 \end{cases} \tag{15}$$

### Coordonnées sphériques

#### 1.2.2. Problème B

$$\begin{cases} x = x(\xi, \eta, \zeta) \\ y = y(\xi, \eta, \zeta) \\ z = z(\xi, \eta, \zeta) \end{cases} \longleftrightarrow \begin{cases} \xi = \xi(x, y, z) \\ \eta = \eta(x, y, z) \\ \zeta = \zeta(x, y, z) \end{cases} \tag{16}$$

$$f(x, y, z) = f(x(\xi, \eta, \zeta), y(\xi, \eta, \zeta), z(\xi, \eta, \zeta)) \tag{17}$$

$$f(\xi, \eta, \zeta) = f(\xi(x, y, z), \eta(x, y, z), \zeta(x, y, z)) \quad (18)$$

$$\nabla f_{xyz} = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} = \frac{\partial f}{\partial x} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \frac{\partial f}{\partial y} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + \frac{\partial f}{\partial z} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \frac{\partial f}{\partial x} \mathbf{i}_x + \frac{\partial f}{\partial y} \mathbf{j}_y + \frac{\partial f}{\partial z} \mathbf{k}_z \quad (19)$$

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} = \begin{bmatrix} \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial x} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial x} \\ \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial y} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial y} \\ \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial z} + \frac{\partial f}{\partial \eta} \frac{\partial \eta}{\partial z} + \frac{\partial f}{\partial \zeta} \frac{\partial \zeta}{\partial z} \end{bmatrix} = \frac{\partial f}{\partial \xi} \begin{bmatrix} \frac{\partial \xi}{\partial x} \\ \frac{\partial \xi}{\partial y} \\ \frac{\partial \xi}{\partial z} \end{bmatrix} + \frac{\partial f}{\partial \eta} \begin{bmatrix} \frac{\partial \eta}{\partial x} \\ \frac{\partial \eta}{\partial y} \\ \frac{\partial \eta}{\partial z} \end{bmatrix} + \frac{\partial f}{\partial \zeta} \begin{bmatrix} \frac{\partial \zeta}{\partial x} \\ \frac{\partial \zeta}{\partial y} \\ \frac{\partial \zeta}{\partial z} \end{bmatrix} \quad (20)$$

$$\begin{aligned} \mathbf{i}_\xi &= \frac{1}{\sqrt{\left(\frac{\partial \xi}{\partial x}\right)^2 + \left(\frac{\partial \xi}{\partial y}\right)^2 + \left(\frac{\partial \xi}{\partial z}\right)^2}} \begin{bmatrix} \frac{\partial \xi}{\partial x} \\ \frac{\partial \xi}{\partial y} \\ \frac{\partial \xi}{\partial z} \end{bmatrix} \\ \mathbf{j}_\eta &= \frac{1}{\sqrt{\left(\frac{\partial \eta}{\partial x}\right)^2 + \left(\frac{\partial \eta}{\partial y}\right)^2 + \left(\frac{\partial \eta}{\partial z}\right)^2}} \begin{bmatrix} \frac{\partial \eta}{\partial x} \\ \frac{\partial \eta}{\partial y} \\ \frac{\partial \eta}{\partial z} \end{bmatrix} \\ \mathbf{k}_\zeta &= \frac{1}{\sqrt{\left(\frac{\partial \zeta}{\partial x}\right)^2 + \left(\frac{\partial \zeta}{\partial y}\right)^2 + \left(\frac{\partial \zeta}{\partial z}\right)^2}} \begin{bmatrix} \frac{\partial \zeta}{\partial x} \\ \frac{\partial \zeta}{\partial y} \\ \frac{\partial \zeta}{\partial z} \end{bmatrix} \end{aligned} \quad (21)$$

$$\begin{aligned}\nabla_{\xi\eta\zeta}f &= \frac{\partial f}{\partial \xi} \sqrt{\left(\frac{\partial \xi}{\partial x}\right)^2 + \left(\frac{\partial \xi}{\partial y}\right)^2 + \left(\frac{\partial \xi}{\partial z}\right)^2} \cdot \mathbf{i}_\xi + \\ &\quad + \frac{\partial f}{\partial \eta} \sqrt{\left(\frac{\partial \eta}{\partial x}\right)^2 + \left(\frac{\partial \eta}{\partial y}\right)^2 + \left(\frac{\partial \eta}{\partial z}\right)^2} \cdot \mathbf{j}_\eta + \\ &\quad + \frac{\partial f}{\partial \zeta} \sqrt{\left(\frac{\partial \zeta}{\partial x}\right)^2 + \left(\frac{\partial \zeta}{\partial y}\right)^2 + \left(\frac{\partial \zeta}{\partial z}\right)^2} \cdot \mathbf{k}_\zeta = \\ &= \begin{bmatrix} \frac{\partial f}{\partial \xi} \sqrt{\left(\frac{\partial \xi}{\partial x}\right)^2 + \left(\frac{\partial \xi}{\partial y}\right)^2 + \left(\frac{\partial \xi}{\partial z}\right)^2} \\ \frac{\partial f}{\partial \eta} \sqrt{\left(\frac{\partial \eta}{\partial x}\right)^2 + \left(\frac{\partial \eta}{\partial y}\right)^2 + \left(\frac{\partial \eta}{\partial z}\right)^2} \\ \frac{\partial f}{\partial \zeta} \sqrt{\left(\frac{\partial \zeta}{\partial x}\right)^2 + \left(\frac{\partial \zeta}{\partial y}\right)^2 + \left(\frac{\partial \zeta}{\partial z}\right)^2} \end{bmatrix} \end{aligned} \tag{22}$$

**Coordonnées cylindriques**

**Coordonnées sphériques**

## **2. Second section**





## **A. First appendix**

