

# Mecânica Quântica Avançada

## Lista 1

Lucas Froguel  
IFT

### List of Exercises

1	Exercise (4.1)	1
2	Exercise (4.4.2 - 1)	1
3	Exercise (4.4.2 - 2)	2
4	Exercise (4.4.2 - 3)	3
5	Exercise (4.4.3)	4
6	Exercise (4.4.4)	4

**Exercise 1** (4.1). *Show that a necessary and sufficient condition for  $|\psi\rangle$  to be an eigenvector of a Hermitian operator  $A$  is that the dispersion (4.8)  $\Delta_\psi A = 0$ .*

**Answer.** Vamos iniciar mostrando que se  $|\psi\rangle = 0$ , então  $\Delta_\psi A = 0$ . Ora, por definição:

$$\Delta_\psi A = \langle \psi | A^2 | \psi \rangle - \langle \psi | A | \psi \rangle^2 \quad (1)$$

É fácil ver que

$$\langle \psi | A^2 | \psi \rangle = \langle \psi | A^\dagger | A | \psi \rangle = a^2 \quad (2)$$

E também

$$\langle \psi | A | \psi \rangle^2 = (a)^2 = a^2 \quad (3)$$

Logo, é trivial que  $\Delta_\psi A = 0$ .

Agora vamos assumir que a dispersão é nula, ou seja,  $\Delta A = 0$ . Então, por definição:

$$0 = \sqrt{\langle A^2 \rangle - \langle A \rangle^2} \quad (4)$$

$$\Rightarrow \langle A^2 \rangle = \langle A \rangle^2 \quad (5)$$

---

**Exercise 2** (4.4.2 - 1). *Let  $|\psi\rangle$  be a vector (not normalized) in the Hilbert space of states and  $H$  be a Hamiltonian. The expectation value  $\langle H \rangle_\psi$  is*

$$\langle \psi \rangle_\psi = \frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle} \quad (6)$$

*Show that if the minimum of this expectation value is obtained for  $|\psi\rangle = |\psi_m\rangle$  and the maximum for  $|\psi\rangle = |\psi_M\rangle$ , then*

$$H |\psi_m\rangle = E_m |\psi_m\rangle, \quad H |\psi_M\rangle = E_M |\psi_M\rangle \quad (7)$$

*where  $E_m$  and  $E_M$  are the smallest and largest eigenvalues.*

**Answer.** É evidente que

$$\langle H \rangle_{\psi_m} = \frac{\langle \psi_m | H | \psi_m \rangle}{\langle \psi_m | \psi_m \rangle} = E_m \quad (8)$$

Portanto, é evidente que se  $\langle H \rangle_{\psi_m}$  for mínimo, então  $E_m$  também é. Vale um raciocínio análogo para  $E_M = \langle H \rangle_{\psi_M}$ .

**Exercise 3** (4.4.2 - 2). *We assume that the vector  $|\varphi\rangle$  depends on a parameter  $\alpha : |\varphi\rangle = |\varphi(\alpha)\rangle$ . Show that if*

$$\left. \frac{\partial \langle H \rangle_{\varphi(\alpha)}}{\partial \alpha} \right|_{\alpha=\alpha_0} = 0, \quad (9)$$

*then  $E_m \leq \langle H \rangle_{\varphi(\alpha_0)}$  if  $\alpha_0$  corresponds to a minimum of  $\langle H \rangle_{\varphi(\alpha)}$ , and  $\langle H \rangle_{\varphi(\alpha_0)} \leq E_M$  if  $\alpha_0$  corresponds to a maximum. This result forms the basis of an approximation method called the variational method (Section 14.1.4).*

**Answer.** Vamos abrir a derivada:

$$\frac{\partial \langle H \rangle_{\varphi(\alpha)}}{\partial \alpha} = \frac{1}{\langle \psi | \psi \rangle} (\langle \psi | H | \partial_\alpha \psi \rangle + (\partial_\alpha \langle \psi |) | H | \psi \rangle) - \frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle^2} ((\partial_\alpha \langle \psi |) | \psi \rangle + \langle \psi | \partial_\alpha \psi \rangle) \quad (10)$$

Então, em  $\alpha_0$

$$\langle \psi | H | \partial_\alpha \psi \rangle + \langle \partial_\alpha \psi | H | \psi \rangle = \frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle} (\langle \partial_\alpha \psi | \psi \rangle + \langle \psi | \partial_\alpha \psi \rangle) \quad (11)$$

Isolando a quantidade de interesse:

$$\langle H \rangle_{\varphi(\alpha_0)} = \frac{\langle \psi | H | \partial_\alpha \psi \rangle + \langle \partial_\alpha \psi | H | \psi \rangle}{\langle \partial_\alpha \psi | \psi \rangle + \langle \psi | \partial_\alpha \psi \rangle} \quad (12)$$

Podemos reescrever, usando que  $H = H^\dagger$ :

$$\langle H \rangle_{\varphi(\alpha_0)} = \frac{\langle \partial_\alpha \psi | H | \psi \rangle^\dagger + \langle \partial_\alpha \psi | H | \psi \rangle}{\langle \partial_\alpha \psi | \psi \rangle^\dagger + \langle \partial_\alpha \psi | \psi \rangle} \quad (13)$$

Podemos expandir qualquer estado usando os autokets do hamiltoniano, que são ortonormais:

$$|\psi\rangle = \sum c_j |\psi_j\rangle \quad (14)$$

De modo que

$$\partial_\alpha |\psi\rangle = \sum \frac{\partial c_j}{\partial \alpha} |\psi_j\rangle \quad (15)$$

Assim, vale que

$$\langle \partial_\alpha \psi | \psi \rangle = \sum c_j \frac{\partial c_j^*}{\partial \alpha} \quad (16)$$

E também

$$\langle \partial_\alpha \psi | H | \psi \rangle = \sum E_j c_j \frac{\partial c_j^*}{\partial \alpha} \quad (17)$$

Considere o denominador:

$$\begin{aligned}
\langle \partial_\alpha \psi | \psi \rangle^\dagger + \langle \partial_\alpha \psi | \psi \rangle &= \sum c_j \frac{\partial c_j^*}{\partial \alpha} + c_j^* \frac{\partial c_j}{\partial \alpha} \\
&= \sum \frac{\partial}{\partial \alpha} (c_j^* c_j) \\
&= \sum \partial_\alpha |c_j|^2 \\
&= \partial_\alpha \sum |c_j|^2
\end{aligned}$$

Considerando, enfim, o numerador e fazendo os mesmos cálculos:

$$\begin{aligned}
\langle \partial_\alpha \psi | H | \psi \rangle^\dagger + \langle \partial_\alpha \psi | H | \psi \rangle &= \sum E_j c_j \frac{\partial c_j^*}{\partial \alpha} + E_j c_j^* \frac{\partial c_j}{\partial \alpha} \\
&= \sum E_j \partial_\alpha |c_j|^2 \\
&= \partial_\alpha \sum E_j |c_j|^2
\end{aligned}$$

É óbvio, então, que

$$E_m \partial_\alpha \sum |c_j|^2 \leq \partial_\alpha \sum E_j |c_j|^2 \leq E_M \partial_\alpha \sum |c_j|^2 \quad (18)$$

Portanto, concluímos que

$$E_m \leq \langle H \rangle_{\varphi(\alpha_0)} \leq E_M \quad (19)$$

**Exercise 4** (4.4.2 - 3). *If  $H$  acts in a two-dimensional space, its most general form is*

$$H = \begin{pmatrix} a+c & b \\ b & a-c \end{pmatrix},$$

*where  $b$  can always be chosen to be real. Parametrizing  $|\varphi(\alpha)\rangle$  as*

$$|\varphi(\alpha)\rangle = \begin{pmatrix} \cos \alpha/2 \\ \sin \alpha/2 \end{pmatrix}$$

*find the values of  $\alpha_0$  by seeking the extrema of  $\langle \varphi(\alpha) | H | \varphi(\alpha) \rangle$ . Rederive (2.35).*

**Answer.** Começamos considerando a equação do valor médio explicitamente:

$$\begin{aligned}
\langle \psi | H | \psi \rangle &= [\cos(\alpha/2) \quad \sin(\alpha/2)] \begin{bmatrix} a+c & b \\ b & a-c \end{bmatrix} \begin{bmatrix} \cos(\alpha/2) \\ \sin(\alpha/2) \end{bmatrix} \\
&= [\cos(\alpha/2) \quad \sin(\alpha/2)] \begin{bmatrix} (a+c) \cos(\alpha/2) + b \sin(\alpha/2) \\ b \cos(\alpha/2) + (a-c) \sin(\alpha/2) \end{bmatrix} \\
&= (a+c) \cos^2(\alpha/2) + b \sin(\alpha/2) \cos(\alpha/2) + b \cos(\alpha/2) \sin(\alpha/2) + (a-c) \sin^2(\alpha/2) \\
&= a(\sin^2(\alpha/2) + \cos^2(\alpha/2)) + c(\cos^2(\alpha/2) - \sin^2(\alpha/2)) + b \sin(\alpha) \\
&= a + b \sin(\alpha) + c \cos(\alpha)
\end{aligned}$$

Vejamos os extremos dessa função:

$$\begin{aligned}
\partial_\alpha \langle \psi | H | \psi \rangle &= b \cos(\alpha) - c \sin(\alpha) = 0 \\
b \cos(\alpha) &= c \sin(\alpha) \\
\tan(\alpha) &= b/c \\
\alpha_0 &= \text{atan}(b/c)
\end{aligned}$$

A Eq.(2.35) se refere aos autovetores e autovalores de  $H$ . Vejamos o caso do nosso  $\psi$ :

$$\begin{aligned} H|\psi\rangle &= \left( a \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \sqrt{b^2 + c^2} \begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{bmatrix} \right) \begin{bmatrix} \cos(\alpha/2) \\ \sin(\alpha/2) \end{bmatrix} \\ &= a \begin{bmatrix} \cos(\alpha/2) \\ \sin(\alpha/2) \end{bmatrix} + \sqrt{b^2 + c^2} \begin{bmatrix} \cos \alpha \cos(\alpha/2) + \sin \alpha \sin(\alpha/2) \\ \sin \alpha \cos(\alpha/2) - \cos \alpha \sin(\alpha/2) \end{bmatrix} \\ &= a + \sqrt{b^2 + c^2} \begin{bmatrix} \cos(\alpha/2) \\ \sin(\alpha/2) \end{bmatrix} \end{aligned}$$

onde usamos a Eq.(2.34) e algumas identidades trigonométricas.

**Exercise 5** (4.4.3). *Let a Hamiltonian  $H$  depend on a parameter  $\lambda : H = H(\lambda)$ . Let  $E(\lambda)$  be a nondegenerate eigenvalue and  $|\varphi(\lambda)\rangle$  be the corresponding normalized eigenvector ( $\|\varphi(\lambda)\|^2 = 1$ ) :*

$$H(\lambda)|\varphi(\lambda)\rangle = E(\lambda)|\varphi(\lambda)\rangle$$

*Demonstrate the Feynman-Hellmann theorem:*

$$\frac{\partial E}{\partial \lambda} = \left\langle \varphi(\lambda) \left| \frac{\partial H}{\partial \lambda} \right| \varphi(\lambda) \right\rangle.$$

**Answer.** Sabemos que podemos escrever

$$E(\lambda) = \langle \psi | H | \psi \rangle \quad (20)$$

Então considere:

$$\partial_\lambda E = \langle \partial_\lambda \psi | H | \psi \rangle + \langle \psi | \partial_\lambda | \psi \rangle + \langle \psi | H | \partial_\lambda \psi \rangle \quad (21)$$

Logo, é suficiente mostrar que

$$\langle \partial_\lambda \psi | H | \psi \rangle + \langle \psi | H | \partial_\lambda \psi \rangle = 0 \quad (22)$$

Considere:

$$\begin{aligned} \langle \partial_\lambda \psi | H | \psi \rangle + \langle \psi | H | \partial_\lambda \psi \rangle &= \langle \partial_\lambda \psi | E | \psi \rangle + \langle \psi | E^\dagger | \partial_\lambda \psi \rangle \\ &= E (\langle \partial_\lambda \psi | \psi \rangle + \langle \psi | \partial_\lambda \psi \rangle) \\ &= E (\partial_\lambda \langle \psi | \psi \rangle) \\ &= E \partial_\lambda 1 \\ &= 0 \end{aligned}$$

onde usamos o fato de  $H$  ser hermitiano e ter autovalores reais e de  $|\psi\rangle$  ser normalizado. Portanto, fica demonstrado o teorema.

**Exercise 6** (4.4.4). *We consider a two-level system with Hamiltonian  $H$  represented by the matrix*

$$H = \hbar \begin{pmatrix} A & B \\ B & -A \end{pmatrix}$$

in the basis

$$|+\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |-\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

According to (2.35), the eigenvalues and eigenvectors of  $H$  are

$$\begin{aligned} E_+ &= \hbar\sqrt{A^2 + B^2}, & |\chi_+\rangle &= \cos \frac{\theta}{2} |+\rangle + \sin \frac{\theta}{2} |-\rangle \\ E_- &= -\hbar\sqrt{A^2 + B^2}, & |\chi_-\rangle &= -\sin \frac{\theta}{2} |+\rangle + \cos \frac{\theta}{2} |-\rangle \end{aligned}$$

with

$$A = \sqrt{A^2 + B^2} \cos \theta, \quad B = \sqrt{A^2 + B^2} \sin \theta, \quad \tan \theta = \frac{B}{A}$$

**Part 1.** The state vector  $|\varphi(t)\rangle$  at time  $t$  can be decomposed on the  $\{|+\rangle, |-\rangle\}$  basis:

$$|\varphi(t)\rangle = c_+(t)|+\rangle + c_-(t)|-\rangle$$

Write down the system of coupled differential equations which the components  $c_+(t)$  and  $c_-(t)$  satisfy.

**Answer.** Começamos escrevendo:

$$|\varphi(t)\rangle = \begin{pmatrix} c_+(t) \\ c_-(t) \end{pmatrix} \quad (23)$$

Agora montamos a equação de Schrödinger:

$$\begin{aligned} H |\varphi(t)\rangle &= -i\hbar \partial_t |\varphi(t)\rangle \\ \hbar \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \begin{pmatrix} c_+(t) \\ c_-(t) \end{pmatrix} &= -i\hbar \begin{pmatrix} \partial_t c_+(t) \\ \partial_t c_-(t) \end{pmatrix} \\ \begin{pmatrix} Ac_+ + Bc_- \\ Bc_+ - Ac_- \end{pmatrix} &= -i \begin{pmatrix} \dot{c}_+(t) \\ \dot{c}_-(t) \end{pmatrix} \end{aligned}$$

Ou seja, temos as equações diferenciais:

$$i\dot{c}_+ = Ac_+ + Bc_- \quad (24)$$

$$i\dot{c}_- = Bc_+ - Ac_- \quad (25)$$

**Part 2.** Let  $|\varphi(t=0)\rangle$  be decomposed on the  $\{|\chi_+\rangle, |\chi_-\rangle\}$  basis:

$$|\varphi(t=0)\rangle = |\varphi(0)\rangle = \lambda |\chi_+\rangle + \mu |\chi_-\rangle, \quad |\lambda|^2 + |\mu|^2 = 1$$

Show that  $c_+(t) = \langle + | \varphi(t) \rangle$  is written as

$$c_+(t) = \lambda e^{-i\Omega t/2} \cos \frac{\theta}{2} - \mu e^{i\Omega t/2} \sin \frac{\theta}{2}$$

with  $\Omega = 2\sqrt{A^2 + B^2}$ . Here  $\hbar\Omega$  is the energy difference of the two levels. Show that  $c_+(t)$  (as well as  $c_-(t)$ ) satisfies the differential equation

$$\ddot{c}_+(t) + \left(\frac{\Omega}{2}\right)^2 c_+(t) = 0.$$

**Answer.** Vamos começar pela segunda parte, mostrando a validade da equação diferencial. Considere as EDOs que obtemos e vamos deriva-las mais uma vez (faremos as contas para  $c_+$ , pois são análogas para  $c_-$ ):

$$i\ddot{c}_+ = A\dot{c}_+ + B\dot{c}_- \quad (26)$$

Mas note, também, que:

$$\begin{aligned} A\dot{c}_+ &= -iA^2c_+ - iABc_- \\ B\dot{c}_- &= -iB^2c_+ + iABc_- \end{aligned}$$

Logo,

$$A\dot{c}_+ + B\dot{c}_- = -i(A^2 + B^2)c_+ \quad (27)$$

Portanto,

$$i\ddot{c}_+ = -i(A^2 + B^2)c_+ \quad (28)$$

que, simplificando, implica na expressão desejada

$$\ddot{c}_+ + \left(\frac{\Omega}{2}\right)^2 c_+ = 0 \quad (29)$$

Vamos considerar agora a outra parte. Ela nos dá condições iniciais:

$$|\varphi_0\rangle = \begin{pmatrix} \lambda \cos(\theta/2) - \mu \sin(\theta/2) \\ \lambda \sin(\theta/2) + \mu \cos(\theta/2) \end{pmatrix} \quad (30)$$

Ou seja,

$$\begin{aligned} c_+(0) &= \lambda \cos(\theta/2) - \mu \sin(\theta/2) \\ c_-(0) &= \lambda \sin(\theta/2) + \mu \cos(\theta/2) \end{aligned}$$

Usando a equação diferencial que deduzimos, é fácil ver que:

$$\begin{aligned} c_+(t) &= A_+ e^{i\Omega t/2} + B_+ e^{-i\Omega t/2} \\ c_-(t) &= A_- e^{i\Omega t/2} + B_- e^{-i\Omega t/2} \end{aligned}$$

Isso obviamente nos diz que:

$$\begin{aligned} A_+ + B_+ &= \lambda \cos(\theta/2) - \mu \sin(\theta/2) \\ A_- + B_- &= \lambda \sin(\theta/2) + \mu \cos(\theta/2) \end{aligned}$$

\*

**Part 3.** We assume that  $c_+(0) = 0$ . Find  $\lambda$  and  $\mu$  up to a phase as well as  $c_+(t)$ . Show that the probability of finding the system in the state  $|+\rangle$  at time  $t$  is

$$p_+(t) = \sin^2 \theta \sin^2 \left( \frac{\Omega t}{2} \right) = \frac{B^2}{A^2 + B^2} \sin^2 \left( \frac{\Omega t}{2} \right).$$

**Answer.** Se  $c_+(0) = 0$ , então precisa ser que

$$\lambda \cos(\theta/2) - \mu \sin(\theta/2) = 0 \quad (31)$$

Logo, é verdade que

$$\lambda = \mu \tan(\theta/2) \quad (32)$$

Mas, pela normalização:

$$\begin{aligned} |\lambda|^2 + |\mu|^2 &= 1 \\ |\mu \tan(\theta/2)|^2 + |\mu|^2 &= 1 \\ |\mu|^2 |\tan(\theta/2)|^2 + |\mu|^2 &= 1 \\ |\mu|^2 &= \frac{1}{1 + |\tan(\theta/2)|^2} \end{aligned}$$

Logo, a menos de uma fase  $e^{ia}$ ,

$$\begin{aligned} \mu &= \frac{1}{\sqrt{1 + |\tan(\theta/2)|^2}} \\ \lambda &= \frac{\tan(\theta/2)}{\sqrt{1 + |\tan(\theta/2)|^2}} \end{aligned}$$

Vamos supor, daqui em diante,  $\theta \in [0, \pi]$  para tirarmos a tangente do módulo. Usando a solução geral de  $c_+(t)$ , sabemos que precisa valer

$$A_+ + B_+ = 0 \quad (33)$$