gamdid: generalized additive models for differential distributions in single-cell proteomics experiments

Lucas Beerland[1], Stijn Vandenbulcke[1], Christophe Vanderaa[1,2,3], Lennart Martens[2,3,4,5], and Lieven Clement*[1]

[1]Department of Mathematics, Ghent University
[2]Department of Biomolecular Medicine, Ghent University
[3]VIB-UGent Center for Medical Biotechnology
[4]BioOrganic Mass Spectrometry Laboratory (LSMBO), IPHC UMR 7178, University of Strasbourg, CNRS, Strasbourg, France
[5]Infrastructure Nationale de Protéomique ProFI-UAR 2048, Strasbourg, France

October 30, 2026

Abstract

Single-cell proteomics (SCP) generates protein abundance measurements across hundreds to thousands of individual cells, offering unprecedented resolution to study cellular heterogeneity. However, existing differential abundance (DA) methods are limited to detecting shifts in mean expression, leaving biologically relevant differences in shape undetected. Indeed, the specific power of SCP is to identify differences between individual cells in a population, which are typically only found as shape differences rather than in mean expression. We therefore present `gamdid` *(generalized additive models for differential distributions), a novel statistical framework and R package for differential distribution (DD) analysis in SCP data.* `gamdid` *is based on generalized additive models (GAMs) to flexibly model heterogeneous distributions, perform inference and provide interpretable visualizations. Through semi-synthetic benchmarking on two SCP datasets,* `gamdid` *demonstrates conservative false discovery rate (FDR) control and substantially outperforms competing methods for differences in shape, while achieving comparable performance for mean shifts. A spike-in case study further demonstrates the utility of* `gamdid` *and its interpretable visualization. Uniquely among DD methods,* `gamdid` *supports omnibus testing across more than two groups, with post-hoc pairwise comparisons via stagewise testing, and is specifically tailored for proteomics abundance data.*

Introduction

High-throughput single-cell technologies have improved the resolution at which biological processes can be studied. A key component of such studies is differential abundance (DA) analysis, which aims to identify features that differ between experimental conditions. However, the majority of established DA single-cell methods primarily detect changes in mean abundance and thus overlook other, biologically very informative, differences in distributions, such as changes in variance, skewness, modality, etc., limiting the advantages of single-cell over bulk analyses. This limiting focus stems from their origin in bulk omics workflows, where analyses beyond the mean are typically not feasible due to the limited number of available samples. Single-cell datasets, however, provide expression measurements for hundreds to thousands of individual cells, making it possible, and likely necessary, to assess changes in the full distribution of expression values.
Indeed, the importance of distribution changes is nicely illustrated in the SCP data from Leduc et al. [1], which compares melanoma cells with monocytes. Figure 1 visualizes the density of three illustrative preprocessed proteins.

*Corresponding author: lieven.clement@ugent.be




Figure 1: ***Motivating examples of distributional heterogeneity in single-cell proteomics*** *Density distributions of three proteins from the Leduc single-cell proteomics dataset, comparing melanoma cells (red) and monocytes (blue). The x-axis shows the normalized log2 protein intensity (log2-transformed, median-centered intensities; see Data in Methods), and the protein accession is given above each panel. The left panel (Q8TCT9) illustrates a conventional mean shift between cell types, which is the type of difference that standard DA methods are designed to detect. The middle panel (P47756) shows a difference in variance (Note that there is also a clear difference in kurtosis). The right panel (Q8TDN6) reveals the emergence of a distinct subpopulation in one cell type, visible as a secondary mode. The latter examples reflect differences in shape rather than mean. These are biologically informative yet systematically missed by conventional differential abundance (DA) approaches. They motivate the need for differential distribution (DD) methods capable of detecting the full spectrum of distributional differences in single-cell data.*

Conventional DA analysis methods are designed to pick up mean shifts, as seen in the left panel in Figure 1, but lack power to prioritize proteins for which the distribution between cell populations mainly differs in shape, as seen in the middle and right panels in Figure 1. These changes in shape may reflect underlying biological processes that remain undetected with conventional DA methods. For example, the increased spread of the abundance values observed in the middle panel may indicate a heterogeneous cellular response, whereas the pattern in the right panel may reflect the emergence of a novel cell state in a subset of cells. To infer such nuanced but relevant effects, more flexible approaches are required, which are referred to as differential distribution (DD) methods.

Several approaches to perform such DD analysis on single cell data already exist, most notably `BASiCS` [2], `scDD` [3] and `distinct` [4]. However, these methods were developed specifically for single-cell RNA-seq data and have been neither tested on nor tailored for proteomics applications. Moreover, they cannot handle large proportions of missing values, a key problem in SCP datasets, and lack interpretability on how the distributions differ. The latter greatly limits the utility of the assessment, as the nature of the difference is an important element in hypothesizing about the possible causes.
Additionally, each approach has methodological limitations: `BASiCS` is restricted to testing for differences in mean and variance, `scDD` can only handle count data, and `distinct` can only be used for two-group comparisons.

Therefore we here present `gamdid`: a novel statistical framework for DD analysis. `gamdid` is capable of detecting any form of distributional difference between experimental conditions and provides an interpretable visualization of these effects. Importantly, `gamdid` also allows users to localize and visualize the specific regions that are differentially distributed, providing a direct link between statistical inference and visualization.
By building on the flexible framework of generalized additive models (GAMs), `gamdid` thus extends well beyond traditional mean-based comparisons while remaining intuitive and highly interpretable.

Results

In this section, we first describe the statistical model and inference procedure underlying `gamdid`. We then evaluate `gamdid` through a simulation study and a case study, both based on real SCP data. The simulation study benchmarks `gamdid` across five distributional scenarios. Subsequently, we demonstrate the practical utility of `gamdid` through a controlled spike-in case study.

Statistical model and inference

A key advantage of SCP analysis over bulk proteomics is its capacity to assess single-cell heterogeneity. This can result in differences in abundance distributions that are more subtle than mean abundance differences, and are thus overlooked by traditional, bulk-oriented differential analysis methods for proteomics data. Instead of modeling the abundance data with a normal distribution, we here adopt a flexible non-parametric approach to estimate the distribution of the protein abundances in single cells in each group by building upon Lindsey's method [5]. Specifically, for each feature the density estimation problem is recast into Poisson regression by discretizing the sample space into bins. In other words, we convert the intensities into histogram bin data, see Figure 2 panel A, and the bin counts are then considered independent Poisson-distributed observations.




Figure 2: ***Overview of the gamdid workflow*** *The complete workflow for fitting GAMs to compare distributions is illustrated using a ribosome biogenesis protein from the Leduc dataset.* ***A****: Raw protein abundance values are discretized into histogram bin counts, recasting the density estimation problem as Poisson regression. This representation preserves distributional shape information that is lost in a mean-based summary.* ***B****: A Poisson generalized additive model (GAM) is fitted to the bin counts as a function of bin midpoints, allowing each group (melanoma and monocyte) to have its own flexible, non-parametric density curve. The GAM framework accommodates any distributional shape without parametric assumptions.* ***C*** *(top): The fitted density curves generated by gamdid's visFit function, with gray shading highlighting regions where the distributions differ significantly between groups.* ***C*** *(bottom): The difference smoother representing the log-ratio of densities between groups along with its simultaneous confidence band. Regions where this band excludes zero identify statistically significant local distributional differences, linking inference to an interpretable visualization. This combination of flexible modeling, hypothesis testing, and interpretable visualization is an innovative feature unique to gamdid.*

This representation allows for density estimation through Poisson generalized additive models, where expected counts are modeled as a function of the bin midpoints and by specifying a group-specific spline,

each group has its own unique density shape. The GAM fits are displayed in Figure 2 panel B.

Inference is subsequently carried out using the GAM framework, with differences between groups assessed using Wald-type tests by comparing group-specific model parameters. This yields a p-value for inferring any type of distributional difference both in location and in shape, referred to as the GAM p-value in the remainder of the manuscript.

To increase the power to detect distributional changes that are strongly driven by differences in the mean, the p-value obtained from the GAM is combined with a p-value derived from the `msqrob2` framework, a conventional mean-based approach for DA analysis [6]. The two p-values are integrated using the harmonic mean p-value (HMP) method [7]. This results in a final aggregated p-value for each feature, providing a single, robust measure of evidence for any distributional change.

Interestingly, `gamdid` also provides an interpretable visualization that is directly linked to the inference, by constructing a difference smoother along with a simultaneous confidence band. Regions where this band does not include zero indicate a statistically significant local distributional difference, as illustrated in Figure 2 panel C.

Simulation study for benchmarking

To benchmark `gamdid` for single cell proteomics applications, we constructed semi-synthetic simulations based on two publicly available mass spectrometry (MS)-based SCP data sets obtained from the R/Bioconductor package `scpdata` [8]: the TMT-labeled single-cell Leduc dataset [1] consisting of 1531 cells and 6706 identified peptides, and the label-free single-cell Petrosius dataset [9] consisting of 564 cells and 5729 identified peptides. For each dataset, preprocessed (see Methods) protein-level summarized assays from one cell type were selected and the corresponding cells were randomly assigned to two mock groups (A and B). Hence, all proteins are non-DD between A and B.

Only proteins with at least 50 non-missing observations per group were retained for benchmarking, consistent with the default setting in the `gamdid` package. This threshold ensures a sufficient number of observations to reliably characterize the distribution of each feature, as fewer observations may yield an incomplete or misleading representation of the underlying density.

Simulation of distributional differences Following the framework established by Korthauer et al. (2016) [3], we introduced five categories of distributional differences by altering protein abundances in cells of group B: differential expression (DE) (a pure mean shift, which corresponds to DA), differential variability (DV), differential both in modality and component means (DB), differential modality (DM), and differential proportion (DP), as illustrated in Figure 3. These effects were introduced in 20% of randomly selected features by systematic shifting of the abundances in subsets of the cells in group B (for the DE, DB, DM and DP scenarios) or by adding controlled random noise (for the DV scenario).




Figure 3: ***Five simulation scenarios used to benchmark gamdid*** *To evaluate gamdid's ability to detect different types of distributional differences, five categories of signal, established by Korthauer et al. (2016) [3], were introduced into real single-cell proteomics data. DE: a pure shift in the mean, equivalent to conventional DA. DV: an increase in variance without a location shift, reflecting e.g. heterogeneous cellular responses. DB: a bimodal distribution induced in one group by shifting subsets of cells in opposite directions, without altering the overall mean. DM: an asymmetric shift in one tail, creating a difference in skewness or the emergence of a subpopulation. DP: a change in the relative weight of a subpopulation shared between groups.*

This semi-synthetic approach leverages real SCP data rather than fully synthetic 'clean' simulations, and thus preserves realistic distributional characteristics inherent to single-cell proteomics measurements.

Method comparison We compare `gamdid` to `distinct` as the primary benchmark method. `BASiCS` and `scDD` were excluded from the comparison as `BASiCS` is limited to detecting shifts in mean and variance, and therefore cannot be applied to the majority of the simulations, while `scDD` requires count data as input, which is incompatible with the continuous abundance measurements obtained in proteomics data.

`distinct` performs group comparisons by computing the average difference between empirical cumulative distribution functions (ECDFs) and can optionally account for subgroup structure in multi-sample multi-cell designs. As our datasets, similar to most public SCP datasets, do not have these multi-sample multi-cell designs, `distinct` was applied without this option.

Note that the current implementation of `gamdid` does not account for subgroup structure.

Method performance is evaluated using false discovery proportion — true positive proportion (FDP-TPP) curves. For a given significance threshold, the FDP represents the proportion of false positives among all features called significant, while the TPP represents the proportion of true positives recovered out of all truly differential features. An FDP-TPP curve allows one to assess the power of a method (TPP) at any given level of false discoveries (FDP). This provides a more complete picture of method performance than a single summary statistic, and allows for a direct visual comparison of the trade-off between sensitivity and false discovery control across competing methods. A method can be considered more powerful when the area under its FDP-TPP curve is larger. The dashed vertical line at FDP = 0.05 serves as a reference for the nominal FDR threshold.

GAM evaluation We first evaluate the performance of the GAM methodology in isolation (without p-value aggregation via `msqrob2`) against `distinct`. The FDP-TPP curves for the Petrosius dataset are presented in Figure 4; those for the Leduc dataset are provided in Supplementary Figure S1.

The GAM method demonstrates conservative false discovery rate control and exhibits substantially superior performance compared to `distinct` when distributional changes do not involve shifts in mean (DV and DB scenarios). Conversely, in the scenarios where mean shifts contribute to the distributional change, `distinct` outperforms the GAM method and is even on par with conventional DA analyses like `msqrob2`. This observation motivated the aggregation of GAM-derived p-values with `msqrob2` p-values, thereby combining the GAM methodology's sensitivity to differences in shape with `msqrob2`'s established performance for detecting mean shifts.

gamdid evaluation Figure 4 also presents the FDP-TPP curves of `gamdid` for the Petrosius dataset (see Supplementary Figure S1 for the Leduc dataset). The corresponding numbers of true and false positives at the 5% FDR threshold are also provided in the supplementary materials (Figures S2 and S3): `gamdid` recovers a comparable or larger number of true positives, while the false positives are similarly low for both methods. `gamdid`'s performance is comparable to `distinct` for pure mean shifts (DE), shows modest improvements for distributional changes that involve location shifts (DM, DP), and shows substantial improvements for distributional changes in shape (DV, DB).

Improvements are more pronounced in the Petrosius simulations than in the Leduc simulations (Supplementary Figure S1), which could be attributable to the smaller sample size in the Petrosius dataset compared to Leduc. To examine this, we subsampled the Leduc dataset such that the average number of non-missing observations per group over all features matched that of the Petrosius dataset. The resulting FDP-TPP curves are provided in the supplementary materials in Figure S4. Indeed, the improvements are more pronounced in the subsampled simulations compared to the Leduc simulations in Supplementary Figure S1. This supports the hypothesis that `gamdid` exhibits higher power under smaller sample conditions compared to `distinct`, which suffers from a reduced resolution of the null distribution due to a limited number of permutations.




Figure 4: ***Performance of gamdid and the standalone GAM method*** *FDP-TPP curves comparing gamdid (combining GAM and msqrob2 p-values via the harmonic mean p-value), the standalone GAM method, msqrob2 and distinct across the five simulation scenarios, based on the protein-level mock data of the Petrosius dataset (269 m2i cells). The corresponding results for the Leduc dataset are shown in Supplementary Figure S1. Each curve traces the trade-off between TPP (sensitivity) and FDP as the significance threshold is varied; the dots mark the 5% FDR threshold and the dashed vertical line marks FDP = 0.05. The standalone GAM method substantially outperforms distinct for distributional changes that do not involve a mean shift (DV, DB), demonstrating its sensitivity to shape differences, whereas distinct and msqrob2 outperform it when mean shifts contribute to the distributional change (DE, DM, DP). This complementarity motivates the aggregation of GAM p-values with msqrob2 p-values: by combining both, gamdid achieves performance that is comparable to distinct for pure mean shifts (DE), shows modest improvements for distributional changes that involve a location component (DM, DP), and demonstrates substantial improvements for pure shape changes (DV, DB), while maintaining conservative FDR control throughout. DB (differential both) denotes a difference in both modality and component means.*

To assess the robustness of the conclusions under less pronounced distributional differences, FDP-TPP curves for simulations with 25% reduced signal magnitudes are provided in the supplementary materials (Figure S5). These results are consistent with the findings above, indicating robust performance.

Interpretation and visualization In contrast to competing methods, `gamdid` provides informative visualizations of features picked up as DD. Figure 5 illustrates this functionality for a significant feature identified under the DV simulation framework. This visualization integrates GAM fits with highlighted differential regions, providing immediate visual evidence of differential variability between groups.




Figure 5: ***gamdid's interpretable visualization for a DV-simulated feature in the Petrosius dataset*** *A feature with induced differential variability (DV simulation), visualized using gamdid's visFit function. The top panel shows the fitted GAM density curves for both groups, overlaid on the observed histograms. The bottom panel displays the difference smoother (log-ratio of densities) along with its simultaneous confidence band; the gray-shaded region identifies the specific abundance range where the distributional difference is statistically significant. Unlike competing DD methods such as distinct, gamdid does not merely return a p-value; this visualization provides immediate visual evidence of differential variability between groups.*

Peptide-level simulations The same simulation framework was applied to peptide-level assays for both the Leduc and Petrosius datasets. The resulting FDP-TPP curves are provided in the supplementary materials in Figure S6 and are consistent with the protein-level results.
Note, however, that the FDR control at the nominal 5% threshold is less conservative at the peptide level due to the correlation structure inherent to peptide-level data, a well-known challenge in peptide-level analyses. Nevertheless, this demonstrates that `gamdid` provides a flexible framework suitable for inferring DD at different quantification levels (precursor, peptide, protein) in MS-based proteomics data.

Three-group simulations A unique advantage of `gamdid` is its support for omnibus testing across more than two groups, followed by post-hoc pairwise comparisons corrected for multiple testing using the stagewise testing procedure [10]. The omnibus test simultaneously assesses whether any distributional difference exists between any of the groups. No competing DD analysis method currently offers this functionality.
To evaluate this functionality, we extended the simulation framework described above to a three-group setting, randomly assigning cells to mock groups A, B, and C, with distributional signal introduced in group B under the same five simulation scenarios as above. The omnibus test was applied to this three-group comparison and the resulting FDP-TPP curves are provided in the supplementary materials in Figure S7. Note that, as none of the competing DD methods supports multi-group testing, a formal benchmark comparison is not possible here.
Interestingly, the three-group setting yields higher power than the two-group comparison under equal signal strength. Note that this might arise from the default minimum observation requirement of 50 cells per group imposed by `gamdid`, which, in the three-group analysis, requires more observations than in the two-group analysis, leading to more powerful inference.

Case study

We next illustrate the utility of `gamdid` in a case study based on the Leduc dataset.
First, we randomly divide the monocyte cells into two mock groups: X (containing 377 cells) and Y (containing 378 cells). In group Y, we spike in 10% of the 697 sub-cluster A melanoma cells (69 cells) to simulate a subpopulation in group Y.
Figure 6 shows the volcano plot of the conventional DA `msqrob2` analysis twice. In panel A, significant features of the DA analysis are marked in red, while in panel B the significant features of DD methods `gamdid` and `distinct` are marked.




Figure 6: ***Case study: gamdid detects distributional changes missed by conventional DA analysis*** *A controlled spike-in experiment based on the Leduc dataset: monocyte cells are split into two mock groups (X: 377 cells; Y: 378 cells), and 10% of the 697 sub-cluster A melanoma cells (69 cells) are spiked into group Y to simulate the emergence of a subpopulation. Panels A and B display the same msqrob2 volcano plot.* ***A****: Features found significant by conventional DA (msqrob2) are highlighted in red.* ***B****: The same volcano plot, now colored by DD method detection: yellow = detected by both distinct and gamdid; orange = detected by distinct only; blue = detected by gamdid only. Both DD methods detect the majority of DA features, while also prioritizing some features missed by conventional DA methods.* ***C****: UpSet plot showing the overlap in significant features between distinct, gamdid and msqrob2. The horizontal bars (left) show the total number of significant features per method (Set size); the vertical bars (top) show the size of each intersection, with the dot matrix below indicating which methods contribute to each intersection. The six features picked up by msqrob2 and missed by gamdid are borderline significant for msqrob2 and borderline non-significant for gamdid (this is not the case for distinct, see Table S1), which means that only gamdid largely preserves the ranking of DA features. Moreover, gamdid prioritizes three features that are missed by both msqrob2 and distinct (see Table S2).*

Figure 6C presents a method comparison for the significant features in the spike-in case study. Overall, the

methods produce similar results with 52 features found significant by all three.

Only six features are picked up by `msqrob2` and missed by `gamdid`. These features are borderline significant for `msqrob2` and borderline non-significant for `gamdid`, which is illustrated in table S1 in supplementary materials.

Furthermore, `gamdid` prioritizes three features that `msqrob2` and `distinct` fail to detect (see Table S2 in supplementary materials). Interestingly, even features detected by all three methods illustrate the utility of `gamdid`. Figure 7 shows the `gamdid` visualization for four representative features (panels A–D), displaying the mock groups (top) and the spike-in study (middle). The boxplot at the bottom of each panel indicates where the spiked-in melanoma cells are located within the distribution of group Y. Figure 7 clearly shows that these features are DD due to the subpopulation of spike-in cells rather than DA. Additionally, it powerfully validates gamdid's localization approach: not only does gamdid detect the distributional change, it correctly identifies its location, providing interpretable visualization that is unavailable in any competing DD method.

The `gamdid` visualizations of two of the four features only detected by `msqrob2` and of the two features only detected by `distinct` are provided in supplementary materials in Figure S8.




Figure 7: ***gamdid visualizations for four representative features detected by all three methods in the spike-in case study***
*Each panel (**A–D**) shows one feature, following the same layout as the panels of Figure S8: the mock comparison (top), the comparison after spike-in (middle), and a boxplot indicating where the spike-in melanoma cells are located (bottom). All four features are detected as significant by msqrob2, gamdid, and distinct, forming part of the 52-feature consensus set (see Figure 6C). In all four cases, gamdid's visualization reveals that the distributional difference extends beyond a simple mean shift: these features are DD due to the subpopulation of spike-in cells rather than DA. Moreover, the spiked-in melanoma cells align closely with the detected differential region (gray shading) in each example. This demonstrates that even within the set of features identified by conventional DA methods, gamdid provides additional interpretive value by characterizing how and where the distributions differ. This information is not accessible from the fold change and p-value from a conventional DA method. These examples underscore the broader argument that accounting for full distributional differences, rather than mean shifts alone, leads to richer biological insights.*

Discussion

Heterogeneity in single-cell data Because of `gamdid`'s flexible approach, it captures the heterogeneity of single-cell data that is, by design, missed by conventional DA methods. In the case study, `gamdid` identifies three features that are not detected by conventional DA methods as the distributional differences do not involve a sufficiently large mean shift.
More importantly, even the 52 features detected by both DA and DD methods further emphasize the need to account for heterogeneity in single-cell data. Indeed, for the majority of these features, the distributional difference extends beyond a pure mean shift. This is illustrated by `gamdid`'s visualization which reveals the important nuances of these more complex distributional patterns.
Four representative examples are provided in Figure 7. Notably, in all four examples the detected differential regions align closely with the regions of the distribution where melanoma cells were spiked in, underscoring the utility of `gamdid`'s visualization.

Ranking of significant features Notably, the ranking of significant features is of increasing interest in proteomics, and `gamdid` offers a key advantage in this regard. For DA features, the `gamdid` ranking is very similar to that of top-performing DA methods (which is not the case for `distinct`, see Table S1). Moreover, `gamdid` can also prioritize interesting features that are completely overlooked by DA methods. This is illustrated in Table S2, which shows that two out of the three features exclusively detected by `gamdid` are assigned high p-values by `msqrob2`.

Conceptual difference between `gamdid` and `distinct` The simulation results reflect a fundamental difference in how `distinct` and the GAM methodology characterize distributional differences between groups. `distinct` operates on empirical cumulative distribution functions (ECDFs) and quantifies the mean difference between the ECDFs of two groups. By design, this mean ECDF difference is sensitive to sustained, global displacements of probability mass. The GAM-based approach, however, models the density function of each group. Because density estimation is inherently local, the GAM is particularly well-suited to detect distributional changes that are concentrated in specific regions of the distribution.
Figure 8A–B illustrates this using a feature from the Petrosius DB simulations that exhibits a local change without substantially displacing the bulk of the probability mass. In this case, the mean difference between the ECDFs of the two groups remains modest, resulting in reduced power for `distinct`, whereas the GAM-based method is able to successfully identify the differential region.

Conversely, distributional changes like mean shifts produce large, sustained differences between the ECDFs of two groups and are therefore readily detected by `distinct`. The GAM-based method, however, exhibits reduced power in such settings, as mean shifts constitute diffuse, global changes that do not concentrate in any particular local region of the distribution. This behavior is illustrated in Figure 8C–D, which presents a feature from the Petrosius DE simulations. This example shows an accumulated difference between the ECDFs, while the GAM method is not able to detect differential regions.
This motivates the aggregation of GAM-derived p-values with those from `msqrob2`, which is specifically designed to detect mean shifts with high sensitivity.




Figure 8: ***Conceptual difference between distinct and gamdid*** *Comparison of how distinct (ECDF curves, top) and gamdid (density visualization, bottom) characterize two features from the Petrosius simulations.* ***A–B****: A feature from the DB scenario, detected by gamdid (FDR adjusted p-value = 0.012) and not by distinct (FDR adjusted p-value = 0.278). The mean ECDF difference (used by distinct) between groups is modest (A) because the induced bimodality does not produce a sustained, global displacement of probability mass, while the gamdid visualization (B) reveals the local bimodality clearly and the difference smoother identifies the specific regions where the densities diverge (gray shading).* ***C–D****: A feature with a pure mean shift (DE), detected by distinct (FDR adjusted p-value = 0.020) and not by the standalone GAM method before aggregation (FDR adjusted p-value = 0.227). The ECDF curves (C) show a large, sustained horizontal displacement between groups, whereas the GAM fits (D) overlap and show no identifiable differential region, reflecting the GAM's reduced power for diffuse, global distributional changes. distinct is thus more sensitive to sustained, global ECDF displacements, while the GAM's density-based, inherently local approach is better suited to detecting concentrated, shape-based differences. This complementarity motivates the aggregation of GAM p-values with msqrob2 p-values: after aggregation, the DE feature (C–D) is detected by gamdid (FDR adjusted p-value = 0.029).*

P-value aggregation The aggregated p-value in `gamdid` combines the local sensitivity of the GAM with the established power of `msqrob2` for mean-based inference, ensuring that `gamdid` is well-suited to detect any form of distributional difference. As expected, a modest loss of power relative to each individual component is observed in scenarios where only one of the two components contributes meaningfully: the DE scenario for the GAM component, and the DV and DB scenarios for `msqrob2`. This is an inherent consequence of p-value aggregation when one of the contributing p-values carries no signal. In scenarios where the distributional change involves both a mean shift and a difference in shape, however, `gamdid` combines the strength of both methods and outperforms each method separately, demonstrating the practical benefit of the aggregation strategy.

FDR control `gamdid` achieves conservative FDR control at the nominal 5% threshold throughout all simulations.

Computational efficiency To evaluate computational efficiency, the runtime of both `gamdid` and `distinct` was measured on the spike-in case study dataset, comprising 1316 features and 824 cells. Each method was executed on a single core, and runtimes were averaged across ten independent runs. The mean runtime for `gamdid` was 16 seconds, whereas `distinct` required 256 seconds on average, indicating that `gamdid` achieved a speedup of more than fifteen-fold (see supplementary Figure S9).

This can be attributed to two key factors. First, `gamdid` does not rely on permutation-based inference as `distinct` does. Second, as `distinct` cannot handle missing values, all missing observations must be removed

on a per-feature basis prior to analysis.

Outlook The development of `gamdid` represents a significant advance toward a more comprehensive interpretation of the information in single-cell proteomics data. By extending differential analysis beyond mean-based comparisons, `gamdid` enables the systematic investigation of distributional changes and thereby facilitates a more detailed characterization of cellular heterogeneity. This functionality uncovers biologically relevant variation that has remained largely unexplored by existing data analysis frameworks.
As single-cell proteomics technologies continue to mature and datasets grow in size and complexity, the ability to detect and interpret subtle distributional differences will become increasingly important. We therefore anticipate that `gamdid` can find applications not only in the comparison of cell types and conditions, but also in the identification of subpopulations, or the characterization of transitional cell states.
More broadly, the framework presented here is not intrinsically limited to proteomics and could, in principle, be adapted to other single-cell modalities, including single-cell transcriptomics, metabolomics and flow cytometry, where distributional heterogeneity likewise constitutes an important source of biological information. As such, `gamdid` has the potential to support a broader methodological shift toward distributional approaches in single-cell data analysis, enabling a more comprehensive characterization of cellular variation beyond changes in average abundance levels.

Methods

Below, we first outline the `gamdid` approach (model equations, test statistics and the simultaneous confidence bands are discussed in more detail in the Supplementary materials). We subsequently describe the covariate adjustment procedure, the datasets, the simulation framework and the implementation.

Overview of the gamdid approach

The analysis is performed for each feature (protein or peptide) separately and consists of five steps (see Figure 2).

Step 1: Estimating a smooth distribution per group For a given feature, the abundance values of all cells are binned into a histogram, using the same bins for every group (see Figure 2, panel A). Following Lindsey [5] and Efron [11], the bin counts, i.e. number of cells with intensities in a specific bin, can be treated as Poisson counts.
We then model the bin counts as a function of the bin midpoints using a Poisson GAM, which allows us to flexibly model the shape of the distribution with a smooth curve. Each group receives its own smooth curve (see Figure 2, panel B). Equation 1 schematically represents the smooth curve for group c and bin k with midpoint mid_k.

$$\log(\text{expected count}_{ck}) = \underbrace{f_c(mid_k)}_{\text{density estimation for group } c} + \text{normalization offset}_c \tag{1}$$

where f_c is a flexible smooth function that captures the shape of the distribution of group c. The counts are normalized using an offset, which allows us to directly interpret the curves f_c as log-transformed densities.

Step 2: Testing whether the distributions differ If two groups have the same distribution, their smooth curves are identical, i.e. $H_0 : f_{c_1} = f_{c_2}$. We test this by assessing if the parameters corresponding to a smooth curve β_{f_c} differ between the groups, i.e. $H_0 : \beta_{f_{c_1}} = \beta_{f_{c_2}}$. We do this by employing a Wald-type test that builds on the work of Wood [12]. The resulting p-value, referred to as the GAM p-value, captures any difference in distribution between the groups: mean shifts, differences in variance, modality, skewness...

Step 3: Combining with a mean-shift test Because the GAM test spreads its power over all possible types of distributional differences, it is less powerful than a dedicated DA method for detecting pure mean shifts. We therefore also compute a p-value with `msqrob2`, a robust linear-model approach designed for detecting mean shifts in proteomics data [6]. The GAM and `msqrob2` p-values are combined into a single 'feature-level aggregated p-value' using the harmonic mean p-value (HMP) method [7]. As a result, `gamdid` largely retains the sensitivity of conventional DA analysis for mean shifts, while additionally detecting

changes in distributional shape that DA methods miss.

Step 4: Multiple testing and pairwise comparisons The feature-level aggregated p-values are corrected for multiple testing across features using the Benjamini-Hochberg FDR procedure [13]. When more than two groups are compared, significant features are followed up with pairwise comparisons between all pairs of groups, using the same combination of GAM and `msqrob2` tests. The feature-level and pairwise hypotheses are jointly controlled for multiple testing by the stagewise testing procedure [10, 14].

Step 5: Detecting differential regions Finally, `gamdid` can indicate where the distributions differ. For this, a log-difference smoother is computed. This smoother displays the difference between the estimated log-densities of two groups, together with a simultaneous confidence band. Regions where this band excludes zero indicate a statistically significant local distributional difference; these are marked by gray shading (Figure 2, panel C).

Covariate adjustment

In addition to the covariate of interest, `gamdid` allows users to account for confounding variables through a nuisance formula, enabling applications such as batch correction in multi-batch experimental designs. Both the covariate of interest and the nuisance formula are combined into one model in the `msqrob2` analysis, and an appropriate contrast is constructed to test for distributional differences in the covariate of interest.
To remove the influence of nuisance covariates from the GAM-based inference, partial residuals with respect to the covariate of interest are computed from the `msqrob2` fit and subsequently used as input to the GAM analysis. This two-step approach is required due to the binning strategy: as a single bin may contain observations that correspond to different groups of the nuisance covariate, direct covariate adjustment within the GAM framework is not feasible.
Note, however, that the uncertainty associated with the regression of the nuisance covariates in the first step is not accounted for, so the resulting p-values should be interpreted with caution.

Data

- `leduc2022_pSCoPE`: These data were acquired using pSCoPE technology: a TMT-18plex protocol with prioritized data-dependent acquisition (DDA) carried out by a Thermo Scientific Q-Exactive mass spectrometer [1]. The MS data were identified and quantified using MaxQuant version 1.6.17 [15]. We performed quality control by filtering low-quality features and filtering cells with a median coefficient of variation greater than 0.6, calculated using the *medianCVperCell* function of the `scp` package [8]. The PSM intensities were log2-transformed for variance-stabilization, median centered and summarized to the protein level using median summarization per MS run. Subsequently, samples were normalized through median centering and proteins with more than 95% missingness were filtered out, resulting in a dataset with 776 melanoma cells and 755 monocytes profiled across 2120 proteins.
- `petrosius2023_mES`: These data originate from a label-free SCP experiment conducted on mouse embryonic stem cells cultured under two distinct conditions: a serum-free 2i condition (m2i) containing cytokine LIF and inhibitors targeting the MEK and GSK3 pathways, and a serum condition (m15) only containing cytokine LIF [9]. Data acquisition was performed using an Orbitrap Eclipse Tribrid mass spectrometer operating in data-independent acquisition (DIA) mode. The resulting mass spectrometry data were subsequently analyzed by the authors using Spectronaut v17 [16]. We performed quality control by removing cells with fewer than 500 detected peptides or a median intensity lower than 11. Features with more than 95% missing values were also removed. The detected intensities were log2-transformed for variance-stabilization, and the intensities of each cell were centered with the median log2-intensity of the corresponding cell. Subsequently, peptides were summarized to the protein level through median polish summarization. The resulting dataset contains 295 m15 and 269 m2i cells with 2492 proteins.

Preprocessed protein-level summarized assays from both datasets served as the foundation for our simulations. From the Petrosius dataset, we selected the 269 m2i-condition cells, while from Leduc, we selected the 755 monocytes (in both cases following removal of non-informative cells during preprocessing). Within each selected subset, no systematic biological differences are expected between cells. Consequently, random assignment of cells to two mock groups (A and B) yields a dataset suitable as a mock basis for a controlled two-group comparative analysis.

Simulation framework

The benchmark simulations are based on protein-level mock data from both the Leduc and Petrosius datasets, as described in the Results section. For each dataset, mock groups A and B were formed by random assignment of cells, and distributional signal was introduced in group B for 20% of randomly selected features. Five types of distributional signal were considered, following the framework of Korthauer et al. (2016) [3]:

- **DE:** A shift equal to 40% of the feature's standard deviation is added to all observations in group B.
- **DV:** Independent Gaussian noise with mean zero and standard deviation equal to 120% of the feature's standard deviation is added to all observations in group B, thereby increasing the spread of the distribution without altering its mean.
- **DB:** For 50% of the observations in group B, a value of 90% of the feature's standard deviation is added; for the remaining 50%, the same value is subtracted. This induces a bimodal distribution in group B.
- **DM:** A value of 90% of the feature's standard deviation is added to 50% of the observations in group B, creating an asymmetric shift concentrated in one tail.
- **DP:** In both groups, observations are split into two subpopulations that are shifted upwards or downwards by 100% of the feature's standard deviation. In group A, 40% of the observations are shifted upwards and the remaining 60% downwards; in group B, 70% are shifted upwards and 30% downwards. Both groups are therefore bimodal with identical component locations but different mixing proportions, which also results in a difference in group means of 0.6 standard deviations. In the three-group simulations, group C is treated identically to group A.

Implementation

The methods for assessing differential distributions are implemented in the `gamdid` R package that is publicly available through GitHub (github.com/statOmics/gamdid) and will be submitted to Bioconductor.
`gamdid` builds upon the CRAN packages `mgcv` [17] and `harmonicmeanp` [18] for fitting generalized additive models and p-value aggregation; and on the R/Bioconductor packages `stageR` [14] for stagewise testing and `msqrob2` for detecting shifts in mean abundance [6]. Specifically, `mgcv` version 1.9.4, `harmonicmeanp` version 3.0.1, `stageR` version 1.32.0 and `msqrob2` version 1.18.0 were used in R version 4.5.3.

Data Availability

All data used in this manuscript are publicly available; no new mass spectrometry data were generated. The Leduc [1] and Petrosius [9] datasets were downloaded using the `scpdata` [8, 19] package from Bioconductor. The `scpdata` package redistributes pubilicly available SCP data using R/BioConductor data containers. The Leduc and Petrosius data were downloaded from the Slavov laboratory website (`https://scp.slavovlab.net/Leduc_et_al_2022`) and Zenodo (`https://zenodo.org/records/8146605`), respectively. The raw data and quantification data can also be found in the massIVE repositories MSV000089159 and MSV000092429.

Code Availability

All code used to prepare and analyze the datasets and produce the figures of this manuscript is available through GitHub (github.com/statOmics/gamdid). Finally, `gamdid` is published as an open source R-package available through GitHub (github.com/statOmics/gamdid) and will be submitted to Bioconductor.

Acknowledgements

This work was supported by grants from Ghent University Special Research Fund [BOF20/GOA/023] (S.V., L.C., C.V.), [BOF21/GOA/033] (L.M.) and [BOF/PDO/2025/018] (L.C., C.V.), Research Foundation Flanders [FWO G062219N] (S.V., L.C.), [G071326N] (L.C.), [G010023N, G028821N] (L.M.) and as WOG [W005325N] (L.M. and L.C.), funding from the European Union's Horizon 2020 Program [101080544 and 101191739] (L.M.).

Supplementary materials




Figure S1: ***Performance of gamdid and the standalone GAM method on the Leduc dataset*** *FDP-TPP curves comparing gamdid, the standalone GAM method, msqrob2 and distinct across the five simulation scenarios, based on the protein-level mock data of the Leduc dataset (755 monocytes). The dots mark the 5% FDR threshold and the dashed vertical line marks FDP = 0.05. The results are consistent with those for the Petrosius dataset (Figure 4): the standalone GAM method outperforms distinct for shape changes (DV, DB) but not when mean shifts are involved (DE, DM, DP), and gamdid combines the strengths of both components while maintaining conservative FDR control. The improvements over distinct are less pronounced than in the Petrosius dataset, which is attributable to the larger sample size: contrary to gamdid, distinct relies on permutation-based inference and suffers from reduced null distribution resolution with fewer observations. DB (differential both) denotes a difference in both modality and component means.*




Figure S2: ***Absolute true positive counts for gamdid and distinct at 5% FDR*** *Bar charts displaying the number of true positives recovered at the nominal 5% FDR threshold for gamdid and distinct across the five simulation scenarios in the Leduc dataset (left) and the Petrosius dataset (right). gamdid recovers a comparable or greater number of true positives in all scenarios, with the largest absolute gains in the DV and DB scenarios (shape-only differences). The false positive counts (Figure S3) are similarly low for both methods.*




Figure S3: ***Absolute false positive counts for gamdid and distinct at 5% FDR*** *Bar charts displaying the number of false positives at the nominal 5% FDR threshold for gamdid and distinct across the five simulation scenarios in the Leduc dataset (left) and the Petrosius dataset (right). Both methods maintain similarly low false positive counts across all scenarios in both datasets, confirming that the gains in true positive recovery demonstrated for gamdid in Figure S2 are not accompanied by inflation of false discoveries.*




Figure S4: ***Performance of gamdid on a subsampled Leduc dataset*** *FDP-TPP curves comparing gamdid and distinct across the five simulation scenarios, based on the protein-level mock data of a subsampled version of the Leduc dataset, in which cells were subsampled such that the average number of non-missing observations per group matched that of the Petrosius dataset. The dots mark the 5% FDR threshold and the dashed vertical line marks FDP = 0.05. This analysis directly tests whether the more pronounced improvements of gamdid over distinct observed in the Petrosius dataset (see Figure 4) are attributable to the smaller sample size. The results support this hypothesis: under reduced sample size, the performance gap between gamdid and distinct increases. distinct likely suffers from reduced null distribution resolution when fewer observations are available. DB (differential both) denotes a difference in both modality and component means.*




Figure S5: ***Performance of gamdid under reduced signal magnitude*** *FDP-TPP curves comparing gamdid and distinct across the five simulation scenarios, based on the protein-level mock data of the Petrosius dataset (top; 269 m2i cells) and the Leduc dataset (bottom; 755 monocytes), under simulation scenarios with 25% reduced signal magnitudes (DE: a shift of 30% instead of 40% of the feature's standard deviation was used; DV: random Gaussian noise with a standard deviation of 90% of the feature's standard deviation was added instead of 120%; DM and DB: a shift of 70% of the feature's standard deviation was used instead of 90%, and for DP 75% instead of 100%). The dots mark the 5% FDR threshold and the dashed vertical line marks FDP = 0.05. This analysis assesses whether gamdid's performance advantages are robust to weaker signals that may be encountered in practice. The results are consistent with the original simulations (Figure 4 and Supplementary Figure S1): gamdid controls the FDR at or below the nominal level and retains its advantages for shape-based distributional changes (DV, DB), while remaining competitive with distinct for scenarios where location shifts are involved. The relative improvements over distinct are more pronounced in the Petrosius dataset, consistent with gamdid's advantage under smaller sample sizes. As expected, the reduced signal does lower the overall power of both methods. DB (differential both) denotes a difference in both modality and component means.*




Figure S6: ***Performance of gamdid on peptide-level data*** *FDP-TPP curves comparing gamdid and distinct across the five simulation scenarios, based on the peptide-level mock data of the Petrosius dataset (top; 269 m2i cells) and the Leduc dataset (bottom; 755 monocytes), rather than the protein-level summarized data used in the main analyses. The dots mark the 5% FDR threshold and the dashed vertical line marks FDP = 0.05. The results are largely consistent with the protein-level findings (Figure 4 and Supplementary Figure S1), and gamdid retains its performance advantages. In the Leduc dataset, FDR control is less conservative at the peptide level than at the protein level for both methods. This reflects the well-known challenge in peptide-level data: multiple peptides mapping to the same protein are correlated. In the Petrosius dataset, gamdid controls the FDR at the peptide level. Despite this, gamdid provides a flexible and applicable framework across all processing levels of proteomics data. DB (differential both) denotes a difference in both modality and component means.*




Figure S7: ***Performance of the gamdid omnibus test in a three-group simulation*** *FDP-TPP curves for gamdid's omnibus test across the five simulation scenarios, based on the protein-level mock data of the Leduc dataset (755 monocytes). Cells were randomly assigned to three mock groups (A, B, C), with distributional signal introduced in group B under the same five simulation scenarios as in the two-group analyses. No competing DD method currently supports multi-group omnibus testing, so no benchmark comparison is possible here; these curves demonstrate gamdid's standalone performance in this setting. The dots mark the 5% FDR threshold and the dashed vertical line marks FDP = 0.05. Notably, the three-group omnibus test achieves higher power than the corresponding two-group comparisons in Supplementary Figure S1. DB (differential both) denotes a difference in both modality and component means.*

msqrob2	gamdid	distinct
0.030*	0.055	0.039*
0.046*	0.081	0.160
0.048*	0.081	0.039*
0.048*	0.084	0.141
0.049*	0.083	0.055
0.049*	0.086	0.128
0.149	0.090	0.035*
0.128	0.197	0.039*

Table S1: ***FDR-corrected p-values for features detected by msqrob2 and/or distinct but not by gamdid in the spike-in case study*** *The eight features that are flagged as significant by msqrob2 and/or distinct but not by gamdid are listed with their FDR-corrected p-values from all three methods. We see that gamdid consistently returns borderline non-significant p-values for the six features detected by msqrob2. In contrast, distinct produces both relatively high and relatively low p-values for these features. This indicates that only gamdid largely preserves the msqrob2 ranking of DA features. Significant FDR-corrected p-values (< 0.05) are marked with an asterisk (*).*

msqrob2	gamdid	distinct
0.931	0.028*	0.499
0.401	0.034*	0.067
0.074	0.045*	0.073

Table S2: ***FDR-corrected p-values for the three features detected exclusively by gamdid in the spike-in case study*** *The three features uniquely identified as significant by gamdid, with FDR-corrected p-values from all three methods. Two of the three features receive very high msqrob2 p-values, confirming that these differences do not involve a detectable mean shift and would be entirely missed by any conventional DA approach. Significant FDR-corrected p-values (< 0.05) are marked with an asterisk (*).*




Figure S8: ***gamdid* visualizations for features detected only by msqrob2 or only by distinct in the spike-in case study** *The output of gamdid's visFit function for two of the four features flagged as significant exclusively by msqrob2 (**A**, **B**) and the two features flagged as significant exclusively by distinct (**C**, **D**). Each panel shows the mock comparison (top), in which the groups show no distributional difference, the comparison after spiking 10% of the 697 sub-cluster A melanoma cells (69 cells) into group Y (middle), and a boxplot indicating where the spike-in melanoma cells are located (bottom). **A–B**: After spike-in, a weak mean shift appears, but no differential region is found by gamdid (no gray shading). This illustrates a feature type where msqrob2's dedicated sensitivity for mean shifts gives it a slight edge over gamdid. **C**: After spike-in, the melanoma cells add a heavier left tail to the distribution of group Y, but no differential region is found by gamdid (no gray shading). Here, distinct's ECDF-based approach accumulates sufficient evidence to reach significance, while gamdid's local density approach does not identify a sufficiently concentrated differential region. **D**: After spike-in, the density of group Y shifts in the right tail, and gamdid's difference smoother identifies a differential region (gray shading), despite the FDR-corrected p-value being non-significant (gamdid FDR adjusted p-value = 0.090).*




Figure S9: ***Computational efficiency of gamdid and distinct*** *Mean runtime for gamdid and distinct on the spike-in case study dataset (1316 features, 824 cells), each run on a single core and averaged across ten independent runs. gamdid completed in 16 seconds on average, whereas distinct required 256 seconds, corresponding to a more than fifteen-fold speedup.*

Supplementary methods: full statistical specification of the gamdid framework

This section provides the detailed statistical specification of the `gamdid` framework introduced in the Methods section. We first detail the density estimation approach and the Poisson generalized additive model used. We subsequently describe the inference procedure and the p-value aggregation strategy. Next we elaborate on the post-hoc pairwise comparisons and stagewise testing procedure, followed by the approach for identifying regions of differential density.

Model and inference The statistical methodology for comparing distributions is developed for a single feature (in this case proteins or peptides). The method is then simply repeated over multiple features and the corresponding p-values are corrected for multiple testing using the conventional Benjamini-Hochberg FDR procedure [13] or stagewise testing [10].

Density estimation through Poisson regression Following the approach of Lindsey [5], we reformulate density estimation in terms of Poisson regression by discretizing the sample space. We partition the sample space $\mathscr{Y}$ into K disjoint bins $\mathscr{Y}_k$ of equal width bw.

$$\mathscr{Y} = \bigcup_{k=1}^{K} \mathscr{Y}_k \tag{2}$$

The data $\boldsymbol{y} = (y_1, y_2, \ldots, y_n)^T$ are summarized into bin counts $\boldsymbol{s} = (s_1, s_2, \ldots, s_K)^T$ where each count s_k represents the number of observations falling into the k-th bin:

$$s_k = \#\{y_i \in \mathscr{Y}_k\} \quad \text{for } k = 1, 2, \ldots K \tag{3}$$

The sum of these counts equals the total number of observations n. Conceptually, this process is equivalent to constructing a histogram of the data, where each bin $\mathscr{Y}_k$ is associated with a count s_k and represented by a midpoint mid_k.

We consider s_k to be independent Poisson observations with expected values μ_k

$$s_k \overset{\text{ind}}{\sim} \text{Po}(\mu_k) \quad \text{for } k = 1, 2, \ldots K \tag{4}$$

Consequently, the problem of density estimation is recast as a spline-based Poisson regression of the bin counts s_k as a function of the bin midpoints mid_k [11].

Poisson generalized additive model The expected bin counts μ_k are modeled using Poisson regression implemented in the existing R package `mgcv` [17] for GAMs.

The linear predictor of a standard GAM with one smoother ($f(mid_k)$) is given by equation 5.

$$g(\mu_k) = \boldsymbol{A}_k \boldsymbol{\gamma} + f(mid_k) \tag{5}$$

with $\boldsymbol{A}_k$ the k-th row of the parametric $K \times G$ model matrix $\boldsymbol{A}$ and $\boldsymbol{\gamma}$ the parametric part of the model parameter vector. For Poisson distributed data, the link function g is a log function.

The relationship between $\log(\mu_k)$ and mid_k, $f(mid_k)$, is modeled using penalized thin plate regression splines (see equation 6) that provide a basis function expansion, i.e.

$$f(mid_k) = \sum_{p=1}^{P} \delta_p \phi_p(mid_k) \tag{6}$$

where $\phi_p(.)$ are thin plate spline basis functions, δ_p are the corresponding smooth basis parameters to be estimated. Let $\boldsymbol{\beta} = (\boldsymbol{\gamma}, \boldsymbol{\delta})$ where $\boldsymbol{\delta} = (\delta_1, \ldots, \delta_P)^T$.

Let $\boldsymbol{\Phi}$ be the $K \times P$ model matrix of the basis functions evaluated at the midpoints: $\phi_p(mid_k)$. We then combine $\boldsymbol{A}$ and $\boldsymbol{\Phi}$ in the $K \times (G + P)$ model matrix $\boldsymbol{X} = (\boldsymbol{A} : \boldsymbol{\Phi})$.

$\boldsymbol{\beta}$ is estimated by maximizing the penalized log-likelihood (see equation 7)

$$\hat{\boldsymbol{\beta}} = \arg\max_{\beta} \left\{ l(\boldsymbol{\beta}|\boldsymbol{s}) - \frac{1}{2}\lambda \boldsymbol{\beta}^{\mathrm{T}} \boldsymbol{S} \boldsymbol{\beta} \right\} \tag{7}$$

The optimization is typically performed using penalized iteratively reweighted least squares (PIRLS). Here, $l(\boldsymbol{\beta}|\boldsymbol{s})$ is the log likelihood of the data given $\boldsymbol{\beta}$, λ is the smoothing parameter that balances goodness-of-fit with smoothness, and $\boldsymbol{S}$ is the penalty matrix quantifying the wiggliness of the basis functions that has zeros on the rows and columns involving parameters for $\boldsymbol{A}$ [20]. We exploit the link between Bayesian statistics, smooth functions, and mixed models to estimate this smoothing parameter λ using the REML method [21, 22].

To approximate the effective degrees of freedom (edf) of the model, we define the influence matrix $\boldsymbol{H}$ in equation 8.

$$\boldsymbol{H} = (\boldsymbol{X}^T \boldsymbol{W} \boldsymbol{X} + \lambda \boldsymbol{S})^{-1} \boldsymbol{X}^T \boldsymbol{W} \boldsymbol{X} \tag{8}$$

with $\boldsymbol{W}$ the diagonal weight matrix from the final PIRLS iteration, where $W_{kk} = \mu_k = \exp(\boldsymbol{X}_k^T \hat{\boldsymbol{\beta}})$ and $\boldsymbol{X}_k$ is the k-th row of $\boldsymbol{X}$.

In the GAM literature $2\boldsymbol{H} - \boldsymbol{H}\boldsymbol{H}$ is the bias-corrected influence matrix (the bias arises due to smoothing), so its trace can be seen as the effective degrees of freedom of the bias-corrected model (see equation 9) [23].

$$edf = 2\,\mathrm{tr}(\boldsymbol{H}) - \mathrm{tr}(\boldsymbol{H}\boldsymbol{H}) \tag{9}$$

Comparing distributions across groups We build on this framework to detect distributional differences across multiple groups. Let c denote a group of the grouping variable C.

Next, the overall sample space $\mathscr{Y}$ is partitioned for all groups c into K disjoint bins $\mathscr{Y}_k$ of equal width bw (as in equation 2), ensuring identical bins (and thus midpoints mid_k) for all groups $c \in C$, as illustrated in Figure 2 panel A. For each group c, a count vector $\boldsymbol{s_c} = (s_{c_1}, s_{c_2}, ..., s_{c_K})$ is obtained, where s_{c_k} is the number of observations from group c in the k-th bin.

The full model allows for different distributions across groups, with the linear predictor specified in equation 10.

$$\log \mu_{ck} = \sum_{c \in C} \gamma_c x_c + \sum_{c \in C} f_c(mid_k) x_c + \log(O_c) \tag{10}$$

where each group c has its own intercept γ_c and smooth function f_c, with identical smoothing parameter λ among all groups.

The offset O_c is defined as

$$O_c = bw \times n_c \tag{11}$$

where n_c is the total number of observations in group c. This allows for normalization of different sample sizes across groups and ensures that the fitted smooth functions $\exp(\gamma_c + f_c(mid_k))$ can be directly interpreted as a probability density.

Conversely, the null model assumes a common density function across all groups of C. The linear predictor is modeled as in equation 12.

$$\log \mu_{ck} = \sum_{c \in C} \gamma_c x_c + f(mid_k) + \log(O_c) \tag{12}$$

where x_c is a dummy variable for the groups of covariate C, μ_{ck} is the expected count for group c in bin k, and the smooth function $f(\text{mid}_k)$ is shared across all groups. O_c is an offset term.

Inference for distributional differences Two approaches were considered for testing whether distributions differ across groups.

- Generalized likelihood ratio test (GLRT)

This approach is possible by comparing the full and the null model. However, it requires fitting two models and does not directly provide pairwise comparisons for a multi-group setting. Therefore, we provide a Wald-type test below and refer the reader to the GLRT section at the end of the supplementary materials.

- Wald-type test

A more practical approach involves Wald-type tests on the parameters of the smooth functions of the full model (see equation 10). This method builds on tests proposed by Simon Wood, which assess whether a smoother significantly deviates from zero, $H_0 : f_c = 0$ [12].

We extend this idea to test whether multiple smooth functions are identical across G groups. Specifically, we test the null hypothesis $H_0 : f_{c_1} = f_{c_2} = \cdots = f_{c_G}$. This hypothesis can be reformulated as a Wald test on the corresponding basis function parameters. Indeed, equality of smooth functions $f_{c_1} = f_{c_2} = \cdots = f_{c_G}$ implies equality of their associated parameter vectors $\boldsymbol{\beta}_{c_1} = \boldsymbol{\beta}_{c_2} = \cdots = \boldsymbol{\beta}_{c_G}$ if basis functions are shared across all groups.

To test this hypothesis, we define contrasts on the full vector of basis function parameters $\hat{\boldsymbol{\beta}}_{\text{full}} = (\hat{\boldsymbol{\beta}}_{c_1}, \ldots, \hat{\boldsymbol{\beta}}_{c_G})$ from the full model in equation 10. The null hypothesis can then be expressed as $H_0 : \boldsymbol{L}\boldsymbol{\beta}_{\text{full}} = 0$ where $\boldsymbol{L}$ is a contrast matrix encoding $G - 1$ independent differences between parameter vectors, i.e. $(\boldsymbol{\beta}_{c_1} - \boldsymbol{\beta}_{c_g} = 0 \text{ for } g = 2, \ldots, G)$.

The corresponding Wald statistic is then given by equation 13.

$$T_W = \hat{\boldsymbol{\beta}}_{\text{full}}^T \boldsymbol{L}^T (\boldsymbol{L}\boldsymbol{V}_{\beta_{\text{full}}}\boldsymbol{L}^T)^{-1} \boldsymbol{L}\hat{\boldsymbol{\beta}}_{\text{full}} \tag{13}$$

where $\boldsymbol{V}_{\beta_{\text{full}}} = (\boldsymbol{X}^T\boldsymbol{W}\boldsymbol{X} + \lambda\boldsymbol{S})^{-1}$ is the covariance matrix for $\boldsymbol{\beta}_{\text{full}}$. T_W is asymptotically χ^2_z distributed.

The degrees of freedom z correspond to the total edf associated with the $G - 1$ smooth functions that are dropped under H_0.

As a conservative estimate, we define z as the sum of the $G - 1$ largest edf values.

The p-value obtained from the χ^2_z distribution of T_W reflects any form of distributional deviation between groups: mean shifts, differences in variance, modality, skewness, etc. In what follows, we refer to this as the GAM p-value.

P-value aggregation While the GAM p-value is sensitive to a wide range of distributional differences, it exhibits limited power in detecting pure mean shifts. To improve sensitivity to such shifts, we complement the GAM p-value with a p-value derived from the msqrob2 framework, which is specifically designed to detect mean shifts in proteomics data using a robust linear regression through a mixed model framework [6].

The GAM p-value and the msqrob2 p-value are aggregated using the HMP method. The HMP method is particularly suited for combining correlated p-values [7]. This is inherently the case since mean shifts are one aspect of overall distributional changes. The resulting aggregated p-value, termed the 'feature-level aggregated p-value', provides a single, comprehensive measure of evidence against the null hypothesis of no distributional difference among groups for that feature.

Post-hoc pairwise comparisons If covariate C comprises more than two groups, post-hoc pairwise comparisons can be conducted. For each of the $\mathcal{C}(\mathcal{C} - 1)/2$ pairs of groups (where $\mathcal{C}$ is the number of groups in C), a pairwise GAM p-value is calculated using the Wald test framework (see equation 13) with an appropriate contrast matrix $\boldsymbol{L}$. Similarly, pairwise msqrob2 p-values are computed. These pairwise p-values are then aggregated for each group-to-group comparison using the HMP method, yielding 'pairwise

aggregated p-values', each representing the evidence against the null hypothesis of identical distributions for that specific pair of groups.

Stagewise testing A stagewise approach is employed to effectively manage the multiple hypotheses arising from feature-level and subsequent pairwise comparisons [10].

- Stage 1 (feature-level FDR control): The feature-level aggregated p-values are adjusted across all features using the Benjamini-Hochberg (BH) procedure to control the FDR at 5%.
- Stage 2 (pairwise FWER control) is only conducted if stage 1 results in a significant p-value. The pairwise aggregated p-values are adjusted for multiple testing across the $\mathcal{C}(\mathcal{C}-1)/2$ pairwise hypotheses within a feature to control the family-wise error rate (FWER) on the BH-adjusted significance level (i.e., the FDR q-value) of the first stage [14].

Identifying regions of differential density For pairs of groups exhibiting significant distributional differences, the specific regions contributing to these differences can be pinpointed. This is achieved by constructing a difference smoother $\hat{d}$, representing the estimated difference in log-densities between the groups, and its associated simultaneous confidence bands.

Consider the comparison between two groups: c_1 and c_2 from the groups of C. The difference smoother $\hat{d}$ is evaluated across a grid of N points: x_n for $n = 1, \ldots, N$ spanning the range of mid_k values (see equation 14).

$$\hat{d}(x_n) = (\hat{\gamma}_{c_1} + \hat{f}_{c_1}(x_n)) - (\hat{\gamma}_{c_2} + \hat{f}_{c_2}(x_n)) \tag{14}$$

Note that $\hat{d}(x_n)$ can also be expressed as a linear combination of the full parameter vector $\hat{\boldsymbol{\beta}}_{\text{full}}$. Let $\boldsymbol{x}_{diff}$ be the contrast row vector for the comparison of c_1 and c_2 so that the difference smoother can be defined as in equation 15.

$$\hat{d}(x_n) = x_{\text{diff},n}\hat{\boldsymbol{\beta}}_{\text{full}} \tag{15}$$

The corresponding pointwise standard error is subsequently computed in equation 16.

$$se(\hat{d}(x_n)) = \sqrt{x_{\text{diff,n}}^T \boldsymbol{V}_{\beta_{\text{full}}} x_{\text{diff,n}}} \tag{16}$$

where $\boldsymbol{V}_{\beta_{\text{full}}}$ is the covariance matrix for $\hat{\beta}_{\text{full}}$.

A $1-\alpha$ pointwise confidence interval for $\hat{d}(x_n)$ is then given by $\hat{d}(x_n) \pm z_{\alpha/2}\, se(\hat{d}(x_n))$ where $z_{\alpha/2}$ is the $1-\alpha/2$ quantile of the standard normal distribution.

To enable valid inference over the entire difference smoother $\hat{d}$ simultaneously, the pointwise confidence intervals need to be adjusted for multiple testing. Essentially, this is equivalent to adjusting for the multiple pointwise tests conducted along the grid. As a computationally efficient approach, we use the Šidák multiple testing correction to transform the pointwise intervals into simultaneous confidence bands [24]. The Šidák correction will be conservative because the points are positively correlated [25].

Regions where the simultaneous band does not encompass zero can then be used to prioritize DD regions.

The $1-\alpha_{\text{overall}}$ simultaneous confidence interval for $\hat{d}(x_n)$ then becomes $\hat{d}(x_n) \pm z^*\, se(\hat{d}(x_n))$, where the adjusted critical value z^* is $z_{1-(1-\alpha_{\text{overall}}/2)^{1/M}}$. M denotes the effective number of independent locations along the grid, which is conservatively estimated as the maximum of the effective degrees of freedom associated with the two smooth functions being compared $\hat{f}_{c_1}$ and $\hat{f}_{c_2}$. This choice is motivated by the equivalence between such grid tests and omnibus Wald-type tests on the basis function parameters (see equation 13).

Similar to the previous section, a stagewise testing approach is used to correct for multiple testing along the features. This means that α_{overall} corresponds to the BH adjusted significance level of the first stage to control the feature-level FDR at 5% [10].

Note, however, that we do not provide simultaneous confidence intervals that account for multiple testing along features as well as within features to address multiple comparisons between groups. Indeed, this would require the extension of the stagewise testing procedure toward multiple stages.

Generalized likelihood ratio test

The null model (equation 12) can be compared to the full model (equation 10) using a GLRT (see equation 17).

$$D = 2\{l(\beta_{full}) - l(\beta_0)\} \tag{17}$$

where $l(\beta_{full})$ and $l(\beta_0)$ are the maximized log-likelihoods for the full and the null models, respectively. Under H_0, D asymptotically follows a chi-squared distribution (see equation 18).

$$D \sim \chi^2_{\Delta\text{edf}} \tag{18}$$

where $\Delta\text{edf} = \text{edf}_{full} - \text{edf}_0$. edf_{full} and edf_0 are derived from equation 9. The p-value is then $P(\chi^2_{\Delta edf} \geq D)$ [26].

This approach requires fitting two models and does not directly provide pairwise comparisons if more than two groups exist.

References

[1] Andrew Leduc et al. "Exploring functional protein covariation across single cells using nPOP". In: *Genome Biology* 23 (2022), p. 261. DOI: 10.1186/s13059-022-02817-5. URL: https://doi.org/10.1186/s13059-022-02817-5.

[2] Catalina A. Vallejos, Sylvia Richardson, and John C. Marioni. "Beyond comparisons of means: understanding changes in gene expression at the single-cell level". In: *Genome Biology* 17 (2016), p. 70. DOI: 10.1186/s13059-016-0930-3. URL: https://doi.org/10.1186/s13059-016-0930-3.

[3] Keegan Korthauer et al. "A statistical approach for identifying differential distributions in single-cell RNA-seq experiments". In: *Genome Biology* 17 (Oct. 2016). DOI: 10.1186/s13059-016-1077-y.

[4] Simone Tiberi et al. "distinct: A novel approach to differential distribution analyses". In: *The Annals of Applied Statistics* 17.2 (June 2023). ISSN: 1932-6157. DOI: 10.1214/22-aoas1689. URL: http://dx.doi.org/10.1214/22-AOAS1689.

[5] J.K. Lindsey and G. Mersch. "Fitting and comparing probability distributions with log linear models". In: *Computational Statistics and Data Analysis* 13.4 (1992), pp. 373–384. ISSN: 0167-9473. DOI: https://doi.org/10.1016/0167-9473(92)90112-S. URL: https://www.sciencedirect.com/science/article/pii/016794739290112S.

[6] Adriaan Sticker et al. "Robust Summarization and Inference in Proteome-wide Label-free Quantification". In: *Molecular & Cellular Proteomics* 19.7 (2020), pp. 1209–1219. DOI: 10.1074/mcp.ra119.001624. URL: https://doi.org/10.1074/mcp.ra119.001624.

[7] DJ Wilson. "Generalized mean p-values for combining dependent tests: comparison of generalized central limit theorem and robust risk analysis [version 1; peer review: 2 approved]". In: *Wellcome Open Research* 5.55 (2020). DOI: 10.12688/wellcomeopenres.15761.1.

[8] Christophe Vanderaa and Laurent Gatto. "The Current State of Single-Cell Proteomics Data Analysis". In: *Current Protocols* 3.1 (2023), e658. DOI: https://doi.org/10.1002/cpz1.658. eprint: https://currentprotocols.onlinelibrary.wiley.com/doi/pdf/10.1002/cpz1.658. URL: https://currentprotocols.onlinelibrary.wiley.com/doi/abs/10.1002/cpz1.658.

[9] Valdemaras Petrosius et al. "Exploration of cell state heterogeneity using single-cell proteomics through sensitivity-tailored data-independent acquisition". In: *Nature Communications* 14 (Sept. 2023), p. 5910. DOI: 10.1038/s41467-023-41602-1.

[10] Ruth Heller et al. "A flexible two-stage procedure for identifying gene sets that are differentially expressed". In: *Bioinformatics* 25.8 (Feb. 2009), pp. 1019–1025. ISSN: 1367-4803. DOI: 10.1093/bioinformatics/btp076. eprint: https://academic.oup.com/bioinformatics/article-pdf/25/8/1019/50287411/bioinformatics_25_8_1019.pdf. URL: https://doi.org/10.1093/bioinformatics/btp076.

[11] Bradley Efron and Robert Tibshirani. "Using Specially Designed Exponential Families for Density Estimation". In: *The Annals of Statistics* 24.6 (1996), pp. 2431–2461. ISSN: 00905364, 21688966. URL: http://www.jstor.org/stable/2242691.

[12] Simon N. Wood. "On p-values for smooth components of an extended generalized additive model". In: *Biometrika* 100.1 (Oct. 2012), pp. 221–228. ISSN: 0006-3444. DOI: 10.1093/biomet/ass048. eprint: https://academic.oup.com/biomet/article-pdf/100/1/221/478658/ass048.pdf. URL: https://doi.org/10.1093/biomet/ass048.

[13] Yoav Benjamini and Yosef Hochberg. "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing". In: *Journal of the Royal Statistical Society. Series B (Methodological)* 57.1 (1995), pp. 289–300. ISSN: 00359246. URL: http://www.jstor.org/stable/2346101.

[14] Koen Van den Berge et al. "stageR: a General Stage-wise Method for Controlling the Gene-level False Discovery Rate in Differential Expression and Differential Transcript Usage". In: *Genome Biology* 18 (2017), p. 151. DOI: 10.1186/s13059-017-1277-0. URL: https://doi.org/10.1186/s13059-017-1277-0.

[15] Jürgen Cox and Matthias Mann. "MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification". In: *Nature Biotechnology* 26.12 (2008), pp. 1367–1372. DOI: 10.1038/nbt.1511.

[16] Oliver Bernhardt et al. *Spectronaut: a fast and efficient algorithm for MRM-like processing of data independent acquisition (SWATH-MS) data*. F1000Posters 5: 1092. Poster presented at the 60th ASMS Conference on Mass Spectrometry and Allied Topics, Vancouver, 20–24 May 2012. 2014. URL: https://f1000research.com/posters/1096450.

[17] Simon N. Wood. "Generalized Additive Models". In: *Annual Review of Statistics and Its Application* 12.Volume 12, 2025 (2025), pp. 497–526. ISSN: 2326-831X. DOI: https://doi.org/10.1146/annurev-statistics-112723-034249. URL: https://www.annualreviews.org/content/journals/10.1146/annurev-statistics-112723-034249.

[18] Daniel J. Wilson. "The harmonic mean p-value for combining dependent tests". In: *Proceedings of the National Academy of Sciences U.S.A.* 116.4 (2019), pp. 1195–1200. URL: https://www.pnas.org/content/116/4/1195.

[19] Christophe Vanderaa and Laurent Gatto. "Replication of single-cell proteomics data reveals important computational challenges". In: *Expert Review of Proteomics* 18.10 (2021). PMID: 34602016, pp. 835–843. DOI: 10.1080/14789450.2021.1988571. eprint: https://doi.org/10.1080/14789450.2021.1988571. URL: https://doi.org/10.1080/14789450.2021.1988571.

[20] Simon N. Wood. "Generalized Additive Models: An Introduction with R, Second Edition". In: 2nd ed. Chapman and Hall/CRC, 2017. Chap. 6, pp. 249–251. DOI: 10.1201/9781315370279.

[21] Simon N. Wood. "Generalized Additive Models: An Introduction with R, Second Edition". In: 2nd ed. Chapman and Hall/CRC, 2017. Chap. 6, pp. 262–263. DOI: 10.1201/9781315370279.

[22] David Ruppert, M. P. Wand, and R. J. Carroll. "Semiparametric Regression". In: Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2003. Chap. 6, pp. 137–142.

[23] Simon N. Wood. "Generalized Additive Models: An Introduction with R, Second Edition". In: 2nd ed. Chapman and Hall/CRC, 2017. Chap. 6, pp. 251–252. DOI: 10.1201/9781315370279.

[24] Zbyněk Šidák. "Rectangular Confidence Regions for the Means of Multivariate Normal Distributions". In: *Journal of the American Statistical Association* 62.318 (1967), pp. 626–633. ISSN: 01621459, 1537274X. URL: http://www.jstor.org/stable/2283989.

[25] Sanat K. Sarkar and Chung-Kuei Chang. "The Simes Method for Multiple Hypothesis Testing with Positively Dependent Test Statistics". In: *Journal of the American Statistical Association* 92.440 (1997), pp. 1601–1608. DOI: 10.1080/01621459.1997.10473682. eprint: https://doi.org/10.1080/01621459.1997.10473682. URL: https://doi.org/10.1080/01621459.1997.10473682.

[26] Simon N. Wood. "Generalized Additive Models: An Introduction with R, Second Edition". In: 2nd ed. Chapman and Hall/CRC, 2017. Chap. 6, pp. 313–315. DOI: 10.1201/9781315370279.