EVERY FINITE GROUP FROM A WALL: LARGE AUTOMORPHISM GROUPS OF SECOND-ORDER CELLULAR AUTOMATA

MICHAEL DAYAH

ABSTRACT. For an elementary rule f, the second-order (Fredkin) automaton F_f on $(\{0,1\}^2)^{\mathbb{Z}}$ is reversible, and the automorphism group of its spacetime subshift is the centralizer of F_f in the automorphism group of the full 4-shift. Many of these automata have walls, patterns that stay fixed or periodic at a fixed position whatever surrounds them; they were listed by Takesue (1989), who observed that no information passes through them. We determine what walls imply for the automorphism group. If a reversible automaton of radius 1 has a word that it maps to itself in every context, and at least two symbols can sit between two copies of it, then permuting the contents of trains of identical segments embeds $\bigoplus_m S_m$, and hence every finite group, in the automorphism group. With Takesue's fixed walls this applies to 32 of the 88 conjugacy classes of second-order elementary automata, including 54R and 37R, and we prove that no other class has a fixed word of any length. Variants of the construction, for walls that are periodic in time or that carry a bit invisible to their neighbours, give the same conclusion in seven further classes. Three affine classes, among them that of 90R, which has no wall and no blocking word, reach it through marker permutations; altogether 42 classes (102 rules) have every finite group in their automorphism group. For four other affine classes the isomorphism rigidity theorem of Kitchens and Schmidt shows that every automorphism is affine, and the automorphism group is $\mathbb{Z}^2$, $\mathbb{Z}^2 \times \mathbb{Z}/2$ or, for rule 150, $\mathbb{Z}^3$. So for affine rules the absence of a blocking word does not force the automorphism group to be virtually $\mathbb{Z}^2$.

1. INTRODUCTION

Let $\mathcal{A}$ be a finite alphabet and σ the left shift on $\mathcal{A}^{\mathbb{Z}}$. By the Curtis–Hedlund–Lyndon theorem [7] the endomorphisms of the full shift are the block maps, and the automorphisms are the invertible block maps. The group $\mathrm{Aut}(\mathcal{A}^{\mathbb{Z}})$ is enormous: it contains every finite group and free groups [5], and every countable, locally finite, residually finite group [23], and its center consists of the powers of the shift [15]. A reversible cellular automaton $F \in \mathrm{Aut}(\mathcal{A}^{\mathbb{Z}})$ determines a two-dimensional subshift, its *spacetime* $X_F \subset \mathcal{A}^{\mathbb{Z}^2}$, whose rows are the configurations $F^n(c)$, $n \in \mathbb{Z}$ [6]. Its automorphism group is the centralizer of F in $\mathrm{Aut}(\mathcal{A}^{\mathbb{Z}})$ (Lemma 2.2), and it always contains $\langle \sigma, F \rangle$. This paper is about automata for which it is much larger.

The automata we consider are the second-order elementary rules, built by the second-order technique of Fredkin [20, 27]; as a family they were introduced and studied systematically by Takesue [18, 19], under the name of elementary reversible cellular automata. For an elementary rule $f\colon \{0,1\}^3 \to \{0,1\}$ define, on the alphabet $\mathcal{A} = \{0,1\}^2$ with cells written (x, y),

$$F_f(c)_i = \big(y_i,\ x_i \oplus f(y_{i-1}, y_i, y_{i+1})\big). \tag{1}$$

Reading y as the current row and x as the previous one, this is the recursion $y_{t+1} = y_{t-1} \oplus f(y_t)$, which is reversible for every f, with $F_f^{-1}(c)_i = \big(y_i \oplus f(x_{i-1}, x_i, x_{i+1}),\ x_i\big)$. Following Takesue [18], one writes fR for this automaton; 37R is Wolfram's standard example of a reversible automaton with complex behaviour [27, pp. 440 and 452–457]. Under reflection and the cellwise complement the 256 rules fall into 88 conjugacy classes, the classes of Takesue [19, Table I], and $\mathrm{Aut}(X_{F_f})$ depends only on the class (Section 2).

Date: November 10, 2026.

For some of these automata the automorphism group is as small as possible. For rule 30, for all non-affine rules permutive in an outer variable, and for the classes of 63 and 187, $\mathrm{Aut}(X_{F_f}) = \langle \sigma, F_f \rangle \cong \mathbb{Z}^2$ [29]; the same holds for the classes of 110, 14, 46 and 79 [28]. The classification of all 88 classes is the subject of [30]. Here we study the opposite behaviour. The spacetime X_F has zero $\mathbb{Z}^2$-entropy, so Hochman's theorem that positive entropy forces every finite group into the automorphism group of a $\mathbb{Z}^2$ shift of finite type [8] never applies to it, and something else must supply the room. In every case we find, it is a local structure.

For most of the classes treated here the local structure is a *wall* in the sense of Takesue [19, Section 4]: a pattern, fixed or periodic in time, that appears at a fixed position irrespective of its surroundings. Takesue relates walls to local conservation laws, exhibits walls in 40 of the 88 classes (the 38 classes of [19, Fig. 2, Table III] and those of 27 and 59), and observes that the two sides of a wall cannot communicate, so that a large system with walls is divided into parts that do not interact and is not mixing. This observation precedes the formalism of blocking words and equicontinuity points [13, 3], of which walls are instances (Remark 6.26). The walls of the second-order elementary automata are therefore known, and we refer to [19] for them. What this paper adds is their consequence for the automorphism group: the embeddings of $\bigoplus_m S_m$ by permutations along trains of walls; a proof that the list of fixed walls is complete for words of every length; a bit hidden in words built from the periodic walls, with the flips that it gives; the census of rule 59R at radius 2; and the results on rule 54. Where there are no walls, it adds the results on the linear and affine rules.

Fixed walls. The basic mechanism is a word that the automaton fixes in every context.

Theorem 1.1. *Let F be a reversible cellular automaton of radius 1 on $\mathcal{A}^{\mathbb{Z}}$ and W a nonempty word such that $F(c)$ contains W at every position at which c contains W. Let $C \subseteq \mathcal{A}$ be the set of symbols s for which $W\,s\,W$ contains W only at its two ends. If $|C| \geq 2$, then $\mathrm{Aut}(X_F)$ contains a copy of $\bigoplus_{m\geq 1} S_m$, hence of every finite group.*

The copies of W act as walls. A symbol between two walls evolves by a fixed permutation of C, whatever lies beyond the walls, so the contents of a run of such cells can be permuted without disturbing the dynamics. For the second-order elementary automata such words are known: the shortest fixed words of the 31 classes of Table 1, and the word 0330 for the class of 133, are the fixed walls (i)–(ix) of Takesue [19, Fig. 2, Table III and p. 391], who examined the words of length at most 6. We verify the hypothesis $|C| \geq 2$ for them, and we prove that no other class has a fixed word of any length (Proposition 5.2).

Corollary 1.2. *For the 81 elementary rules f in the 31 classes listed in Table 1, $\mathrm{Aut}(X_{F_f})$ contains every finite group. These include 54, 37, 22 and 73. For the class $\{133, 161\}$ the same holds, through the fixed word 0330 of length 4. No other elementary rule has a word of any length fixed by F_f in every context.*

Rule 54. Rule 54 is, with rule 110, one of the two elementary rules for which Wolfram reports class 4 behaviour, rule 110 as the clear example and rule 54 as a possible one [27, pp. 693–697]. The two second-order automata behave in opposite ways: $\mathrm{Aut}(X_{F_{110}}) = \langle \sigma, F_{110} \rangle$ [28], while for 54R the pattern 33 (writing 3 for the cell $(1, 1)$) is a wall, the wall (iv) of [19, Table III], and the walls give the following.

Theorem 1.3. *For every $m \geq 1$ there is an injective homomorphism $\Psi_m \colon S_m \to \mathrm{Aut}(X_{F_{54}})$ whose image consists of block maps of radius at most $3m + 3$, and the images for different m commute, so $\bigoplus_{m\geq 1} S_m$ embeds in $\mathrm{Aut}(X_{F_{54}})$.*

The relabellings of the segments between walls are automorphisms of smaller radius: a permutation of the contents of the segments of a given length L is an automorphism whenever

it commutes with the permutation that F induces on these contents, and these maps give an embedding of the direct sum over L of the centralizers of the induced permutations (Theorem 4.2). The smallest of them is D, of radius 2 and order 3, which acts as the 3-cycle $0 \to 1 \to 2 \to 0$ on every cell that is a single-cell segment between two walls and as the identity elsewhere. At radius 2 an exhaustive computation shows that the walls account for everything: the automorphisms of radius at most 2 are exactly the 13 maps $\sigma^a F^b$ with $|a| + |b| \le 2$ and the 10 maps $F^b D^k$ with $|b| \le 2$, $k \in \{1, 2\}$ (Theorem 4.4). At radius 3 this is no longer so: there is an involution τ of radius 3 that commutes with F, moves the boundary between a long wall and a single cell next to it, and does not commute with D (Theorem 4.6). A partial enumeration of the automorphisms of radius 3 (Section 4.6) finds that every automorphism it produces differs from some $\sigma^a F^b$ only on windows that contain a wall, and that most of them, like τ, do not preserve walls.

Walls that are not fixed. A wall need not be fixed in every context. Takesue's list contains walls that are periodic in time and not fixed, in the classes of 27, 59, 179, 127, 223, 126 and 219 [19, Section 4, (4.1)–(4.3), Fig. 2 and Table III, walls (xii), (xiii), (xv), (xvi)]. None of these seven classes has a word fixed in every context, so Theorem 1.1 does not apply to them. In Section 6 we give a general mechanism (Theorem 6.6) for walls whose image never depends on the context, except at cells that do not matter, and which carry one bit that F cannot see. Permuting these hidden bits along chains of walls gives every finite group again. The mechanism applies to words built from Takesue's periodic walls in the classes of 59, 27, 127, 179 and 223 (Theorems 6.14 and 6.20, Proposition 6.19). For 219R and 126R the walls rotate through the alphabet in every context, and permuting the cells between them again embeds every finite group (Theorem 6.23, Proposition 6.25).

The affine rules 90 and 165. Rule 90 has no blocking word [13, 3], hence no wall of any of the kinds above (Corollary 7.13), and the automorphisms of $X_{F_{90}}$ of radius at most 2 are all affine (Section 7). Nevertheless F_{90} has two factor maps onto the binary full shift that carry F to $\sigma^{\pm 1}$, with matching embeddings (Theorem 7.2), and these carry marker automorphisms of the binary full shift into $X_{F_{90}}$: by Theorem 7.10, $(\bigoplus_m S_m)^2$ embeds in $\mathrm{Aut}(X_{F_{90}})$ and in $\mathrm{Aut}(X_{F_{165}})$. Theorem 7.2 also describes the block maps of radius at most 3 that commute with F_{90}, invertible or not: they are the sums of an affine map and of maps that factor through the two factor maps.

The other affine rules. For the affine rules whose characteristic polynomial $t^2 + pt + 1$ is irreducible and whose spacetime action is mixing, the isomorphism rigidity theorem of Kitchens and Schmidt [11] forces every automorphism to be affine (Lemma 8.4), and the automorphism group becomes a unit group, which we compute. Write κ for the cellwise complement.

Theorem 1.4. *Every automorphism of X_{F_f} is affine for the rules f of the four classes below, and:*

1. *for $f \in \{170, 240\}$, $\mathrm{Aut}(X_{F_f}) = \langle F, h\rangle \cong \mathbb{Z}^2$ with $h = \mathrm{id} + F$, and $h^2 = \sigma^{\pm 1} F$, so $\langle \sigma, F\rangle$ has index 2; for $f \in \{15, 85\}$ the same holds with $h(c) = c + F(c) + (1, 1)^{\mathbb{Z}}$;*
2. *for $f \in \{60, 102\}$, $\mathrm{Aut}(X_{F_f}) = \langle \sigma, F\rangle \times \langle \kappa\rangle \cong \mathbb{Z}^2 \times \mathbb{Z}/2$;*
3. *for $f \in \{153, 195\}$, $\mathrm{Aut}(X_{F_f}) = \langle \sigma, F\rangle \times \langle \kappa\rangle \cong \mathbb{Z}^2 \times \mathbb{Z}/2$;*
4. *for $f = 150$, $\mathrm{Aut}(X_{F_f}) = \langle \sigma, F, h\rangle \cong \mathbb{Z}^3$ with $h = \mathrm{id} + \sigma F$; for $f = 105$ the same holds with $h(c) = c + \sigma F(c) + (1, 1)^{\mathbb{Z}}$.*

For 170 the spacetime, with the action of F and $\mathrm{id} + F$, is conjugate to Ledrappier's three-dot shift, with $\langle \sigma, F\rangle$ acting as a subgroup of index 2 of its shift action (Remark 8.6). The hypotheses fail exactly for the other affine classes, those of 0, 51, 90, 165 and 255, and each of them has non-affine automorphisms (Remark 8.7). For 255, F adds 1 modulo 4 in every cell, and

$\mathrm{Aut}(X_{F_{255}}) \cong \mathbb{Z}/4 \times \mathcal{G}_4$, where $\mathcal{G}_4$ is a group of automorphisms of the full shift on $\mathbb{Z}/4$ that contains $\bigoplus_m S_m$ (Proposition 8.10).

Summary. Collecting these results, the following 42 classes of second-order elementary automata, comprising 102 rules, have the property that $\bigoplus_{m\geq 1} S_m$ embeds in $\mathrm{Aut}(X_{F_f})$, so that $\mathrm{Aut}(X_{F_f})$ contains every finite group and is not virtually $\mathbb{Z}^n$ for any n:

- the 31 classes (81 rules) of Table 1, with a word of length at most 3 fixed in every context, and the class of 133 (2 rules), with one of length 4 (Theorem 1.1, Corollary 1.2);
- the classes of 59 and 27 (8 rules), with hidden-bit walls (Theorems 6.14 and 6.20);
- the classes of 127, 179 and 223 (6 rules), with time-periodic walls (Proposition 6.19);
- the classes of 219 and 126 (2 rules), with rotating walls (Theorem 6.23, Proposition 6.25);
- the affine classes of 90 and 165 (2 rules), where $(\bigoplus_m S_m)^2$ embeds (Theorem 7.10);
- the affine class of 255 (1 rule), through marker permutations of the difference sequence (Proposition 8.10).

The first four items comprise 39 classes. Their walls are those of Takesue or are built from them, and with the class of 255, which also has walls in his list, they are the 40 classes in which he exhibits walls [19, Section 4]. Apart from the description $\mathrm{Aut}(X_{F_{255}}) \cong \mathbb{Z}/4 \times \mathcal{G}_4$, we do not determine the whole automorphism group in these cases. For the other 46 classes, the status of each is recorded in [30]: for the class of 95 the automorphism group is determined in [31], and it also contains $\bigoplus_{m\geq 1} S_m$, through automorphisms that permute moving particles and not through a wall; 11 classes have $\mathrm{Aut}(X_{F_f}) = \langle \sigma, F_f \rangle \cong \mathbb{Z}^2$ [28, 29, 31], the four affine classes of Theorem 1.4 have the groups given there, two further classes have the cellwise symmetry κ, and for 28 classes an exhaustive search finds no automorphism outside $\langle \sigma, F_f \rangle$ of radius at most 3.

Blocking words and linearity. In the nonlinear examples of this paper the automorphisms outside $\langle \sigma, F \rangle$ all come from blocking words with a degree of freedom next to the blocking column (Remark 6.26). The following remark records what the affine rules say about the converse.

Remark 1.5. Without a hypothesis on f, the absence of a blocking word does not force $\mathrm{Aut}(X_{F_f})$ to be virtually $\mathbb{Z}^2$. The affine rules 90 and 165 have no blocking word (Corollary 7.13), yet $(\bigoplus_m S_m)^2$ embeds in their automorphism groups (Theorem 7.10). Rule 150 is a counterexample of a different kind. It has no blocking word, and its commuting block maps of radius at most 3 are all linear (Proposition 7.4); but $\mathrm{id} + \sigma F_{150}$ is an automorphism outside $\langle \sigma, F_{150} \rangle$, and $\langle \sigma, F_{150}, \mathrm{id} + \sigma F_{150} \rangle \cong \mathbb{Z}^3$ (Proposition 7.5), so $\mathrm{Aut}(X_{F_{150}})$ is not virtually $\mathbb{Z}^2$; in fact $\mathrm{Aut}(X_{F_{150}}) = \langle \sigma, F_{150}, \mathrm{id} + \sigma F_{150} \rangle$ (Theorem 1.4). In all three the room comes from linearity, not from a local structure.

Nor does the absence of a blocking word force $\mathrm{Aut}(X_{F_f})$ to be virtually $\mathbb{Z}^2$ when f is not affine: the rule 95 is not affine, F_{95} has no blocking word, and $\mathrm{Aut}(X_{F_{95}})$ contains every finite group [31]. There the room comes from automorphisms that permute particles moving at light speed, and neither from a local structure nor from linearity. We do not know another example of this kind among the elementary rules.

Question 1.6. Let f be an elementary rule that is not affine and does not belong to the class of 95. If the second-order automaton F_f has no blocking word [13, 3], is $\mathrm{Aut}(X_{F_f})$ virtually $\mathbb{Z}^2$?

Having no blocking word is stronger than having no word fixed in every context; the rules of Section 6 show that the difference matters (Remark 6.26). For 110R and 30R, whose automorphism groups are $\mathbb{Z}^2$ [28, 29], a single changed cell produces a difference that spreads in both directions at linear speed, whereas in 54R it spreads only a few cells (Table 2).

Related work. *Large automorphism groups.* The automorphism group of a mixing one-dimensional shift of finite type contains every finite group [5]. In two dimensions the same holds under strong irreducibility [21] and under positive entropy [8]. Hochman also notes that zero entropy does not prevent a large automorphism group, with the example of a shift whose columns are constant; that shift is the spacetime of the identity automaton. Theorems 1.1 and 1.3 are of the same kind, but for nonlinear automata of infinite order, where the room comes from a conserved local structure. For linear automata the algebraic theory of Kitchens and Schmidt applies [10]; their isomorphism rigidity theorem [11] is what determines the automorphism groups in Theorem 1.4, and for Ledrappier's shift [14] the automorphism group is $\mathbb{Z}^2$ [21], [1, Theorem 7], [16, Example 8.20].

Centralizers and blocking words. Ryan's theorem identifies the centralizer of the whole automorphism group of a full shift with the powers of the shift [15], and finitary versions find finite sets of automorphisms with the same centralizer [22, 12]; a spacetime automorphism group is the centralizer of a single automaton, which these results do not cover. The term blocking word is due to Blanchard and Tisseur [3]; the relation of blocking words to equicontinuity is in the proof of [13, Theorem 3]. For the second-order elementary automata, that a wall separates the two sides of a configuration was observed by Takesue [18], [19, Section 4], before these notions were introduced.

Rule 54. The particles and collisions of the elementary rule 54 were catalogued by Boccara, Nasser and Roger [4] and by Martínez, Adamatzky, Chen and Chua [24]. A different reversible automaton, introduced by Bobenko, Bordemann, Gunn and Pinkall [25] and studied extensively as an integrable model of statistical mechanics [26], is also called Rule 54 because it uses the same local function $b \oplus (a \vee c)$, but it updates the two sublattices alternately in place. In the present notation it is one of the two sublattices of a spacetime diagram of 250R [26], which is the conjugate of 95R under the cellwise complement: it is the half system of the class of 95, whose automorphism group is determined in [31]. It is not the second-order automaton F_{54} studied here, which has the extra term from the previous row; its solitons move at speed ± 1, while in F_{54} the structures that carry the automorphisms are the stationary walls.

Organisation. Section 2 fixes notation, proves Lemma 2.2, and shows that $\langle \sigma, F_f \rangle \cong \mathbb{Z}^2$ unless f reads only its middle argument. Section 3 describes the SAT encoding of the commutation condition, the invertibility test and the verification. Section 4 treats rule 54: it proves Theorem 1.3, and it states and proves Theorems 4.2 and 4.6. Section 5 proves Theorem 1.1 and Corollary 1.2, including the class of 133. Section 6 treats walls with hidden bits, rules 59 and 27, the time-periodic walls of rules 127, 179 and 223, the rotating walls of rules 219 and 126, and blocking words. Section 7 treats the linear rules 90 and 150 (Theorem 7.2, Proposition 7.4) and proves that $(\bigoplus_m S_m)^2$ embeds in the automorphism groups of 90R and 165R. Section 8 proves Theorem 1.4 and treats rule 255. Section 9 lists open questions. The appendix describes the computations.

2. Preliminaries

Throughout, $\mathcal{A} = \{0,1\}^2$, and a cell $(x,y) \in \mathcal{A}$ is also written as the integer $2x+y \in \{0,1,2,3\}$; thus $3 = (1,1)$ and $0 = (0,0)$. The left shift is $\sigma(c)_i = c_{i+1}$. A *block map of radius* R is a map $g\colon \mathcal{A}^{\mathbb{Z}} \to \mathcal{A}^{\mathbb{Z}}$ of the form $g(c)_i = \gamma(c_{i-R}, \dots, c_{i+R})$ for a local rule $\gamma\colon \mathcal{A}^{2R+1} \to \mathcal{A}$; we identify g with the lookup table of γ, which has 4^{2R+1} entries of two bits each. A block map of radius R is also a block map of every larger radius. By Curtis–Hedlund–Lyndon, $\mathrm{End}(\mathcal{A}^{\mathbb{Z}})$ is the monoid of block maps and $\mathrm{Aut}(\mathcal{A}^{\mathbb{Z}})$ the group of invertible ones. We write

$$\mathcal{C}_R(F) = \{g \text{ a block map of radius} \le R : \ g \circ F = F \circ g\}, \qquad \mathcal{C}(F) = \bigcup_R \mathcal{C}_R(F),$$

for the centralizer of F in $\mathrm{End}(\mathcal{A}^{\mathbb{Z}})$, and $\mathcal{C}_R^{\times}(F)$, $\mathcal{C}^{\times}(F)$ for the invertible elements. For a configuration c and integers $i < j$ we write $c_{[i,j)} = c_i c_{i+1} \cdots c_{j-1}$, and $c_{[i,j]} = c_{[i,j+1)}$.

For an elementary rule f, identified with its Wolfram number $\sum_{a,b,c} f(a,b,c)2^{4a+2b+c}$, the automaton F_f is defined by (1). It has radius 1, and F_f^b has radius at most $|b|$, so $\sigma^a F_f^b \in \mathcal{C}_{|a|+|b|}(F_f)$.

Definition 2.1. The *spacetime subshift* of a reversible cellular automaton F on $\mathcal{A}^{\mathbb{Z}}$ is

$$X_F = \left\{(F^n(c))_{n\in\mathbb{Z}} :\ c \in \mathcal{A}^{\mathbb{Z}}\right\} \subset \mathcal{A}^{\mathbb{Z}^2},$$

with the $\mathbb{Z}^2$-action generated by the horizontal shift σ and the vertical shift τ, $(\tau z)_n = z_{n+1}$. An *automorphism* of X_F is a homeomorphism of X_F commuting with both.

Lemma 2.2. *For reversible F the map $c \mapsto (F^n(c))_{n\in\mathbb{Z}}$ is a homeomorphism $\mathcal{A}^{\mathbb{Z}} \to X_F$ conjugating (σ, F) to (σ, τ), and it induces an isomorphism*

$$\mathrm{Aut}(X_F) \ \cong\ \mathcal{C}^{\times}(F) = \{g \in \mathrm{Aut}(\mathcal{A}^{\mathbb{Z}}) :\ gF = Fg\}.$$

Under this isomorphism $\langle\sigma,\tau\rangle$ corresponds to $\langle\sigma, F\rangle$.

Proof. The map is continuous, injective (the 0-th row recovers c), and surjective onto X_F by definition, hence a homeomorphism of compact spaces; it carries σ to σ and F to τ since $F^n(F(c)) = F^{n+1}(c)$. A homeomorphism of $\mathcal{A}^{\mathbb{Z}}$ commuting with σ is a block map by Curtis–Hedlund–Lyndon, and commuting with τ means commuting with F. □

So determining $\mathrm{Aut}(X_F)$ means determining which invertible block maps commute with F. The non-invertible elements of $\mathcal{C}(F)$ are the endomorphisms of X_F that are not automorphisms; the constant map onto a fixed point of F is always one. For every rule with $f(0,0,0) = f(1,1,1) = 0$, which includes 54 and 90, both $(0,0)^{\mathbb{Z}}$ and $(1,1)^{\mathbb{Z}}$ are fixed by F_f; for rule 150, which has $f(1,1,1) = 1$, only $(0,0)^{\mathbb{Z}}$ is.

Conjugacies. Let $\mathrm{rev}(c)_i = c_{-i}$ be the reflection and $\kappa(x,y) = (1-x, 1-y)$ the cellwise complement. By (1), $\mathrm{rev}\, F_f\, \mathrm{rev} = F_{f'}$ with $f'(a,b,c) = f(c,b,a)$, and $\kappa F_f \kappa = F_{f''}$ with $f''(a,b,c) = f(1-a, 1-b, 1-c)$. Both rev and κ are involutions that act cellwise up to reflection, so conjugation by them maps $\mathcal{C}_R(F_f)$ bijectively onto $\mathcal{C}_R(F_{f'})$, respectively $\mathcal{C}_R(F_{f''})$, preserves invertibility and maps $\langle\sigma, F_f\rangle$ onto the corresponding group. So $\mathrm{Aut}(X_{F_f})$, and every statement below about block maps of bounded radius, depends only on the class of f under these two operations. There are 88 classes [18], [19, Table I]; one representative of each appears in Table 1. Wolfram's complement $1 \oplus f(1-a, 1-b, 1-c)$ is not a conjugacy of second-order automata, so these are not the classes of the elementary rules [19, p. 377 and footnote 3].

2.1. **The group generated by the shift and the automaton.** The automorphisms constructed below lie outside $\langle\sigma, F_f\rangle$. To make statements about the group they generate together with σ and F_f we need the structure of $\langle\sigma, F_f\rangle$ itself. For $c \in \mathcal{A}^{\mathbb{Z}}$ let $Y_c(t,i)$ be the y-bit of $F^t(c)_i$; then the x-bit of $F^t(c)_i$ is $Y_c(t-1,i)$, and by (1)

$$Y(t+1,i) \oplus Y(t-1,i) = f\big(Y(t,i-1), Y(t,i), Y(t,i+1)\big) \qquad \text{for all } (t,i).$$

A block map g *depends on* a bit of the cell c_i (for the output bit $y(g(c)_0)$, say) if two configurations that differ only in that bit have different outputs. For a nonconstant block map g let $r^+(g)$ be the largest and $-r^-(g)$ the smallest i such that $g(c)_0$ depends on c_i. Then $r^{\pm}(\sigma^a g) = r^{\pm}(g) \pm a$, and $r^{\pm}(F_f^n) \le |n|$ because $F_f^n(c)_0$ depends only on $c_{[-|n|,|n|]}$.

Lemma 2.3 (free staircases)**.** *Let f be any elementary rule and $n \ge 1$. For every assignment of bits to the $2n$ points $(k, n-k-2)$ and $(k, n-k-1)$, $0 \le k < n$, there is a configuration c whose diagram Y_c takes the assigned values at these points.*

Proof. The points lie on the two diagonals $d_m = \{(t,i) : t+i = m\}$ with $m = n-2$ and $m = n-1$. Put $K = 2n+2$, $\tau(m) = m-n+1+K$, and let Ω be the set of (t,i) with $n-1-K \le t+i \le n-1$ and $-1 \le t \le \tau(t+i)$; so $\Omega \cap d_m = \{(t, m-t) : -1 \le t \le \tau(m)\}$ for $n-1-K \le m \le n-1$, and $\tau(m+1) = \tau(m)+1$. We define $Y\colon \Omega \to \{0,1\}$, one diagonal at a time, from right to left. On d_{n-1} and d_{n-2} let Y take the assigned values at the $2n$ points, which lie in Ω because $k \le n-1 < \tau(n-2)$, and the value 0 elsewhere. For $m = n-3, n-4, \dots, n-1-K$ in turn put $Y(\tau(m), m-\tau(m)) = 0$ and then, for $t = \tau(m), \tau(m)-1, \dots, 0$,

$$(2)\quad Y(t-1, m-t+1) = Y(t+1, m-t+1) \oplus f\big(Y(t,m-t),\, Y(t,m-t+1),\, Y(t,m-t+2)\big).$$

The point $(t, m-t)$ of d_m has just been defined, and $(t, m-t+1) \in d_{m+1}$, $(t+1, m-t+1)$ and $(t, m-t+2) \in d_{m+2}$ have been defined before, because $t \le \tau(m) < \tau(m+1)$ and $t+1 \le \tau(m+2)$. So every point of Ω receives a value, and (2) is the spacetime relation at the point $(t, m-t+1)$.

Conversely, suppose that the five points $(t \pm 1, i)$, $(t, i \pm 1)$ of the relation at (t,i) all lie in Ω, and put $m = t+i-1$. Then $(t, i-1) \in d_m$ and $(t, i+1) \in d_{m+2}$ give $n-1-K \le m \le n-3$ and $t \le \tau(m)$, and $(t-1, i) \in \Omega$ gives $t \ge 0$. So the relation at (t,i) is (2) for this m and t, and it holds. Thus Y satisfies every instance of the spacetime relation whose five points lie in Ω.

Let c be any configuration with $c_i = (Y(-1,i), Y(0,i))$ for $n-K \le i \le n-1$; these points lie in Ω. We claim that $Y_c(t,i) = Y(t,i)$ for $t \in \{-1, 0\}$ and $n-K \le i \le n-1$, and for $t \ge 1$ and $n-K+t \le i \le n-1-t$. For $t \le 0$ this is the choice of c. For $t \ge 1$, $Y_c(t,i) = Y_c(t-2,i) \oplus f(Y_c(t-1,i-1), Y_c(t-1,i), Y_c(t-1,i+1))$, the four points on the right lie in the ranges already covered for the rows $t-2$ and $t-1$, and the five points of the relation at $(t-1,i)$ lie in Ω (their diagonals are between $t+i-2 \ge n-K$ and $t+i \le n-1$, and their rows between $t-2 \ge -1$ and $t \le \tau(t+i)$, the last since $i \ge n-K+t > n-1-K$); so Y satisfies the same relation there, and the claim follows by induction on t. The assigned points $(k, n-k-2)$, $(k, n-k-1)$ lie in these ranges: for $k = 0$ in row 0, and for $k \ge 1$ because $n-K+k \le n-k-2$, that is $K \ge 2k+2$. So Y_c takes the assigned values there. □

Theorem 2.4. *Let f be an elementary rule, $F = F_f$ and $n \ge 1$.*

1. *If f depends on its third argument, then $y(F^n(c)_0)$ depends on $y(c_n)$, and $x(F^{-n}(c)_0)$ depends on $x(c_n)$. So $r^+(F^n) = r^+(F^{-n}) = n$.*
2. *If f depends on its first argument, then $y(F^n(c)_0)$ depends on $y(c_{-n})$, and $x(F^{-n}(c)_0)$ depends on $x(c_{-n})$. So $r^-(F^n) = r^-(F^{-n}) = n$.*
3. *The map $(a,b) \mapsto \sigma^a F^b$ is an isomorphism $\mathbb{Z}^2 \to \langle \sigma, F\rangle$ if and only if f is not a function of its middle argument alone.*

Proof. (1) Choose (a^*, b^*) with $f(a^*, b^*, 0) \ne f(a^*, b^*, 1)$. By Lemma 2.3 there is a configuration c with $Y_c(k, n-k-2) = a^*$ and $Y_c(k, n-k-1) = b^*$ for $0 \le k < n$. Let c' be c with $y(c_n)$ complemented, and $\Delta = Y_c \oplus Y_{c'}$. We show by induction on k, $0 \le k \le n$, that $\Delta(k,i) = 0$ for $i < n-k$, $\Delta(k, n-k) = 1$, and $\Delta(k-1, i) = 0$ for $i < n-k+1$. For $k = 0$ this holds because c and c' differ only in $y(c_n)$. For the step from $k < n$ to $k+1$, the relation at (k,i) gives $\Delta(k+1, i) = 0$ for $i < n-k-1$, since all four inputs $\Delta(k-1,i)$, $\Delta(k, i-1)$, $\Delta(k,i)$, $\Delta(k, i+1)$ vanish. At $i = n-k-1$ the inputs $\Delta(k-1, n-k-1)$, $\Delta(k, n-k-2)$ and $\Delta(k, n-k-1)$ vanish, so both diagrams have the values a^*, b^* at $(k, n-k-2)$, $(k, n-k-1)$, while $\Delta(k, n-k) = 1$; hence $\Delta(k+1, n-k-1) = f(a^*, b^*, 0) \oplus f(a^*, b^*, 1) = 1$. For $k = n$ this says $Y_c(n, 0) \ne Y_{c'}(n, 0)$, that is $y(F^n(c)_0) \ne y(F^n(c')_0)$. For F^{-n} use the time reversal $\rho(x,y) = (y,x)$, which satisfies $\rho F \rho = F^{-1}$ for every f [18, p. 2500]: $x(F^{-n}(c)_0) = y(F^n(\rho c)_0)$ depends on $y((\rho c)_n) = x(c_n)$. With $r^+(F^{\pm n}) \le n$ this gives $r^+(F^{\pm n}) = n$.

(2) The reflection conjugates F_f to $F_{f'}$ with $f'(a,b,c) = f(c,b,a)$, and f' depends on its third argument; apply (1) to f'.

(3) Suppose f depends on its third argument. By (1), $r^+(\sigma^a F^b) = a + |b|$ for all (a, b). If $\sigma^a F^b = \mathrm{id}$, then also $\sigma^{-a} F^{-b} = \mathrm{id}$, so $a + |b| = 0 = -a + |b|$ and $a = b = 0$. Since σ and F commute, $(a, b) \mapsto \sigma^a F^b$ is a homomorphism onto $\langle \sigma, F \rangle$, and it is injective. If f depends on its first argument, the same argument with $r^-(\sigma^a F^b) = |b| - a$ applies. If f depends on neither, then $f(a, b, c) = g(b)$, and $F(c)_i = (y_i, x_i \oplus g(y_i))$ applies one permutation of $\mathcal{A}$ to every cell, so F has finite order and the map is not injective. □

Proposition 2.5. *For $f \in \{126, 219\}$ the cellwise complement κ commutes with F_f and $\kappa \notin \langle \sigma, F_f \rangle$, so $\langle \sigma, F_f, \kappa \rangle = \langle \sigma, F_f \rangle \times \langle \kappa \rangle \cong \mathbb{Z}^2 \times \mathbb{Z}/2$.*

Proof. Since $\kappa F_f \kappa = F_{f''}$ with $f''(a, b, c) = f(1 - a, 1 - b, 1 - c)$, κ commutes with F_f if and only if $f(1 - a, 1 - b, 1 - c) = f(a, b, c)$ for all a, b, c, that is, if and only if f is self-conjugate in the sense of [19, Appendix A], which holds for 126 (which vanishes exactly on 000 and 111) and for 219 (which vanishes exactly on 010 and 101). For 126, $f(0, 0, 0) = 0$, so $(0, 0)^{\mathbb{Z}}$ is fixed by σ and F_f, hence by every element of $\langle \sigma, F_f \rangle$, while κ maps it to $(1, 1)^{\mathbb{Z}}$. For 219, the orbit of $(01)^\infty$ under $\langle \sigma, F_{219} \rangle$ consists of the four configurations of period 2 with words $01, 10, 02, 20$, and κ maps $(01)^\infty$ to $(32)^\infty$, which is not among them. Since an element of $\langle \sigma, F_f \rangle$ maps each configuration into its orbit, $\kappa \notin \langle \sigma, F_f \rangle$. Both rules depend on their first and third arguments, so $\langle \sigma, F_f \rangle \cong \mathbb{Z}^2$ by Theorem 2.4(3). As κ is an involution commuting with σ and F_f, and $\langle \sigma, F_f \rangle \cong \mathbb{Z}^2$ is torsion-free, $\langle \kappa \rangle \cap \langle \sigma, F_f \rangle = \{\mathrm{id}\}$, which gives the direct product. □

3. The computation

Several statements below are finite computations of $\mathcal{C}_R(F)$ or $\mathcal{C}_R^\times(F)$ for small R. This section describes the method; Appendix A lists what was computed.

3.1. **Commutation as a finite constraint.** Let g be a block map of radius R with local rule γ, and $F = F_f$. Both $g \circ F$ and $F \circ g$ are block maps of radius $R + 1$, so they are equal if and only if their local rules agree on every window $w \in \mathcal{A}^{2R+3}$. Write $n = 2R + 1$ and, for a window $w = (w_0, \ldots, w_{2R+2})$, let e_1, e_2, e_3 be the three subwords of length n starting at positions $0, 1, 2$ and let $e_L = F(w)$ be the word of length n obtained by applying the local rule of F to w. Then

$$(g \circ F)(w) = \gamma(e_L), \qquad (F \circ g)(w) = \Big(\gamma(e_2)_y,\ \gamma(e_2)_x \oplus f\big(\gamma(e_1)_y, \gamma(e_2)_y, \gamma(e_3)_y\big)\Big),$$

where $\gamma(e)_x, \gamma(e)_y$ are the two output bits. Writing X_e, Y_e for the two unknown bits of table entry e, each window contributes the two constraints

$$X_{e_L} = Y_{e_2}, \qquad Y_{e_L} \oplus X_{e_2} = f(Y_{e_1}, Y_{e_2}, Y_{e_3}). \tag{3}$$

The first is two binary clauses; the second is a constraint on at most five variables, encoded by its truth table in at most sixteen clauses. Altogether there are $2 \cdot 4^{2R+1}$ variables and at most $18 \cdot 4^{2R+3}$ clauses: 17,680 clauses at $R = 1$ and 2.9×10^5 at $R = 2$.

3.2. **Enumeration and verification.** All solutions are enumerated by repeated solving with blocking clauses, using PySAT [9] with CaDiCaL 1.5.3 [2]. Every table returned by the solver is re-checked, independently of the CNF, by evaluating $g \circ F$ and $F \circ g$ on all 4^{2R+3} windows. Each solution is then classified: it is matched against the tables of the maps $\sigma^a F^b$ with $|a| + |b| \le R$, which are built directly by composing local rules, and against the constant maps. The program asserts that every $\sigma^a F^b$ with $|a| + |b| \le R$ occurs among the solutions, which checks the encoding against the only maps known in advance to commute.

3.3. **Invertibility.** A block map is an automorphism of $\mathcal{A}^{\mathbb{Z}}$ if and only if it is injective. Injectivity of a block map of radius R is decided exactly by the pair graph: its vertices are the pairs (u, v) of words of length $2R$, with an edge $(u, v) \to (u', v')$ labelled $(a, b) \in \mathcal{A}^2$ whenever u' and v' are ua and vb with the first letter dropped, and $\gamma(ua) = \gamma(vb)$. Then g fails to be injective if and only if a bi-infinite path passes through an off-diagonal vertex, which is decided by iteratively deleting vertices without a predecessor or without a successor. The test was implemented twice, in pure Python and vectorised, and the two implementations were checked to agree at $R = 1$; at $R \le 2$ every map found injective was also checked to be injective on all spatially periodic configurations of period at most $2R + 4$.

When $\mathcal{C}_R(F)$ is too large to enumerate, only its invertible part is enumerated. An invertible block map is surjective, and a surjective block map of radius R is *balanced* [7, Theorem 5.4]: each output symbol occurs exactly 4^{2R} times in the table. Balance is added to the CNF as four cardinality constraints, the balanced commuting maps are enumerated, and each is submitted to the pair-graph test. Balance is necessary and the pair-graph test is exact, so this yields $\mathcal{C}_R^{\times}(F)$ exactly.

3.4. **Linear rules.** When f is linear over $\mathbb{F}_2$, the constraints (3) are homogeneous linear equations in the table bits and $\mathcal{C}_R(F)$ is an $\mathbb{F}_2$-subspace of table space. Its dimension is computed by Gaussian elimination: the equalities $X_{e_L} = Y_{e_2}$ are merged by union–find, which reduces the $2 \cdot 4^{2R+1}$ bits to 4^{2R} classes, and the remaining equations are reduced by bit-packed elimination. For $R \le 2$ the result was cross-checked against the SAT enumeration of all $2^{\dim}$ solutions.

4. Rule 54: walls

Rule 54 is $f(a, b, c) = b \oplus (a \vee c)$; its values with $b = 1$ are $f(0, 1, 0) = 1$ and $f(a, 1, c) = 0$ whenever $a \vee c = 1$. We need only

$$f(\,\cdot\,, 1, 1) = f(1, 1, \,\cdot\,) = 0, \qquad f(1, 0, \,\cdot\,) = f(\,\cdot\,, 0, 1) = 1. \tag{4}$$

4.1. **Walls and segments.** Call a pair of adjacent cells $(c_i, c_{i+1}) = (3, 3)$ a *wall*; it is the wall (iv) of Takesue [19, Fig. 2, Table III], whose list for this wall contains 54R. A *segment* of a configuration c is a maximal interval $[i, j]$ such that $c_{i-2} = c_{i-1} = 3$, $c_{j+1} = c_{j+2} = 3$, $c_i \ne 3$, $c_j \ne 3$, and c contains no wall inside $[i, j]$; its *content* is the word $c_i \cdots c_j \in \mathcal{A}^L$, $L = j - i + 1$, and $S_L \subset \mathcal{A}^L$ denotes the set of words that can occur as contents, namely those with first and last letter different from 3 and without 33 as a factor. Note that a segment may contain isolated 3's.

Lemma 4.1. *Let $F = F_{54}$.*

(i) *$F(c)$ has a wall at $(i, i + 1)$ if and only if c does. The same holds for F^{-1}.*

(ii) *If $[i, j]$ is a segment of c then $[i, j]$ is a segment of $F(c)$, and the content of $F(c)$ on $[i, j]$ depends only on the content of c on $[i, j]$. Writing $F_L \colon S_L \to S_L$ for the induced map, F_L is a permutation.*

(iii) *F_1 is the 3-cycle $\pi \colon 0 \mapsto 1 \mapsto 2 \mapsto 0$ on $S_1 = \{0, 1, 2\}$.*

Proof. (i) If $c_i = c_{i+1} = 3$ then $y_i = y_{i+1} = 1$ and $x_i = x_{i+1} = 1$, so by (1) and (4), $F(c)_i = (1, 1 \oplus f(y_{i-1}, 1, 1)) = 3$ and $F(c)_{i+1} = (1, 1 \oplus f(1, 1, y_{i+2})) = 3$. Conversely, if $F(c)_i = F(c)_{i+1} = 3$ then $y_i = y_{i+1} = 1$, hence $f(y_{i-1}, 1, 1) = f(1, 1, y_{i+2}) = 0$, and the second coordinates force $x_i = x_{i+1} = 1$. The statement for F^{-1} follows by applying the forward statement to $F^{-1}(c)$.

(ii) By (i) the walls of $F(c)$ are those of c, so the maximal wall-free intervals between walls are the same, and the condition $c_i \ne 3$, $c_j \ne 3$ holds for $F(c)$ as well, since otherwise $F(c)$ would have a wall at $(i - 1, i)$ or $(j, j + 1)$ that c lacks. The new value at a position $k \in [i, j]$ depends on c_{k-1}, c_k, c_{k+1}, and the only positions outside $[i, j]$ involved are $i - 1$ and $j + 1$, where c equals

3 regardless of the configuration; so the content of $F(c)$ on $[i,j]$ is a function of the content of c on $[i,j]$. This function maps S_L into S_L by the first part, and it is injective since F^{-1} induces its inverse by the same argument.

(iii) For a segment $3\,s\,3$ with $s=(x,y)\neq 3$, (1) gives $F(c)$ at that position $=(y,x\oplus f(1,y,1))=(y,x\oplus y\oplus 1)$, which sends $0\mapsto 1\mapsto 2\mapsto 0$. □

4.2. **Relabelling automorphisms.**

Theorem 4.2. *Let $F=F_{54}$. The pattern 33 is a wall: for every configuration c, $F(c)$ has 33 at positions $(i,i+1)$ if and only if c does, and F acts on each maximal wall-free segment of c as a function of that segment alone. For $L\geq 1$ let S_L be the set of admissible contents of a walled segment of length L, let $F_L\colon S_L\to S_L$ be the induced permutation and C_L its centralizer in $\mathrm{Sym}(S_L)$. Then there is an injective homomorphism*

$$\Phi\colon \bigoplus_{L\geq 1} C_L \longrightarrow \mathrm{Aut}(X_F),$$

and $\Phi(\varphi)$ has radius at most $L+1$ for $\varphi\in C_L$. Each C_L contains F_L, which has order $o_L\geq 2$, and $C_2\cong(\mathbb{Z}/3)^3\rtimes S_3$ is nonabelian. Consequently $\mathrm{Aut}(X_{F_{54}})$ contains an infinitely generated subgroup, so it is not virtually $\mathbb{Z}^d$ for any d, and it is not abelian.

Definition 4.3. For $L\geq 1$ and a permutation φ of S_L, let $\Phi_L(\varphi)\colon \mathcal{A}^{\mathbb{Z}}\to\mathcal{A}^{\mathbb{Z}}$ be the map that replaces the content s of every segment of length exactly L by $\varphi(s)$ and leaves every other cell unchanged.

Whether the cell at position k belongs to a segment of length L, and if so which content and which position in it, is determined by $c_{k-L-1},\dots,c_{k+L+1}$: one needs at most $L-1$ cells on one side to reach the end of the segment and two more to see the wall. So $\Phi_L(\varphi)$ is a block map of radius at most $L+1$.

Proof of Theorem 4.2. Let $\varphi\in C_L$, the centralizer of F_L in $\mathrm{Sym}(S_L)$, and $g=\Phi_L(\varphi)$. Since $\varphi(s)\in S_L$ has no wall inside and does not begin or end with 3, the walls of $g(c)$ are the walls of c, so $g(c)$ has the same segments as c, and $\Phi_L(\varphi^{-1})$ inverts g. Hence g is an automorphism of $\mathcal{A}^{\mathbb{Z}}$ and Φ_L is an injective homomorphism $\mathrm{Sym}(S_L)\to\mathrm{Aut}(\mathcal{A}^{\mathbb{Z}})$; injectivity is seen on a configuration consisting of one wall and one segment of length L repeated periodically.

To see that g commutes with F, compare $F(g(c))$ and $g(F(c))$ cell by cell. Both configurations have the walls of c. At a position inside a segment of length L with content s, the first is $F_L(\varphi(s))$ and the second is $\varphi(F_L(s))$ at the corresponding position, by Lemma 4.1(ii) applied to $g(c)$ and to c respectively, and these agree because φ commutes with F_L. At a position k inside a segment of another length, or inside a wall, or not in any segment, g fixes c_k and $F(c)_k$; moreover $F(c)_k$ and $F(g(c))_k$ agree, because the only cells where $g(c)$ differs from c are inside segments of length L, and such a cell is adjacent to k only if k is a wall cell, whose image under F is 3 regardless of its neighbours by (4).

For $L\neq L'$ the maps in $\Phi_L(C_L)$ and $\Phi_{L'}(C_{L'})$ act on disjoint sets of cells and commute, so the Φ_L assemble to a homomorphism Φ on the restricted direct sum $\bigoplus_L C_L$. It is injective: suppose $\Phi(\varphi_{L_1},\dots,\varphi_{L_m})=\mathrm{id}$ with $\varphi_{L_i}\in C_{L_i}$. Fix i and $s\in S_{L_i}$, and let c be the periodic configuration $(3\,3\,s)^\infty$. Every segment of c has length L_i, so the components φ_{L_j} with $j\neq i$ act on c as the identity, and $\Phi(\varphi_{L_1},\dots,\varphi_{L_m})(c)=\Phi_{L_i}(\varphi_{L_i})(c)=(3\,3\,\varphi_{L_i}(s))^\infty$. This equals c only if $\varphi_{L_i}(s)=s$; as s was arbitrary, $\varphi_{L_i}=\mathrm{id}$. Every $\Phi_L(\varphi)$ commutes with σ, being a block map.

Finally, $F_L\in C_L$, and $F_L\neq\mathrm{id}$ because the content 0^L is not fixed: by (4) its first cell becomes $(0,0\oplus f(1,0,0))=(0,1)$. So $o_L\geq 2$ and $\Phi(\bigoplus_L\langle F_L\rangle)\cong\bigoplus_L\mathbb{Z}/o_L$ is an infinite direct sum of nontrivial finite cyclic groups; it is not finitely generated, since any finite set of elements lies

in finitely many summands and generates a finite group. Every subgroup of a virtually $\mathbb{Z}^d$ group is finitely generated, since a subgroup H meets the finite-index copy of $\mathbb{Z}^d$ in a finitely generated group of finite index in H. Hence $\mathrm{Aut}(X_F)$ is not virtually $\mathbb{Z}^d$ for any d. For $L = 2$, both cells of a segment are adjacent to a wall, and by (4) each evolves as $(x, y) \mapsto (y, x \oplus y \oplus 1)$ independently of the other, so $F_2 = \pi \times \pi$ on $S_2 = \{0, 1, 2\}^2$, a product of three 3-cycles, and $C_2 \cong \mathbb{Z}/3 \wr S_3 \cong (\mathbb{Z}/3)^3 \rtimes S_3$, which is nonabelian. So $\mathrm{Aut}(X_F)$ is not abelian, and in particular is not $\langle \sigma, F \rangle$. □

The element $D = \Phi_1(F_1) = \Phi_1(\pi)$ is the radius-2 block map that sends each window $3\,3\,s\,3\,3$ with $s \neq 3$ to $\pi(s)$ and acts as the identity on every other window. Its table differs from that of the identity on exactly 3 of the 1024 entries. The orders o_L of F_L for $L = 1, \ldots, 7$ are $3, 3, 6, 90, 2520, 102702600$ and 4354384834800; the content 0^L has period $3, 3, 6, 9, 18, 21, 18, 21, 30, 33$ for $L = 1, \ldots, 10$. More automorphisms come from the rest of C_L: for $L = 2$, S_2 has 9 elements on which F_2 acts as three 3-cycles, so $C_2 \cong (\mathbb{Z}/3)^3 \rtimes S_3$ has order 162, and $\Phi_2(C_2)$ consists of 162 automorphisms of radius 3, among them the elements of order 2 that swap two of the three cycles.

4.3. **The census at radius 2.**

Theorem 4.4. *$\mathcal{C}_2^{\times}(F_{54})$ has exactly 23 elements: the 13 maps $\sigma^a F^b$ with $|a| + |b| \leq 2$ and the 10 maps $F^b D^k$ with $|b| \leq 2$ and $k \in \{1, 2\}$. At radius 1, $\mathcal{C}_1(F_{54})$ has 13 elements: the 5 maps $\sigma^a F^b$, the 2 constants, and 6 non-injective maps $g_{\pm,k}$, $k = 0, 1, 2$, where $g_{+,k}(c)_i = \pi^k(c_{i-1})$ if $(c_i, c_{i+1}) = (3, 3)$ and $g_{+,k}(c)_i = 3$ otherwise, and $g_{-,k}$ is the mirror image. These record, on each wall, the cell just outside it; they commute with F because by (4) the cell adjacent to a wall evolves by π whatever its other neighbour is.*

Proof. The radius-1 centralizer was enumerated completely; the 13 tables are listed in the data accompanying the paper, and each can be checked by hand against (3). At radius 2 the full centralizer exceeded the enumeration cap of 2×10^4 solutions (it contains, for instance, the compositions of the $g_{\pm,k}$ with the maps Φ_L and many other non-injective endomorphisms), so the balanced commuting maps were enumerated instead: there are exactly 23, every one of them passed the pair-graph test, 13 matched the tables of $\sigma^a F^b$, and each of the remaining 10 differs from the table of some F^b on exactly the three windows $3\,3\,s\,3\,3$, $s \in \{0, 1, 2\}$, where it applies π^k to the value that F^b would produce. That is the table of $F^b D^k = D^k F^b$. □

The count is what the wall mechanism predicts. Maps $\Phi_L(\varphi)$ with $L \geq 2$ have radius at least 3, and $\sigma^a D^k$ with $a \neq 0$ has radius 3, so the only elements of $\langle \sigma, F \rangle \cdot \Phi(\bigoplus_L C_L)$ of radius at most 2 are the $13 + 10$ listed. At radius 3 the wall relabellings stop being the whole story.

4.4. **Every finite group.** Call a maximal run of the symbol 3 of length at least 2 a *wall of length* equal to the run length. By Lemma 4.1, F maps every wall onto itself and every maximal wall-free interval onto itself, so walls and their lengths are invariant. A *car* is a cell $c_i \neq 3$ with walls immediately on both sides, that is $c_{i-2} = c_{i-1} = c_{i+1} = c_{i+2} = 3$. By (4) a car evolves by π, whatever lies beyond the walls. A *train with m cars* is a word

$$W\ s_1\,3\,3\,s_2\,3\,3 \cdots 3\,3\,s_m\ W'$$

in which every $s_j \neq 3$ is a car, consecutive cars are separated by walls of length exactly 2, and the outer walls W, W' have length at least 3. Trains are disjoint, and their positions, lengths and number of cars are invariant under F.

Proof of Theorem 1.3. For $\sigma \in S_m$ let $\Psi_m(\sigma)$ replace the cars $(s_1, \ldots, s_m)$ of every train with exactly m cars by $(s_{\sigma^{-1}(1)}, \ldots, s_{\sigma^{-1}(m)})$ and leave every other cell unchanged. Whether a cell is the j-th car of an m-car train is decided by the cells within distance $3m + 3$, so $\Psi_m(\sigma)$ is a block

map of radius at most $3m+3$. It maps trains to trains with the same cars in other places, so it preserves the set of trains, $\Psi_m(\sigma)\Psi_m(\rho) = \Psi_m(\sigma\rho)$, and $\Psi_m(\sigma^{-1})$ inverts $\Psi_m(\sigma)$. It commutes with F: both $F\Psi_m(\sigma)(c)$ and $\Psi_m(\sigma)F(c)$ carry the walls and non-train cells of $F(c)$, because $\Psi_m(\sigma)$ changes only car cells and a car's neighbours are wall cells whose images are 3 regardless, and in each m-car train both carry $(\pi(s_{\sigma^{-1}(1)}), \dots, \pi(s_{\sigma^{-1}(m)}))$, since F applies the same π to every car. The homomorphism is injective: on the periodic configuration whose period is one m-car train with a single car equal to 1 in position j and all other cars 0, followed by a wall of length 3, the map $\Psi_m(\sigma)$ moves the 1 to position $\sigma(j)$, so σ is determined by its action. Maps for different m act on disjoint sets of cells and commute. Finally every finite group G embeds in $S_{|G|}$ by Cayley's theorem. □

Since $\bigoplus_m S_m$ is not finitely generated and contains nonabelian finite groups, Theorem 1.3 also implies the qualitative conclusions of Theorem 4.2; the relabellings Φ remain of interest because they have small radius and account exactly for the automorphisms of radius 2.

Remark 4.5. A group that contains every finite group is not virtually abelian. Indeed, if A is an abelian normal subgroup of finite index n, then a finite subgroup $H \cong A_m$ with $m \geq 5$ and $m!/2 > n$ meets A in an abelian normal subgroup of H of index at most n, which is impossible because A_m is simple and not abelian. So $\mathrm{Aut}(X_{F_{54}})$ is not virtually abelian, and the same holds for every class of this paper in which $\bigoplus_m S_m$ embeds.

4.5. **Trading wall cells for segment cells.** A singleton segment and a two-cell segment of the form $(\pi(s), s)$ both evolve as a 3-cycle, so the boundary between a long wall and the singleton next to it can be moved by one cell without disturbing the dynamics.

Theorem 4.6. *Let $F = F_{54}$, π the 3-cycle $0 \mapsto 1 \mapsto 2 \mapsto 0$ on $\{0,1,2\}$, and let $\tau\colon \mathcal{A}^{\mathbb{Z}} \to \mathcal{A}^{\mathbb{Z}}$ be the map that, in every occurrence of the pattern $3\,3\,3\,s\,3\,3$ with $s \neq 3$, replaces the third 3 by $\pi(s)$, in every occurrence of $3\,3\,a\,s\,3\,3$ with $s \neq 3$ and $a = \pi(s)$ replaces a by 3, and leaves every other cell unchanged. Then τ is a block map of radius 3, $\tau^2 = \mathrm{id}$, τ commutes with F, τ does not preserve walls, and $\tau D \neq D\tau$. Hence $\tau \in \mathrm{Aut}(X_F)$ lies outside the subgroup generated by σ, F and the wall relabellings, and $\langle \tau, D\rangle$ is a nonabelian subgroup of $\mathrm{Aut}(X_F)$.*

Proof. For a configuration c let $P_1(c)$ be the set of positions i with $c_{i-3} = c_{i-2} = c_{i-1} = 3$, $c_i \neq 3$, $c_{i+1} = c_{i+2} = 3$ (a singleton whose left wall has length at least 3), and $P_2(c)$ the set of positions i with $c_{i-3} = c_{i-2} = 3$, $c_{i-1} = \pi(c_i)$, $c_i \neq 3$, $c_{i+1} = c_{i+2} = 3$ (a two-cell segment of the form $(\pi(s), s)$). By definition $\tau(c)_{i-1} = \pi(c_i)$ for $i \in P_1(c)$, $\tau(c)_{i-1} = 3$ for $i \in P_2(c)$, and $\tau(c)_j = c_j$ at every other position j. Whether j is modified, and how, depends on $c_{j-2}, \dots, c_{j+3}$, so τ is a block map of radius 3. The sets $P_1(c)$ and $P_2(c)$ are disjoint, since $c_{i-1} = 3$ in the first and $c_{i-1} \neq 3$ in the second, and distinct i give distinct modified positions $i-1$.

Involution. Modifying position $i-1$ for $i \in P_1(c) \cup P_2(c)$ changes no other membership. Indeed, a position $j \neq i$ in P_1 or P_2 reads the cells $j-3, \dots, j+2$, and $i-1$ lies among them only for $j \in \{i-3, i-2, i, i+1, i+2\}$; but $i+1$ and $i+2$ are wall cells in both patterns, $i-2$ is a wall cell, and $i-3$ is a wall cell in P_1 and in P_2, while membership of j in $P_1 \cup P_2$ requires $c_j \neq 3$. So the modifications at different positions are independent, $P_1(\tau(c)) = P_2(c)$ and $P_2(\tau(c)) = P_1(c)$, and the two modifications undo each other. Hence $\tau^2 = \mathrm{id}$, and τ is a bijection, so $\tau \in \mathrm{Aut}(\mathcal{A}^{\mathbb{Z}})$.

Commutation. By Lemma 4.1, F preserves walls. A cell adjacent to a wall evolves by $(x, y) \mapsto (y, x \oplus y \oplus 1)$, which is π on $\{0,1,2\}$, whatever its other neighbour is, by (4). For $i \in P_1(c)$ the cell c_i is adjacent to a wall, so $F(c)_i = \pi(c_i) \neq 3$ and $i \in P_1(F(c))$. For $i \in P_2(c)$ both cells $c_{i-1} = \pi(s)$ and $c_i = s$ are adjacent to walls, so $F(c)$ carries $(\pi^2(s), \pi(s)) = (\pi(\pi(s)), \pi(s))$ there and $i \in P_2(F(c))$. Thus $P_1(F(c)) = P_1(c)$ and $P_2(F(c)) = P_2(c)$. We compare $\tau(F(c))$ and $F(\tau(c))$ cell by cell. At $i-1$ with $i \in P_1(c)$: the first is $\pi(F(c)_i) = \pi^2(c_i)$; in $\tau(c)$ the cell $i-1$ holds $\pi(c_i)$ and has the wall $c_{i-3}c_{i-2} = 3\,3$ on its left, so it evolves by π and the second is $\pi^2(c_i)$

as well. At i with $i \in P_1(c)$: both equal $\pi(c_i)$, since c_i is adjacent to the wall on its right in c and in $\tau(c)$. At $i-1$ with $i \in P_2(c)$: the first is 3; in $\tau(c)$ the cell $i-1$ is 3 with 3 3 on its left, hence part of a wall, and stays 3. At i with $i \in P_2(c)$: both equal $\pi(c_i)$. At any other position j, τ changes neither c_j nor $F(c)_j$, and $F(\tau(c))_j = F(c)_j$ because the only cells where $\tau(c)$ differs from c are the positions $i-1$, whose neighbours are i (already treated) and $i-2$, a wall cell, whose image under F is 3 regardless of its right neighbour by (4). So $\tau F = F\tau$.

The rest. τ sends $\cdots 3\,3\,3\,0\,3\,3\cdots$ to $\cdots 3\,3\,1\,0\,3\,3\cdots$, so it does not preserve walls, and therefore lies outside the subgroup generated by σ, F and the wall relabellings, all of which preserve walls. For $c = \cdots 3\,3\,3\,0\,3\,3\,3\cdots$ one has $\tau D(c) = \tau(\cdots 3\,3\,3\,1\,3\,3\,3\cdots) = \cdots 3\,3\,2\,1\,3\,3\,3\cdots$, whereas $\tau(c) = \cdots 3\,3\,1\,0\,3\,3\,3\cdots$ contains no singleton, so $D\tau(c) = \tau(c) \neq \tau D(c)$. □

4.6. **Radius 3.** The enumeration of $\mathcal{C}_3^{\times}(F_{54})$ needs more than the balance filter, because the balanced commuting maps at radius 3 number more than 2×10^4 and almost all of them are not injective. The computation adds to the CNF the injectivity of g on all spatially periodic configurations of period at most 5, and then learns further necessary conditions lazily: whenever the solver returns a non-injective map, a local witness is extracted, either a diamond (two words of length 13 with the same prefix and suffix of length 6 and equal images of length 7, which by Hedlund's theorem no surjective map admits) or a pair of periodic configurations of period at most 10 with equal images, and the clause forbidding that witness is added and shared among all worker processes. Every surviving map is decided by the pair-graph test. This enumeration is not complete, and we give no count of $\mathcal{C}_3^{\times}(F_{54})$ (Appendix A.1). Every automorphism of radius 3 that it has produced differs from some $\sigma^a F^b$, $|a| + |b| \leq 3$, only on windows of length 7 that contain a wall, and most of them, like τ, do not preserve walls. The wall relabellings of Theorem 4.2 and the wall-trading involutions of Theorem 4.6 are two instances of one principle: between two walls, F acts as a finite permutation, and a local, reversible re-encoding of the walls and segments that intertwines these permutations commutes with F.

Remark 4.7. Nothing in Lemma 4.1(i),(ii) uses more of rule 54 than the three values $f(0,1,1) = f(1,1,0) = f(1,1,1) = 0$, so 33 is a wall for the 32 elementary rules with these values, among them rules 0–7, 16–23, 32–39 and 48–55, and each such second-order automaton admits the relabelling automorphisms $\Phi_L(\varphi)$ for φ in the centralizer of its own F_L. Rule 110 has $f(0,1,1) = 1$, rule 30 has $f(0,1,1) = 1$, and rule 90 has $f(0,1,1) = 1$; none of them has 33 as a wall. For rules 110 and 30 the automorphism groups are $\langle \sigma, F \rangle \cong \mathbb{Z}^2$ [28, 29].

5. Walls in general

In this section F is a reversible cellular automaton of radius 1 on a finite alphabet $\mathcal{A}$. A nonempty word W is *fixed in every context* if $F(c)$ contains W at every position at which c contains W.

Lemma 5.1. *Under the hypothesis of Theorem 1.1, F neither creates nor destroys occurrences of W: $F(c)$ contains W at position i if and only if c does.*

Proof. Fix n and let P_n be the finite set of configurations of spatial period n. F maps P_n bijectively to itself, being injective and commuting with σ^n. Write $N(c)$ for the number of positions $i \in \{0, \ldots, n-1\}$ at which c contains W. By hypothesis $N(F(c)) \geq N(c)$ for every $c \in P_n$, and $\sum_{c \in P_n} N(F(c)) = \sum_{c \in P_n} N(c)$ because F permutes P_n. Hence $N(F(c)) = N(c)$, so on P_n the occurrences of W in $F(c)$ are exactly those of c. If some configuration c had an occurrence of W in $F(c)$ at a position i where c has none, the same would hold for the periodic configuration obtained by repeating a long enough window of c around i, since both properties depend only on $c_{i-1}, \ldots, c_{i+|W|}$. □

TABLE 1. Second-order elementary automata with a word W fixed in every context, one rule per conjugacy class, with the shortest such W (cells written 0–3). The words are the fixed walls (i)–(viii) of Takesue [19, Fig. 2, Table III], and the 31 classes are those of the rules listed there for these walls. Classes are taken under reflection and the cellwise complement $\kappa(x, y) = (1 - x, 1 - y)$, which conjugates F_f to $F_{f'}$ with $f'(a, b, c) = f(1 - a, 1 - b, 1 - c)$, as in [19, Table I]; Wolfram's complement $1 \oplus f(1 - a, 1 - b, 1 - c)$ is not a conjugacy of second-order automata [19, p. 377], and for instance 110R and 137R have different numbers of periodic points. The property is constant on each class. For each listed rule $|C| \geq 2$. Of the 57 classes in the last row, only $133 = \{133, 161\}$ has a fixed word of greater length ($W = 0330$, $k = 4$, the wall (ix) of [19]; Section 5.1).

$W = 0$: 0, 4, 12 $W = 3$: 1, 2, 3, 18, 19, 33, 35, 50, 51
$W = 30$: 9, 10, 11, 131, 139 $W = 03$: 129 $W = 030$: 73 $W = 303$: 147 $W = 00$: 36
$W = 33$: 5, 6, 7, 22, 23, 37, 38, 39, 54, 55
no fixed word of length ≤ 3 (57 classes): 13–15, 24–31, 41, 43, 45–47, 57–63, 75, 77, 79, 90, 91, 94, 95, 105, 107, 109, 111, 123, 126, 127, 133, 135, 141, 143, 151, 153, 155, 157, 159, 165, 167, 175, 179, 183, 187, 189, 191, 219, 223, 255

Proof of Theorem 1.1. Let $k = |W|$. A *car* of c is a position i with c containing W at $i - k$ and at $i + 1$ and $c_i \in C$; by the definition of C no occurrence of W covers i. By Lemma 5.1 the occurrences of W in $F(c)$ are those of c, and the cells of each occurrence keep their values, so the neighbours of a car keep their values and the car evolves by a map $\varphi\colon C \to \mathcal{A}$, $\varphi(s) = F(W\, s\, W)$ at the middle position, independent of the rest of c. The image $\varphi(s)$ lies in C, because an occurrence of W in $W\, \varphi(s)\, W$ other than at its ends would be an occurrence created by F. And φ is injective, because F is injective on the periodic configurations $(W\, s)^\infty$, which differ only at their cars. So φ is a permutation of C, and F maps cars to cars. Conversely every car of $F(c)$ is a car of c: its position is flanked by the same occurrences of W, and if the symbol of c there were outside C, then c would contain an occurrence of W covering it, which F would preserve. So cars are invariant, and F acts on every car by the same φ; this uses that the neighbours of a car lie in occurrences of W, whose images under F do not depend on the context, which is where the hypothesis "fixed in every context" enters. A *train with m cars* is a chain of cars $i_1 < \cdots < i_m$ with $i_{j+1} = i_j + k + 1$, such that c contains W at $i_1 - 2k$ and at $i_m + k + 1$ (two copies of W with no car between them at each end), and that is maximal with these properties. Trains are determined by the occurrences of W and the cars, so they are invariant under F. As in the proof of Theorem 1.3, permuting the cars of every m-car train by $\sigma \in S_m$ defines a block map $\Psi_m(\sigma)$ of radius at most $m(k + 1) + 2k + 1$, which is an automorphism commuting with F because F applies the same φ to every car. With two distinct symbols $a, b \in C$, the periodic configuration whose period is one m-car train with cars $(b, \ldots, b, a, b, \ldots, b)$ shows that σ is recovered from $\Psi_m(\sigma)$. The map $\Psi_m(\sigma)$ preserves the occurrences of W, since any occurrence covering a car lies inside $W\, s\, W$ and the new symbol is again in C; hence it preserves cars and trains. Maps for different m act on disjoint cells and commute, and the resulting homomorphism of $\bigoplus_m S_m$ is injective, as one sees by evaluating on periodic configurations $(W\, W\, s_1\, W\, s_2 \cdots W\, s_m)^\infty$ whose only trains have m cars. □

The train definition here, bounded by two adjacent copies of W, differs from the one used for rule 54 in Theorem 1.3, where outer walls of length at least 3 suffice; both definitions work, and the general one is what is needed for words that overlap themselves.

TABLE 2. Spread of a single changed cell after 300 steps, mean over 40 random configurations of length 801, and number of trials in which the difference disappeared.

rule	leftward spread	rightward spread	healed
54R	15	21	0
110R	195 (min 167)	41 (min 22)	0
30R	98 (min 72)	300 (min 299)	0

Proof of Corollary 1.2. For each rule of Table 1 the listed word W is fixed in every context, and the set C of Theorem 1.1 has at least two elements; both are finite checks of the local rule over the 4^2 contexts of W and the four symbols s in $W\,s\,W$. Theorem 1.1 applies, and by the conjugacy invariance of Section 2 it applies to every rule of the 31 classes, 81 rules in all. The class of 133 is Proposition 5.2(1), and the last statement is Proposition 5.2(3). □

The walls also confine the spreading of differences: in random configurations of 54R a single changed cell produces a difference that spreads only a few cells, while in 110R and 30R it spreads in both directions at linear speed (Table 2).

5.1. **The class of 133: a wall of length 4.** Table 1 lists the fixed words of length at most 3. One more class has a fixed word, of length 4. It is the wall (ix) of Takesue, who notes that it is not detected with fewer than four cells [19, Table III and p. 391]. In the next proposition the fixed word of part (1) is his; the automorphisms of parts (1) and (2) and the completeness statement (3) are what is added. The examination in [19, Section 4] covers the words of at most 6 cells, and it is accompanied by the conjecture, stated there without proof, that 4 cells suffice to detect a local conservation law of a second-order elementary automaton. Part (3) concerns fixed words only, and for them it gives a proof for every length.

Proposition 5.2 (the fixed word of 133R: [19, p. 391])**.** *Let $f \in \{133, 161\}$ and $F = F_f$.*

(1) *The word $W = 0330$ is fixed by F_{133} in every context, and $W = 3003$ is fixed by F_{161} in every context. In both cases the set C of Theorem 1.1 is all of $\mathcal{A}$, so $\bigoplus_m S_m$ embeds in $\mathrm{Aut}(X_F)$.*

(2) *For $f = 133$, let D be the block map of radius 4 that replaces the middle cell s of every occurrence of $W\,s\,W$ by $F(W\,s\,W)$ at that cell and fixes every other cell. Then $D \in \mathrm{Aut}(X_F)$, D acts on each such cell as the 4-cycle $0 \mapsto 1 \mapsto 3 \mapsto 2 \mapsto 0$, $D^4 = \mathrm{id} \neq D^2$, and $D \notin \langle \sigma, F \rangle$.*

(3) *Every word fixed by some $F_{f'}$ in every context lies in $\{0,3\}^k$. Among the 57 classes with no such word of length at most 3, the class of 133 is the only one with such a word of any length.*

Proof. (1) A cell of a word fixed in every context keeps its value, so by (1) its two bits agree and $f(y_{j-1}, y_j, y_{j+1}) = 0$ at each of its positions j, the outer neighbour bits being arbitrary. For 133, $f(a,b,c) = 1$ exactly on 000, 010 and 111. The y-row of $W = 0330$ is 0110, and the triples $(u,0,1)$, $(0,1,1)$, $(1,1,0)$ and $(1,0,v)$, $u, v \in \{0,1\}$, all avoid 000, 010 and 111; so W is fixed in all 16 contexts. Since $161 = \kappa(133)$ and κ maps 0330 to 3003, the statement for 161 follows. Finally $W\,s\,W$ contains W only at its two ends for each $s \in \mathcal{A}$, so $C = \mathcal{A}$. Theorem 1.1 applies.

(2) The neighbours of the middle cell s of $W\,s\,W$ have y-bit 0, so it evolves by $(x,y) \mapsto (y, x \oplus f(0,y,0)) = (y, x \oplus 1)$, the 4-cycle $0 \to 1 \to 3 \to 2 \to 0$. So D is the map that applies the car permutation φ of the proof of Theorem 1.1 to every car. As there, every length-4 window covering a car lies in $W\,s\,W$, and $W\,\varphi(s)\,W$ contains W only at its ends, so D preserves the

occurrences of W and the cars, $D^{-1} = D^3$, and D commutes with F because F applies φ to every car and the neighbours of a car lie in occurrences of W, whose images do not depend on the context. Since φ is a 4-cycle, $D^4 = \mathrm{id} \neq D^2$. Suppose $D = \sigma^a F^b$. The configuration $0^{\mathbb{Z}}$ contains no W, so D fixes it; F acts on constant configurations as the 4-cycle $0 \to 1 \to 3 \to 2 \to 0$, and σ fixes them, so $4 \mid b$. On $c = (0330\,0)^\infty$, every fifth cell is a car, so $D(c) = F(c)$, and therefore $\sigma^a F^{b-1} c = c$. But $F^{b-1}c = (0330\,s)^\infty$ with $s = \varphi^{b-1}(0) \neq 0$, as $b - 1 \not\equiv 0 \pmod 4$, and no shift of it equals c. So $D \notin \langle \sigma, F \rangle$.

(3) The first statement is the first sentence of the proof of (1). With w the y-row, a fixed word of length $k \geq 2$ is a word $w \in \{0,1\}^k$ with $f(u, w_0, w_1) = 0$ and $f(w_{k-2}, w_{k-1}, v) = 0$ for $u, v \in \{0,1\}$ and $f = 0$ on its interior triples: a walk through $k-1$ states in a graph on the 4 pairs (w_j, w_{j+1}), from a set of allowed initial pairs to a set of allowed final pairs. If such a walk exists, a shortest one visits each state at most once, so some fixed word has length at most 5. A search of all lengths $k \leq 40$ for all 256 rules therefore decides the question for every length; it finds fixed words for exactly the 81 rules of Table 1 at length at most 3, and, among the other rules, only for 133 and 161, at the lengths 4, 7, 8 and every k with $10 \leq k \leq 40$. A brute-force search over all of $\mathcal{A}^k$, $k \leq 4$, agrees with the reduction to $\{0,3\}^k$. The search is a finite computation, described in Appendix A.2. □

6. Walls with hidden bits

The walls of Sections 4 and 5 are words that F fixes in every context, and the automorphisms of Theorems 1.3 and 1.1 permute the cars that sit between them. A wall need not be fixed for this to work. Walls that are periodic in time and not fixed are in the list of Takesue [19, Section 4]: for 27R and 59R the runs of cells in equal phase given by the local conservation law [19, (4.1)–(4.3)], of period 3; for 179R the pairs of cells (xii) and (xiii) of [19, Fig. 2, Table III], of period 3; for 219R and 223R the pairs (xv), and for 126R and 127R the pairs (xvi), of period 4. The walls of this section are built from these, and we say in each case how. What is added here is the bit that such a wall can carry, the flips, and the embeddings that result. In this section the wall is a word that moves under F, but whose image never depends on the context, except perhaps at cells that do not matter, and which carries one bit that F cannot see. Permuting these hidden bits along chains of walls gives every finite group again. The main example is rule 59R, which has no word of any length fixed in every context (Lemma 6.10) and whose walls are not fixed by any power of F that we have checked; the general mechanism also covers walls that are fixed by a power of F, which we call time-periodic, and these occur for rules 127R, 179R and 223R. A variant of the rule-59 walls with windows of length 6 gives the same conclusion for rule 27R (Theorem 6.20). Finally, for rules 219R and 126R the walls are runs of cells bound by a relation that F preserves, and they rotate through the alphabet in every context; we call them rotating walls (Section 6.6). All seven classes discussed here, 27, 59, 126, 127, 179, 219 and 223, lie in the last row of Table 1: none of them has a word of length at most 3 fixed in every context.

Throughout, F is a reversible cellular automaton of radius 1 on $\mathcal{A}^{\mathbb{Z}}$, and for a configuration c and integers $i < j$ we write $c_{[i,j)} = c_i c_{i+1} \cdots c_{j-1}$.

6.1. **The general mechanism.** Fix $k \geq 3$, a set of words $W \subseteq \mathcal{A}^k$, a position j with $1 \leq j \leq k-2$, and an involution $\iota \colon W \to W$ such that $\iota(w)$ differs from w exactly at position j, for every $w \in W$. For a configuration c let

$$\mathrm{Occ}(c) = \{ i \in \mathbb{Z} : \; c_{[i,i+k)} \in W \}$$

be its set of *occurrences*; for $i \in \mathrm{Occ}(c)$ write $w_i(c) = c_{[i,i+k)}$, and let $\iota_i c$ be the configuration obtained from c by replacing $w_i(c)$ by $\iota(w_i(c))$. Only the cell $i + j$, the *flip cell* of the occurrence,

changes. We consider the following hypotheses. Each is stated at position 0; since F commutes with σ, it then holds at every position.

(W1) If $0 \in \mathrm{Occ}(c)$ then $0 \in \mathrm{Occ}(F(c))$, for every c.
(H1) If $0 \in \mathrm{Occ}(c)$ then $F(\iota_0 c) = \iota_0 F(c)$.
(H2) If $0, i \in \mathrm{Occ}(c)$ with $i \neq 0$, then $i \in \mathrm{Occ}(\iota_0 c)$, and $\iota_0 \iota_i c = \iota_i \iota_0 c$.
(H3) There are $\beta\colon W \to \mathbb{Z}/2$ and $\varepsilon \in \mathbb{Z}/2$ such that $\beta \circ \iota = \beta + 1$ and $\beta(w_0(F(c))) = \beta(w_0(c)) + \varepsilon$ whenever $0 \in \mathrm{Occ}(c)$.
(H3′) There are $\beta\colon W \to \mathbb{Z}/2$, a surjection $\kappa\colon W \to K$ onto a finite set, and maps $\theta\colon K \to K$, $\varepsilon\colon K \to \mathbb{Z}/2$ such that $\kappa \circ \iota = \kappa$, $\beta \circ \iota = \beta + 1$, and whenever $0 \in \mathrm{Occ}(c)$,

$$\kappa(w_0(F(c))) = \theta(\kappa(w_0(c))), \qquad \beta(w_0(F(c))) = \beta(w_0(c)) + \varepsilon(\kappa(w_0(c))).$$

(H4) If $0, i \in \mathrm{Occ}(c)$ with $i \neq 0$, then $\beta(w_i(\iota_0 c)) = \beta(w_i(c))$, and under (H3′) also $\kappa(w_i(\iota_0 c)) = \kappa(w_i(c))$.

We call $\beta(w_i(c))$ the *bit* and $\kappa(w_i(c))$ the *class* of the occurrence i of c. Hypothesis (H3) is the case $|K| = 1$ of (H3′), and we work with (H3′) throughout. In (H2), $\iota_i \iota_0 c$ is defined by the first half of (H2), and $\iota_0 \iota_i c$ by the first half with 0 and i exchanged. The bit is "hidden" in the sense of (H1): flipping it commutes with F, so the rest of the configuration evolves as if it were not there.

Lemma 6.1. *Under* (W1), $\mathrm{Occ}(F(c)) = \mathrm{Occ}(c)$ *and* $\mathrm{Occ}(F^{-1}(c)) = \mathrm{Occ}(c)$ *for every configuration* c.

Proof. This is the proof of Lemma 5.1, with "c contains W at i" read as $i \in \mathrm{Occ}(c)$. On the finite set of configurations of spatial period n, which F permutes, the number $N(c)$ of occurrences in a period satisfies $N(F(c)) \geq N(c)$ by (W1), and summing over the set gives equality, so $\mathrm{Occ}(F(c)) = \mathrm{Occ}(c)$ there. For general c, whether $i \in \mathrm{Occ}(c)$ and whether $i \in \mathrm{Occ}(F(c))$ depend only on $c_{[i-1,i+k+1)}$, so a counterexample would yield a periodic one. □

For c and $I \subseteq \mathrm{Occ}(c)$ define $\iota_I c$ cellwise by

$$(\iota_I c)_{i+j} = \iota(w_i(c))_j \quad (i \in I), \qquad (\iota_I c)_n = c_n \quad (n \notin I + j).$$

This is well defined because distinct occurrences have distinct flip cells. Note that $\iota_I c$ is computed from the words of c, all at once.

Lemma 6.2. *Assume* (W1) *and* (H2)*, and let* $I, J \subseteq \mathrm{Occ}(c)$.

(a) $\mathrm{Occ}(\iota_I c) = \mathrm{Occ}(c)$*; in particular flips neither destroy nor create occurrences.*
(b) $\iota_J \iota_I c = \iota_{I \triangle J} c$*. In particular* $\iota_I \iota_I c = c$.
(c) *If also* (H3′) *and* (H4) *hold, then for every* $i \in \mathrm{Occ}(c)$ *the class of* i *in* $\iota_I c$ *is its class in* c*, and the bit of* i *in* $\iota_I c$ *is its bit in* c *plus* $[i \in I]$.
(d) *If also* (H1) *holds, then* $F(\iota_I c) = \iota_I F(c)$.

Proof. One flip. Let $i \in \mathrm{Occ}(c)$. The word of $\iota_i c$ at i is $\iota(w_i(c))$, since ι changes only position j; hence $\iota_i \iota_i c = c$. By (H2), every $i' \in \mathrm{Occ}(c)$ lies in $\mathrm{Occ}(\iota_i c)$. Conversely, if $i' \in \mathrm{Occ}(\iota_i c)$ with $i' \neq i$, then (H2) applied to the configuration $\iota_i c$, which has occurrences at i and i', shows that $i' \in \mathrm{Occ}(\iota_i \iota_i c) = \mathrm{Occ}(c)$. So $\mathrm{Occ}(\iota_i c) = \mathrm{Occ}(c)$: flips do not create occurrences, because ι is an involution.

Finitely many flips. Let $I = \{i_1, \ldots, i_r\}$ be finite. By the first step every intermediate configuration has occurrence set $\mathrm{Occ}(c)$, so the composition $\iota_{i_1} \circ \cdots \circ \iota_{i_r}(c)$ is defined, and by (H2), applied to intermediate configurations, it does not depend on the order. It equals $\iota_I c$: the cell $i_s + j$ is changed only by ι_{i_s}, and ordering the composition so that ι_{i_s} acts first shows that this cell becomes $\iota(w_{i_s}(c))_j$; cells outside $I + j$ are never changed. This proves (a) for finite I,

and (b) follows, since $\iota_J\iota_I c$ is then a composition of single flips in which those at $I \cap J$ occur twice and cancel. For (c), a single flip at i_s changes the word at i_s to its ι-partner, which has the same class and the other bit, and by (H4), applied to the intermediate configuration, it leaves the bit and class of every other occurrence unchanged. For (d), if $i \in \mathrm{Occ}(y)$ then $F(\iota_i y) = \iota_i F(y)$ by (H1), and $\mathrm{Occ}(F(y)) = \mathrm{Occ}(y)$ by Lemma 6.1; induction on r gives $F(\iota_I c) = \iota_I F(c)$.

Infinite sets. For a finite set U of cells, $\iota_I c$ agrees on U with $\iota_{I_U} c$, where $I_U = \{i \in I : i+j \in U\}$ is finite. Each of (a)–(d) is a statement about finitely many cells at a time (an occurrence reads k cells, a flip reads one window, F reads three cells), so each follows from the finite case applied to suitable finite subsets of I and J. □

Corollary 6.3. *Under* (W1), (H1) *and* (H2), *the map* $\tau_W(c) = \iota_{\mathrm{Occ}(c)} c$, *which flips every occurrence, is an involutive automorphism of* $\mathcal{A}^{\mathbb{Z}}$ *of radius at most* $\max(j, k-1-j)$ *commuting with* F.

Proof. Cell n of $\tau_W(c)$ depends only on $c_{[n-j,n-j+k)}$. By Lemma 6.2(a),(b), $\tau_W(\tau_W(c)) = \iota_{\mathrm{Occ}(c)}\iota_{\mathrm{Occ}(c)} c = c$, and by Lemma 6.2(d) and Lemma 6.1,

$$F(\tau_W(c)) = \iota_{\mathrm{Occ}(c)} F(c) = \iota_{\mathrm{Occ}(F(c))} F(c) = \tau_W(F(c)).$$

□

Definition 6.4. Fix a gap $d \geq 1$ and assume (H3′). A *train of length* m of c is a set $\{i, i+d, \ldots, i+(m-1)d\} \subseteq \mathrm{Occ}(c)$ of occurrences that all have the same class x, such that neither $i-d$ nor $i+md$ is an occurrence of class x. Its *bit vector* is $(b_1, \ldots, b_m)$, where b_r is the bit of $i+(r-1)d$. For $\sigma \in S_m$ let $I_\sigma(c)$ be the set of occurrences $i+(r-1)d$, taken over all trains of length m, with $b_{\sigma^{-1}(r)} \neq b_r$, and define

$$\Psi_m(\sigma)(c) = \iota_{I_\sigma(c)} c.$$

Every occurrence lies in at most one train: the occurrences of a given class x whose positions are congruent modulo d split into maximal arithmetic progressions of step d, and the trains are the finite ones among them. An occurrence that lies in an infinite progression, one-sided or two-sided, belongs to no train; nothing below needs these, since Ψ_m touches only trains of length exactly m. By Lemma 6.2(c), $\Psi_m(\sigma)$ replaces the bit vector b of every train of length m by $b \circ \sigma^{-1} = (b_{\sigma^{-1}(1)}, \ldots, b_{\sigma^{-1}(m)})$ and changes no other bit and no class.

Lemma 6.5. *Assume* (W1), (H1), (H2), (H3′) *and* (H4), *and let* $d \geq 1$. *Then for every* $m \geq 1$:

(i) $\Psi_m(\sigma)$ *is a block map of radius at most* $md + \max(j, k-1-j)$;
(ii) $\Psi_m(\sigma)\Psi_m(\rho) = \Psi_m(\sigma\rho)$ *and* $\Psi_m(\mathrm{id}) = \mathrm{id}$, *so each* $\Psi_m(\sigma)$ *is an automorphism of* $\mathcal{A}^{\mathbb{Z}}$;
(iii) $F\Psi_m(\sigma) = \Psi_m(\sigma)F$, *so* $\Psi_m(\sigma) \in \mathrm{Aut}(X_F)$ *by Lemma 2.2;*
(iv) $\Psi_m(\sigma)$ *and* $\Psi_{m'}(\rho)$ *commute for* $m \neq m'$, *and* $\Psi_{m'}(\rho)$ *does not change the bits of the trains of length* m.

Proof. (i) Cell n can change only if $n - j =: i$ is an occurrence. Whether i is the r-th member of a train of length exactly m, and the bits of that train, are decided by the words at $i + td$, $-m \leq t \leq m$, and these lie in $c_{[n-j-md,\, n-j+md+k)}$. The construction is the same at every position, so $\Psi_m(\sigma)$ commutes with σ.

We show that trains are invariant. By Lemma 6.2(a),(c), $\iota_I c$ has the same occurrences, with the same classes, as c; so it has the same trains. For F: the map θ is a permutation of K. Indeed, given $x \in K$ choose c with $w_0(c)$ of class x; then $0 \in \mathrm{Occ}(F^{-1}(c))$ by Lemma 6.1, and θ maps the class of 0 in $F^{-1}(c)$ to x, so θ is onto, hence bijective. Since $\mathrm{Occ}(F(c)) = \mathrm{Occ}(c)$ and every class is replaced by its image under the injective map θ, two occurrences have the same class in $F(c)$ if and only if they do in c. Hence $F(c)$ has the same trains as c, and by (H3′) the bit vector

of a train of class x changes from b to $b+\varepsilon(x)(1,\dots,1)$. The same constant is added to every bit of a train, so the set of r with $b_{\sigma^{-1}(r)} \neq b_r$ is unchanged: $I_\sigma(F(c)) = I_\sigma(c)$.

(iii) By Lemma 6.2(d), $F\Psi_m(\sigma)(c) = F(\iota_{I_\sigma(c)}c) = \iota_{I_\sigma(c)}F(c) = \iota_{I_\sigma(F(c))}F(c) = \Psi_m(\sigma)F(c)$.

(ii) Let $c' = \Psi_m(\rho)(c) = \iota_{I_\rho(c)}c$. It has the trains of c, and the bit vector b of each train of length m has become $b\circ\rho^{-1}$. Then $\Psi_m(\sigma)(c') = \iota_{I'}c'$, where I' consists of the members at which $b\circ\rho^{-1}\circ\sigma^{-1}$ differs from $b\circ\rho^{-1}$. By Lemma 6.2(b), $\Psi_m(\sigma)\Psi_m(\rho)(c) = \iota_{I_\rho(c)\triangle I'}c$, and $I_\rho(c)\triangle I'$ is the set of members at which $b\circ(\sigma\rho)^{-1}$ differs from b, which is $I_{\sigma\rho}(c)$. Clearly $I_{\mathrm{id}}(c) = \emptyset$. So $\Psi_m(\sigma^{-1})$ inverts $\Psi_m(\sigma)$, and an invertible block map is an automorphism of $\mathcal{A}^{\mathbb{Z}}$.

(iv) $\Psi_{m'}(\rho)$ flips only members of trains of length m', so by Lemma 6.2(c) it leaves the trains and the bits of the trains of length m unchanged; hence $I_\sigma(\Psi_{m'}(\rho)(c)) = I_\sigma(c)$, and symmetrically. By Lemma 6.2(b) both composites equal $\iota_{I_\sigma(c)\cup I_\rho(c)}c$. □

Theorem 6.6. *Let F be a reversible cellular automaton of radius 1 on $\mathcal{A}^{\mathbb{Z}}$, and let $W \subsetneq \mathcal{A}^k$, j and ι be as above. Assume* (W1), (H1), (H2), (H4), *and either* (H3) *or* (H3′). *Then for every $d \geq k$ and $m \geq 1$ the maps Ψ_m of Definition 6.4 form an injective homomorphism $S_m \to \mathrm{Aut}(X_F)$ whose image consists of block maps of radius at most $md + \max(j, k-1-j)$; the images for different m commute, and together they give an embedding of $\bigoplus_{m\geq 1} S_m$ in $\mathrm{Aut}(X_F)$. In particular $\mathrm{Aut}(X_F)$ contains every finite group and is not virtually $\mathbb{Z}^n$ for any n.*

Proof. Everything except injectivity is Lemma 6.5, and (iv) there shows that the Ψ_m assemble to a homomorphism Ψ of the restricted direct sum. Let $g = \Psi((\sigma_{m'})_{m'})$ with $\sigma_m \neq \mathrm{id}$ for some m, and choose r with $\sigma_m(r) \neq r$. Pick $w \in W$, a word $z \in \mathcal{A}^k \setminus W$ and a letter $e \in \mathcal{A}$, and let $u_t = \iota(w)$ for $t = r$ and $u_t = w$ for $t \neq r$, $1 \leq t \leq m$. Let c be the configuration of spatial period $(m+1)d$ whose period, starting at position 0, is

$$u_1\, e^{d-k}\, u_2\, e^{d-k} \cdots u_m\, e^{d-k}\, z\, e^{d-k};$$

this uses $d \geq k$. The positions $0, d, \dots, (m-1)d$ are occurrences, all of the class $\kappa(w)$ because $\kappa\circ\iota = \kappa$, and md and $-d$ are not occurrences because $c_{[md,md+k)} = c_{[-d,-d+k)} = z$. So these positions form a train of length m whose bit vector b has b_t equal to $\beta(w) + [t = r]$. Other occurrences, created by the fillers, may form further trains; this does not matter. By Lemma 6.5(iv) and the remark before it, the bit vector of this train in $g(c)$ is $b\circ\sigma_m^{-1}$, which differs from b at $\sigma_m(r)$. Since bits are functions of the configuration, $g(c) \neq c$. So Ψ is injective. Every finite group embeds in some S_m by Cayley's theorem. Finally $\bigoplus_m S_m$ is not finitely generated, and every subgroup of a virtually $\mathbb{Z}^n$ group is finitely generated, as in the proof of Theorem 4.2. □

Remark 6.7. Hypothesis (H4) cannot be dropped from the argument. (H2) guarantees that a flip at 0 preserves an overlapping occurrence at i and commutes with the flip there, but the flip cell j may be another letter of the word at i, so the word at i can move to a different ι-pair, with a different bit or class. Then the bit vectors after $\Psi_m(\rho)$ need not be $b\circ\rho^{-1}$, and both the homomorphism law and the train structure under (H3′) can fail; commutation with F is unaffected, since it uses only $I_\sigma(F(c)) = I_\sigma(c)$. Rule 59 below shows that such overlaps occur: a flip changes the protector of the next wall. There (H4) holds because the bit does not read the protector; it would fail if the class were taken to be the ι-pair of the whole word. Alternatively, (H4) can be avoided by building trains only from occurrences that overlap no other occurrence. That set is preserved by F and by flips, since both preserve Occ, and a flip at such an occurrence changes no other word; the injectivity argument then needs a configuration in which the members of the train are of this kind.

Remark 6.8. The proof uses about $\sigma \in S_m$ only that $b \mapsto b\circ\sigma^{-1}$ is a permutation of $(\mathbb{Z}/2)^m$ commuting with $b \mapsto b + (1,\dots,1)$, which is how F acts on the bit vector of a train when $\varepsilon = 1$.

For every permutation π of $(\mathbb{Z}/2)^m$ commuting with $b \mapsto b + (1, \dots, 1)$, replacing the bit vector b of every train of length m by $\pi(b)$ defines an automorphism $\Psi_m(\pi)$ commuting with F, by the same proof with $I_\pi(c)$ the set of members of trains of length m at which $\pi(b)$ and b differ, and $\pi \mapsto \Psi_m(\pi)$ is a homomorphism. For injectivity the witness must be a train with an arbitrary bit vector, rather than the vector with one odd bit used for S_m. The construction in the proof of Theorem 6.6 gives one: choose each $u_t \in \{w, \iota(w)\}$ freely, so that the train at $0, d, \dots, (m-1)d$ has bit vector b with $b_t = \beta(u_t)$, which is arbitrary. Given $\pi \neq \mathrm{id}$, choose b with $\pi(b) \neq b$; then the bit vector of this train in $\Psi_m(\pi)(c)$ is $\pi(b) \neq b$, so $\Psi_m(\pi)(c) \neq c$. For rule 59 the witness of Theorem 6.14 serves in the same way, with the centres C_t chosen freely. So the centralizer of $b \mapsto b + (1, \dots, 1)$ in $\mathrm{Sym}((\mathbb{Z}/2)^m)$, which contains $(\mathbb{Z}/2)^m \rtimes S_m$, embeds in $\mathrm{Aut}(X_F)$.

Theorem 6.6 generalizes the mechanism of Theorem 1.1, though not its statement, which it does not contain. In Theorem 1.1 the walls are fixed and the permuted information sits in cars between them; here the information is a letter inside the wall, and the rest of the wall moves. The common point is that every cell next to the permuted information has a trajectory that does not depend on the context.

The simplest case is a wall that is a single periodic orbit of F in every context.

Corollary 6.9. *Let $P = 2h$ be even and let $w_0, \dots, w_{P-1} \in \mathcal{A}^k$ be distinct words, indices modulo P, such that:*

(a) *if $c_{[0,k)} = w_s$ then $F(c)_{[0,k)} = w_{s+1}$, for every c and s;*
(b) *w_{s+h} differs from w_s exactly at a position j with $1 \le j \le k-2$, the same for all s;*
(c) *no configuration has occurrences of $W = \{w_0, \dots, w_{P-1}\}$ at two positions at distance δ with $1 \le \delta \le \max(j, k-1-j)$.*

Then $\bigoplus_{m \ge 1} S_m$ embeds in $\mathrm{Aut}(X_F)$. In particular each w_s is fixed by F^P in every context, and the flip-all map τ_W of Corollary 6.3 is an involutive automorphism of radius $\max(j, k-1-j)$.

Proof. Let $\iota(w_s) = w_{s+h} = F^h$ on W. (W1) is (a). For (H1), let $c_{[0,k)} = w_s$. The configurations $\iota_0 c$ and c differ only at cell j, so $F(\iota_0 c)$ and $F(c)$ differ at most at cells $j-1, j, j+1$, which lie in $[0,k)$ because j is interior. On $[0,k)$, $F(\iota_0 c)$ carries w_{s+h+1} by (a), which is what $\iota_0 F(c)$ carries there; outside $[0,k)$ both agree with $F(c)$. So $F(\iota_0 c) = \iota_0 F(c)$. For (H3′) take $K = \mathbb{Z}/h$, $\kappa(w_s) = s \bmod h$, $\beta(w_s) = \lfloor s/h \rfloor$ for $0 \le s < P$, $\theta(x) = x + 1$ and $\varepsilon(x) = [x = h-1]$: then $\kappa \circ \iota = \kappa$, $\beta \circ \iota = \beta + 1$, and passing from w_s to w_{s+1} changes $\lfloor s/h \rfloor$ exactly when $s \equiv h-1 \pmod h$. For (H2) and (H4): if $0, i \in \mathrm{Occ}(c)$ with $i \neq 0$, then $|i| > \max(j, k-1-j)$ by (c), so the flip cell of each of the two occurrences lies outside the window of the other. Hence ι_0 leaves $w_i(c)$ unchanged, the new value it writes at cell j depends only on $w_0(c)$, which ι_i leaves unchanged, and so (H2) and (H4) hold. Finally $W \neq \mathcal{A}^k$ by (c). Theorem 6.6 and Corollary 6.3 apply. □

Condition (b) contains the requirement that w_s and w_{s+h} agree at their end positions 0 and $k-1$; this is sufficient for (H1) but not necessary, since a disagreement at an end letter that the outside neighbour's update ignores would not break (H1). Conditions (a)–(c) are finite checks, but (H2)–(H4) do not follow from (a) and (b) alone, which is why (c) is assumed.

6.2. **Rule 59R: clocks.** Rule 59 is $f(a,b,c) = \neg b \vee (\neg a \wedge c)$. Call a cell i *triggered* in c if $y_{i-1} = 0$ and $y_{i+1} = 1$, that is $c_{i-1} \in \{0, 2\}$ and $c_{i+1} \in \{1, 3\}$. Since $f(a, 0, c) = 1$ and $f(a, 1, c) = [a = 0,\ c = 1]$, (1) gives, for $F = F_{59}$:

$$0 \mapsto 1, \qquad 2 \mapsto 0, \qquad 1 \mapsto \begin{cases} 3 & \text{triggered} \\ 2 & \text{otherwise,} \end{cases} \qquad 3 \mapsto \begin{cases} 2 & \text{triggered} \\ 3 & \text{otherwise.} \end{cases} \tag{5}$$

So every cell is a clock $0 \to 1 \to 2 \to 0$ of period 3, whose y-bit is 1 exactly at phase 1; the state 3 is phase 1 held, a trigger at phase 1 holds the clock, and a later trigger releases it. A cell

can only be triggered while its y-bit is 1, and only by the pattern "left neighbour silent, right neighbour firing". In particular two adjacent cells in the same phase never disturb each other.

Say that a word u of length n is *fixed in every context* if $F(c)_{[0,n)} = u$ for every c with $c_{[0,n)} = u$.

Lemma 6.10. *For $F = F_{59}$, no word of any length is fixed in every context.*

Proof. By (5), a cell with value 0, 1 or 2 changes its value in every context, so a word fixed in every context would be 3^n. Let $c_{[0,n)} = 3^n$, $c_{-1} = 0$ and $c_n = 1$. The left neighbour of cell 0 has $y = 0$, and its right neighbour (the next 3, or the outside cell $c_1 = 1$ when $n = 1$) has $y = 1$. So cell 0 is triggered, and $F(c)_0 = 2$. □

For $p \in \{0,1,2\}$, expressions such as $p+1$ and $p+2$ are taken modulo 3 and denote cells in $\{0,1,2\}$. Let

$$X_A = \{\, a\,p\,C\,(p+1) \;:\; p \in \{0,1,2\},\ a \in \{p,3\},\ C \in \{p+1,3\} \,\},$$
$$X_B = \{\, a\,p\,C\,(p+2) \;:\; p \in \{0,1,2\},\ a \in \{p,3\},\ C \in \{p+2,3\} \,\},$$

two disjoint sets of 12 words of length 4. In an *A-wall* the right side is one phase ahead of the left, in a *B-wall* one phase behind. We say c has an A-wall (B-wall) at i if $c_{[i,i+4)} \in X_A$ ($\in X_B$), and call c_i, c_{i+1}, c_{i+2}, c_{i+3} its *protector* a, its *left cell* $L = p$, its *centre* C and its *right cell* R; the pair (L, C) is its *core*. Write $\delta_A = 1$ and $\delta_B = 2$, so that $R = L + \delta$ and $C \in \{L + \delta, 3\}$.

In this notation the local conservation law of 27R and 59R found by Takesue [19, (4.1), (4.2)] says that the set of pairs $a\,p$ with $p \in \{0,1,2\}$ and $a \in \{p,3\}$ is invariant: such a pair remains one in every context. His wall is a run $a\,p\,p\cdots p$ of at least three cells with $a \in \{p,3\}$, in which every cell but the first passes from p to $p+1$ in every context [19, (4.3)]. A word of $X_A \cup X_B$ consists of two such pairs, the protector with the left cell and the centre with the right cell, whose phases differ by δ. The cycles of the cores, the bit carried by the centre and the flips (Lemmas 6.11–6.13) are not in [19]. For 59R the invariance of the pairs is contained in the proof of Lemma 6.11, in the images of the protector and of the left cell, where it is needed together with the images of the other cells.

Lemma 6.11. *Let $F = F_{59}$ and $X \in \{A, B\}$. If c has an X-wall at i with left cell p, then $F(c)$ has an X-wall at i with left cell $p+1$, and $F(c)_{i+1}, F(c)_{i+2}, F(c)_{i+3}$ depend only on the core of c at i, not on the rest of c. The cores run through the 6-cycles*

$$A: \quad (0,1) \to (1,3) \to (2,3) \to (0,3) \to (1,2) \to (2,0) \to (0,1),$$
$$B: \quad (0,2) \to (1,0) \to (2,1) \to (0,3) \to (1,3) \to (2,3) \to (0,2).$$

Moreover $F(c)$ has an X-wall at i if and only if c does, and the same holds for F^{-1}.

Proof. By (5), cells with value 0 or 2 advance regardless of their neighbours, so only cells with value 1 or 3 need attention. The six cases are:

	(a, L, C, R)	(a', L', C', R')	$C \mapsto C'$	triggered
A, $p = 0$	$(\{0,3\}, 0, \{1,3\}, 1)$	$(\{1,3\}, 1, \{3,2\}, 2)$	$1 \mapsto 3,\ 3 \mapsto 2$	C
A, $p = 1$	$(\{1,3\}, 1, \{2,3\}, 2)$	$(\{2,3\}, 2, \{0,3\}, 0)$	$2 \mapsto 0,\ 3 \mapsto 3$	possibly a
A, $p = 2$	$(\{2,3\}, 2, \{0,3\}, 0)$	$(\{0,3\}, 0, \{1,3\}, 1)$	$0 \mapsto 1,\ 3 \mapsto 3$	none
B, $p = 0$	$(\{0,3\}, 0, \{2,3\}, 2)$	$(\{1,3\}, 1, \{0,3\}, 0)$	$2 \mapsto 0,\ 3 \mapsto 3$	none
B, $p = 1$	$(\{1,3\}, 1, \{0,3\}, 0)$	$(\{2,3\}, 2, \{1,3\}, 1)$	$0 \mapsto 1,\ 3 \mapsto 3$	possibly a
B, $p = 2$	$(\{2,3\}, 2, \{1,3\}, 1)$	$(\{0,3\}, 0, \{2,3\}, 2)$	$1 \mapsto 3,\ 3 \mapsto 2$	C

We justify the entries. If $L = 1$ (case $p = 1$), its left neighbour $a \in \{1,3\}$ has $y = 1$, so L is not triggered and $L' = 2$. If $R = 1$ (cases A, $p = 0$ and B, $p = 2$), its left neighbour $C \in \{1,3\}$ has $y = 1$, so $R' = 2$. The centre, when it equals 1 or 3, is triggered exactly when $y_L = 0$ and $y_R = 1$, that is when $p \neq 1$ and $R = 1$: this happens in the cases A, $p = 0$ and B, $p = 2$

and in no other case (in the other cases $p = 1$, or $R \in \{0, 2\}$). The protector: $a = 0 \mapsto 1$, $a = 2 \mapsto 0$, $a = 1 \mapsto \{2, 3\}$, and $a = 3$ is triggered only if $y_L = 1$, so $a = 3 \mapsto 3$ when $p \in \{0, 2\}$ and $a = 3 \mapsto \{2, 3\}$ when $p = 1$. In every case $a' \in \{p + 1, 3\}$. Thus each row maps the family into itself with p replaced by $p + 1$, and the images of L, C, R are determined by (L, C), since $R = L + \delta$; only a' depends on the context. Reading off the core images gives the two 6-cycles. The last statement follows from Lemma 6.1 applied to $W = X_A$ and to $W = X_B$. □

Only the protector depends on the context, through its own left neighbour. The left cell L of a wall is therefore a column whose whole trajectory, forwards and backwards, is determined by the wall (backwards because the core map is a bijection), and the same holds for C and R. Cells to the left of the wall feel the right side only through the y-sequence of L, and cells to the right feel the left side only through that of R.

At a wall the centre takes one of two values, $C \in \{L + \delta, 3\}$. Let ι exchange them; on each family it is a fixed-point-free involution changing only position $j = 2$. On the core, ι is the shift by three steps of the 6-cycle, so ι acts on the wall as F^3 acts on its core, while leaving every other cell alone. Define

$$\beta_A(a\,L\,C\,R) = [C = 3] + [L = 2], \qquad \beta_B(a\,L\,C\,R) = [C = 3] + [L = 1] \qquad (\mathrm{mod}2).$$

Lemma 6.12. *Let $F = F_{59}$, $X \in \{A, B\}$, and let c have an X-wall at i. Then $F(\iota_i c) = \iota_i F(c)$ and $F^{-1}(\iota_i c) = \iota_i F^{-1}(c)$. Moreover $\beta_X \circ \iota = \beta_X + 1$, and $\beta_X(F(c)_{[i,i+4)}) = \beta_X(c_{[i,i+4)}) + 1$ in every context.*

Proof. The configurations $\iota_i c$ and c differ only at the centre $i + 2$, so $F(\iota_i c)$ and $F(c)$ can differ only at $i + 1$, $i + 2$, $i + 3$. The images of L and R computed in the proof of Lemma 6.11 do not depend on which of the two values C has: L and R read C only through its y-bit, and either $y_C = 1$ for both values (A, $p = 0$; B, $p = 2$), or the cell reading C is $L = 1$ with $y_a = 1$ and is never triggered ($p = 1$), or L and R are in $\{0, 2\}$ and ignore their neighbours. The centre's update is, by the table, a bijection from $\{L + \delta, 3\}$ onto $\{L' + \delta, 3\}$, the centre pair of the image wall. Hence $F(\iota_i c)_{i+2}$ is the element of that pair other than $F(c)_{i+2}$, that is, $(\iota_i F(c))_{i+2}$, and $F(\iota_i c) = \iota_i F(c)$. Applying this to $F^{-1}(c)$, which has an X-wall at i by Lemma 6.11, gives the statement for F^{-1}. The partners under ι are $(0, 1) \leftrightarrow (0, 3)$, $(1, 3) \leftrightarrow (1, 2)$, $(2, 3) \leftrightarrow (2, 0)$ for A and $(0, 2) \leftrightarrow (0, 3)$, $(1, 0) \leftrightarrow (1, 3)$, $(2, 1) \leftrightarrow (2, 3)$ for B, and along the A-cycle β_A takes the values $0, 1, 0, 1, 0, 1$, along the B-cycle β_B takes the values $0, 1, 0, 1, 0, 1$; both statements about β follow, since β_X depends only on the core. □

In spacetime the centre column of an A-wall has y-history $1, 1, 1, 1, 0, 0$ repeated, and ι replaces it by its shift by half a period; the neighbours read that column only at times when the two histories agree. By Lemma 6.12, $F^n(\iota_i c) = \iota_i F^n(c)$ for all $n \in \mathbb{Z}$, so flipping the bit of one wall changes exactly one column of the spacetime diagram, at all times.

Lemma 6.13. *Let $F = F_{59}$ and $W = X_A$, with $k = 4$ and $j = 2$.*

(i) *No configuration has A-walls at i and $i + 1$.*

(ii) *If c has A-walls at 0 and $i \neq 0$, then $\iota_0 c$ has an A-wall at i, $\iota_0 \iota_i c = \iota_i \iota_0 c$, and the bit β_A of the wall at i is the same in c and in $\iota_0 c$.*

Hence (W1), (H1), (H2), (H3) *with $\varepsilon = 1$, and* (H4) *hold for X_A, ι and β_A.*

Proof. (i) Suppose c has A-walls at i and $i + 1$, and let $p = c_{i+1}$. The left cell of the second wall is c_{i+2}, the centre of the first, which must then lie in $\{0, 1, 2\}$; so $c_{i+2} = p + 1$. The protector of the second wall is $c_{i+1} = p$, which must lie in $\{p + 1, 3\}$; impossible.

(ii) The flip ι_0 changes only cell 2, writing the element of $\{c_1 + 1, 3\}$ other than c_2; it reads only c_1 and c_2. The window $[i, i + 4)$ contains cell 2 only if $-1 \leq i \leq 2$, so by (i) only if $i = 2$.

In that case cell 2 is the protector of the wall at 2, whose left cell is $c_3 = c_1 + 1$, so its allowed protectors are $\{c_1 + 1, 3\}$, the same pair; the wall at 2 survives. In all cases the wall at i survives, and its core (c_{i+1}, c_{i+2}) is untouched, since $i + 1 \neq 2$ by (i) and $i + 2 \neq 2$. So its bit is unchanged. The same argument with 0 and i exchanged shows that ι_i does not touch c_1 or c_2; as each flip reads only the core of its own wall, the two flips commute. The hypotheses: (W1) is Lemma 6.11, (H1) and (H3) are Lemma 6.12, (H2) and (H4) are (ii). □

The same holds for walls of both families together; this is Lemma 6.15(i) below. Overlapping walls then occur only at distances 2 and 3, and by computation on all words of length at most 9 both distances occur for all four type pairs AA, AB, BA, BB.

Theorem 6.14. *For every $m \geq 1$ there is an injective homomorphism $\Psi_m \colon S_m \to \mathrm{Aut}(X_{F_{59}})$ whose image consists of block maps of radius at most $2m + 2$, and the images for different m commute. So $\bigoplus_{m \geq 1} S_m$ embeds in $\mathrm{Aut}(X_{F_{59}})$, which therefore contains every finite group and is not virtually $\mathbb{Z}^n$ for any n. The same holds for the rules* 115*,* 206 *and* 220*, which are conjugate to* 59 *by reflection and complement.*

Here a train of length m is a maximal chain of A-walls at gap 2: A-walls at $i, i+2, \ldots, i+2(m-1)$ and none at $i - 2$ or $i + 2m$. Consecutive walls overlap, the centre of each being the protector of the next, so a train looks like a staircase $a\ p\ C_1\ (p+1)\ C_2\ (p+2) \cdots C_m\ (p+m)$. This is not literally an instance of Theorem 6.6, which requires the gap d to be at least the word length $k = 4$, so that the members of a train can be written down independently; at gap 2 they overlap, and the injectivity witness has to be built by hand. (With $d = 4$, Theorem 6.6 does apply to X_A, by Lemma 6.13, with the weaker radius bound $4m + 2$.)

Proof. By Lemma 6.13, the hypotheses of Lemma 6.5 hold for $W = X_A$ with $|K| = 1$ and $\varepsilon = 1$. Applying that lemma with $d = 2$, the maps $\Psi_m(\sigma)$ are automorphisms of radius at most $2m + \max(2, 1) = 2m + 2$, commuting with F, forming homomorphisms of the S_m whose images commute for different m. It remains to prove injectivity, as in the proof of Theorem 6.6 but with a different witness. Fix m, $p \in \{0, 1, 2\}$ and choices $C_t \in \{p + t, 3\}$ for $1 \leq t \leq m$, and let

$$c = \cdots 3\,3\,3\ p\ C_1\ (p+1)\ C_2\ (p+2) \cdots C_m\ (p+m)\ 3\,3\,3 \cdots ,$$

with $c_{-1} = p$, $c_{2t-2} = C_t$ and $c_{2t-1} = p + t$, and all other cells equal to 3. For $1 \leq t \leq m$ the window $c_{[2t-4,2t)}$ is $(c_{2t-4},\ p + t - 1,\ C_t,\ p + t)$, with protector $c_{-2} = 3$ for $t = 1$ and $C_{t-1} \in \{p + t - 1, 3\}$ for $t \geq 2$; so it is an A-wall. There is no A-wall at -4, since $c_{-3} = 3$ is not a possible left cell, and none at $2m - 2$, since $c_{2m+1} = 3$ is not a possible right cell. So the walls at $-2, 0, \ldots, 2m - 4$ form a train of length m, and its bits $\beta_A = [C_t = 3] + [p + t - 1 = 2]$ can be prescribed arbitrarily by the choice of the C_t. (By direct check, for $m \leq 7$ these are the only walls of c of either family, but this is not needed.) Given $(\sigma_{m'})_{m'}$ with $\sigma_m(r) \neq r$, choose the bits to be 1 at r and 0 elsewhere; as in the proof of Theorem 6.6, the bit vector of this train in $\Psi((\sigma_{m'}))(c)$ is $b \circ \sigma_m^{-1} \neq b$. The statement for the conjugate rules follows because conjugation by the reflection or by the cellwise complement induces an isomorphism of the automorphism groups of the spacetime subshifts. □

By Remark 6.8, $\mathrm{Aut}(X_{F_{59}})$ also contains the centralizer of $b \mapsto b + (1, \ldots, 1)$ in $\mathrm{Sym}((\mathbb{Z}/2)^m)$ for every m.

6.3. **The radius-2 census of rule 59R.** For $X \in \{A, B\}$, call an X-wall at i *level* if $c_{i+4} = c_{i+3}$ (the cell after R is in phase with R), and *doubly protected* if $c_i = c_{i+1}$ and $c_{i-1} \in \{c_{i+1}, 3\}$ (the protector is itself protected). For a property Q of X-walls, flipping the centre of every X-wall with property Q, all at once, defines a block map; we write $\tau_{X,=}$, $\tau_{X,\neq}$ for the maps that flip the level and the non-level X-walls, $\lambda_{X,\mathrm{pp}}$, $\lambda_{X,\neg\mathrm{pp}}$ for those that flip the doubly protected and the

other X-walls, and $\tau_X = \tau_{X,=}\tau_{X,\neq} = \lambda_{X,\mathrm{pp}}\lambda_{X,\neg\mathrm{pp}}$ for the flip of all X-walls, which is the map τ_W of Corollary 6.3 for $W = X_A$ or $W = X_B$. The maps $\tau_{X,=}$ and $\tau_{X,\neq}$ have radius 2 and differ from the identity on 12 and 36 entries of the table for $X = A$; the maps λ have radius 3. Let

$$T_R = \langle \tau_{A,=}, \tau_{A,\neq}, \tau_{B,=}, \tau_{B,\neq} \rangle, \qquad T_L = \langle \lambda_{A,\mathrm{pp}}, \lambda_{A,\neg\mathrm{pp}}, \lambda_{B,\mathrm{pp}}, \lambda_{B,\neg\mathrm{pp}} \rangle, \qquad T = T_R T_L.$$

The *class* of an X-wall at i is the triple (X, level or not, doubly protected or not); there are 8 possible classes. The class reads the cells $i-1$, i, $i+1$, $i+3$, $i+4$ (the family is determined by L and R), but not the centre $i+2$. Let $W = X_A \cup X_B$, with $k = 4$, $j = 2$, and ι exchanging the two values of the centre in each family. The flip ι_i of a wall at i writes only cell $i+2$, and reads only cells $i+1$, $i+2$, $i+3$, since the new centre is the other element of $\{L+\delta, 3\}$ and $\delta = R - L$. Let $Y = \{a\,p\,p :\ p \in \{0,1,2\},\ a \in \{p,3\}\}$. An X-wall at i is doubly protected if and only if $c_{[i-1,i+2)} \in Y$, since $L = c_{i+1} \in \{0,1,2\}$. It is level if and only if $c_{[i+2,i+5)} \in Y$, since the centre always lies in $\{R,3\}$.

Lemma 6.15. *Let $F = F_{59}$ and $W = X_A \cup X_B$.*

(i) *No configuration has walls, of either family, at two positions at distance* 1. *The pair (W,ι) satisfies* (W1), (H1) *and* (H2), *so Lemma 6.2(a), (b), (d) applies to it.*
(ii) *If c has walls at i and i', possibly $i = i'$, then the wall at i has the same class in $\iota_{i'}c$ as in c. Hence for every set I of walls of c, the configuration $\iota_I c$ has the same walls as c, with the same classes.*
(iii) *$F(c)$ has a wall at i of class x if and only if c does.*
(iv) *All* 8 *classes occur.*

Proof. (i) Suppose c has walls at i and $i+1$, and let $p = c_{i+1}$ and δ belong to the first wall. The left cell of the second wall is c_{i+2}, the centre of the first, so it lies in $\{0,1,2\}$, and hence $c_{i+2} = p + \delta$. The protector of the second wall is $c_{i+1} = p$, which must lie in $\{p+\delta, 3\}$. This is impossible. (W1) is Lemma 6.11 and (H1) is Lemma 6.12, applied to each family. For (H2), let c have walls at 0 and $i \neq 0$, so $|i| \geq 2$. The cell 2 written by ι_0 lies in the window $[i, i+4)$ only if $-1 \leq i \leq 2$, hence only if $i = 2$. In that case cell 2 is the protector of the wall at 2 and the centre of the wall at 0. The centre pair of the wall at 0 is $\{R_0, 3\} = \{c_3, 3\}$, and c_3 is the left cell of the wall at 2, so this pair is exactly the set of allowed protectors of that wall. In every case the wall at i survives, with its family. The flip ι_i writes cell $i+2$, which is not among the cells 1, 2, 3 read by ι_0, since $|i| \neq 1$ and $i \neq 0$. Symmetrically, ι_0 writes no cell that ι_i reads, so the two flips commute.

(ii) If $i' = i$, the flip changes only the centre, which the class does not read. Otherwise $|i' - i| \geq 2$. The cell $i'+2$ written by $\iota_{i'}$ is one of the cells $i-1$, i, $i+4$ read by the class only if $i' \in \{i-3, i-2, i+2\}$, because $i'+2 \in \{i+1, i+3\}$ would give $|i'-i| = 1$. The wall at i survives by (i), and we check these three cases.

- $i' = i+2$: cell $i+4$ is the centre of the wall at $i+2$, whose left cell is c_{i+3}. Both possible values of that centre differ from its left cell. So the wall at i is not level, before or after the flip.
- $i' = i-2$: the wall at $i-2$ has left cell $c_{i-1} \in \{0,1,2\}$ and right cell $c_{i+1} \neq c_{i-1}$. So $c_{i-1} \notin \{c_{i+1}, 3\}$, and the wall at i is not doubly protected, before or after the flip. The flip writes c_i but not c_{i-1}.
- $i' = i-3$: the flip exchanges c_{i-1}, the centre of the wall at $i-3$, between c_i (its right cell) and 3. If $c_i = c_{i+1}$, both values lie in $\{c_{i+1}, 3\}$, so the wall at i is doubly protected before and after the flip. If $c_i \neq c_{i+1}$, it is doubly protected in neither.

For a set I, the walls and the class of a given wall read finitely many cells, so the second statement follows from the first, Lemma 6.2(a), and induction, as in the proof of Lemma 6.2.

(iii) The family is preserved by Lemma 6.11. By the remark before the lemma, it remains to show that F and F^{-1} preserve the occurrences of Y. By Lemma 6.1 it suffices to check (W1) for Y and F. Let $c_{[0,3)} = a\,p\,p$ with $a \in \{p, 3\}$. Cells 1 and 2 have left neighbours in $\{p, 3\}$. For $p \in \{0, 2\}$ they advance to $p+1$ by (5). For $p = 1$ their left neighbours have $y = 1$, so they are not triggered and also become $2 = p + 1$. The protector $a = p$ becomes $p + 1$, or 3 when $p = 1$ and it is triggered. The protector $a = 3$ can be triggered only if $y_1 = 1$, that is only if $p = 1$, and then it becomes $2 = p + 1$. So $F(c)_{[0,3)} \in Y$.

(iv) Let X have δ, and take a window $c_{[i-1,i+5)} = u\,a\,p\,C\,(p+\delta)\,v$ with $C \in \{p+\delta, 3\}$. With $a = u = p$ the wall at i is doubly protected; with $a = 3$ it is not, since then $c_i \neq c_{i+1}$. With $v = p + \delta$ it is level; with $v = 3$ it is not. For instance 000111, 000113, 030111, 030113 give the four A-classes, and 000222, 000223, 030222, 030223 give the four B-classes, each for the wall at position 1. □

Proposition 6.16. *Let $F = F_{59}$. The 64 elements of T are distinct automorphisms commuting with F and with each other, and each is an involution or the identity; $T_R \cong T_L \cong (\mathbb{Z}/2)^4$, $T_R \cap T_L = \{1, \tau_A, \tau_B, \tau_A\tau_B\}$, and $T \cong (\mathbb{Z}/2)^6$. The group $\mathcal{C}_2^\times(F)$ has exactly 133 elements, and each of them has exactly one expression of one of the following forms:*

(i) *$\sigma^a F^b$ with $|a| + |b| \leq 2$ (13 maps);*
(ii) *$F^b t$ with $|b| \leq 2$ and $1 \neq t \in T_R$ (75 maps);*
(iii) *$\sigma F^b t$ with $|b| \leq 1$ and $1 \neq t \in T_L$ (45 maps).*

Proof. The group T. This part is exact. For a set U of classes, let $I_U(c)$ be the set of walls of c whose class lies in U, and $\varphi_U(c) = \iota_{I_U(c)}c$. This is a block map of radius 3, since cell n changes only if there is a wall at $n-2$, and its class is read from $c_{[n-3,n+3)}$. By Lemma 6.15(ii), $I_V(\iota_J c) = I_V(c)$ for every set J of walls of c. So Lemma 6.2(b) gives

$$\varphi_V\varphi_U(c) = \iota_{I_V(c)}\iota_{I_U(c)}c = \iota_{I_U(c)\triangle I_V(c)}c = \varphi_{U\triangle V}(c),$$

and $U \mapsto \varphi_U$ is a homomorphism from the group of subsets of the 8 classes under $\triangle$, which is $(\mathbb{Z}/2)^8$. In particular each φ_U is an involution or the identity, hence an automorphism, and any two commute. By Lemma 6.15(iii), $I_U(F(c)) = I_U(c)$, so $F\varphi_U = \varphi_U F$ by Lemma 6.2(d). The homomorphism is injective. If $U \neq \emptyset$, pick $x \in U$ and, by Lemma 6.15(iv), a configuration c with a wall of class x; then φ_U changes the centre of that wall, so $\varphi_U \neq \mathrm{id}$.

Write $R_{X,=}$, $R_{X,\neq}$ for the sets of the two level and the two non-level classes of family X, and $L_{X,\mathrm{pp}}$, $L_{X,\neg\mathrm{pp}}$ for the sets of the two doubly protected and the two other classes. Then $\tau_{X,=} = \varphi_{R_{X,=}}$, $\tau_{X,\neq} = \varphi_{R_{X,\neq}}$, $\lambda_{X,\mathrm{pp}} = \varphi_{L_{X,\mathrm{pp}}}$ and $\lambda_{X,\neg\mathrm{pp}} = \varphi_{L_{X,\neg\mathrm{pp}}}$. Hence the map from $(\mathbb{Z}/2)^8$, indexed by the eight generators, onto T that sends s to the product of the generators with exponents s sends s to $\varphi_{U(s)}$. Here the class (X, l, q) lies in $U(s)$ if and only if the exponents of $\tau_{X,l}$ and $\lambda_{X,q}$ have sum 1. So s lies in the kernel if and only if, for each X, the four exponents of $\tau_{X,=}$, $\tau_{X,\neq}$, $\lambda_{X,\mathrm{pp}}$, $\lambda_{X,\neg\mathrm{pp}}$ are equal. The kernel has dimension 2, so $|T| = 2^6 = 64$ and $T \cong (\mathbb{Z}/2)^6$. The four sets $R_{X,l}$ are disjoint and nonempty, so $T_R \cong (\mathbb{Z}/2)^4$, and likewise $T_L \cong (\mathbb{Z}/2)^4$. A set of classes that is a union of sets $R_{X,l}$ and also a union of sets $L_{X,q}$ contains, for each X, either none or all four classes of X. So $T_R \cap T_L = \{\varphi_\emptyset, \varphi_A, \varphi_B, \varphi_{A\cup B}\} = \{1, \tau_A, \tau_B, \tau_A\tau_B\}$, where φ_X flips all X-walls and so equals τ_X. This also gives $\tau_X = \tau_{X,=}\tau_{X,\neq} = \lambda_{X,\mathrm{pp}}\lambda_{X,\neg\mathrm{pp}}$.

The census. This part is a computation. $\mathcal{C}_2^\times(F)$ was enumerated as in Section 3, and every solution passed the pair-graph test. The tables of the maps in (i)–(iii) were computed by exact composition, and each of the 133 tables was matched against them; each matched exactly one, so the three lists are disjoint. In (iii), $\sigma\lambda$ has radius 2 although λ has radius 3, because λ reads three cells to the left of the flip cell and one to the right. □

The census is consistent with the picture of Theorem 6.14: radius 2 is too small for any train of length $m \geq 2$, and what appears instead are flips of whole classes of walls, defined by properties of the walls and their immediate surroundings that F and the flips preserve.

Proposition 6.17. *Let $F = F_{59}$ and $G = \langle \sigma, F, \mathcal{C}_2^{\times}(F) \rangle$. Then:*

(i) $G = \langle \sigma, F \rangle\, T$, *and* $T \cong (\mathbb{Z}/2)^6$ *is central in* G;
(ii) $\langle \sigma, F \rangle \cap T = \{1\}$;
(iii) $G \cong \mathbb{Z}^2 \times (\mathbb{Z}/2)^6$.

Every $t \neq 1$ in T has order 2, and every $\sigma^a F^b t$ with $(a, b) \neq (0, 0)$ has infinite order.

Proof. (i) By Proposition 6.16 the generators of G are of the form $\sigma^a F^b t$ with $t \in T$. The elements of T are automorphisms commuting with F, hence with σ and F, and with each other; so T is central and $G = \langle \sigma, F \rangle T$.

(ii) If $\sigma^a F^b = t \in T$, then $\sigma^{2a} F^{2b} = t^2 = 1$, so $a = b = 0$ by Theorem 2.4(3); rule 59 depends on its first and third arguments.

(iii) By Theorem 2.4(3), $\langle \sigma, F \rangle \cong \mathbb{Z}^2$. By (ii) and since T is central, $G \cong \langle \sigma, F \rangle \times T \cong \mathbb{Z}^2 \times (\mathbb{Z}/2)^6$.

For the orders: $(\sigma^a F^b t)^2 = \sigma^{2a} F^{2b}$, which has infinite order if $(a, b) \neq (0, 0)$. □

Corollary 6.18. *F_{59} has infinite order, and $\langle \sigma, F_{59}, \mathcal{C}_2^{\times}(F_{59}) \rangle \cong \mathbb{Z}^2 \times (\mathbb{Z}/2)^6$.*

Proof. Theorem 2.4 and Proposition 6.17(iii). □

Some orbits of F_{59} move slowly. Let c_0 be the configuration of spatial period 4000 with period $(021)^{666}\,02\,0^{2000}$; it has no wall of either family, and a direct computation shows $F^n(c_0) \neq c_0$ for $1 \leq n \leq 400{,}000$. At the interface between the staircase $(021)^{666}02$ and the uniform block 0^{2000} of c_0, F behaves like a binary odometer: the disturbance first reaches depth k at times close to 2^{k+2} (at $t = 501$, 1014, 2037, …, 131,055 for $k = 7, \ldots, 15$), so the period of c_0 under F, which is finite because c_0 is spatially periodic, is very large. Apart from walls, the dynamics of rule 59R is rather tame. A single changed cell on random rings stayed, in all trials, between the cores of the nearest walls on either side, and after transients most columns are periodic in time.

The walls of rule 59R are not time-periodic in the sense of Corollary 6.9, because the protector depends on the context. For $p \not\equiv 0 \pmod 6$ no word of $X_A \cup X_B$ is fixed by F^p in every context, since F^p moves the core along its 6-cycle; for $p \in \{6, 12, 30, 60, 120\}$ a computer search found, for every word of $X_A \cup X_B$, a context in which F^p changes the protector. We have not proved this for all multiples of 6; nothing depends on it, since Lemma 6.5 applies to the family X_A as it stands.

6.4. **Time-periodic walls: rules 127R, 179R and 223R.** For these three rules the walls of Takesue are pairs of adjacent cells whose images do not depend on the context: for 127R the pairs $b\,\kappa(b)$, $b \in \mathcal{A}$, and for 223R the pairs $b\,b$, on which F acts by the 4-cycle $0 \mapsto 1 \mapsto 3 \mapsto 2 \mapsto 0$ in both cells, so that they have period 4; and for 179R the pairs of the two cycles $01 \to 12 \to 20 \to 01$ and $10 \to 21 \to 02 \to 10$, of period 3 [19, Fig. 2 and Table III, walls (xii), (xiii), (xv), (xvi)]. Each word of the next proposition consists of two such pairs with one cell between them. The cell in the middle is the flip cell, and with it the words have period 8, 8 and 6, twice the period of the pairs.

Proposition 6.19. *For $f \in \{127, 179, 223\}$ the following cycles of words of length 5 satisfy the hypotheses of Corollary 6.9 for $F = F_f$, with $k = 5$, $j = 2$ and $\iota = F^{P/2}$ on W:*

$$127: \quad 03112 \to 12230 \to 30021 \to 21103 \to 03312 \to 12330 \to 30221 \to 21003 \quad (P = 8),$$

$$179: \quad 01110 \to 12321 \to 20302 \to 01310 \to 12221 \to 20002 \quad (P = 6),$$

$$223: \quad 00122 \to 11300 \to 33211 \to 22133 \to 00322 \to 11200 \to 33011 \to 22033 \quad (P = 8).$$

Hence $\bigoplus_{m\geq 1} S_m$ embeds in $\mathrm{Aut}(X_{F_f})$, and so in $\mathrm{Aut}(X_{F_{f'}})$ for the conjugate rules $f' = 254$, 205 and 251. Each word is fixed by F^P in every context.

Proof. (a) The images of the three interior cells of a word of length 5 depend only on the word, and the arrows above are the result of evaluating (1) on them and on the end cells. It remains to see that the images of the end cells do not depend on the outside neighbours, which enter only through their y-bits. For $f = 127$, $f(a, b, c) = \neg(a \wedge b \wedge c)$ depends on a only when $b = c = 1$ and on c only when $a = b = 1$; for $f = 179$ the same is true, since $f(0, b, c) = f(1, b, c)$ unless $b = c = 1$ and $f(a, b, 0) = f(a, b, 1)$ unless $a = b = 1$; so for these two rules it suffices that the y-bits of the first two and of the last two letters of each word are not both 1, which one reads off the words (y is the parity of the letter: for 127 the y-words are 01110, 10010, 10001, 01101, 01110, 10110, 10001, 01001, and for 179 they are 01110, 10101, 00100, 01110, 10001, 00000). For $f = 223$, $f(a, b, c) = \neg(a \wedge \neg b \wedge c)$ depends on a only when $(b, c) = (0, 1)$ and on c only when $(a, b) = (1, 0)$, and the y-words 00100, 11100, 11011, 00111, 00100, 11000, 11011, 00011 never begin with 01 or end with 10. (b) In each list the words $P/2$ apart differ exactly at position 2, as one reads off: for instance 03112 and 03312, 01110 and 01310, 00122 and 00322. (c) Comparing suffixes with prefixes shows that two words of W can overlap only at offsets 3 and 4 for 127 and 223, and only at offset 4 for 179; in particular never at offsets 1 or $2 = \max(j, k - 1 - j)$. Corollary 6.9 applies. The rules 254, 205, 251 are conjugate to 127, 179, 223 by reflection and complement. □

For 179, where $h = P/2 = 3$ is odd, one can also take $|K| = 1$: $\beta(w_s) = s \bmod 2$ satisfies (H3) with $\varepsilon = 1$. For 127 and 223, where $h = 4$, (H3) with a constant ε is impossible, since it would give $\beta(\iota w) = \beta(F^4 w) = \beta(w) + 4\varepsilon = \beta(w)$; so the classes of (H3′) are genuinely needed there, and trains consist of walls in the same phase modulo h. By Corollary 6.9 the flip of all walls is in each case an involutive automorphism of radius 2 commuting with F; this was also confirmed by composing tables exactly at radius 4 and by the pair-graph test. Rings of length 1500 seeded with trains at gaps 5, 6 and 8, with walls of other classes mixed in, confirmed commutation with F and F^{-1} and the homomorphism law, with no failures. We have not computed $\mathcal{C}_2^{\times}(F_f)$ for these three rules.

6.5. **Rule 27R.** The class $\{27, 83, 202, 216\}$ carries a version of the rule-59 mechanism with windows of length 6. Rule 27 has $f(a, 0, c) = \neg(a \wedge c)$ and $f(a, 1, c) = \neg a \wedge c$, so it differs from rule 59 only at $(a, b, c) = (1, 0, 1)$. Call a cell i *pinched* if $y_{i-1} = y_{i+1} = 1$. By (1), F_{27} acts on cells with values 1 and 3 as in (5), and

$$0 \mapsto \begin{cases} 0 & \text{pinched} \\ 1 & \text{otherwise,} \end{cases} \qquad 2 \mapsto \begin{cases} 1 & \text{pinched} \\ 0 & \text{otherwise.} \end{cases} \tag{6}$$

Let

$$W_{27} = \{\, a\,p\,p\,C\,(p+1)\,(p+1) \;:\; p \in \{0,1,2\},\ a \in \{p,3\},\ C \in \{p+1,3\} \,\},$$

twelve words. As for rule 59, call a the *protector*, C the *centre* and (p, C) the *core*. Let ι exchange the two values of C, at position $j = 3$, and let

$$\beta(a\,p\,p\,C\,(p+1)\,(p+1)) = [C = 3] + [p = 2] \pmod 2.$$

Theorem 6.20. *Let $F = F_{27}$. The family W_{27}, with $k = 6$, $j = 3$, ι and β as above, satisfies* (W1), (H1), (H2), (H3) *with $\varepsilon = 1$, and* (H4). *Consequently, for every $d \geq 6$ and $m \geq 1$ the maps Ψ_m of Definition 6.4 form an injective homomorphism $S_m \to \mathrm{Aut}(X_{F_{27}})$ whose image consists of block maps of radius at most $md + 3$, and the images for different m commute. So $\bigoplus_{m\geq 1} S_m$ embeds in $\mathrm{Aut}(X_{F_f})$ for $f \in \{27, 83, 202, 216\}$; each of these groups contains every finite group and is not virtually $\mathbb{Z}^n$ for any n. The map that flips every occurrence of W_{27} is an involutive automorphism of radius 3 commuting with F_{27}.*

Proof. (W1) and the core. Let $c_{[0,6)} = a\,p\,p\,C\,(p+1)\,(p+1)$. The cells $1, \ldots, 4$ read only cells of the word. Cells 0 and 5 read the outside neighbours, and only through their y-bits. By (5) and (6):

	(a, L, L, C, R, R)	(a', L', L', C', R', R')	$C \mapsto C'$	context enters
$p = 0$	$(\{0,3\}, 0, 0, \{1,3\}, 1, 1)$	$(\{1,3\}, 1, 1, \{3,2\}, 2, 2)$	$1 \mapsto 3,\ 3 \mapsto 2$	nowhere
$p = 1$	$(\{1,3\}, 1, 1, \{2,3\}, 2, 2)$	$(\{2,3\}, 2, 2, \{0,3\}, 0, 0)$	$2 \mapsto 0,\ 3 \mapsto 3$	at a
$p = 2$	$(\{2,3\}, 2, 2, \{0,3\}, 0, 0)$	$(\{0,3\}, 0, 0, \{1,3\}, 1, 1)$	$0 \mapsto 1,\ 3 \mapsto 3$	nowhere

We justify the entries.

- $p = 0$. Cells 1 and 2 have value 0 and are neighbours of each other with $y = 0$, so they are not pinched and become 1. The centre has $y_2 = 0$ and $y_4 = 1$, so it is triggered: $1 \mapsto 3$, $3 \mapsto 2$. Cells 4 and 5 have value 1 and left neighbours with $y = 1$, so they are not triggered and become 2. The protector has $y_1 = 0$, so it is neither pinched nor triggered: $0 \mapsto 1$, $3 \mapsto 3$.
- $p = 1$. Cells 1 and 2 have value 1 and left neighbours with $y = 1$, so they become 2. The centre $C = 2$ has $y_4 = 0$, so it is not pinched and becomes 0; the centre $C = 3$ has $y_2 = 1$, so it is not triggered and stays 3. Cells 4 and 5 have value 2 and are neighbours of each other with $y = 0$, so they become 0. The protector has $y_1 = 1$ and is triggered exactly when $y_{-1} = 0$, so $a' \in \{2, 3\}$ depends on the context.
- $p = 2$. Cells 1 and 2 have value 2 and are neighbours of each other with $y = 0$, so they become 0. The centre $C = 0$ has $y_2 = 0$, so it becomes 1; the centre $C = 3$ has $y_4 = 0$, so it stays 3. Cells 4 and 5 have value 0 and are neighbours of each other with $y = 0$, so they become 1. The protector has $y_1 = 0$: $2 \mapsto 0$, $3 \mapsto 3$.

In each case $F(c)_{[0,6)} \in W_{27}$ with p replaced by $p + 1$. This is (W1), and by Lemma 6.1 F and F^{-1} preserve the occurrences. The cells $1, \ldots, 5$ of the image depend only on the core, which runs through the 6-cycle

$$(0,1) \to (1,3) \to (2,3) \to (0,3) \to (1,2) \to (2,0) \to (0,1),$$

the A-cycle of Lemma 6.11.

(H3). The map ι changes $[C = 3]$ and not p, so $\beta \circ \iota = \beta + 1$. Along the cycle, β takes the values $0, 1, 0, 1, 0, 1$. Since β depends only on the core, $\beta(w_0(F(c))) = \beta(w_0(c)) + 1$ in every context, which is (H3) with $\varepsilon = 1$.

(H1). The configurations $\iota_0 c$ and c differ only at cell 3, so $F(\iota_0 c)$ and $F(c)$ can differ only at cells 2, 3, 4. In the justifications above, the images of cells 2 and 4 do not depend on which of the two values C has. For $p = 0$, cell 2 is decided by cell 1, and cell 4 reads C only through $y_C = 1$. For $p = 1$ and $p = 2$, cell 2 is decided by cell 1 and cell 4 by cell 5. The centre map $C \mapsto C'$ is a bijection from $\{p + 1, 3\}$ onto $\{p + 2, 3\}$, the centre pair of the image word. So $F(\iota_0 c)_3$ is the element of that pair other than $F(c)_3$, and $F(\iota_0 c) = \iota_0 F(c)$.

Overlaps. Suppose c has occurrences at 0 and $\delta \in \{1, 2\}$, with $c_1 = p$. If $\delta = 1$, then $c_2 = p$ and $c_3 = C$ are both the left cell of the second word, so $C = p$, which contradicts $C \in \{p + 1, 3\}$. If $\delta = 2$, then $c_3 = C$ and $c_4 = p + 1$ are both the left cell of the second word, so its protector

$c_2 = p$ must lie in $\{p+1, 3\}$, which is impossible. So distinct occurrences are at distance at least 3. Distances 3, 4 and 5 do occur: at distance 3 the centre of the first word is the protector of the second.

(H2) and (H4). The flip ι_0 writes only cell 3. It reads only cells 1 and 3, since the new value is the element of $\{c_1+1, 3\}$ other than c_3. The bit β reads positions 1 and 3 of its word. Let $i \neq 0$ be another occurrence, so $|i| \geq 3$. The window $[i, i+6)$ contains cell 3 only if $-2 \leq i \leq 3$, hence only if $i = 3$. Then cell 3 is the protector of the word at 3, whose left cell is $c_4 = c_1 + 1$. So its allowed protectors are $\{c_1+1, 3\}$, the pair between which ι_0 exchanges c_3. Hence the occurrence at i survives in $\iota_0 c$ in every case. Its positions 1 and 3, the cells $i+1$ and $i+3$, are not cell 3, so its bit is unchanged; this is (H4). The flip ι_i writes cell $i+3$, which is neither 1 nor 3 since $i \notin \{-2, 0\}$. Symmetrically, ι_0 writes no cell read by ι_i. So the two flips commute, which is (H2).

Conclusion. $W_{27} \neq \mathcal{A}^6$. Theorem 6.6 applies with $d \geq k = 6$ and gives the radius bound $md + \max(3, 2) = md + 3$. Corollary 6.3 gives the statement about the flip-all map. The conjugate rules 83, 202, 216 follow as in the proof of Theorem 6.14. □

Machine checks of each step of the proof are listed in Appendix A.4. At gap 3 consecutive walls overlap, as at gap 2 for rule 59; Theorem 6.6 does not cover this case, and we do not need it.

As for 59, the protector depends on the context; for instance the image of 1 1 1 2 2 2 is 2 2 2 0 0 0 or 3 2 2 0 0 0 depending on the left neighbour. The flip-all map has radius 3, and the maps Ψ_m have larger radius. This is consistent with the census of [30], in which the block maps of radius at most 2 that commute with F_{27} are the 13 maps $\sigma^a F^b$ and one constant map. A computer search (Appendix A.3) looked for invariant families with a hidden involution satisfying (W1) and (H1), with windows of length at most 6, among the 57 classes with no fixed word of length at most 3. It considered the closure of each word under F and F^{-1} in all contexts, if that closure has at most 300 words. It found such families only in the classes of 27, 59, 127, 179 and 223, all of which have periodic walls in [19, Section 4]. It says nothing about longer windows or larger families.

6.6. **Rotating walls: rules 219R and 126R.** In the walls of Sections 6.1–6.5, the cells next to the permuted information have context-free trajectories because they belong to a word with that property. For rules 219 and 126 the same effect comes from a relation between adjacent cells that F preserves. The cells that satisfy it all evolve in every context by one permutation of the alphabet, of order 4, so the walls they form stay in place but rotate through the symbols. We call them *rotating walls*. Throughout this subsection π is the cell permutation $\pi(x, y) = (y, 1-x)$, the 4-cycle $0 \mapsto 1 \mapsto 3 \mapsto 2 \mapsto 0$, so that $\pi^2 = \kappa$ and $\pi\kappa = \kappa\pi$. The pairs of equal cells of Lemma 6.21, which rotate by π, are the wall (xv) of Takesue for 219R, and the pairs $b\,\kappa(b)$ used for 126R below are his wall (xvi), both of period 4 [19, Fig. 2, Table III]. The source lists these walls without a proof, and it does not state that F creates no such pair; so Lemma 6.21 is proved here. The cars, the trains and the automorphisms are what is added.

Lemma 6.21 (bonds; cf. [19, Table III, wall (xv)])**.** *Let $F = F_{219}$, and say that c has a* bond *at $(i, i+1)$ if $c_i = c_{i+1}$.*

(a) *$F(c)$ has a bond at $(i, i+1)$ if and only if c does, and the same holds for F^{-1}.*
(b) *If c has a bond at $(i-1, i)$ or at $(i, i+1)$, then $F(c)_i = \pi(c_i)$, whatever the other cells of c are.*

Proof. The rule 219 vanishes exactly on 010 and 101, so $f(a, b, c) = 1$ whenever $a = b$ or $b = c$. If $c_i = c_{i+1}$, then $y_i = y_{i+1}$, so $f = 1$ on the triples (y_{i-1}, y_i, y_{i+1}) and (y_i, y_{i+1}, y_{i+2}), and by (1) $F(c)_i = (y_i, 1 - x_i) = \pi(c_i)$ and $F(c)_{i+1} = \pi(c_{i+1})$. This proves (b) and one half of (a). Conversely, if $F(c)_i = F(c)_{i+1}$, the first coordinates give $y_i = y_{i+1}$, hence $f = 1$ at i and at $i+1$

as before, and then the second coordinates give $1 - x_i = 1 - x_{i+1}$; so $c_i = c_{i+1}$. The statement for F^{-1} follows by applying this to $F^{-1}(c)$. □

So every maximal run of at least two equal cells keeps its position and length, and its symbol runs through the 4-cycle π in every context. A *car* of c is a position p with $c_{p-2} = c_{p-1} = c_{p+1} = c_{p+2} \neq c_p$; the common value s of its four neighbours is its *wall symbol*. For $s \in \mathcal{A}$ and $a \neq s$ let $\varphi_s(a)$ be the middle letter of $F(s\,s\,a\,s\,s)$, which by (1) depends only on s and a.

Lemma 6.22. *Let $F = F_{219}$.*

(a) *The cars of $F(c)$ are exactly the cars of c. A car p of c with wall symbol s is a car of $F(c)$ with wall symbol $\pi(s)$, and $F(c)_p = \varphi_s(c_p)$; and φ_s is a bijection $\mathcal{A} \setminus \{s\} \to \mathcal{A} \setminus \{\pi(s)\}$.*

(b) *If c' is obtained from c by changing the values of some cars, each to a value different from its wall symbol, then c' has the same bonds, the same cars and the same wall symbols as c.*

Proof. (a) Let p be a car of c with wall symbol s. The cells $p \pm 1$, $p \pm 2$ carry bonds, so they become $\pi(s)$ by Lemma 6.21(b), and $F(c)_p = \varphi_s(c_p)$ because $F(c)_p$ reads only c_{p-1}, c_p, c_{p+1}. By Lemma 6.21(a) there is no bond at $(p-1, p)$ or $(p, p+1)$ in $F(c)$, so $\varphi_s(c_p) \neq \pi(s)$ and p is a car of $F(c)$. The same argument for F^{-1}, under which bonded cells evolve by π^{-1}, shows that every car of $F(c)$ is a car of c, and that the corresponding map $\mathcal{A} \setminus \{\pi(s)\} \to \mathcal{A} \setminus \{s\}$ is inverse to φ_s. (Explicitly, $\varphi_0 = (1 \mapsto 2,\ 2 \mapsto 0,\ 3 \mapsto 3)$, $\varphi_1 = (0 \mapsto 0,\ 2 \mapsto 1,\ 3 \mapsto 2)$, $\varphi_2 = (0 \mapsto 1,\ 1 \mapsto 2,\ 3 \mapsto 3)$ and $\varphi_3 = (0 \mapsto 0,\ 1 \mapsto 3,\ 2 \mapsto 1)$.)

(b) Let p be a changed car with wall symbol s. The bonds at $(p-2, p-1)$ and $(p+1, p+2)$ do not involve p, and there is no bond at $(p-1, p)$ or $(p, p+1)$ before or after, since the new value differs from s. Whether a position q is a car depends on $c_{[q-2,q+2]}$. For $q \in \{p \pm 1, p \pm 2\}$ the cell q carries a bond in c and in c', so it is not a car of either; for $|q - p| \geq 3$ the window of q does not contain p. Wall symbols are cells next to cars, which are not changed. □

Two cars at distance 3 share the pair of wall cells between them, so they have the same wall symbol. A *train of length m* of c is a set $\{p, p+3, \ldots, p+3(m-1)\}$ of cars of c such that neither $p-3$ nor $p+3m$ is a car; its *car vector* is $(c_p, c_{p+3}, \ldots, c_{p+3(m-1)})$. Every car lies in at most one train (it may also lie in an infinite progression of cars, which is not a train). By Lemma 6.22(a), $F(c)$ has the same trains as c, and F applies the same bijection φ_s to every entry of the car vector of a train with wall symbol s. By Lemma 6.22(b), changing car values within $\mathcal{A} \setminus \{s\}$ does not change the trains.

Theorem 6.23. *Let $F = F_{219}$. For $\sigma \in S_m$ let $\Psi_m(\sigma)$ replace the car vector $(a_1, \ldots, a_m)$ of every train of length m by $(a_{\sigma^{-1}(1)}, \ldots, a_{\sigma^{-1}(m)})$ and leave every other cell unchanged. Then $\Psi_m(\sigma)$ is a block map of radius at most $3m + 2$ that commutes with F, and the Ψ_m assemble to an injective homomorphism $\bigoplus_{m \geq 1} S_m \to \mathrm{Aut}(X_F)$. In particular $\mathrm{Aut}(X_{F_{219}})$ contains every finite group and is not virtually $\mathbb{Z}^n$ for any n.*

Proof. Locality. Whether q is the r-th member of a train of length exactly m depends on whether the positions $q + 3t$, $-r \leq t \leq m - r + 1$, are cars, which is decided by $c_{[q-3r-2,\, q+3(m-r+1)+2]}$; the value written at q is one of the cars of that train. So $\Psi_m(\sigma)$ is a block map of radius at most $3m + 2$, and it commutes with σ.

Commutation. Let $c' = \Psi_m(\sigma)(c)$. By Lemma 6.22(b), c' has the bonds, cars, wall symbols and trains of c, and it differs from c only at members of trains of length m. Let q be the r-th member of such a train, with wall symbol s and car vector a. Then $F(c')_q = \varphi_s(a_{\sigma^{-1}(r)})$ by Lemma 6.22(a), and the same train in $F(c)$ has car vector $(\varphi_s(a_1), \ldots, \varphi_s(a_m))$, so $\Psi_m(\sigma)(F(c))_q = \varphi_s(a_{\sigma^{-1}(r)})$ as well. Let q be any other position. Then $\Psi_m(\sigma)(F(c))_q = F(c)_q$, because $F(c)$ has the trains of c. If q carries a bond in c, it carries one in c', and $F(c')_q = \pi(c'_q) = \pi(c_q) = F(c)_q$ by Lemma 6.21(b).

If q carries no bond, its neighbours $q \pm 1$ are not cars, because a neighbour of a car carries a bond; so c' agrees with c on $[q-1, q+1]$ and $F(c')_q = F(c)_q$. Hence $F\Psi_m(\sigma) = \Psi_m(\sigma)F$.

Group law. Since $\Psi_m(\rho)$ preserves trains and replaces each car vector a of length m by $a \circ \rho^{-1}$, $\Psi_m(\sigma)\Psi_m(\rho)$ replaces it by $a \circ \rho^{-1} \circ \sigma^{-1} = a \circ (\sigma\rho)^{-1}$; so Ψ_m is a homomorphism, each $\Psi_m(\sigma)$ is invertible with inverse $\Psi_m(\sigma^{-1})$, and $\Psi_m(\sigma) \in \mathrm{Aut}(X_F)$ by Lemma 2.2. For $m \neq m'$, $\Psi_m(\sigma)$ and $\Psi_{m'}(\rho)$ change disjoint sets of cells and preserve each other's trains and car vectors, so they commute, and the Ψ_m assemble to a homomorphism Ψ of $\bigoplus_m S_m$.

Injectivity. Let $g = \Psi((\sigma_{m'})_{m'})$ with $\sigma_m \neq \mathrm{id}$, and choose r with $\sigma_m(r) \neq r$. Let c be the configuration with $c_{3t} = 1$ for $t = r$, $c_{3t} = 2$ for $t \in \{1, \dots, m\} \setminus \{r\}$, and $c_q = 0$ at every other position q. Every cell 0 of c carries a bond, so the cars of c are exactly $3, 6, \dots, 3m$, which form one train of length m with wall symbol 0. Then $g(c)$ has the value 1 at $3\sigma_m(r) \neq 3r$, so $g(c) \neq c$. The last statement follows as in the proof of Theorem 4.2. □

Neither Theorem 1.1 nor Corollary 6.9 applies to 219R as stated. The walls are not fixed, since their symbol rotates, and the words $s\,s\,a\,s\,s$ with $a \neq s$, whose images are context-free by Lemma 6.21, have orbits of period 12 under F in which no word differs from the word half a period later at the car alone.

Proposition 6.24. *Let $F = F_{219}$, and for $s \in \mathcal{A}$ let $R_s = \varphi_{\pi^3(s)} \circ \varphi_{\pi^2(s)} \circ \varphi_{\pi(s)} \circ \varphi_s$, the permutation of $\mathcal{A} \setminus \{s\}$ by which F^4 acts on a car with wall symbol s. Let D be the map that replaces the value a of every car with wall symbol s by $R_s(a)$ and leaves every other cell unchanged. Then D is an automorphism of radius 2 commuting with F, $D^3 = \mathrm{id} \neq D$, and $D \notin \langle \sigma, F\rangle \times \langle \kappa \rangle$.*

Proof. D has radius 2, since being a car and the wall symbol are read from $c_{[p-2,p+2]}$. Since $\pi^4 = \mathrm{id}$, $R_{\pi(s)} \circ \varphi_s = \varphi_s \circ R_s$, so F intertwines the action of D on a car with wall symbol s and on the same car, with wall symbol $\pi(s)$, in $F(c)$; at all other cells the commutation argument in the proof of Theorem 6.23 applies verbatim, with Lemma 6.22(b). A direct computation gives $R_0 = (1 \mapsto 3 \mapsto 2 \mapsto 1)$, $R_1 = (0 \mapsto 2 \mapsto 3 \mapsto 0)$, $R_2 = (0 \mapsto 3 \mapsto 1 \mapsto 0)$ and $R_3 = (0 \mapsto 1 \mapsto 2 \mapsto 0)$, all 3-cycles; so $D^3 = \mathrm{id} \neq D$, and D is invertible. By Proposition 2.5, $\langle \sigma, F\rangle \times \langle \kappa \rangle \cong \mathbb{Z}^2 \times \mathbb{Z}/2$, which has no element of order 3. □

The map D is the analogue of the relabelling D of 54R. A variant of Ψ_2 with trains bounded by runs of at least three wall cells, which acts on the windows $s\,s\,s\,a\,s\,s\,b\,s\,s\,s$ ($a, b \neq s$) by exchanging a and b, has radius 6; it commutes with F on all 4^{15} windows and is an involution (computer-assisted; Appendix A.3).

For rule 126 the relation is $c_{i+1} = \kappa(c_i)$, and the walls are runs that alternate between a symbol and its complement. The two rules are related by the correspondence of [19, (A.5)–(A.7)]: the cellwise map θ that complements the y-bit of the cells at odd sites and the x-bit of the cells at even sites satisfies $\theta F_{126} = F_{219}\,\sigma\theta\sigma^{-1}$, and it maps the pairs $b\,\kappa(b)$ to pairs of equal cells. It does not commute with σ, so it does not identify the two automorphism groups, and the two rules are treated separately. Here Theorem 6.6 applies verbatim.

Proposition 6.25. *Let $F = F_{126}$ and*

$$W = \{\, b\,\kappa(b)\,q\,b\,\kappa(b) \;:\; b \in \mathcal{A},\ q \in \mathcal{A} \setminus \{b, \kappa(b)\} \,\} \subsetneq \mathcal{A}^5,$$

with $j = 2$ and ι replacing the letter q at position 2 by $\kappa(q)$. Then (W1), (H1), (H2), (H4) *and* (H3′) *hold, with the class of an occurrence $w = b\,\kappa(b)\,q\,b\,\kappa(b)$ equal to $b \in K = \mathcal{A}$, $\theta = \pi$, bit $\beta(w) = [q = \pi(b)]$ and $\varepsilon \equiv 0$. Hence $\bigoplus_{m \geq 1} S_m$ embeds in $\mathrm{Aut}(X_F)$. The flip-all map $E = \tau_W$ of Corollary 6.3, which replaces c_p by $\kappa(c_p)$ whenever $c_{[p-2,p+2]} \in W$, is an involutive automorphism of radius 2, and $E \notin \langle \sigma, F\rangle \times \langle \kappa \rangle$.*

Proof. The rule 126 vanishes exactly on 000 and 111. In every cell of $w \in W$ the y-bit differs from that of a neighbour inside w: for the cells b and $\kappa(b)$ because $y(\kappa(b)) = 1 - y(b)$, and for q because its two neighbours $\kappa(b)$ and b have different y-bits. So $f = 1$ at every cell of an occurrence, whatever the outside neighbours, and F maps the occurrence w to the cellwise image $\pi(w) = \pi(b)\,\kappa(\pi(b))\,\pi(q)\,\pi(b)\,\kappa(\pi(b))$, which lies in W because π commutes with κ. This is (W1), and it also gives the class map $\theta = \pi$. For (H1), $\iota_0 c$ again has an occurrence at 0, which F maps to $\pi(\iota(w)) = \iota(\pi(w))$; outside $[0, 5)$ the configurations $F(\iota_0 c)$ and $F(c)$ agree, since ι_0 changes only cell 2. For (H3′), $q \in \mathcal{A} \setminus \{b, \pi^2(b)\} = \{\pi(b), \pi^3(b)\}$, and $\kappa = \pi^2$ exchanges these two values, so $\beta \circ \iota = \beta + 1$; and $[\pi(q) = \pi(\pi(b))] = [q = \pi(b)]$, so $\varepsilon \equiv 0$. Two occurrences cannot lie at distance 1 or 2: at distance 1 the second would force $q = b$, and at distance 2 it would force $q = \kappa(b)$. As in the proof of Corollary 6.9, this gives (H2) and (H4). Theorem 6.6 and Corollary 6.3 apply. Finally, $E \neq \mathrm{id}$, and $E \neq \kappa$ since E fixes $(0, 0)^{\mathbb{Z}}$, which contains no occurrence. By Proposition 2.5, $\langle \sigma, F \rangle \times \langle \kappa \rangle \cong \mathbb{Z}^2 \times \mathbb{Z}/2$, whose only element of order 2 is κ. □

The trains of Theorem 6.6 here consist of occurrences $b\,\kappa(b)\,q\,b\,\kappa(b)$ in the same phase b, and the walls between their cars are alternating runs, which F rotates by π in every context. In both rules the commuting block maps of radius 2 are too many to enumerate, so the census of [30] does not list these automorphisms of radius 2.

6.7. **Blocking words.**

Remark 6.26. For a cellular automaton of radius 1, a word u is *blocking* [3] if there is a position q in u such that $F^n(x)_q = F^n(x')_q$ for all $n \geq 0$ and all x, x' that contain u at the same place; such a column separates the dynamics on its two sides, and by the argument in the proof of [13, Theorem 3] a cellular automaton has a blocking word if and only if it has equicontinuity points, that is, if and only if it is not sensitive [13, Corollary 2]. For the second-order elementary automata Takesue observed that the two sides of a wall cannot communicate and that a system with walls is not mixing [19, Section 4]. Every word of $X_A \cup X_B$ is blocking for F_{59}, with q the left cell (or the right cell), and also for F_{59}^{-1}, by Lemma 6.11. There are even blocking words of length 3: the six words $a\,p\,p$ with $a \in \{p, 3\}$, which are the walls of [19, (4.3)] on three cells, form a family invariant under F_{59} and F_{59}^{-1} in which the last two cells have context-free images $p + 1$, $p + 1$, as follows from (5) (each of the two in-phase cells has its left neighbour in the same phase, hence is never triggered). The time-periodic walls of Proposition 6.19 are blocking as well, since their cells have context-free trajectories, and so are the words of W_{27} for F_{27}, whose cells $1, \ldots, 5$ have context-free trajectories by the proof of Theorem 6.20. Words fixed in every context are blocking, but the converse fails, and the rules of this section show that the gap matters for automorphisms: rules 27, 59, 127, 179 and 223 have no word of length at most 3 fixed in every context (Table 1), and rule 59 has none of any length (Lemma 6.10), yet they have blocking words and their automorphism groups contain every finite group. This is why Question 1.6 is phrased in terms of blocking words rather than fixed words. In the nonlinear examples of this paper, the automorphisms outside $\langle \sigma, F \rangle$ all come from blocking words with a degree of freedom next to the blocking column: cars between fixed walls in Theorem 1.1, a hidden bit inside a moving wall in Theorems 6.6 and 6.14, and cars between rotating walls in Theorem 6.23 and Proposition 6.25. For the affine rules 90 and 165 this fails: they have no blocking word, yet their automorphism groups contain every finite group (Corollary 7.13).

7. The linear rules 90 and 150

For a linear rule $f(a,b,c) = \alpha a \oplus \beta b \oplus \gamma c$ the automaton F_f is $\mathbb{F}_2$-linear, and in terms of the ring $\Lambda = \mathbb{F}_2[s, s^{-1}]$ acting on $\mathbb{F}_2^{\mathbb{Z}}$ by $s = \sigma$ it is the matrix

$$M = \begin{pmatrix} 0 & 1 \\ 1 & p \end{pmatrix}, \qquad p = \alpha s^{-1} + \beta + \gamma s,$$

acting on column vectors $(x,y)^{\mathsf{T}}$; indeed $f(y_{i-1}, y_i, y_{i+1}) = (py)_i$. Rule 90 has $p = s + s^{-1}$ and rule 150 has $p = s + 1 + s^{-1}$. The characteristic polynomial of M is $t^2 + pt + 1$.

Lemma 7.1. *Let $K = \mathbb{F}_2(s)$. If $t^2 + pt + 1$ has distinct roots in K, or is irreducible over K, then the matrices in $M_2(\Lambda)$ commuting with M are exactly $aI + bM$ with $a, b \in \Lambda$. Suppose moreover that $p = s^{-1} + \beta + s$ with $\beta \in \mathbb{F}_2$, as for rules 90 and 150, and let $R \geq 1$. Then the matrices $aI + bM$ whose entries all lie in the span of $s^{-R}, \dots, s^R$ are those with $a \in \operatorname{span}(s^{-R}, \dots, s^R)$ and $b \in \operatorname{span}(s^{-R+1}, \dots, s^{R-1})$, an $\mathbb{F}_2$-space of dimension $4R$. For $R = 0$ they are 0 and I.*

Proof. In both cases M is not scalar and its minimal polynomial over K has degree 2, so its commutant in $M_2(K)$ is $K[M] = \{aI + bM : a, b \in K\}$. If $aI + bM = \left(\begin{smallmatrix} a & b \\ b & a+bp \end{smallmatrix}\right)$ has entries in Λ then $a, b \in \Lambda$. For the radius count, the entries a and b must lie in the span of $s^{-R}, \dots, s^R$, and then $a + bp$ lies in that span if and only if bp does. Since p has the terms s and s^{-1}, the terms of bp of degree $R+1$ and $-(R+1)$ are those of b of degree R and $-R$ multiplied by s and s^{-1}, so the condition is that b has no term $s^{\pm R}$. For $R \geq 1$ this leaves $2R - 1$ coefficients of b and $2R + 1$ of a, all free; for $R = 0$ it forces $b = 0$, and $a \in \{0, 1\}$. □

For rule 90, $t^2 + (s + s^{-1})t + 1 = (t + s)(t + s^{-1})$ has the distinct roots $s^{\pm 1}$. For rule 150 the polynomial is irreducible over K: a root would be integral over Λ, hence in Λ, and the product of the two roots being 1 makes it a unit s^k, but then $s^k + s^{-k} = s + 1 + s^{-1}$ is impossible.

For the linear rules the centralizer is a vector space and can be computed by Gaussian elimination over $\mathbb{F}_2$.

Theorem 7.2. *Let $F = F_{90}$, let $\Lambda = \mathbb{F}_2[s, s^{-1}]$ act on $\mathbb{F}_2^{\mathbb{Z}}$ through $s = \sigma$, and let $M = \left(\begin{smallmatrix} 0 & 1 \\ 1 & p \end{smallmatrix}\right)$ with $p = s + s^{-1}$, so that $F(x,y) = M(x,y)^{\mathsf{T}}$.*

- (a) *The affine block maps commuting with F are the maps $c \mapsto (aI + bM)c + k$ with $a, b \in \Lambda$ and $k \in \{(0,0)^{\mathbb{Z}}, (1,1)^{\mathbb{Z}}\}$. For $R \geq 1$, those of radius at most R form an $\mathbb{F}_2$-space of dimension $4R + 1$; those of radius 0 are the four maps $c \mapsto ac + k$ with $a \in \{0, 1\}$.*
- (b) *The maps $u = x + s^{-1}y$ and $w = x + sy$ are factor maps onto the binary full shift with $u \circ F = \sigma^{-1} \circ u$ and $w \circ F = \sigma \circ w$; the maps $\iota(v) = (v, s^{-1}v)$ and $\iota'(v) = (v, sv)$ are embeddings with $F \circ \iota = \iota \circ \sigma^{-1}$ and $F \circ \iota' = \iota' \circ \sigma$. For every block map h of the binary full shift, $\iota \circ h \circ u$ and $\iota' \circ h \circ w$ commute with F.*
- (c) *For every $R \geq 1$ the $\mathbb{F}_2$-span of the affine commuting maps of radius at most R and of the maps in (b) of radius at most R has dimension exactly $2^{2R} + 1$. For $R \leq 3$ this span is the whole space of block maps of radius at most R commuting with F, so that space has dimension 5, 17 and 65 for $R = 1, 2, 3$.*

Proof. (a) An affine block map has the form $c \mapsto Nc + k$ with N a linear block map and $k = g(0)$ a constant configuration, $k \in \{0^{\mathbb{Z}}, 1^{\mathbb{Z}}, 2^{\mathbb{Z}}, 3^{\mathbb{Z}}\}$. It commutes with $F = M$ if and only if $NM = MN$ and $Mk = k$. For a constant $k = (x,y)^{\mathbb{Z}}$ one has $F(k) = (y, x \oplus f(y,y,y))^{\mathbb{Z}} = (y,x)^{\mathbb{Z}}$ since $f(y,y,y) = 0$ for rule 90, so k is fixed if and only if $x = y$, that is, $k \in \{(0,0)^{\mathbb{Z}}, (1,1)^{\mathbb{Z}}\}$. Combined with Lemma 7.1 this gives dimension $4R + 1$ for $R \geq 1$, and for $R = 0$ the linear parts 0 and I with the two constants.

(b) The row vector $(1, s^{-1})$ satisfies $(1, s^{-1})M = (s^{-1}, 1 + s^{-1}p) = (s^{-1}, s^{-2}) = s^{-1}(1, s^{-1})$, which is $u \circ F = \sigma^{-1} \circ u$ for $u(x,y) = x + s^{-1}y$; similarly $(1, s)M = s(1, s)$. The column vector

$(1, s^{-1})^{\mathsf{T}}$ satisfies $M(1, s^{-1})^{\mathsf{T}} = (s^{-1}, 1 + ps^{-1})^{\mathsf{T}} = s^{-1}(1, s^{-1})^{\mathsf{T}}$, which is $F \circ \iota = \iota \circ \sigma^{-1}$, and similarly for ι'. Both u and w are surjective onto $\mathbb{F}_2^{\mathbb{Z}}$ (choose $y = 0$). For a block map h of the binary full shift, which commutes with $\sigma^{\pm 1}$,

$$(\iota h u)F = \iota h \sigma^{-1} u = \iota \sigma^{-1} h u = F \iota h u,$$

and likewise for $\iota' h w$.

(c) Fix $R \geq 1$ and let A_R be the space in (a), U_R the span of the maps $\iota h u$ with h a polynomial in the variables u_j, $j \in J_U = [-(R-2), R]$, and W_R the span of the $\iota' h w$ with h a polynomial in w_k, $k \in J_W = [-R, R-2]$; the index ranges are those for which $\iota h u$ and $\iota' h w$ have radius at most R, since $(\iota h u)(c)_i = (h(u)_i, h(u)_{i-1})$ and $u_{i+j} = x_{i+j} \oplus y_{i+j-1}$. The linear forms u_j ($j \in J_U$) and w_k ($k \in J_W$) on the $2(2R+1)$ coordinates of a window are jointly linearly independent: a relation $\sum a_j u_j + \sum b_k w_k = 0$ gives $a_m = b_m$ from the coefficient of x_m and $a_{m+1} = b_{m-1}$ from that of y_m, with a, b extended by zero outside their ranges; so $a_{m+1} = a_{m-1}$ for all $m \in \mathbb{Z}$. Since $R \notin J_W$ and $R - 1 \notin J_W$, we have $b_R = b_{R-1} = 0$, hence $a_R = b_R = 0$ and $a_{R-1} = b_{R-1} = 0$; descending with $a_{m-1} = a_{m+1}$ gives $a_m = 0$ for all $m \leq R$, hence for all m, and then $b_m = a_m = 0$ for all m. Consequently distinct monomials in the u_j give linearly independent functions, $\dim U_R = 2^{|J_U|} = 2^{2R-1}$, likewise $\dim W_R = 2^{2R-1}$, and a polynomial in the u's equals a polynomial in the w's only if both are constant, so $U_R \cap W_R$ is spanned by the constant map onto $(1,1)^{\mathbb{Z}}$ and $\dim(U_R + W_R) = 2^{2R} - 1$. An element $\iota h u + \iota' h' w$ is affine if and only if h and h' are affine, again by independence of the variables, so $A_R \cap (U_R + W_R) = (A_R \cap U_R) + (A_R \cap W_R)$, where $A_R \cap U_R$ consists of the $\iota h u$ with h affine and has dimension $|J_U| + 1 = 2R$, likewise $\dim(A_R \cap W_R) = 2R$, and the two meet in the constants. Therefore

$$\dim(A_R + U_R + W_R) = (4R+1) + (2^{2R} - 1) - (4R - 1) = 2^{2R} + 1.$$

This is a lower bound for $\dim \mathcal{C}_R(F)$, since all three families commute with F. The Gaussian elimination of Section 3 gives $\dim \mathcal{C}_R(F) = 5, 17, 65$ for $R = 1, 2, 3$, and a direct rank computation on the tables of the three families, each table re-verified to commute, gives the same numbers, so the span is everything for $R \leq 3$. □

Conjecture 7.3. $\dim_{\mathbb{F}_2} \mathcal{C}_R(F_{90}) = 2^{2R} + 1$ *for every* $R \geq 1$*; equivalently, every block map commuting with* F_{90} *is a sum of an affine commuting map, a map* $\iota h u$ *and a map* $\iota' h' w$*.*

The invertible elements are few. At $R = 1$ the 32 commuting maps contain 10 invertible ones, all affine, namely the 5 maps $\sigma^a F^b$ and the 5 further maps $(aI + bM)c + k$ whose determinant $a^2 + abp + b^2$ is a unit of Λ. The maps $\iota h u$ and $\iota' h w$ are never injective, since they factor through u or w. (The maps $\mathrm{id} + \iota h u$ can be injective; see Section 7.1.) At $R = 2$ the enumeration of all 2^{17} commuting maps, with the pair-graph test applied to the 98,222 balanced ones, finds exactly 26 invertible maps, all affine: the 13 maps $\sigma^a F^b$ and the 13 further maps $(aI + bM)c + k$ with unit determinant. So every automorphism of $X_{F_{90}}$ of radius at most 2 is affine. This does not persist: by Theorem 7.10, $X_{F_{90}}$ has non-affine automorphisms, and its automorphism group contains every finite group.

Proposition 7.4. *For* $1 \leq R \leq 3$*,* $\mathcal{C}_R(F_{150})$ *consists exactly of the* 2^{4R} *linear maps* $(aI + bM)c$ *of Lemma 7.1.*

Proof. Lemma 7.1 gives 2^{4R} linear commuting maps, and the only constant configuration fixed by F_{150} is $(0,0)^{\mathbb{Z}}$: since $f(y, y, y) = y$, F_{150} maps $(x, y)^{\mathbb{Z}}$ to $(y, x \oplus y)^{\mathbb{Z}}$, which equals $(x, y)^{\mathbb{Z}}$ only for $x = y = 0$. So there is no further constant. Gaussian elimination gives $\dim \mathcal{C}_R(F_{150}) = 4, 8, 12$ for $R = 1, 2, 3$. □

So the non-affine commuting maps of F_{90} are explained entirely by the two eigen-factors onto the binary shift, and the rule without eigen-factors has none, at least through radius 3. Both

TABLE 3. Block maps of radius at most R commuting with F_{90} and F_{150}. "Named" are the maps $\sigma^a F^b$ with $|a| + |b| \le R$; "other" are maps that are neither named nor constant; the last two columns count the invertible commuting maps, that is, the automorphisms of radius at most R, and those outside $\langle \sigma, F \rangle$; for 150R, $h = \mathrm{id} + \sigma F$ as in Proposition 7.5.

rule	R	commuting	named	constants	other	invertible	outside $\langle \sigma, F \rangle$
90R	1	2^5	5	2	25	10	5, all affine
90R	2	2^{17}	13	2	rest	26	13, all affine
90R	3	2^{65}	25	2	rest	—	—
150R	1	2^4	5	1	10	9	4, all in $\langle \sigma, F, h \rangle$
150R	2	2^8	13	1	rest	43	30, all in $\langle \sigma, F, h \rangle$
150R	3	2^{12}	25	1	rest	115	90, all in $\langle \sigma, F, h \rangle$

results also confirm the encoding: for the linear rules the SAT enumeration at $R \le 2$ and the elimination agree, and the $2^{2R} + 1$ and $4R$ patterns are exactly what the algebra predicts.

The commuting maps of F_{150} are all linear through radius 3, but its automorphism group is still larger than $\langle \sigma, F_{150} \rangle$. In $M_2(\Lambda)$ the shift is the scalar matrix sI and F_{150} is M; under the ring isomorphism $\Lambda[M] \cong \Lambda[t]/(t^2 + pt + 1)$ the commutant of Lemma 7.1 becomes a subring of the field $L = K[t]/(t^2 + pt + 1)$, a quadratic extension of $K = \mathbb{F}_2(s)$, and composition of commuting linear maps becomes multiplication in L.

Proposition 7.5. *Let $F = F_{150}$ and $h = \mathrm{id} + \sigma F$. Then $h \in \mathrm{Aut}(X_F)$, $h \notin \langle \sigma, F \rangle$, and $(a, b, e) \mapsto \sigma^a F^b h^e$ is an isomorphism $\mathbb{Z}^3 \to \langle \sigma, F, h \rangle$. Moreover F has no blocking word. In particular $\mathrm{Aut}(X_{F_{150}})$ is not virtually $\mathbb{Z}^2$.*

Proof. The map h has matrix $I + sM$, so it is a block map commuting with F and σ. It is invertible, because $\det(I + sM) = 1 + sp + s^2 = 1 + s(s + 1 + s^{-1}) + s^2 = s$ is a unit of Λ, so the inverse matrix has entries in Λ. By Lemma 2.2, $h \in \mathrm{Aut}(X_F)$.

For independence, identify $\sigma^a F^b h^e$ with the element $s^a t^b (1 + st)^e$ of $L^\times$, where t is the image of M; the identification is an injective homomorphism on the commuting linear invertible maps, since I and M are linearly independent over Λ. Suppose $s^a t^b (1 + st)^e = 1$. Let $N \colon L^\times \to K^\times$ be the norm, which is the determinant: $N(s) = s^2$, $N(t) = \det M = 1$ and $N(1 + st) = s$. So $s^{2a+e} = 1$, and $e = -2a$. Put $u = (1 + st)^2 s^{-1}$; the relation becomes $t^b = u^a$. Since $t^2 = pt + 1$, we have $t^{-1} = t + p$, and a direct computation with $sp = s^2 + s + 1$ and $(s^2 + s + 1)p = s^3 + s + s^{-1}$ gives

$$u = s^{-1} + s + (s^2 + s + 1)t = s^3 + (s^2 + s + 1)t^{-1}, \qquad ut = s^3 t + s^2 + s + 1.$$

Let v be a valuation of L extending a valuation of K with $v(s) = e_0 > 0$. Then $v(p) = -e_0$, and since $t \cdot t^{-1} = 1$ and $t + t^{-1} = p$, one of $v(t)$, $v(t^{-1})$ is $-e_0$ and the other e_0. If $v(t) = -e_0$, the second expression gives $v(u) = \min(3e_0, e_0) = e_0$; if $v(t) = e_0$, the first gives $v(u) = -e_0$, since $v(s^{-1}) = -e_0$ and the other terms have positive valuation. In both cases $v(u) = -v(t)$, and $t^b = u^a$ gives $b = -a$, so $(ut)^a = 1$. The element ut is not a root of unity: a root of unity in L is algebraic over $\mathbb{F}_2$, so it lies in the algebraic closure of $\mathbb{F}_2$ in L, a finite field $\mathbb{F}_q$ with $[\mathbb{F}_q : \mathbb{F}_2] \le [L : K] = 2$; every element of $\mathbb{F}_4$ has trace over K in $\mathbb{F}_2$, while the trace of ut is $s^3(t + t^{-1}) = s^3 p = s^4 + s^3 + s^2$. Hence $a = 0$, and then $b = e = 0$. So the homomorphism $\mathbb{Z}^3 \to \langle \sigma, F, h \rangle$ is injective. In particular $h \notin \langle \sigma, F \rangle$.

For the blocking words, let u be a word, q a position in it, and x a configuration containing u on $[0, |u|)$. For $N \ge |u|$ let x' be x with the y-bit of the cell N complemented, and let

$\Delta = Y_x \oplus Y_{x'}$. By linearity Δ satisfies $\Delta(t+1,i) = \Delta(t-1,i) \oplus \Delta(t,i-1) \oplus \Delta(t,i) \oplus \Delta(t,i+1)$, with $\Delta(-1,\cdot) = 0$ and $\Delta(0,\cdot)$ the unit vector at N. By induction on $t \geq 0$, $\Delta(t,i) = 0$ for $i < N - t$ and $\Delta(t, N-t) = 1$: at $i < N - t - 1$ all four terms vanish, and at $i = N - t - 1$ only $\Delta(t, N-t) = 1$ is nonzero. For $t = N - q$ this gives $F^{N-q}(x)_q \neq F^{N-q}(x')_q$, so u is not blocking with column q.

Finally, if $\mathrm{Aut}(X_F)$ had a finite-index subgroup isomorphic to $\mathbb{Z}^2$, it would meet $\langle \sigma, F, h\rangle \cong \mathbb{Z}^3$ in a subgroup of finite index in $\mathbb{Z}^3$, hence of rank 3, inside $\mathbb{Z}^2$; this is impossible. □

In fact $\mathrm{Aut}(X_{F_{150}}) = \langle \sigma, F, h\rangle$: every automorphism of $X_{F_{150}}$ is affine (Lemma 8.4), so it is a map $aI + bM$ whose determinant $a^2 + abp + b^2$ is a unit of Λ, and these maps are exactly the elements $s^a t^b (1+st)^e$ (Lemma 8.5 and Theorem 1.4(4)).

Remark 7.6. For $M \geq 0$ let Λ_M be the light-speed window graph of F: its vertices are the words $\omega \in \mathcal{A}^{M+1}$, and for every configuration $\tilde{c}$ with $\tilde{c}_{[p,p+M]} = \omega$ it has an edge from ω to the word $F(\tilde{c})_{[p+1,p+M+1]}$, which depends only on ω and on $\tilde{c}_{p+M+1}, \tilde{c}_{p+M+2}$. Its strong connectivity for every M is one of the hypotheses of the rigidity theorems of [29]. A direct construction of Λ_M from the local rule (Appendix A.3) finds it strongly connected for $M \leq 4$ for the affine rules 15, 60, 105, 150, 195 and 240, and not strongly connected for 90 and 165. So for 150 the extra automorphisms of Proposition 7.5 are not detected by these graphs; they come from linearity.

7.1. **Every finite group for 90R and 165R.** The maps $\iota h u$ of Theorem 7.2(b) are never invertible, but a map $\mathrm{id} + \iota h u$ can be. Here we show that such maps carry two commuting copies of $\bigoplus_m S_m$, for rule 90 and for its complement 165.

Throughout this subsection F is F_{90} or F_{165}, and $\varepsilon = 0$ or 1 accordingly. Since $165(a,b,c) = 1 \oplus a \oplus c$, we have $F_{165}(c) = F_{90}(c) + \varepsilon(0,1)^{\mathbb{Z}}$, with cellwise addition in $\mathbb{F}_2^2$. Binary configurations $v \in \mathbb{F}_2^{\mathbb{Z}}$ are added cellwise, $\mathbf{1} = 1^{\mathbb{Z}}$, $\bar{v} = v + \mathbf{1}$, and $(\sigma v)_n = v_{n+1}$. Define

$$d(c)_n = x_{n+1} + y_{n+2}, \qquad e(c)_n = y_n + x_{n+1}, \qquad \iota(v)_i = (v_{i-1}, v_i), \qquad \iota'(v)_i = (v_{i+1}, v_i);$$

thus ι is the embedding of Theorem 7.2(b), and $d = \sigma w$, $e = \sigma u$ for its factor maps w and u.

Lemma 7.7. (a) *d and e are $\mathbb{F}_2$-linear block maps $\mathcal{A}^{\mathbb{Z}} \to \mathbb{F}_2^{\mathbb{Z}}$ commuting with σ, d reads the cells $n+1, n+2$ and e the cells $n, n+1$, and both are surjective.*

(b) *$d(F(c)) = \sigma d(c) + \varepsilon\mathbf{1}$ and $e(F(c)) = \sigma^{-1} e(c) + \varepsilon\mathbf{1}$.*

(c) *$F_{90} \circ \iota = \iota \circ \sigma$ and $F_{90} \circ \iota' = \iota' \circ \sigma^{-1}$.*

(d) *$d \circ \iota = \mathrm{id} + \sigma^2$, $e \circ \iota = 0$, $d \circ \iota' = 0$ and $e \circ \iota' = \mathrm{id} + \sigma^2$.*

Proof. (a) is clear; for surjectivity take $y = 0$. Write $F(c)_i = (y_i,\ x_i + y_{i-1} + y_{i+1} + \varepsilon)$. Then $d(F(c))_n = y_{n+1} + x_{n+2} + y_{n+1} + y_{n+3} + \varepsilon = d(c)_{n+1} + \varepsilon$, and $e(F(c))_n = x_n + y_{n-1} + y_{n+1} + \varepsilon + y_{n+1} = e(c)_{n-1} + \varepsilon$, which is (b). For (c), $F_{90}(\iota v)_i = (v_i,\ v_{i-1} + v_{i-1} + v_{i+1}) = (v_i, v_{i+1}) = \iota(\sigma v)_i$ and $F_{90}(\iota' v)_i = (v_i,\ v_{i+1} + v_{i-1} + v_{i+1}) = (v_i, v_{i-1}) = \iota'(\sigma^{-1} v)_i$. For (d), $d(\iota v)_n = v_n + v_{n+2}$, $e(\iota v)_n = v_n + v_n = 0$, $d(\iota' v)_n = v_{n+2} + v_{n+2} = 0$ and $e(\iota' v)_n = v_n + v_{n+2}$. □

Definition 7.8. Let $\mathcal{G}$ be the set of automorphisms ψ of the binary full shift with $\psi(0^{\mathbb{Z}}) = 0^{\mathbb{Z}}$ for which there is a block map $J\colon \mathbb{F}_2^{\mathbb{Z}} \to \mathbb{F}_2^{\mathbb{Z}}$ with $J(0^{\mathbb{Z}}) = 0^{\mathbb{Z}}$ and

$$\psi(z) = z + J(z) + \sigma^2 J(z) \qquad (z \in \mathbb{F}_2^{\mathbb{Z}}),$$

that is, $\psi = \mathrm{id} + (\mathrm{id} + \sigma^2)J$. We call $J = J_\psi$ the *integral* of ψ. Let $\mathcal{G}^\kappa \subseteq \mathcal{G}$ consist of the ψ with $J_\psi(\bar{z}) = J_\psi(z)$ for all z; such ψ satisfy $\psi(\bar{z}) = \overline{\psi(z)}$.

Lemma 7.9. *The integral is unique. $\mathcal{G}$ is a group, with $J_{\psi\chi} = J_\chi + J_\psi \circ \chi$ and $J_{\psi^{-1}} = J_\psi \circ \psi^{-1}$, and $\mathcal{G}^\kappa$ is a subgroup.*

Proof. Let K be a block map of radius r with $K(0^{\mathbb{Z}}) = 0^{\mathbb{Z}}$ and $(\mathrm{id}+\sigma^2)K = 0$, so that every $K(z)$ is invariant under σ^2. Given z and n, let z' agree with z on $[n-r, n+r]$ and vanish on $[n+2N-r, n+2N+r]$ for some large N. Then $K(z)_n = K(z')_n = K(z')_{n+2N} = K(0^{\mathbb{Z}})_n = 0$. So $K = 0$, which gives uniqueness. For $\psi, \chi \in \mathcal{G}$, $(\mathrm{id}+\sigma^2)(J_\chi + J_\psi \circ \chi) = (\chi+\mathrm{id})+(\psi\chi+\chi) = \psi\chi+\mathrm{id}$, and $J_\chi + J_\psi \circ \chi$ is a block map vanishing at $0^{\mathbb{Z}}$; so $\psi\chi \in \mathcal{G}$ with the stated integral. Similarly $(\mathrm{id}+\sigma^2)(J_\psi \circ \psi^{-1}) = (\psi+\mathrm{id}) \circ \psi^{-1} = \mathrm{id} + \psi^{-1}$, so $\psi^{-1} \in \mathcal{G}$. If J_ψ and J_χ are invariant under $z \mapsto \bar{z}$, then ψ and χ commute with it, since $\psi(\bar{z}) = \bar{z} + (\mathrm{id}+\sigma^2)J_\psi(z) = \overline{\psi(z)}$, and hence so are $J_\chi + J_\psi \circ \chi$ and $J_\psi \circ \psi^{-1}$. □

Theorem 7.10. *For $\psi \in \mathcal{G}$ define*

$$\Phi_\psi(c) = c + \iota\big(J_\psi(d(c))\big), \qquad \Phi'_\psi(c) = c + \iota'\big(J_\psi(e(c))\big).$$

Let $H = \mathcal{G}$ if $F = F_{90}$ and $H = \mathcal{G}^\kappa$ if $F = F_{165}$. Then $(\psi, \chi) \mapsto \Phi_\psi \Phi'_\chi$ is an injective homomorphism $H \times H \to \mathrm{Aut}(X_F)$. The group $\bigoplus_{m\geq 1} S_m$ embeds in $\mathcal{G}^\kappa$ (Lemma 7.11), so $(\bigoplus_{m\geq 1} S_m)^2$ embeds in $\mathrm{Aut}(X_{F_{90}})$ and in $\mathrm{Aut}(X_{F_{165}})$. In particular both groups contain every finite group, are nonabelian, and are not virtually $\mathbb{Z}^n$ for any n.

Proof. Locality. Φ_ψ and Φ'_ψ are composites and sums of block maps commuting with σ, so they are block maps.

The coordinates. By linearity of d and e and Lemma 7.7(d),

$$d \circ \Phi_\psi = \psi \circ d, \qquad e \circ \Phi_\psi = e, \qquad d \circ \Phi'_\psi = d, \qquad e \circ \Phi'_\psi = \psi \circ e.$$

For instance $d(\Phi_\psi(c)) = d(c) + (\mathrm{id}+\sigma^2)J_\psi(d(c)) = \psi(d(c))$.

Commutation. Write $F = F_{90} + k$ with $k = \varepsilon(0,1)^{\mathbb{Z}}$, and recall that J_ψ commutes with σ. By Lemma 7.7(c), $F(\Phi_\psi(c)) = F(c) + F_{90}\iota J_\psi(d(c)) = F(c) + \iota\sigma J_\psi(d(c))$. By Lemma 7.7(b), $\Phi_\psi(F(c)) = F(c) + \iota J_\psi(\sigma d(c) + \varepsilon\mathbf{1})$, and $J_\psi(\sigma d(c) + \varepsilon\mathbf{1}) = J_\psi(\sigma d(c)) = \sigma J_\psi(d(c))$: for $\varepsilon = 0$ trivially, and for $\varepsilon = 1$ because $\psi \in \mathcal{G}^\kappa$. So $F\Phi_\psi = \Phi_\psi F$. The same computation with ι', e and σ^{-1} gives $F\Phi'_\psi = \Phi'_\psi F$.

Homomorphism and invertibility. By the coordinate identities, the linearity of ι and Lemma 7.9,

$$\Phi_\psi(\Phi_\chi(c)) = c + \iota J_\chi(d(c)) + \iota J_\psi(\chi(d(c))) = c + \iota J_{\psi\chi}(d(c)) = \Phi_{\psi\chi}(c),$$

and $\Phi_{\mathrm{id}} = \mathrm{id}$ since $J_{\mathrm{id}} = 0$. So Φ_ψ is invertible with inverse $\Phi_{\psi^{-1}}$; here the invertibility of $\psi = \mathrm{id} + (\mathrm{id}+\sigma^2)J_\psi$ is what makes Φ_ψ invertible. With the commutation, $\Phi_\psi \in \mathrm{Aut}(X_F)$ by Lemma 2.2, and $\psi \mapsto \Phi_\psi$ is a homomorphism $H \to \mathrm{Aut}(X_F)$. The same holds for $\psi \mapsto \Phi'_\psi$.

The two movers commute. Since $d \circ \Phi'_\chi = d$ and $e \circ \Phi_\psi = e$,

$$\Phi_\psi \Phi'_\chi(c) = c + \iota' J_\chi(e(c)) + \iota J_\psi(d(c)) = \Phi'_\chi \Phi_\psi(c).$$

So $(\psi, \chi) \mapsto \Phi_\psi \Phi'_\chi$ is a homomorphism $H \times H \to \mathrm{Aut}(X_F)$.

Injectivity. If $\Phi_\psi \Phi'_\chi = \mathrm{id}$, then applying d gives $\psi(d(c)) = d(c)$ for every c, so $\psi = \mathrm{id}$ because d is surjective; applying e gives $\chi = \mathrm{id}$.

The remaining statements follow from Lemma 7.11, with $\mathcal{G}^\kappa \subseteq \mathcal{G}$, and from the argument in the proof of Theorem 4.2. □

It remains to produce automorphisms of the binary full shift with integrals. They come from marker permutations whose changes have zero sum on the even and on the odd positions of each segment.

Lemma 7.11. *For $L \geq 1$ let U_L be the set of words $u \in \{0,1\}^L$ with $u_0 = u_{L-1} = 0$, with no factor 11 and no factor 000, and with $\sum_{k \text{ even}} u_k \equiv \sum_{k \text{ odd}} u_k \equiv 0 \pmod 2$. A* frame *of $z \in \mathbb{F}_2^{\mathbb{Z}}$ of length L at p is a position p with $z_{[p,p+L+6)} = 111\,u\,111$ (a positive frame) or $z_{[p,p+L+6)} = 000\,\bar{u}\,000$ (a negative frame) for some $u \in U_L$, its* content*; its* segment *is $[p+3, p+L+2]$. For a family*

$\rho = (\rho_L)_{L\geq 1}$ of permutations $\rho_L \in \mathrm{Sym}(U_L)$, all but finitely many equal to the identity, let $\psi_\rho(z)$ be obtained from z by replacing the content u of every frame of length L by $\rho_L(u)$, that is, writing $\rho_L(u)$ on the segment of a positive frame and $\overline{\rho_L(u)}$ on that of a negative one. Then $\rho \mapsto \psi_\rho$ is an injective homomorphism $\bigoplus_L \mathrm{Sym}(U_L) \to \mathcal{G}^\kappa$. Moreover $|U_{4N+11}| \geq N+1$ for $N \geq 0$, so $\bigoplus_{m\geq 1} S_m$ embeds in $\mathcal{G}^\kappa$.

Proof. (i) *No run of three equal symbols meets a segment.* In a positive frame every 1 of the segment has two neighbours 0, since u has no 11 and starts and ends with 0; and the cells next to the segment are 1, so a run 000 meeting the segment would lie inside it, which u excludes. Negative frames are the complements.

(ii) *Segments of distinct frames are disjoint, at least three cells apart.* Two frames at the same position have the same sign, given by z_p, and if their lengths were $L < L'$, then the cell $p+L+3$, which lies in a run of three, would lie in the segment $[p+3, p+L'+2]$ of the longer one, contradicting (i). Let $p < q$ be frames, of lengths L_p and L_q. The run $z_{[q,q+3)}$ is disjoint from the segment of p by (i), and $q+2 > p+2$, so $q \geq p+L_p+3$. Hence the segment of q starts at $q+3 \geq p+L_p+6$, at least three cells after the end $p+L_p+2$ of the segment of p.

(iii) *ψ_ρ preserves frames.* Let $z' = \psi_\rho(z)$. By (i) and (ii), ψ_ρ changes only cells of segments and leaves the runs of three equal symbols in place, so every frame of z is a frame of z', of the same length and sign, with content $\rho_L(u)$; note that $\rho_L(u) \in U_L$ again satisfies (i). Conversely let p be a frame of z'. If its segment meets the segment of a frame q of z, then q is also a frame of z', and $p = q$ by (ii) applied to z'. Otherwise $z = z'$ on the segment of p, and also on the two runs of p: a cell of one of them that lay in a segment of some frame of z would lie in a segment of the same frame of z', contradicting (i) for z'. So p is a frame of z. Hence z and z' have the same frames, with the same lengths and signs, and the contents of z' are the images under ρ of those of z.

(iv) *Group law.* By (iii), $\psi_\rho\psi_{\rho'} = \psi_{\rho\rho'}$ and $\psi_{\mathrm{id}} = \mathrm{id}$; and ψ_ρ is a block map commuting with σ, since a cell reads the frame containing it in its segment, of length at most $\max\{L : \rho_L \neq \mathrm{id}\} + 6$. So each ψ_ρ is an automorphism. It fixes $0^{\mathbb{Z}}$, which has no frames, and if $\rho_L(u) \neq u$ then ψ_ρ moves the configuration $0^{\mathbb{Z}}$ with $111\,u\,111$ written at 0; so $\rho \mapsto \psi_\rho$ is injective. The frames of $\bar{z}$ are those of z with the signs exchanged and the same contents, so $\psi_\rho(\bar{z}) = \overline{\psi_\rho(z)}$.

(v) *The integral.* Let $\delta = z + \psi_\rho(z)$. It vanishes outside segments, and on the segment $[a, a+L-1]$ of a frame with content u it is $\delta_{a+k} = u_k + \rho_L(u)_k$, for either sign. Define $J(z)_{a+k} = \sum\{\delta_{a+k'} : \ k \leq k' \leq L-1, \ k' \equiv k \pmod 2\}$ for $0 \leq k \leq L-1$ on each segment, and $J(z)_n = 0$ outside segments. This is a block map with $J(0^{\mathbb{Z}}) = 0^{\mathbb{Z}}$, and $J(\bar{z}) = J(z)$, since $\bar{z}$ has the same segments and the same δ. We check $J(z)_n + J(z)_{n+2} = \delta_n$ for every n. If n and $n+2$ lie in one segment, this is the definition. If $n = a+k$ lies in a segment and $n+2$ does not, then $k \geq L-2$, $J(z)_n = \delta_n$, and $n+2$ lies in no segment by (ii), so $J(z)_{n+2} = 0$. If n lies in no segment, then $\delta_n = J(z)_n = 0$, and either $n+2$ lies in no segment, or by (ii) $n+2 \in \{a, a+1\}$ for the start a of a segment; then $J(z)_{n+2}$ is the sum of δ over the even, respectively odd, positions of that segment, which is 0 because u and $\rho_L(u)$ both lie in U_L. Thus $\psi_\rho = \mathrm{id} + (\mathrm{id} + \sigma^2)J$, and $\psi_\rho \in \mathcal{G}^\kappa$.

(vi) *Counting.* For $0 \leq i \leq N$ the word $0\,(10)^{2i+1}\,100\,10\,100\,(10)^{2(N-i)}$ has length $4N+11$, starts and ends with 0, and has no factor 11 or 000. Its 1's lie at the odd positions $1, 3, \ldots, 4i+1$, then at $4i+3$, $4i+6$ and $4i+8$, and then at the $2(N-i)$ odd positions $4i+11, \ldots$; so $4N+2$ of them are at odd and 2 at even positions, and the word lies in U_{4N+11}. These $N+1$ words are distinct. Embedding S_m in $\mathrm{Sym}(U_{4m+7})$ for each m gives $\bigoplus_m S_m \hookrightarrow \bigoplus_L \mathrm{Sym}(U_L) \hookrightarrow \mathcal{G}^\kappa$. □

Example 7.12. A smaller instance uses frames $111\,u\,111$ and $000\,\bar{u}\,000$ with u in $U = \{0010110, 0101010, 0110100\}$, which are not in U_7 but all have even-position sum 0 and odd-position sum 1. For $\rho \in S_3$, ψ_ρ replaces u_i by $u_{\rho(i)}$, and J_ρ is defined on each segment by the formula in Lemma 7.11(v). Exhaustive checks on the binary full shift (computer-assisted;

Appendix A.3) show that distinct frames share no segment cell (two frames can overlap only in a shared run, at offset 10), that ψ_ρ preserves the frames and permutes the contents, on all 2^{31} windows of length 31, that $d + J_\rho(d) + \sigma^2 J_\rho(d) = \psi_\rho(d)$ on all 2^{21} windows, and that $J_\rho(\bar{d}) = J_\rho(d)$. So $\psi_\rho \in \mathcal{G}^\kappa$, and $\rho \mapsto \Phi_{\psi_\rho}$ embeds S_3 in $\mathrm{Aut}(X_{F_{90}})$ and in $\mathrm{Aut}(X_{F_{165}})$ by maps of cell radius 11. The maps $\Phi_{\psi_{(01)}}$ and $\Phi_{\psi_{(12)}}$ do not commute.

The maps Φ_ψ have the form $\mathrm{id} + \iota \circ h \circ u'$ with $u' = d$ a factor map and $h = J_\psi$. In the classification of [30], rules 90 and 165, which are affine, are listed among the classes whose automorphism group contains every finite group.

Corollary 7.13. *For $f \in \{90, 165\}$, F_f has no blocking word, and in particular no word fixed in every context, and $\mathrm{Aut}(X_{F_f})$ is not virtually $\mathbb{Z}^2$.*

Proof. Let u be a word, q a position in it, and x a configuration containing u on $[0, |u|)$. For $N \geq |u| + q$ let $e_N \in \mathbb{F}_2^{\mathbb{Z}}$ be the unit vector at N, and $x' = x + \iota(e_N)$, which differs from x only at the cells N and $N+1$ and so also contains u on $[0, |u|)$. Since F^n differs from F_{90}^n by a constant, Lemma 7.7(c) gives $F^n(x') - F^n(x) = F_{90}^n(\iota(e_N)) = \iota(\sigma^n e_N) = \iota(e_{N-n})$, whose y-bit at q is 1 for $n = N - q$. So $F^{N-q}(x')_q \neq F^{N-q}(x)_q$, and u is not blocking with column q. A word fixed in every context is blocking. The last statement is Theorem 7.10. □

8. Affine rules

The 16 affine elementary rules $f(a, b, c) = \varepsilon \oplus \alpha a \oplus \beta b \oplus \gamma c$ form nine classes:

$$\{0\},\ \{255\},\ \{51, 204\},\ \{90\},\ \{165\},\qquad \{15, 85, 170, 240\},\ \{60, 102\},\ \{153, 195\},\ \{105, 150\}.$$

In each of the first five, $\bigoplus_m S_m$ embeds in $\mathrm{Aut}(X_{F_f})$: for 0 and 51 through fixed walls (Table 1), for 90 and 165 by Theorem 7.10, and for 255 by Proposition 8.10 below. For the last four we prove Theorem 1.4: every automorphism is affine, and the automorphism group is determined. The proof uses the isomorphism rigidity theorem of Kitchens and Schmidt [11], which we state in Section 8.2; everything else is proved here.

8.1. **Affine commuting maps.** Let f be affine, with $\Lambda = \mathbb{F}_2[s, s^{-1}]$, $p = \alpha s^{-1} + \beta + \gamma s$ and $M = \left(\begin{smallmatrix} 0 & 1 \\ 1 & p \end{smallmatrix}\right)$ as in Section 7. Then $F_f(c) = Mc + k_0$ with $k_0 = (0, \varepsilon)^{\mathbb{Z}}$. Put $q = t^2 + pt + 1 \in \Lambda[t]$ and

$$\mathcal{O} = \Lambda[t]/(q).$$

Since $t(t + p) = 1$ in $\mathcal{O}$, t is a unit, and $\mathcal{O}$ is also the quotient of $\mathbb{F}_2[s^{\pm 1}, t^{\pm 1}]$ by q. The ring $\Lambda[M] = \{aI + bM : a, b \in \Lambda\}$ is free over Λ with basis I, M, since the entries of $aI + bM$ are a, b, b, $a + bp$, and $M^2 = pM + I$; so $a + bt \mapsto aI + bM$ is a ring isomorphism $\mathcal{O} \cong \Lambda[M]$, under which σ corresponds to s and the linear part of F_f to t. We use it to identify the two rings. The proof of Lemma 7.1 uses only that M is not scalar, so for every affine rule the matrices in $M_2(\Lambda)$ commuting with M are the elements of $\Lambda[M]$; in particular they commute with each other. On a constant configuration k an element $a \in \Lambda$ acts as the sum $a(1)$ of its coefficients, and $N \in \Lambda[M]$ acts as the 2×2 matrix $N(1)$ over $\mathbb{F}_2$ obtained by putting $s = 1$.

Lemma 8.1. *Let f be affine, $N \in M_2(\Lambda)$ and k a constant configuration. The block map $\varphi(c) = Nc + k$ commutes with F_f if and only if $N \in \Lambda[M]$ and $(N + I)k_0 = (M + I)k$. The automorphisms of X_{F_f} that are affine, that is, of this form, form a group $\mathrm{Aut}_{\mathrm{aff}}(X_{F_f})$, and $\varphi \mapsto N$ is a homomorphism from it to the abelian group $\Lambda[M]^\times$ whose kernel consists of the translations $c \mapsto c + k$ with $(M + I)k = 0$, at most four maps. In particular $\mathrm{Aut}_{\mathrm{aff}}(X_{F_f})$ is metabelian.*

Proof. $\varphi(F_f(c)) = NMc + Nk_0 + k$ and $F_f(\varphi(c)) = MNc + Mk + k_0$; these agree for all c if and only if $NM = MN$ (compare the parts depending on c) and $Nk_0 + k = Mk + k_0$. If φ is invertible, then so is the continuous additive map $c \mapsto Nc = \varphi(c) + k$, which commutes with σ; its inverse is again continuous, additive and commutes with σ, so by Curtis–Hedlund–Lyndon it is a block map with an $\mathbb{F}_2$-linear local rule, that is, a matrix $N' \in M_2(\Lambda)$ with $NN' = N'N = I$. So N is invertible in $M_2(\Lambda)$, the inverse $c \mapsto N'(c+k)$ of φ is again affine and commutes with F_f, and $N' = N^{-1} \in \Lambda[M]$. The composite of $c \mapsto Nc+k$ and $c \mapsto N'c+k'$ has linear part NN', and the map with linear part I and constant k commutes with F_f if and only if $(M + I)k = 0$. □

So a class with $\bigoplus_m S_m \leq \mathrm{Aut}(X_{F_f})$ always has automorphisms that are not affine, since $\bigoplus_m S_m$ is not solvable. This applies to the first five classes above.

8.2. **The rigidity theorem.** Let β be an action of $\mathbb{Z}^d$ by continuous automorphisms of a compact abelian group G, with normalised Haar measure λ_G. The action is *irreducible* if every closed β-invariant subgroup $H \subsetneq G$ is finite, and *mixing* if it is mixing with respect to λ_G. A *measurable conjugacy* between two such actions (G, β) and (G', β') is a measure-preserving map $\varphi\colon (G, \lambda_G) \to (G', \lambda_{G'})$, invertible modulo null sets, with $\varphi \circ \beta^v = \beta'^v \circ \varphi$ almost everywhere for every $v \in \mathbb{Z}^d$.

Theorem 8.2 (Kitchens–Schmidt [11, Theorem 1.1]; see also [16, Theorem 8.4])**.** *Let $d \geq 2$, and let (G, β) and (G', β') be irreducible and mixing actions of $\mathbb{Z}^d$ by continuous automorphisms of compact zero-dimensional abelian groups. Then every measurable conjugacy φ between them is affine almost everywhere: there are a continuous surjective group homomorphism $A\colon G \to G'$ and $g' \in G'$ with $\varphi(g) = A(g) + g'$ for λ_G-almost every g.*

We apply it with $d = 2$ and $G = G' = \mathcal{A}^{\mathbb{Z}} = (\mathbb{F}_2^2)^{\mathbb{Z}}$. The dual group $\widehat{G}$ consists of the characters $\chi_{\xi,\eta}(x, y) = (-1)^{\sum_i (\xi_i x_i + \eta_i y_i)}$ with $\xi, \eta \in \mathbb{F}_2^{\mathbb{Z}}$ finitely supported, and we identify ξ with $\sum_i \xi_i s^i \in \Lambda$.

Lemma 8.3. *Let f be affine with linear part M. Under $\chi \mapsto \chi \circ \sigma$ and $\chi \mapsto \chi \circ M$, the group $\widehat{G}$ is an $\mathbb{F}_2[s^{\pm 1}, t^{\pm 1}]$-module, with s acting as $\chi \mapsto \chi \circ \sigma$ and t as $\chi \mapsto \chi \circ M$, and it is isomorphic to $\mathcal{O}$ with s and t acting by multiplication.*

Proof. For finitely supported ξ and any x, $\sum_i \xi_i x_{i+1} = \sum_i \xi_{i-1} x_i$, so $\chi_{\xi,\eta} \circ \sigma = \chi_{s\xi, s\eta}$; and since $(py)_i = \alpha y_{i-1} + \beta y_i + \gamma y_{i+1}$, $\sum_i \eta_i (py)_i = \sum_i (p\eta)_i y_i$. Hence $\chi_{\xi,\eta}(M(x, y)) = \chi_{\xi,\eta}(y, x + py) = \chi_{\eta, \xi + p\eta}(x, y)$, so $\chi \circ M$ corresponds to the matrix M acting on the column $(\xi, \eta)^{\mathsf{T}} \in \Lambda^2$. The vector $e_1 = (1, 0)^{\mathsf{T}}$ has $Me_1 = (0, 1)^{\mathsf{T}}$, so $a + bt \mapsto (aI + bM)e_1 = (a, b)^{\mathsf{T}}$ is a bijection $\mathcal{O} \to \Lambda^2$ compatible with s and t. □

8.3. **Every automorphism is affine.**

Lemma 8.4. *Let f be affine with linear part M, and suppose:*

- (R1) *$q = t^2 + pt + 1$ is irreducible over $K = \mathbb{F}_2(s)$;*
- (R2) *$s^a t^b \neq 1$ in $\mathcal{O}$ for $(a, b) \neq (0, 0)$;*
- (R3) *there is $m \geq 1$ with $F_f^m = M^m$ such that the subring $S = \Lambda[t^m, t^{-m}]$ of $\mathcal{O}$ has finite index in $\mathcal{O}$ as an additive group.*

Then every automorphism of X_{F_f} is affine: it has the form $c \mapsto Nc+k$ with $N \in \Lambda[M]^\times = \mathcal{O}^\times$ and k a constant configuration with $(N + I)k_0 = (M + I)k$. Consequently $\mathrm{Aut}(X_{F_f}) = \mathrm{Aut}_{\mathrm{aff}}(X_{F_f})$.

For a linear rule (R3) holds with $m = 1$ and $S = \mathcal{O}$.

Proof. Let β be the $\mathbb{Z}^2$-action on $G = \mathcal{A}^{\mathbb{Z}}$ generated by σ and $F_f^m = M^m$, which act by continuous automorphisms. By Lemma 8.3, $\widehat{G}$ is $\mathcal{O}$, with the generators of β acting by multiplication by s and t^m. By (R1), $\mathcal{O}$, which is free over Λ with basis $1, t$, embeds in the field $K[t]/(q)$, so it is a domain.

Irreducibility. The map $t \mapsto t + p$ is a ring automorphism of $\mathcal{O}$, since $t + p$ is again a root of q. For $n = a + bt \neq 0$ put $\bar{n} = a + b(t + p)$; then $n\bar{n} = a^2 + abp + b^2 =: \nu$ lies in Λ and is nonzero, as $\mathcal{O}$ is a domain. So $\nu\mathcal{O} \subseteq n\mathcal{O}$, and $\mathcal{O}/\nu\mathcal{O} \cong (\Lambda/\nu\Lambda)^2$ is finite, since $\Lambda/\nu\Lambda \cong \mathbb{F}_2[s]/(\nu_0)$ for the polynomial ν_0 obtained from ν by removing powers of s. Hence $\mathcal{O}/n\mathcal{O}$ is finite, and so is $n\mathcal{O}/nS \cong \mathcal{O}/S$ by (R3); so $\mathcal{O}/nS$ is finite. Now let $H \subsetneq G$ be closed and β-invariant. Its annihilator $H^\perp \subseteq \widehat{G}$ is nonzero, and it is invariant under $s^{\pm 1}$ and $t^{\pm m}$, hence an S-submodule of $\mathcal{O}$. Choosing $0 \neq n \in H^\perp$ we get $nS \subseteq H^\perp$, so $\widehat{H} \cong \mathcal{O}/H^\perp$ is finite, and H is finite.

Mixing. For characters χ, ψ, $\int (\chi \circ \beta^v)\,\overline{\psi}\, d\lambda_G$ is 1 if $\chi \circ \beta^v = \psi$ and 0 otherwise. If $\chi \neq 1$ corresponds to $0 \neq n \in \mathcal{O}$ and $\chi \circ \beta^{(a,b)} = \chi \circ \beta^{(a',b')}$, then $(s^{a-a'}t^{m(b-b')} - 1)s^{a'}t^{mb'}n = 0$, so $s^{a-a'}t^{m(b-b')} = 1$ and $(a, b) = (a', b')$ by (R2). So for $\chi \neq 1$ the equation $\chi \circ \beta^v = \psi$ has at most one solution v, while for $\chi = 1$ it holds for all v or none according as $\psi = 1$. Hence $\int (\chi \circ \beta^v)\overline{\psi}\, d\lambda_G \to \int \chi\, d\lambda_G \int \overline{\psi}\, d\lambda_G$ as $|v| \to \infty$, for all characters. By linearity this holds for trigonometric polynomials, which are dense in $L^2(\lambda_G)$, and since each β^v preserves λ_G it holds for all L^2 functions; this is mixing.

Haar measure. Haar measure on G is the uniform Bernoulli measure. Let $\varphi \in \mathrm{Aut}(X_{F_f})$, a block map of some radius r that is onto. By Hedlund [7, Theorem 5.4], for an onto block map of the full 4-shift every word of length n has exactly 4^{2r} preimages of length $n + 2r$ under the local rule; so the preimage of a cylinder of length n is a union of 4^{2r} cylinders of length $n + 2r$ and has the same measure 4^{-n}. Hence φ preserves λ_G. It commutes with σ and F_f, hence with β, so it is a measurable conjugacy of (G, β) with itself.

Conclusion. By Theorem 8.2 there are a continuous surjective homomorphism $A\colon G \to G$ and $k \in G$ with $\varphi = A + k$ almost everywhere. Both sides are continuous, and a nonempty open set has positive measure, so $\varphi = A + k$ everywhere. Then $k = \varphi(0^{\mathbb{Z}})$ is fixed by σ, hence constant, and $A = \varphi - k$ commutes with σ and is bijective, since φ is. A continuous additive map of G commuting with σ is a block map with an $\mathbb{F}_2$-linear local rule (Curtis–Hedlund–Lyndon), that is, a matrix $N \in M_2(\Lambda)$. Lemma 8.1 gives $N \in \Lambda[M]$, N invertible there, and the condition on k. □

8.4. **The unit groups.**

Lemma 8.5. (1) *For $p = s^{\pm 1}$ (rules 170 and 240), $\mathcal{O} \cong \mathbb{F}_2[t, t^{-1}, (1 + t)^{-1}]$ with $s^{\pm 1} = (1 + t)^2 t^{-1}$, and $\mathcal{O}^\times = t^{\mathbb{Z}}(1 + t)^{\mathbb{Z}} \cong \mathbb{Z}^2$.*

(2) *For $p = 1 + s^{\pm 1}$ (rules 102 and 60), $\mathcal{O} \cong \mathbb{F}_2[t, t^{-1}, (1 + t + t^2)^{-1}]$ with $s^{\pm 1} = (1 + t + t^2)t^{-1}$, and $\mathcal{O}^\times = s^{\mathbb{Z}}t^{\mathbb{Z}} \cong \mathbb{Z}^2$.*

(3) *For $p = s + 1 + s^{-1}$ (rule 150), $\mathcal{O}^\times = s^{\mathbb{Z}}t^{\mathbb{Z}}(1 + st)^{\mathbb{Z}} \cong \mathbb{Z}^3$.*

In all three cases (R1) *and* (R2) *hold.*

Proof. (1) Take $p = s$; the case $p = s^{-1}$ follows by the automorphism $s \mapsto s^{-1}$ of Λ. Let $B = \mathbb{F}_2[t, t^{-1}, (1 + t)^{-1}]$ and $u = (1 + t)^2 t^{-1} \in B$. The homomorphism $\Lambda[t] \to B$ with $s \mapsto u$, $t \mapsto t$ is onto, since $(1+t)^{-1} = (1+t)t^{-1}u^{-1}$, and it kills q, since $t^2 + ut + 1 = t^2 + (1+t)^2 + 1 = 0$. The induced map $\mathcal{O} \to B$ sends $a + bt$ to $a(u) + b(u)t$, and it is injective: u is transcendental over $\mathbb{F}_2$ and $[\mathbb{F}_2(t) : \mathbb{F}_2(u)] = 2$, so 1 and t are linearly independent over $\mathbb{F}_2(u)$. So $\mathcal{O} \cong B$. The ring B is a localisation of $\mathbb{F}_2[t]$, hence a unique factorisation domain whose units are the products of powers of the inverted primes t and $1 + t$. Since $\mathcal{O}$ is a domain, q is prime in $\Lambda[t]$, and being monic it is irreducible over K by Gauss's lemma; this is (R1). In B, $s^a t^b = (1 + t)^{2a} t^{b-a}$, which is 1 only for $a = b = 0$; this is (R2).

(2) The same argument with $u = (1+t+t^2)t^{-1}$, which satisfies $t^2+(1+u)t+1=0$, and the prime $1+t+t^2$ of $\mathbb{F}_2[t]$; here $(1+t+t^2)^{-1} = t^{-1}u^{-1}$, the units are $t^{\mathbb{Z}}(1+t+t^2)^{\mathbb{Z}} = t^{\mathbb{Z}}(st)^{\mathbb{Z}} = s^{\mathbb{Z}}t^{\mathbb{Z}}$, and $s^at^b = (1+t+t^2)^at^{b-a}$.

(3) Irreducibility (R1) was shown in Section 7. Multiplying q by s, $\mathcal{O}$ is the coordinate ring of the affine curve $s^2t+st^2+st+s+t=0$ in the torus $st\neq 0$. Let $E\subset\mathbb{P}^2$ be the cubic

$$s^2t+st^2+stz+sz^2+tz^2=0.$$

Its partial derivatives in characteristic 2 are t^2+tz+z^2, s^2+sz+z^2 and st; if all vanish, then $s=0$ or $t=0$, and either case forces $s=t=z=0$. So E is smooth, hence an elliptic curve over $\mathbb{F}_2$, geometrically irreducible, with function field the fraction field of $\mathcal{O}$; in particular $\mathbb{F}_2$ is algebraically closed in it, and a function on E without poles is a constant in $\mathbb{F}_2$. The points of E over $\overline{\mathbb{F}}_2$ outside the affine torus $\{z\neq 0,\ s\neq 0,\ t\neq 0\}$ are, since $s=0$ forces $tz^2=0$, $t=0$ forces $sz^2=0$, and $z=0$ forces $st(s+t)=0$,

$$P_0=(0:0:1),\quad P_1=(1:0:0),\quad P_2=(0:1:0),\quad P_3=(1:1:0);$$

these are the four points of $E(\mathbb{F}_2)$. Put $g=1+st$. Its norm is $(1+st)(1+st^{-1}) = 1+sp+s^2 = s$, so $g\in\mathcal{O}^\times$. The divisors of $s=s/z$, $t=t/z$ and $g=(z^2+st)/z^2$ are

$$\operatorname{div} s = P_0+P_2-P_1-P_3,\qquad \operatorname{div} t = P_0+P_1-P_2-P_3,\qquad \operatorname{div} g = P_1+P_2-2P_3.$$

Indeed, the line $s=0$ meets E where $tz^2=0$, that is at P_0 simply and at P_2 doubly; the line $t=0$ meets E at P_0 simply and at P_1 doubly; the line $z=0$ meets E simply at P_1,P_2,P_3. For g: g has no zero or pole on the affine part, it equals 1 at P_0, and its numerator z^2+st is 1 at P_3, so g has a double pole at P_3. At P_1, in the chart $s=1$ the curve is $t(1+t+z+z^2)=z^2$, so z is a local parameter, $t=z^2w^{-1}$ with $w=1+t+z+z^2$, and $z^2+t=z^2(1+w)w^{-1}$ with $1+w=t+z+z^2$ of order 1; so $g=(z^2+t)/z^2$ has a simple zero at P_1, and by the symmetry $s\leftrightarrow t$ of E and g also at P_2.

Let $\mathcal{D}$ be the group of divisors of degree 0 supported on $\{P_0,\dots,P_3\}$, which is free of rank 3, and define $\theta\colon\mathcal{D}\to\mathbb{Z}/4$ by $\theta(\sum n_iP_i) = 2n_0+n_1-n_2$. It vanishes on the three divisors above (on $\operatorname{div} t$ the value is 4), and it is onto. The three divisors span a subgroup $L\subseteq\ker\theta$ of index $|\det| = 4$ in $\mathcal{D}$, in the basis $P_0-P_3, P_1-P_3, P_2-P_3$, where they are $(1,-1,1)$, $(1,1,-1)$ and $(0,1,1)$. Since $\ker\theta$ also has index 4, $L=\ker\theta$. In particular s,t,g generate a free abelian group W of rank 3, and $\operatorname{div}\colon W\to L$ is an isomorphism.

Now let $u\in\mathcal{O}^\times$. Both u and u^{-1} are regular on the affine curve, so $D=\operatorname{div} u\in\mathcal{D}$. The values $\theta(P_i-P_3)$ are $2,1,3,0$ for $i=0,1,2,3$, so there is an i with $D-(P_i-P_3)\in\ker\theta = L$, say $D-(P_i-P_3)=\operatorname{div} w$ with $w\in W$. Then $P_i-P_3=\operatorname{div}(u/w)$ is principal, and on an elliptic curve a divisor $P-P'$ is principal only if $P=P'$ [17, Lemma III.3.3]. So $i=3$, u/w has no zeros or poles, and $u/w\in\mathbb{F}_2^\times=\{1\}$. Hence $\mathcal{O}^\times = W$. Finally (R2) holds since $\operatorname{div}(s^at^b) = (a+b,\,b-a,\,a-b)$ in the basis above vanishes only for $a=b=0$. □

8.5. **The automorphism groups.** For $p(1)=1$, that is, when f has an odd number of linear terms, conjugation by the cellwise complement $\kappa(c)=c+(1,1)^{\mathbb{Z}}$ relates the two members of a pair $\{f,255-f\}$: on constants $M(1)=\left(\begin{smallmatrix}0&1\\1&1\end{smallmatrix}\right)$, so $(M+I)(1,1)^{\mathbb{Z}}=(0,1)^{\mathbb{Z}}$, and hence

$$F_{255-f} = \kappa\, F_f\, \kappa \qquad \text{when } f \text{ is linear and } p(1)=1. \tag{7}$$

Since κ is a homeomorphism commuting with σ, $\varphi\mapsto\kappa\varphi\kappa$ is an isomorphism $\operatorname{Aut}(X_{F_f})\to\operatorname{Aut}(X_{F_{255-f}})$. This relates 170 to 85, 240 to 15 and 150 to 105.

Proof of Theorem 1.4. (1) Let $f\in\{170,240\}$. By Lemma 8.5(1), Lemma 8.4 applies with $m=1$. Since $p(1)=1$, $M(1)+I=\left(\begin{smallmatrix}1&1\\1&0\end{smallmatrix}\right)$ is invertible, so the only constant with $(M+I)k=0$ is $0^{\mathbb{Z}}$, and $\varphi\mapsto N$ is an isomorphism $\operatorname{Aut}(X_F)\to\mathcal{O}^\times = t^{\mathbb{Z}}(1+t)^{\mathbb{Z}}$, under which $F\mapsto t$ and $h=\mathrm{id}+F\mapsto 1+t$. So $(a,b)\mapsto F^ah^b$ is an isomorphism $\mathbb{Z}^2\to\operatorname{Aut}(X_F)$. As $s^{\pm1}=(1+t)^2t^{-1}$,

$h^2 = \sigma^{\pm 1}F$, so $\langle \sigma, F\rangle = \langle F, h^2\rangle$ has index 2. For $f \in \{15, 85\}$ conjugate by κ, using (7): $\kappa h \kappa(c) = c + Mc + M(1,1)^{\mathbb{Z}}$, which is $c + F_f(c) + (1,1)^{\mathbb{Z}}$ because $F_f(c) = Mc + (0,1)^{\mathbb{Z}}$ and $M(1,1)^{\mathbb{Z}} = (1,0)^{\mathbb{Z}}$.

(2) Let $f \in \{60, 102\}$. By Lemma 8.5(2), Lemma 8.4 applies with $m = 1$, and $\mathcal{O}^\times = s^{\mathbb{Z}}t^{\mathbb{Z}}$ consists of the linear parts of the maps $\sigma^a F^b$. Since $p(1) = 0$, $M(1) = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, and the constants with $(M + I)k = 0$ are $0^{\mathbb{Z}}$ and $(1,1)^{\mathbb{Z}}$. So $\varphi \mapsto N$ is onto $\mathcal{O}^\times$, with kernel $\{\mathrm{id}, \kappa\}$, and it is split by $s^a t^b \mapsto \sigma^a F^b$. The involution κ commutes with every automorphism, since $N(1,1)^{\mathbb{Z}} = (1,1)^{\mathbb{Z}}$ for $N = s^a t^b$; so $\mathrm{Aut}(X_F) = \langle \sigma, F\rangle \times \langle \kappa\rangle$.

(3) Let $f \in \{153, 195\}$, with linear parts 102 and 60, so $\mathcal{O}$ and $\mathcal{O}^\times = s^{\mathbb{Z}}t^{\mathbb{Z}}$ are as in (2). On constants $M(1)$ is the transposition of the two bits and $k_0 = (0,1)^{\mathbb{Z}}$, so $(I + M + M^2 + M^3)k_0 = 0$ and $F^4 = M^4$. In $\mathcal{O}$, $t^4 = p^3 t + p^2 + 1$ and $t^{-4} = (t + p)^4 = p^3 t + p^4 + p^2 + 1$, so $S = \Lambda[t^4, t^{-4}]$ is contained in $\Lambda \oplus \Lambda p^3 t$; the latter is a subring, as $(p^3 t)^2 = p^4 \cdot p^3 t + p^6$, and it is generated by $t^{\pm 4}$ over Λ, so $S = \Lambda \oplus \Lambda p^3 t$ and $\mathcal{O}/S \cong \Lambda/p^3\Lambda$, a group of order 8. So Lemma 8.4 applies with $m = 4$. For $N = s^a t^b$, $(N + I)k_0 = (M(1)^b + I)(0,1)$ is 0 for even b and $(1,1)$ for odd b, and $(M + I)(x,y)^{\mathbb{Z}} = (x + y, x + y)^{\mathbb{Z}}$; so for each N there are exactly two constants k, differing by $(1,1)^{\mathbb{Z}}$, and the automorphisms with linear part $s^a t^b$ are $\sigma^a F^b$ and $\kappa\sigma^a F^b$. The involution κ commutes with F (Section 2), and $\kappa \notin \langle \sigma, F\rangle$ since its linear part is I. As in (2), $\mathrm{Aut}(X_F) = \langle \sigma, F\rangle \times \langle \kappa\rangle$.

(4) Let $f = 150$. By Lemma 8.5(3), Lemma 8.4 applies with $m = 1$; as in (1) there is no nonzero constant with $(M + I)k = 0$, so $\varphi \mapsto N$ is an isomorphism onto $\mathcal{O}^\times = s^{\mathbb{Z}}t^{\mathbb{Z}}(1 + st)^{\mathbb{Z}}$, under which $\sigma \mapsto s$, $F \mapsto t$ and $h = \mathrm{id} + \sigma F \mapsto 1 + st$. For $f = 105$ conjugate by κ: $\kappa h \kappa(c) = c + \sigma(Mc) + (1,0)^{\mathbb{Z}} = c + \sigma F_{105}(c) + (1,1)^{\mathbb{Z}}$.

In all four classes every automorphism is affine by Lemma 8.4. The statements for the remaining members follow by reflection, which conjugates F_{170} to F_{240}, F_{85} to F_{15}, F_{102} to F_{60} and F_{153} to F_{195}. □

Remark 8.6. For $f = 170$ the ring $\mathcal{O} \cong \mathbb{F}_2[t^{\pm 1}, (1 + t)^{-1}]$ is also $\mathbb{F}_2[u_1^{\pm 1}, u_2^{\pm 1}]/(1 + u_1 + u_2)$, with $u_1 = t$ and $u_2 = 1 + t$, the dual module of Ledrappier's shift $\{z \in \mathbb{F}_2^{\mathbb{Z}^2} : z_v + z_{v+e_1} + z_{v+e_2} = 0\}$ [14]. By duality, $X_{F_{170}}$ with the action of F and $h = \mathrm{id} + F$ is conjugate to Ledrappier's shift with its two coordinate shifts; explicitly, $c \mapsto (y(F^a h^b(c)_0))_{(a,b)\in\mathbb{Z}^2}$ satisfies the three-dot relation because $h^{b+1} = h^b + Fh^b$. Under this conjugacy $\langle \sigma, F\rangle$ is the restriction of the Ledrappier action to the subgroup of index 2 generated by e_1 and $2e_2 - e_1$, and Theorem 1.4(1) says that $\mathrm{Aut}(X_{F_{170}})$ is the full Ledrappier action.

Remark 8.7. The hypotheses of Lemma 8.4 fail for the five other affine classes, and each of them has automorphisms that are not affine (Lemma 8.1 and the remark after it). For 90 and 165, $q = (t + s)(t + s^{-1})$ is reducible; for 0 and 255, $p = 0$ and $q = (t + 1)^2$; for 51 and 204, $p = 1$ and $q = t^2 + t + 1$ is irreducible over K, but $t^3 = 1$ in $\mathcal{O}$, so (R2) fails and the action is not mixing. So among the affine rules, (R1) and (R2) hold exactly for the classes whose automorphisms are all affine.

Remark 8.8. Lemma 8.4 concerns automorphisms. Whether every block map commuting with F_f, invertible or not, is affine for the classes of Theorem 1.4 is open. By Lemma 8.1 the affine commuting block maps of radius at most R, for $R \geq 1$, form an affine $\mathbb{F}_2$-space of dimension $4R + 1$ for 170, $4R + 2$ for 102, $4R + 1$ for 153 and $4R$ for 150: the linear parts $aI + bM$ of radius at most R have a of degree in $[-R, R]$ and b of degree in $[-R, R - 1]$ for $p = s$ and $p = 1 + s$, and in $[-R + 1, R - 1]$ for 150, as a direct check of the entries a, b, b, $a + bp$ shows; the constants are $0^{\mathbb{Z}}$ only for 170 and 150, and $0^{\mathbb{Z}}$, $(1,1)^{\mathbb{Z}}$ for 102; and for 153 the condition $(N + I)k_0 = (M + I)k$ cuts the linear parts by one affine condition, $a(1) + b(1) = 1$, and leaves two constants for each. For $R \leq 3$ the elimination of Section 3 gives exactly these dimensions

for the space of all commuting block maps (Appendix A.1), so for these rules every commuting block map of radius at most 3 is affine.

8.6. **Rule 255.** For $f = 255$, $F(x, y) = (y, 1 \oplus x)$ permutes the cells as $(0,0) \mapsto (0,1) \mapsto (1,1) \mapsto (1,0) \mapsto (0,0)$. Relabel the cells by $\mathbb{Z}/4$, writing $0, 1, 2, 3$ for these four values in this order; then a configuration becomes $x \in (\mathbb{Z}/4)^{\mathbb{Z}}$, and $F(x) = x + \mathbf{1}$ adds 1 in every cell. So $\mathrm{Aut}(X_{F_{255}})$ is the group of automorphisms of the full shift on $\mathbb{Z}/4$ that commute with adding the constant $\mathbf{1}$, and $\langle \sigma, F\rangle = \langle\sigma\rangle \times \langle F\rangle \cong \mathbb{Z} \times \mathbb{Z}/4$. Let $\Delta\colon (\mathbb{Z}/4)^{\mathbb{Z}} \to (\mathbb{Z}/4)^{\mathbb{Z}}$, $\Delta(x)_i = x_{i+1} - x_i$, that is $\Delta = \sigma - \mathrm{id}$. It is an onto block map whose fibres are the cosets of the four constant configurations.

Definition 8.9. Let $\mathcal{G}_4$ be the set of automorphisms ψ of the full shift $(\mathbb{Z}/4)^{\mathbb{Z}}$ for which there is a block map J of $(\mathbb{Z}/4)^{\mathbb{Z}}$ with $\psi = \mathrm{id} + (\sigma - \mathrm{id})J$.

Every $\psi \in \mathcal{G}_4$ preserves the sum over a period of every spatially periodic configuration, since $(\sigma - \mathrm{id})J(z)$ has sum 0 over a period of z.

Proposition 8.10. *Let $F = F_{255}$. For $\varphi \in \mathrm{Aut}(X_F)$ there is a unique $\tilde\varphi \in \mathcal{G}_4$ with $\Delta \circ \varphi = \tilde\varphi \circ \Delta$, and a unique $\epsilon(\varphi) \in \mathbb{Z}/4$ with $\varphi(0^{\mathbb{Z}}) = \epsilon(\varphi)^{\mathbb{Z}}$. The map $\varphi \mapsto (\epsilon(\varphi), \tilde\varphi)$ is an isomorphism*

$$\mathrm{Aut}(X_{F_{255}}) \;\cong\; \mathbb{Z}/4 \times \mathcal{G}_4,$$

with $\mathbb{Z}/4 = \langle F\rangle$. Moreover $\bigoplus_{m\geq 1} S_m$ embeds in $\mathcal{G}_4$, so $\mathrm{Aut}(X_{F_{255}})$ contains every finite group.

Proof. The map $\varphi \mapsto \tilde\varphi$. Since $\varphi(x + \mathbf{c}) = \varphi(x) + \mathbf{c}$ for every constant $\mathbf{c}$, the configuration $\varphi(x) - x$ depends only on the coset $x + \{\text{constants}\}$, that is on Δx: there is a map J with $\varphi(x) = x + J(\Delta x)$. It commutes with σ, and $J(z)_0 = \varphi(x)_0 - x_0$ depends on $x_{[-r,r]}$ only up to adding a constant, that is on $z_{[-r,r-1]}$; so J is a block map. Then $\Delta\varphi(x) = \Delta x + (\sigma - \mathrm{id})J(\Delta x)$, so $\tilde\varphi = \mathrm{id} + (\sigma - \mathrm{id})J$ satisfies $\Delta\varphi = \tilde\varphi\Delta$, and it is unique because Δ is onto. It is onto because φ is, and injective: if $\tilde\varphi(\Delta x) = \tilde\varphi(\Delta x')$, then $\varphi(x') = \varphi(x) + \mathbf{c} = \varphi(x + \mathbf{c})$ for a constant $\mathbf{c}$, so $x' = x + \mathbf{c}$ and $\Delta x' = \Delta x$. So $\tilde\varphi \in \mathcal{G}_4$, and $\varphi \mapsto \tilde\varphi$ is a homomorphism, again because Δ is onto.

Kernel. If $\tilde\varphi = \mathrm{id}$, then $(\sigma - \mathrm{id})J = 0$, so every $J(z)$ is constant. If J has radius r, then for any z, z' and large n a configuration z'' agreeing with z on $[-r, r]$ and with z' on $[n - r, n + r]$ gives $J(z)_0 = J(z'')_0 = J(z'')_n = J(z')_n = J(z')_0$, so J is a constant $\mathbf{c}$ and $\varphi = F^c$. So the kernel is $\langle F\rangle \cong \mathbb{Z}/4$.

Onto $\mathcal{G}_4$. Let $\psi = \mathrm{id} + (\sigma - \mathrm{id})J \in \mathcal{G}_4$ and $\varphi(x) = x + J(\Delta x)$. This is a block map commuting with σ and with adding constants, and $\Delta\varphi = \psi\Delta$. It is injective: $\varphi(x) = \varphi(x')$ gives $\psi(\Delta x) = \psi(\Delta x')$, so $x' = x + \mathbf{c}$, and then $\varphi(x') = \varphi(x) + \mathbf{c}$ forces $\mathbf{c} = 0$. It is onto: given w, choose x with $\Delta x = \psi^{-1}(\Delta w)$; then $\Delta\varphi(x) = \Delta w$, so $\varphi(x) = w + \mathbf{c}$ and $\varphi(x - \mathbf{c}) = w$. So $\varphi \in \mathrm{Aut}(X_F)$ and $\tilde\varphi = \psi$.

Splitting. $\varphi(0^{\mathbb{Z}})$ is fixed by σ, hence constant, and $\varphi\chi(0^{\mathbb{Z}}) = \varphi(\epsilon(\chi)^{\mathbb{Z}}) = \varphi(0^{\mathbb{Z}}) + \epsilon(\chi)^{\mathbb{Z}}$, so ϵ is a homomorphism to $\mathbb{Z}/4$; it maps F^c to c, so it is an isomorphism on the kernel $\langle F\rangle$, which is central. Hence $\mathrm{Aut}(X_F) = \langle F\rangle \times \ker\epsilon$, and $\varphi \mapsto \tilde\varphi$ maps $\ker\epsilon$ isomorphically onto $\mathcal{G}_4$.

Markers. For $L \geq 1$ let V_L be the set of words $u \in \{1,2,3\}^L$ with $\sum_k u_k \equiv 0 \pmod 4$. Call a *segment* of $z \in (\mathbb{Z}/4)^{\mathbb{Z}}$ of length L an interval $[a, a + L - 1]$ on which z has no 0, with $z_{a-1} = z_{a+L} = 0$; distinct segments are disjoint and separated by a 0. For $\rho = (\rho_L)_L$ with $\rho_L \in \mathrm{Sym}(V_L)$, all but finitely many the identity, let $\psi_\rho(z)$ replace the content u of every segment of length L with $u \in V_L$ by $\rho_L(u)$. It changes only nonzero cells to nonzero values, so it preserves the set of zeros, hence the segments, their lengths and the property $u \in V_L$. So $\psi_\rho\psi_{\rho'} = \psi_{\rho\rho'}$, each ψ_ρ is an automorphism of the full shift, a block map of radius at most $\max\{L : \rho_L \neq \mathrm{id}\}$, and $\rho \mapsto \psi_\rho$ is injective, since ψ_ρ moves the configuration $0^\infty u\, 0^\infty$ when $\rho_L(u) \neq u$. Let $\delta = \psi_\rho(z) - z$; on a segment $[a, a + L - 1]$ it has sum 0 modulo 4, because u and $\rho_L(u)$ lie in V_L, and it vanishes off segments. Put $J(z)_{a+j} = \sum_{0\leq k<j} \delta_{a+k}$ for $0 \leq j \leq L - 1$ on every segment, and $J(z)_n = 0$

off segments. This is a block map, and $J(z)_{n+1} - J(z)_n = \delta_n$ for every n: inside a segment by definition, at its last cell because the sum of δ over the segment is 0, and at the 0 before a segment because J vanishes at the first cell of a segment. So $\psi_\rho = \mathrm{id} + (\sigma - \mathrm{id})J \in \mathcal{G}_4$, and $\bigoplus_L \mathrm{Sym}(V_L)$ embeds in $\mathcal{G}_4$. The 2^m words of length $2m$ made of the blocks 13 and 31 lie in V_{2m}, so S_m embeds in $\mathrm{Sym}(V_{2m})$, and $\bigoplus_m S_m$ embeds in $\mathcal{G}_4$. □

So $\mathrm{Aut}(X_{F_{255}})$ contains every finite group and is not virtually $\mathbb{Z}^n$ for any n. Through $\varphi \mapsto \tilde{\varphi}$, its automorphism group is, up to the central factor $\langle F \rangle$, the group $\mathcal{G}_4$ of automorphisms of the full shift on $\mathbb{Z}/4$ whose displacement $\psi - \mathrm{id}$ is a difference $(\sigma - \mathrm{id})J$ of a block map.

9. Open questions

(1) Is $\mathrm{Aut}(X_{F_{54}})$ generated by σ, F, the wall relabellings of Theorem 4.2 and the wall-trading involutions of the kind in Theorem 4.6, of all radii? At radius 2 the relabellings suffice (Theorem 4.4); at radius 3 they do not, by Theorem 4.6. A complete count of $\mathcal{C}_3^\times(F_{54})$ would be a first test.

(2) Determine $\mathrm{Aut}(X_{F_{90}})$. Its elements of radius at most 2 are affine, but it contains $(\bigoplus_m S_m)^2$ (Theorem 7.10). Is it generated by the affine automorphisms and the maps Φ_ψ, Φ'_χ of that theorem?

(3) For the affine classes of Theorem 1.4, is every block map commuting with F_f affine? Every automorphism is (Lemma 8.4), and every commuting block map of radius at most 3 is (Remark 8.8).

(4) In every example of this paper with a nonlinear rule, the automorphisms outside $\langle \sigma, F \rangle$ come from a blocking word with a degree of freedom next to the blocking column. For a rule that is not affine, the absence of blocking words does not force the automorphism group to be virtually $\mathbb{Z}^2$: the class of 95 is a counterexample [31]. Question 1.6 asks whether it is the only one among the elementary rules.

Appendix A. Computations

This appendix describes the finite computations used in the paper. Every result marked as computer-assisted in the text is the output of a terminating exact search over a finite set; where random sampling is used, this is said, and such runs serve only as checks. The code and its output accompany the paper. Section A.1 describes the censuses of commuting block maps, Section A.2 the computations on which proofs rely, Section A.3 the computations whose results are stated in the text without a proof by hand, and Section A.4 computational confirmations of results that are proved by hand. Section A.5 lists running times.

A.1. **The censuses.** The censuses of Theorem 4.4 and Proposition 6.16, and the dimension counts of Theorem 7.2(c), Proposition 7.4 and Remark 8.8, use the encoding, the enumeration, the re-verification and the invertibility test of Section 3.

Rule 54 *at radius* 2. The full centralizer $\mathcal{C}_2(F_{54})$ exceeds the enumeration cap of 2×10^4 solutions, so the balanced commuting maps were enumerated, with the balance condition as native cardinality constraints; there are 23, and all pass the pair-graph test.

Rule 54 *at radius* 3. The enumeration of $\mathcal{C}_3^\times(F_{54})$ described in Section 4.6 adds to the CNF the injectivity of g on all spatially periodic configurations of period at most 5 and learns necessary conditions lazily (diamonds of length 13, and collisions of periodic configurations of period at most 10), shared among the worker processes; the balanced commuting maps number more than 2×10^4. The enumeration is not complete, and the paper uses from it only the qualitative statements of Section 4.6.

Rule 59 *at radius* 2 (Proposition 6.16). $\mathcal{C}_2^{\times}(F_{59})$ was enumerated as in Section 3; each of the 133 solutions passed the pair-graph test, and each was matched against the tables of the maps (i)–(iii) of the proposition, computed by exact composition.

Rule 90 *at radius* 2. The invertibility of the 2^{17} commuting maps was decided with the balance condition and the pair-graph test, in 5,851 seconds: 98,222 of them are balanced, and 26 are invertible, all affine.

Linear rules. The dimensions of $\mathcal{C}_R(F_{90})$ and $\mathcal{C}_R(F_{150})$ for $R \leq 3$ were computed by the elimination of Section 3, and for $R \leq 2$ cross-checked against the SAT enumeration of all solutions. For Theorem 7.2(c) the tables of the three families were built directly, each re-verified to commute, and their span was computed by a rank computation over $\mathbb{F}_2$.

Affine rules (Remark 8.8). For $f \in \{170, 102, 153, 150\}$ and $R \leq 3$ the commutation condition, an affine system over $\mathbb{F}_2$ in the $2 \cdot 4^{2R+1}$ table bits (with a constant term for 153), was solved by elimination in an implementation independent of the SAT encoding. The solution spaces are consistent and have dimensions $5, 9, 13$ for 170, $6, 10, 14$ for 102, $5, 9, 13$ for 153 and $4, 8, 12$ for 150, at $R = 1, 2, 3$, which are the dimensions $4R + 1$, $4R + 2$, $4R + 1$ and $4R$ of the affine commuting maps. By reflection the same holds for 240, 60 and 195, and by the conjugation (7) for 85, 15 and 105.

A.2. **Computations used in proofs.** *Fixed words of all lengths* (Proposition 5.2(3) and Corollary 1.2). For each of the 256 rules and each $k \leq 40$, the words $w \in \{0,1\}^k$ that are fixed in every context are found as walks in the graph on the four pairs (w_j, w_{j+1}), as in the proof. Fixed words exist exactly for the 81 rules of Table 1 at length at most 3, and, among the other rules, only for 133 and 161, at the lengths 4, 7, 8 and every k with $10 \leq k \leq 40$. A brute-force search over all of $\mathcal{A}^k$, $k \leq 4$, agrees with the reduction to $\{0,3\}^k$. For each rule of Table 1, the set C of Theorem 1.1 for the listed word has at least two elements.

A.3. **Computer-assisted statements.** The following results are stated in the text as computations. For each we give the method in brief.

Walls and searches (Sections 6.2 and 6.5). Words fixed or time-periodic in every context, and invariant families of type (W1), are found by computing, for each word, its images under $F^{\pm p}$ in all contexts. The search for invariant families with a hidden involution satisfying (W1) and (H1), with windows of length at most 6, considers the closure of each word under F and F^{-1} in all contexts when that closure has at most 300 words. For 59R and $p \in \{6, 12, 30, 60, 120\}$, a search finds, for every word of $X_A \cup X_B$, a context in which F^p changes the protector. The overlaps of walls of rule 59 at distances 2 and 3 were found on all words of length at most 9.

The marker example (Example 7.12). The statements about frames are checked exhaustively on all binary words of length 31 and 21, as stated there.

Rule 219 (after Proposition 6.24). The radius-6 variant of Ψ_2 is checked to commute with F on all 4^{15} windows and to be an involution.

Linear rules (Remark 7.6). The graphs Λ_M are built directly from the local rule and tested for strong connectivity, for $M \leq 4$ for the affine left-permutive rules listed there, and for $M \leq 8$ for rule 90.

Rule 54 (after Theorem 4.2 and Table 2). The orders o_L of F_L and the periods of the content 0^L were computed by iterating F_L on S_L. The spreading data of Table 2 come from simulation on random configurations, as stated in its caption.

A.4. **Confirmations of results proved by hand.** The following computations check results that the paper proves by hand. None of them is used in a proof.

Walls. The construction of Theorem 1.3 was checked on 3,000 random cyclic configurations of length about 60 containing trains of 2 to 4 cars: $\Psi_m(\sigma)$ commuted with F, was inverted by

$\Psi_m(\sigma^{-1})$ and composed as a homomorphism, and it acted nontrivially on 476 of them. For each rule of Table 1, the construction of Theorem 1.1 was checked on 1,500 random cyclic configurations. The table of τ (Theorem 4.6) differs from that of the identity on exactly 24 of the 16,384 entries, commutes with F on all 4^9 windows, composes with itself to the identity on all 4^{13} windows of length 13, and differs from $D\tau$ on 147,456 of them; the pair-graph test reports τ injective, and τ is injective on all spatially periodic configurations of period at most 10. The automorphism D of Proposition 5.2 commutes with F on all 4^{11} windows of length 11.

Hidden and rotating walls. For Proposition 6.16, the elements of T_R were composed exactly on windows of radius 4, and the eight generators were checked to commute with F exhaustively at radius 3. For 59R (Theorem 6.14), on rings of length 2000 seeded with trains of length up to 6, with all $m = 2, \dots, 6$ permuted at once, $F\Psi = \Psi F$, $F^{-1}\Psi = \Psi F^{-1}$ and the homomorphism law held for 60 steps, and the radius bound $2m + 2$ held when all cells outside the window were re-randomized. For 127R, 179R and 223R (Proposition 6.19), the flip of all walls was composed exactly at radius 4 and passed the pair-graph test, and rings of length 1500 seeded with trains at gaps 5, 6 and 8, with walls of other classes mixed in, showed no failure of commutation with F and F^{-1} or of the homomorphism law. For 27R (Theorem 6.20), (W1) and (H1) were checked for the 12 words in all 16 contexts, for F and F^{-1}; (H2) and (H4) on all pairs of occurrences at distances 3, 4, 5, that is, on words of length at most 11; over all 16 contexts, β and $\beta + 1$ are the only maps $W_{27} \to \mathbb{Z}/2$ with $\beta \circ \iota = \beta + 1$ that satisfy (H3) for some ε; and simulations on rings with trains at gaps 3, 6 and 7 showed no failure of commutation or of the homomorphism law. For 219R and 126R (Theorem 6.23 and Proposition 6.25), the hypotheses were checked on all contexts by two implementations; the bond invariance, the π-evolution of bonded cells, the maps φ_s and R_s, and the commutation of D and E with F on all 4^7 windows were checked, D and E are injective by the pair-graph test, and Ψ_1, Ψ_2, Ψ_3 were tested on planted rings.

Markers for $90R$ *and* $165R$ (Theorem 7.10). The maps Φ and Φ' were checked on all cyclic configurations of periods 10 and 11 and on planted random rings for commutation with F, the group law and the commutation of the two movers, with no failures; as a negative control, for 165R the frames $111\,u\,111$ alone, whose integral is not complement-invariant, give commutation failures. The construction of Lemma 7.11 with $N = 2$, three words of length 19, was checked in the same way.

Theorem 2.4 and Lemma 2.3. Parts (1) and (2) were checked by exhaustion for all 256 rules and $n \le 4$, and the construction of Lemma 2.3 with random staircase data for all 240 rules reading c and $n \le 31$.

A.5. **Running times and reproducibility.** The censuses were run on one core of a machine with 32 GB of memory under Linux, using Python 3.12 or later, NumPy and PySAT 1.9 with CaDiCaL 1.5.3 and MiniCard. Table 4 lists instance sizes and times. The 54R radius-2 invertible census used MiniCard's native cardinality constraints for the balance condition. The code for all computations of this appendix accompanies this paper, with its logs and with the solution tables of the complete enumerations for 54R at radius 1, for the 23 balanced maps of 54R at radius 2, and for 59R; the larger solution sets are regenerated by the programs.

TABLE 4. Instance sizes and running times (encode / enumerate / classify, in seconds).

instance	variables	clauses	solutions	time (s)
54R, $R = 2$, balanced	$2{,}048 + 4{,}096$	2.9×10^5+card.	23	3 / 11 / –
90R, $R = 2$, pair-graph test	—	—	2^{17} maps tested	5,851
90R, $R = 3$, elimination	4,096 classes	2.6×10^5 eqs	2^{65}	4

Use of AI tools. Large language models (Anthropic's Claude models) were used extensively in this work: to propose and develop proofs, to design, write and run the computer searches and their independent re-implementations, to check proofs and computations, and to draft the manuscript. The models are not authors. The author has reviewed the content and takes full responsibility for it. All code and computational output accompany the paper (Appendix A).

References

[1] M. Baake, J. A. G. Roberts and R. Yassawi, Reversing and extended symmetries of shift spaces, *Discrete Contin. Dyn. Syst.* **38** (2018), 835–866.

[2] A. Biere, K. Fazekas, M. Fleury and M. Heisinger, CaDiCaL, Kissat, Paracooba, Plingeling and Treengeling entering the SAT Competition 2020, in *Proc. SAT Competition 2020*, Univ. Helsinki, 2020, 51–53.

[3] F. Blanchard and P. Tisseur, Some properties of cellular automata with equicontinuity points, *Ann. Inst. H. Poincaré Probab. Statist.* **36** (2000), 569–582.

[4] N. Boccara, J. Nasser and M. Roger, Particlelike structures and their interactions in spatiotemporal patterns generated by one-dimensional deterministic cellular-automaton rules, *Phys. Rev. A* **44** (1991), 866–875.

[5] M. Boyle, D. Lind and D. Rudolph, The automorphism group of a shift of finite type, *Trans. Amer. Math. Soc.* **306** (1988), 71–114.

[6] V. Cyr, J. Franks and B. Kra, The spacetime of a shift endomorphism, *Trans. Amer. Math. Soc.* **371** (2019), 461–488.

[7] G. A. Hedlund, Endomorphisms and automorphisms of the shift dynamical system, *Math. Systems Theory* **3** (1969), 320–375.

[8] M. Hochman, On the automorphism groups of multidimensional shifts of finite type, *Ergodic Theory Dynam. Systems* **30** (2010), 809–840.

[9] A. Ignatiev, A. Morgado and J. Marques-Silva, PySAT: a Python toolkit for prototyping with SAT oracles, in *Theory and Applications of Satisfiability Testing (SAT 2018)*, Lecture Notes in Comput. Sci. 10929, Springer, 2018, 428–437.

[10] B. Kitchens and K. Schmidt, Automorphisms of compact groups, *Ergodic Theory Dynam. Systems* **9** (1989), 691–735.

[11] B. Kitchens and K. Schmidt, Isomorphism rigidity of irreducible algebraic $\mathbb{Z}^d$-actions, *Invent. Math.* **142** (2000), 559–577.

[12] J. Kopra, Glider automata on all transitive sofic shifts, *Ergodic Theory Dynam. Systems* **42** (2022), 3716–3744.

[13] P. Kůrka, Languages, equicontinuity and attractors in cellular automata, *Ergodic Theory Dynam. Systems* **17** (1997), 417–433.

[14] F. Ledrappier, Un champ markovien peut être d'entropie nulle et mélangeant, *C. R. Acad. Sci. Paris Sér. A-B* **287** (1978), A561–A563.

[15] J. P. Ryan, The shift and commutativity, *Math. Systems Theory* **6** (1972), 82–85.

[16] K. Schmidt, Algebraic $\mathbb{Z}^d$-actions, PIMS Distinguished Chair Lectures, University of Victoria, 2002.

[17] J. H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd ed., Grad. Texts in Math. 106, Springer, New York, 2009.

[18] S. Takesue, Reversible cellular automata and statistical mechanics, *Phys. Rev. Lett.* **59** (1987), 2499–2502.

[19] S. Takesue, Ergodic properties and thermodynamic behavior of elementary reversible cellular automata. I. Basic properties, *J. Stat. Phys.* **56** (1989), 371–402.

[20] T. Toffoli and N. Margolus, Invertible cellular automata: a review, *Phys. D* **45** (1990), 229–253.

[21] T. Ward, Automorphisms of $\mathbb{Z}^d$-subshifts of finite type, *Indag. Math. (N.S.)* **5** (1994), 495–504.

[22] V. Salo, Transitive action on finite points of a full shift and a finitary Ryan's theorem, *Ergodic Theory Dynam. Systems* **39** (2019), 1637–1667.

[23] K. H. Kim and F. W. Roush, On the automorphism groups of subshifts, *Pure Math. Appl. Ser. B* **1** (1990), no. 4, 203–230.

[24] G. J. Martínez, A. Adamatzky, F. Chen and L. Chua, On soliton collisions between localizations in complex elementary cellular automata: rules 54 and 110 and beyond, *Complex Systems* **21** (2012), 117–142; arXiv:1301.6258.

[25] A. Bobenko, M. Bordemann, C. Gunn and U. Pinkall, On two integrable cellular automata, *Comm. Math. Phys.* **158** (1993), 127–134.

[26] B. Buča, K. Klobas and T. Prosen, Rule 54: exactly solvable model of nonequilibrium statistical mechanics, *J. Stat. Mech.* (2021), 074001.

[27] S. Wolfram, *A New Kind of Science*, Wolfram Media, Champaign, IL, 2002.

[28] M. Dayah, Reversible rule 110 has no hidden symmetries: six families of second-order cellular automata settled by edge moves, in preparation.
[29] M. Dayah, Fronts at the speed of light: rigidity of permutive second-order cellular automata, in preparation.
[30] M. Dayah, The 88 second-order elementary cellular automata: a classification by automorphism group, in preparation.
[31] M. Dayah, The cellular automata that commute with Rule 54, and rigidity of the second-order automaton 94R, in preparation.

Email address: michael@dayah.com