GLOBAL HEALTH ANALYSIS

Sexually Transmitted Diseases Dashboard

Global Analysis Report | 2015 – 2025

HIV/AIDs | HPV | Hepatitis B | Syphilis

Towards Awareness, Prevention and Control

Contents

01 **Objectives**
What we set out to analyze and why it matters

02 **Global Overview**
Total cases, deaths and new infections 2015–2025

03 **Disease Breakdown**
HIV/AIDs · Hepatitis B · HPV · Syphilis

04 **Regional Distribution**
Burden across Africa, Asia, Americas, Europe & Oceania

05 **Trends Over Time**
Annual new cases and death trends

06 **Treatment Coverage**
How well are patients being reached?

07 **Key Insights**
Data-driven findings from the analysis

08 **Recommendations**
Strategic actions for stakeholders

Objectives

What This Report Sets Out to Achieve

1. **Analyze Disease Burden**
   Quantify total cases, new infections and deaths across 4 major STDs from 2015 to 2025 across the globe.

2. **Identify High-Risk Regions**
   Pinpoint which regions and countries carry the highest disease burden and why action is most urgent.

3. **Understand Trends Over Time**
   Track how new cases and mortality rates have shifted year-over-year to identify patterns and turning points.

4. **Evaluate Treatment Reach**
   Assess treatment coverage rates to determine how effectively health systems are responding to each disease.

5. **Inform Strategic Action**
   Provide data-driven recommendations for policymakers, health organizations and stakeholders to reduce impact.

Global Overview | 2015 - 2025

Key Figures at a Glance

336.9M
Total Cases Recorded

15.7M
Total Deaths

38.9M
Total New Infections

5
Regions Covered

4
Diseases Tracked

11
Years of Data

~78%
Avg Treatment Coverage

Key Takeaway: Despite ongoing global health efforts, STDs continue to impose a massive burden. With over 336 million cumulative cases recorded across 5 regions, the data underscores the urgent need for sustained investment in testing, treatment and prevention programmes.

Disease Breakdown

Four Major STDs Under the Microscope

HIV / AIDs

Highest total cases & deaths

Cases: 102.9M **Deaths: 4.99M**

The largest STD burden globally. HIV remains most prevalent in Africa and Asia, with Sub-Saharan Africa carrying a disproportionate share. Treatment coverage averages 77.9%, leaving millions without access to antiretroviral therapy.

HPV

Most common among youth

Cases: 85.2M **Deaths: 3.29M**

The most common viral STD. HPV disproportionately affects young females aged 15–24. Vaccines are available but coverage varies significantly across regions, particularly in low-income settings. Treatment coverage: 78.5%.

Hepatitis B

2nd highest disease burden

Cases: 96.0M **Deaths: 4.91M**

Close behind HIV in scale. Hepatitis B is highly prevalent in Asia where vaccination coverage remains inconsistent. Untreated, it leads to liver disease and cancer. Average treatment coverage: 78.4%.

Syphilis

Lowest burden but high risk

Cases: 52.8M **Deaths: 2.55M**

Though lowest in total cases, Syphilis remains a serious concern due to complications if untreated, including neurological damage and increased HIV risk. Treatment coverage averages 77.9% globally.

Regional Distribution of STD Cases

Which Regions Carry the Highest Burden?



Why Asia Leads
Asia's high burden is driven by large populations in India and China with dense urban centers and varying access to healthcare and STD screening.

Africa's Challenge
Africa faces high HIV/AIDs rates, limited treatment infrastructure and stigma around STD testing, particularly in Sub-Saharan nations.

The Americas
The Americas sees high HPV and Syphilis rates in Brazil and the USA, offset by stronger healthcare systems and broader vaccination access.

New Cases & Death Trends | 2015 - 2025

How Have Infections & Mortality Shifted Over the Decade?

Annual New Cases (millions)



Annual Deaths (millions)



▲ 2016 & 2024 saw the highest new case spikes — possible surveillance surges or outbreak events.

▼ 2020 recorded the lowest deaths (1.17M) — likely due to COVID-era healthcare disruptions reducing reported STD deaths.

● New cases show no consistent decline, signaling that prevention efforts need to be significantly scaled up.

Treatment Coverage & Health System Response

Are Patients Getting the Care They Need?

77.9%	78.5%	78.4%	77.9%
HIV / AIDs	**HPV**	**Hepatitis B**	**Syphilis**
Treatment Coverage	Treatment Coverage	Treatment Coverage	Treatment Coverage

TREATMENT GAP ANALYSIS

~22%	~22%	~22%	~22%
untreated	untreated	untreated	untreated
≈22.6M people not receiving treatment globally	Vaccine access is limited in low-income regions	Despite effective antivirals, coverage gaps persist	Penicillin-based treatment is cheap yet underutilised

Key Insights from the Data

01 HIV/AIDs Remains the Primary Threat

With over 102.9M total cases and 4.99M deaths, HIV/AIDs accounts for the largest share of the global STD burden. Africa and Asia together represent the core epicentres of this epidemic.

02 No Consistent Downward Trend in New Cases

Annual new infections fluctuated between 2.9M and 4.5M across the decade, with 2016 and 2024 being peak years. This signals that current prevention strategies have not achieved sustained reduction.

03 Asia Carries the Highest Regional Burden

Asia recorded 108.5M total cases — nearly 32% of the global total — driven by densely populated countries like India and China where healthcare access and STD awareness remain uneven.

04 Treatment Coverage Plateaus at ~78%

All four diseases show treatment coverage hovering around 78%, suggesting a structural ceiling. The remaining 22% gap represents millions of untreated individuals who continue to drive transmission.

05 2020 COVID Effect on Reported Deaths

Deaths in 2020 dropped to their lowest point (1.17M), likely reflecting disrupted healthcare reporting during the COVID-19 pandemic rather than a genuine reduction in STD mortality.

06 HPV Disproportionately Impacts Young Females

HPV cases are concentrated in the 15–24 female age group globally. Vaccine availability is unequal, with high-income nations achieving far greater coverage than low-income regions.

Recommendations

Strategic Actions for Stakeholders & Policymakers

1 Scale Up Testing Infrastructure

Governments must invest in free, accessible STD testing in high-burden regions especially Africa and Asia. Mobile testing units and community clinics can reach underserved populations.

2 Close the Treatment Coverage Gap

A 22% treatment gap across all diseases must be urgently addressed. Subsidised antiretroviral therapy, Hepatitis B antivirals and Syphilis treatments must reach last-mile communities.

3 Expand HPV Vaccination in Youth

National vaccination programmes must prioritise adolescent girls aged 9–14. School-based vaccination initiatives in low-income countries can dramatically reduce future HPV burden.

4 Launch Targeted Awareness Campaigns

Combat stigma around STD testing through culturally sensitive education. Digital platforms, community leaders and health workers are key delivery channels for awareness messages.

5 Improve Cross-Border Data Reporting

Inconsistent reporting, especially during global disruptions like COVID-19, masks the true burden. A standardised WHO-aligned reporting system should be adopted across all regions.

6 Fund STD Research & Innovation

Sustained investment in next-generation vaccines, rapid diagnostics and drug resistance research is critical to getting ahead of STD transmission curves over the next decade.

Thank You

Together We Can Reduce the STD Burden

Data Source: Global Health Dataset | Period: 2015 – 2025 | Diseases: HIV/AIDs, HPV, Hepatitis B, Syphilis

336.9M	15.7M	~78%
Total Cases	Total Deaths	Avg Treatment Coverage

Towards Awareness, Prevention and Control