HEALTH ANALYTICS

Smoking Health Risk Analysis

A comprehensive assessment of smoking-related health impacts across 2,500 patients — presented for executive review

2,500
Total Patients

35.9%
Damaged Organs

29.3%
Current Smokers

This analysis examines the health outcomes of 2,500 patients across multiple organ systems, evaluating the relationship between smoking history, duration, and disease burden — providing critical insights for targeted intervention strategies.

2,500

Patients Analyzed

Avg Age 54 • Avg BMI 29.6

35.9%

Organ Damage Rate

897 patients affected

29.3%

Current Smokers

733 active users

25.2%

High BP Risk

629 at elevated risk

KEY FINDINGS

- Current smokers average 22.2 years of smoking history and 16.8 cigarettes per day — indicating deeply entrenched dependency.
- Organ damage is significantly concentrated among smoking populations, with lungs, heart, and kidneys at highest risk.
- Nearly 30% of patients have a positive family history risk, compounding the impact of active smoking behavior.
- Both male and female patients show similar smoking distribution — the health risk burden is gender-neutral.

SMOKING STATUS DISTRIBUTION

% of Smoking Status



Never Smoked Former Smoker Current Smoker

BREAKDOWN BY CATEGORY

Never Smoked
1136 patients
45.4%

Former Smoker
631 patients
25.2%

Current Smoker
733 patients
29.3%

Insight
Over half the dataset has never smoked or has quit — suggesting interventions for former smokers could yield measurable health recovery benefits.

SMOKING BY GENDER & AGE GROUP





Male – Current

374

29.6% of all males

Female – Current

359

29.0% of all females

Avg Smoking Duration

22.2 yrs

Among current smokers

Avg Cigarettes/Day

16.8

Among current smokers

ORGAN DAMAGE & HEALTH RISK PROFILE





35.9%
Organ Damage Rate
897 out of 2,500 patients

25.2%
High BP Risk
629 patients with elevated BP

29.7%
Family History Risk
742 patients — compounding factor

45.4%
Healthy & Never Smoked
Protection benchmark group

⚠ CRITICAL RISK FINDING:

Current smokers have the highest organ damage concentration. Patients with both active smoking and family history risk represent a compounded vulnerability tier requiring immediate clinical intervention.

STRATEGIC RECOMMENDATIONS

Based on our analysis of 2,500 patients, we recommend the following actionable steps:

01 Launch Targeted Cessation Programs

With 733 current smokers averaging 22 years of use and 16.8 cigarettes/day, structured behavioral therapy and pharmacological support are urgently needed.

02 Prioritize High-Risk Demographic Screening

Patients with both active smoking and family history (≈30%) face compounded organ damage risk. These patients should receive quarterly organ function assessments.

03 Support Former Smoker Recovery Pathways

631 former smokers show partial recovery potential. Structured nutritional, pulmonary, and cardiac rehab programs can meaningfully reduce residual damage.

04 Implement Population-Wide BP Monitoring

With 25.2% carrying high BP risk and smoking as a compounding factor, routine blood pressure monitoring and early statin/ACE inhibitor intervention are recommended.

Thank You

for your attention

Dataset: 2,500 patients | Variables: 13 | Analysis: Smoking & Health Outcomes

Questions? Let's discuss the data, findings, and next steps for your health strategy.