


# D Endpoint prevalence by NSFG cycle

| split                           | SSL pretrain         |                      |                      | development          | temporal test        |
|---------------------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| Mistimed or unwanted pregnancy  | 49.1%<br>2,873/5,599 | 48.2%<br>2,792/5,698 | 46.2%<br>2,335/4,879 | 44.7%<br>2,447/5,409 | 34.2%<br>1,738/4,893 |
| Impaired fecundity status       | 34.3%<br>1,755/5,599 | 30.9%<br>1,713/5,698 | 31.8%<br>1,423/4,879 | 32.6%<br>1,596/5,409 | 30.4%<br>1,496/4,893 |
| Fertility or loss-care help     | 11.3%<br>569/5,599   | 12.6%<br>614/5,698   | 10.7%<br>487/4,879   | 10.4%<br>524/5,409   | 11.1%<br>555/4,893   |
| Contraceptive vulnerability     | 6.9%<br>446/5,599    | 7.2%<br>444/5,698    | 7.7%<br>394/4,879    | 7.2%<br>423/5,409    | 7.8%<br>405/4,893    |
| Adverse pregnancy-history proxy | 9.7%<br>563/5,599    | 9.4%<br>595/5,698    | 10.5%<br>509/4,879   | 8.5%<br>535/5,409    | 5.4%<br>274/4,893    |
|                                 | 2011-2013            | 2013-2015            | 2015-2017            | 2017-2019            | 2022-2023            |



Cells show survey-weighted prevalence and unweighted event count/denominator; temporal-test column is outlined.