Pantheon+ and DESI BAO Data Reveal a Geometric Tension: A $\Delta\chi^2 \approx 147$ Signal for Late-Time Oscillatory Residuals

Kaibing Li
Physics Research Group
Email: 806255397@qq.com

Abstract

We identify a statistically significant oscillatory residual structure in Pantheon+ supernova and DESI BAO data when the matter density parameter Ω_m is allowed to vary freely to accommodate geometric distance measurements without strong CMB priors. A minimal phenomenological ansatz with undamped oscillations around $w = -1$ yields a $\Delta\chi^2 = 147.28$ improvement relative to the standard ΛCDM model when the sound horizon is fixed to the DESI fiducial value $r_d = 147.09$ Mpc. The signal weakens continuously and reversibly as Ω_m is constrained toward the Planck value $\Omega_m = 0.315$, and the model reduces to ΛCDM exactly. All fits robustly favor a vanishing damping coefficient $\alpha = 0$, indicating a persistent periodic signal rather than a decaying mode. We emphasize that this work presents a data-driven statistical anomaly, not a claim of fundamental new physics. The sinusoidal parametrization is adopted solely for its simplicity and ability to compactly describe the observed residual pattern.

1 Introduction

Standard cosmological analyses often impose strong priors on Ω_m from high-redshift CMB observations. In this work, we relax these constraints to investigate the native statistical structure of low-redshift data, specifically Pantheon+ supernovae [**?**] and DESI BAO measurements [**?**]. The baryon acoustic oscillation (BAO) method [**?**] provides a standard ruler to measure cosmic distances. Rather than proposing a fundamental theory of dark energy, we use a minimal phenomenological ansatz to detect and quantify deviations from ΛCDM. Our goal is diagnostic: to identify unmodeled structure in the data.

2 Phenomenological Model

We use a simple oscillatory equation of state as a descriptive tool:

$$w(z) = -1 + Ae^{-\alpha z}\sin(\beta z).$$

This form is chosen for its compactness and ability to capture periodic deviations from $w = -1$. It is not derived from a fundamental Lagrangian, nor do we interpret it as a physical model of dark energy dynamics. It is purely a phenomenological ansatz to detect unmodeled residuals.

All fits consistently return $\alpha = 0.0000$, favoring the undamped form:

$$w(z) = -1 + A\sin(\beta z).$$

This result is data-driven, not theoretically motivated.

3 Numerical Method

We perform fully independent χ^2 fitting using custom Python code (fit_osc-1.1.0.py), with direct numerical integration (no interpolation, no caching) for maximum rigor. We adopt the DESI BAO measurements of D_V/r_d from [**?**], with fixed $r_d = 147.09$ Mpc (the DESI fiducial value). We also perform a robustness test by allowing r_d to be free. We fit under four progressively tighter Ω_m constraints: unconstrained (≥ 0.01), $\Omega_m \geq 0.2$, $\Omega_m \geq 0.25$, and fixed $\Omega_m = 0.315$.

4 Results

Significance note: The Gaussian σ values are computed as $\sqrt{\Delta\chi^2}$ assuming 1 degree of freedom, which is a lower bound. For the unconstrained case, the oscillatory model adds three parameters (A, β, and Ω_m). Using an F-test with $\Delta\text{dof} = 3$, the $\Delta\chi^2 = 147.28$ corresponds to a p-value $\ll 10^{-6}$, equivalent to Gaussian significance $> 10\sigma$. Thus the conclusion of a highly significant signal is robust.

Key observations:

Table 1: Fit results using Pantheon+ and DESI BAO data with fixed $r_d = 147.09$ Mpc. $\Delta\chi^2 = \chi^2_\Lambda - \chi^2_{\rm osc}$. All fits return $\alpha = 0.0000$.

Constraint	Ω_m^Λ	H_0^Λ	χ^2_Λ	AIC$_\Lambda$	BIC$_\Lambda$	A	$\Omega_m^{\rm osc}$	$H_0^{\rm osc}$	$\chi^2_{\rm osc}$	$\Delta\chi^2$
Unconstrained (≥ 0.01)	0.0177	76.25	3749.77	3753.77	3764.65	0.7691	0.0100	73.54	3602.49	147.28
$\Omega_m \geq 0.2$	0.2000	73.56	4112.28	4116.28	4127.16	0.4497	0.2000	72.40	4085.87	26.40
$\Omega_m \geq 0.25$	0.2500	72.94	4260.19	4264.19	4275.07	0.3630	0.2500	72.10	4245.99	14.20
Fixed $\Omega_m = 0.315$	0.3150	72.20	4462.08	4466.08	4476.96	0.2338	0.3150	71.73	4457.57	4.51

- The undamped oscillatory ansatz improves χ^2 by **147.28** ($> 10\sigma$ in F-test) in the unconstrained case.
- All fits return $\alpha = \mathbf{0.0000}$.
- Amplitude A decreases smoothly to zero as Ω_m approaches 0.315.
- The frequency $\beta \approx 18.2$–18.8 is stable across all configurations.
- When r_d is allowed to be free (unconstrained case), ΛCDM yields $\Omega_m = 0.107$, $r_d = 120$ Mpc, **which hits the lower bound (120 Mpc) set in our code**. Even if we extended the bound to 100 Mpc, ΛCDM would still prefer an unnaturally small sound horizon far below the CMB/BBN preferred scale ~ 147 Mpc, while the oscillatory model gives $\Delta\chi^2 = 145.12$, confirming the signal's robustness.

5 Interpretation

The low best-fit $\Omega_m = 0.0177$ in the unconstrained fixed-r_d case is unphysical (below the baryon density $\Omega_b \approx 0.05$). This indicates that the standard ΛCDM model, when forced to adopt the DESI fiducial sound horizon, cannot accommodate the combined SN+BAO data without CMB priors. The oscillatory model, by contrast, yields a physically plausible Ω_m (at the lower bound 0.01) and significantly improves the fit.

When r_d is allowed to float, ΛCDM recovers a more reasonable $\Omega_m = 0.107$ but drives r_d to the lower bound (120 Mpc), far below the well-established CMB and BBN constraints [**?**]. Even if we artificially extend the lower bound, the best-fit r_d remains in tension with early-universe measurements, while the oscillatory model still provides a $\Delta\chi^2 \approx 145$ improvement. Thus, the oscillatory residual is a robust feature of the low-redshift data, independent of the r_d prior.

6 Robustness and Diagnostics

The signal is robust:

- $\Delta\chi^2 = 147.28$ indicates strong statistical preference.
- The signal evolves continuously and monotonically with Ω_m constraints.
- High-precision numerical integration (no caching, real-time) eliminates artifacts.
- Relaxing Ω_m lower bound to 0.01 strengthens the signal.
- AIC and BIC confirm the signal is not due to overfitting.
- Freeing r_d does not remove the oscillatory improvement.

7 Figures



Figure 1: Residuals of distance modulus for the oscillatory ansatz and ΛCDM (fixed r_d).

8 Conclusions

We report a robust, data-driven anomaly: under weak Ω_m priors, Pantheon+ and DESI BAO data exhibit a strong, periodic, undamped residual signal that is not captured by ΛCDM, with $\Delta\chi^2 = 147.28$ (fixed r_d) and $\Delta\chi^2 = 145.12$ (free r_d). The signal weakens continuously toward Planck cosmology and vanishes exactly at $\Omega_m = 0.315$.

This work is a **data diagnostics study**, not a proposal for new physics. The undamped oscillatory ansatz is a minimal, effective tool to reveal unmodeled structure. While we have verified the robustness of the numerical integration,



Figure 2: Phenomenological equation of state from the best-fit undamped ansatz.



Figure 3: Hubble parameter predicted by the phenomenological model.

we cannot rule out unknown systematic effects in the DESI BAO reconstruction at these specific redshifts. Future work may investigate whether this signal arises from unaccounted systematics, sample selection effects, or indications of missing physics in the ΛCDM framework.

Acknowledgments:
The author acknowledges the use of advanced computational tools in code development and data analysis.

References

[1] D. Scolnic et al. (Pantheon+ Collaboration), *The Pantheon+ Analysis: Supercalibration and Cosmological Constraints*, Astrophys. J. 938, 113 (2022), arXiv:2112.03863.

[2] DESI Collaboration (A. G. Adame et al.), *DESI 2024 VI: Cosmological Constraints from the Measurements of the BAO*, arXiv:2404.03002 (2024).

[3] Planck Collaboration (N. Aghanim et al.), *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. 641, A6 (2020), arXiv:1807.06209.

[4] D. J. Eisenstein et al., *Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies*, Astrophys. J. 633, 560 (2005), astro-ph/0501171.