

Fundamentals of Artificial Intelligence

Exercise 5: First Order Logic

Florian Lercher

Technical University of Munich

Friday 8th December, 2023

Problem 5.1: Syntax of first-order logic

Recall the formal syntax of first-order logic, in particular the definition of terms and sentences.

Reminder: Basic Elements

Syntactic element	Examples
Variables	x, y, a, b, \dots
Connectives	$\wedge, \vee, \neg, \Rightarrow, \Leftrightarrow$
Equality	$=$
Quantifiers	\forall, \exists

Reminder: Basic Elements

Syntactic element	Representation of	Examples
Constants	Objects	<i>KingJohn</i> , 2, <i>TUM</i> , ...
Predicates	Relations	<i>Brother</i> , $>$, ...
Functions	Functions	<i>Sqrt</i> , <i>LeftLegOf</i> , ...

Reminder: Terms and sentences

Backus-Naur Form of terms

$\text{Term} ::= \text{Function}(\text{Term}, \dots) \mid \text{Constant} \mid \text{Variable}$

Backus-Naur Form of atomic sentences

$\text{AtomicSentence} ::= \text{Predicate} \mid \text{Predicate}(\text{Term}, \dots) \mid \text{Term} = \text{Term}$

Backus-Naur Form of complex sentences

$\text{ComplexSentence} ::= (\text{Sentence}) \mid [\text{Sentence}] \mid \neg \text{Sentence} \mid$
 $\text{Sentence} \wedge \text{Sentence} \mid \text{Sentence} \vee \text{Sentence} \mid$
 $\text{Sentence} \Rightarrow \text{Sentence} \mid \text{Sentence} \Leftrightarrow \text{Sentence} \mid$
 $\text{Quantifier Variable}, \dots \text{Sentence}$

Problem 5.1: Syntax of first-order logic

Problem 5.1.1: Let $\mathcal{F} = \{d, f, g\}$ be the set of symbols, where d is a constant, f a function symbol with two arguments, and g a function symbol with three arguments. Which of the following expressions are terms over \mathcal{F} ? If an expression is not a term, give a reason why. (You may assume that x, y, z are variables.)

$f(x, y)$ $g(x, y, z)$

1. $g(\underline{d}, d)$ ✗
2. $f(x, \underline{g(y, z)}, d)$ ✗
3. $g(x, f(y, z), d)$ ✓
4. $g(x, \underline{k(y, z)}, d)$ ✗

Problem 5.1: Syntax of first-order logic

Problem 5.1.2: Let m be a constant, h a function symbol with one argument, and S and B two predicate symbols, each with two arguments. Which of the following expressions are sentences in first-order logic? If an expression is not a sentence, give a reason why. (You may assume that x, y, z are variables.)

1. $S(m, x)$ ✓ *terms* (arrows pointing to m and x)
2. $B(m, h(m))$ ✓ *x term* (arrow pointing to $h(m)$)
3. $B(B(m, x), y)$ ✗
4. $B(x, y) \Rightarrow [\exists z S(z, y)]$ ✓
5. $S(x, y) \Rightarrow S(y, h(h(x)))$ ✓

$S(x, y)$ $B(x, y)$

Problem 5.2: Universal and existential quantifier

Let us abbreviate the predicate “ x is taking the Bus” by $B(x)$, and “ x has a Ticket” by $T(x)$. Suppose that we only consider the universe of discourse where there are only three people: Alice, Bob, and Charlie.

x	$B(x)$	$T(x)$
Alice	True	True
Bob	False	True
Charlie	False	False

Problem 5.2.1: For each of our protagonists, the table above lists whether they have a ticket, and whether they take the bus. Suppose that we want to formalize the sentence “All people who take the bus have a ticket.” Given the characteristics in the table above, do you think the sentence evaluates to true? Or false?

True

Problem 5.2: Universal and existential quantifier

Problem 5.2.2: Now, consider the formula

$$\forall x \quad B(x) \wedge T(x).$$

By using the extended interpretation for universal quantifiers (in other words, the transformation to propositional logic), decide whether this formula is true or false in our universe of discourse. Do you think this formula is a faithful formalization of “All people who take the bus have a ticket”? If not, what would be a better formula?

$$\begin{aligned} & \forall x B(x) \wedge T(x) \\ \equiv & [B(A) \wedge T(A)] \wedge [B(B) \wedge T(B)] \\ & \quad \frac{}{\top} \qquad \frac{}{\text{F}} \wedge [B(C) \wedge T(C)] \\ & \quad \frac{}{\text{F}} \\ \equiv & \text{False} \end{aligned}$$

Problem 5.2: Universal and existential quantifier

x	$B(x)$	$T(x)$
Alice	<i>False</i>	<i>True</i>
Bob	<i>False</i>	<i>True</i>
Charlie	<i>False</i>	<i>False</i>

Problem 5.2.3: Now consider the changed truth assignments in the table above. Suppose that we want to formalize the sentence “Some people who take the bus have a ticket”¹. Given the characteristics in the table above, do you think the sentence evaluates to true? Or false?

False

¹This has to be understood as “There is **at least** one person that takes the bus, who has a ticket”.

Problem 5.2: Universal and existential quantifier

Problem 5.2.4: Now, consider the formula

$$\exists x \quad B(x) \Rightarrow T(x).$$

By using the extended interpretation for existential quantifiers, decide whether this formula is true or false in our universe of discourse. Do you think this formula is a faithful formalization of “Some people who take the bus have a ticket”? If not, what would be a better formula?

$$\equiv [B(A) \Rightarrow T(A)] \vee [B(B) \Rightarrow T(B)] \vee [B(C) \Rightarrow T(C)]$$

$\models \quad \models \quad \models$

$$\equiv \top$$

Learnings from Problem 5.2

Limiting the domain of quantification

To limit a quantifier to objects satisfying a predicate P , use ...

- $\forall x \ P(x) \Rightarrow \dots$ for universal quantification, and
- $\exists x \ P(x) \wedge \dots$ for existential quantification.

Example:

- "Some doors are locked" $\rightsquigarrow \exists x \ Door(x) \wedge Locked(x)$
- "Everything that glitters is gold" $\rightsquigarrow \forall x \ Glitters(x) \Rightarrow Gold(x)$

Problem 5.3: Formalization to First Order Logic

Problem 5.3.1: Formalize the next sentences using the following predicates and constant:

$A(x, y)$: x admires y

$P(x)$: x is a professor

d : Dan

1. Dan admires every professor.

$$\forall x \ P(x) \Rightarrow A(d, x)$$

2. Some professor admires Dan.

$$\exists x \ P(x) \wedge A(x, d)$$

3. Dan admires himself.

$$A(d, d)$$

Problem 5.3: Formalization to First Order Logic

Problem 5.3.2: Formalize the next sentences using the following predicates:

$A(x, y)$: x attended y

$S(x)$: x is a student

$L(x)$: x is a lecture

1. No student attended every lecture.

$$\neg \exists x S(x) \wedge \forall y L(y) \Rightarrow A(x, y)$$

2. No lecture was attended by every student.

$$\neg \exists x L(x) \wedge \forall y S(y) \Rightarrow A(y, x)$$

3. No lecture was attended by any student.

$$\neg \exists x L(x) \wedge \exists y S(y) \wedge A(y, x)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

Let the constant h stand for Holmes (Sherlock Holmes) and m for Moriarty (Professor Moriarty). Let us abbreviate " x can trap y " with the predicate $T(x, y)$. Give symbolic rendition of the following statements:

1. Holmes can trap everyone who can trap Moriarty.
2. Holmes can trap everyone whom Moriarty can trap.
3. Holmes can trap everyone who can be trapped by Moriarty.
4. If someone can trap Moriarty, then Holmes can.
5. If everyone can trap Moriarty, then Holmes can.
6. Anyone who can trap Holmes can trap Moriarty.
7. No one can trap Holmes unless that person can trap Moriarty.
8. Everyone can trap someone who cannot trap Moriarty.
9. Everyone who can trap Holmes can trap everyone whom Holmes can trap.

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

1. Holmes can trap everyone who can trap Moriarty.

$$\forall(x) T(x, M) \Rightarrow T(H, x)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

2. Holmes can trap everyone whom Moriarty can trap.

$$\forall x \quad T(M, x) \Rightarrow T(H, x)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

3. Holmes can trap everyone who can be trapped by Moriarty.

$$\forall x \ T(M, x) \Rightarrow T(H, x)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

4. If someone can trap Moriarty, ^{\Rightarrow} then Holmes can.

$$[\exists x \ T(x, m)] \Rightarrow T(h, m)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

5. If everyone can trap Moriarty, then Holmes can.

$$[\forall x \ T(x, m)] \Rightarrow T(H, m)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

6. Anyone who can trap Holmes can trap Moriarty.

$$\forall x \ T(x, H) \Rightarrow T(x, M)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

7. No one can trap Holmes unless that person can trap Moriarty.

$$\neg \exists x T(x, H) \wedge \neg T(x, M)$$

$$\equiv \forall x T(x, H) \Rightarrow T(x, M)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty

8. Everyone can trap someone who cannot trap Moriarty.

$$\forall x \exists y \neg T(y, M) \wedge T(x, y)$$

Problem 5.3.3: Sherlock Holmes and Professor Moriarty


9. Everyone who can trap Holmes can trap everyone
whom Holmes can trap.

$$\forall x T(x, H) \Rightarrow T(x, y)$$

$$\forall y T(H, y)$$

$$\forall x T(x, H) \Rightarrow \forall y T(H, y) \Rightarrow T(x, y)$$

Intuition for the Resolution Rule



"If $A \vee B$ and $\neg B \vee C$, then also $A \vee C$."

$$\frac{A \vee B \quad \neg B \vee C \leftarrow \text{Premises}}{A \vee C \leftarrow \text{Conclusion}}$$

But why? Let's do a case distinction on the truth assignment of B :

Case 1: B is assigned to **True**

- $A \vee B$ is satisfied, because B holds
- B holds $\leadsto \neg B$ does not hold
- To satisfy $\neg B \vee C$ we have to assign C to **True**

Case 2: B is assigned to **False**

- $\neg B \vee C$ is satisfied, because $\neg B$ holds
- To satisfy $A \vee B$ we have to assign A to **True**

\leadsto Since we cannot assign B to True and False, we must assign A to True or C to True

$\hookrightarrow A \vee C$ has to hold (assigning both A and B to True is also fine)