

Word count:

Student number:

Promoters

Supervisors:

A dissertation submitted to Ghent University in partial fulfilment of the requirements for the degree of
Master of

Academic Year:

Preface

Contents

Preface	v
1 Abstract	1
2 Introduction	3
2.1 The occurrence of ferroptosis	3
2.2 Ferroptosis in cancer therapy	5
2.3 Oxford Nanopore Episequencing	5
3 Aims	7
4 Methods	9
4.1 Introduction	9
4.2 Samples	10
4.3 Gene Whitelist	10
4.4 RNA-seq	10
4.5 ONT-seq	10
4.6 Differential Gene Expression Analysis	11
4.7 Feature Reduction	11
5 Results	13

6 Discussion	15
A Appendix	17

List of Figures

2.1 Figure depicting the various regulatory pathways associated with ferroptosis (Zhang et al., 2022a). 4

4.1 Overview workflow. 9

List of Tables

4.1	Cell Line Metadata	10
-----	------------------------------	----

List of Abbreviations and Symbols

Chapter 1

Abstract

Cancer cells exhibit a characteristic ability to evade programmed cell death, a feature critical for their uncontrolled proliferation. An emerging strategy in cancer therapy involves targeting alternative forms of cell death, distinct from apoptosis, as evidenced by recent studies (Tait, Ichim, and Green, 2014). Non-apoptotic cell death mechanisms, including necroptosis and pyroptosis, offer promising avenues to circumvent apoptosis resistance in tumors, thus enhancing the efficacy of anti-cancer therapies (Ye et al., 2018).

One such non-apoptotic pathway, ferroptosis, reliant on iron and reactive oxygen species (ROS), represents a novel approach in cancer treatment. However, inherent resistance to ferroptosis induction poses a challenge in its therapeutic application (Zou and Schreiber, 2020). Detecting drug resistance before treatment initiation could significantly impact treatment outcomes (Volm and Efferth, 2015).

Epigenetic modifications, such as DNA methylation, RNA methylation, and non-coding RNAs, have emerged as key regulators of cellular processes. Notably, DNA methylation alterations in genes associated with ferroptosis, including GPX4, FSP1, and NRF2, influence their expression levels, potentially modulating ferroptosis susceptibility (X. Wang et al., 2023).

Bisulfite genomic sequencing is widely utilized for DNA methylation detection, despite its limitations. Challenges such as incomplete bisulfite conversion or assay failures may compromise data accuracy (Li and Tollefsbol, 2011; Heiss and Just,

2018).

Recent advancements in sequencing technologies, such as nanopore sequencing, offer improved accuracy and resolution compared to traditional methods. Nanopore episequencing, in particular, shows promise in refining genetic associations and identifying novel loci previously undetected by microarray platforms (Gombert et al., 2023; Flynn et al., 2022).

In this project, we will aim to develop a machine learning model which, based on a nanopore episequencing profile, will be able to predict the likelihood of ferroptosis resistance for a given tumor strain. The model's capacity to generalize to other (un-trained) strains will be tested. Lastly, we will identify specific methylation changes brought about by ferroptosis-inducing drugs by sequencing sensitized strains.

Chapter 2

Introduction

2.1 The occurrence of ferroptosis

Ferroptosis is a complex process with many contributing, interconnected factors. At its core, it is an intracellular iron-dependent form of cell death that is distinct from apoptosis, necrosis, and autophagy (Yan et al., 2021). As the name suggests, this phenomenon is principally driven by iron molecules absorbed by cells from the surrounding extracellular environment. One such receptor is transferrin receptor 1 (TFR1), which mediates endocytosis of diferric transferrin (S. Wang et al., 2019). This intracellular iron is often transported and stored in storage proteins such as ferritin, aiding in maintaining iron homeostasis and metabolism (Knovich et al., 2009).

Labile iron, however, which refers to loosely bound or chelatable iron, becomes a critical player in the ferroptotic process. Labile iron participates in the Fenton reaction, leading to the generation of reactive oxygen species (ROS), particularly hydroxyl radicals (Cabantchik, 2014). ROS, in turn, contributes to oxidative stress within the cell.

Polyunsaturated fatty acids in cellular membranes are highly susceptible to oxidative damage by ROS (Ayala, Muñoz, and Argüelles, 2014). Lipid peroxides are formed as a result of the oxidation of these fatty acids. Accumulation of lipid peroxides disrupts the integrity of cellular membranes, particularly the lipid bilayer

(Gaschler and Stockwell, 2017). Membrane damage leads to the loss of compartmentalization and function within the cell, including the mitochondrion (Ademowo et al., 2017).

Healthy mammalian cells stave off this threat in a number of ways. The primary cellular mechanism of protection against oxytosis/ferroptosis is mediated by glutathione peroxidase 4 (GPX4), a glutathione-dependent hydroperoxidase that converts lipid peroxides into non-toxic lipid alcohols (Xie et al., 2023). Glutathione itself, which is produced from cysteine, is able to neutralize ROS present within the cell directly, playing the role of antioxidant (Averill-Bates, 2023).

Even when glutathione peroxidase functionality is uncompromised, research has shown that the mitochondrion plays a substantial role in ferroptosis regulation and induction, being the major compartment for cellular iron metabolism (Liu et al., 2023). Mitochondrial shrinkage and rupture are also common observations when a cell undergoes ferroptosis, since this organelle is enveloped in a membrane (Gao et al., 2019).




Figure 2.1: Figure depicting the various regulatory pathways associated with ferroptosis (Zhang et al., 2022a).

2.2 Ferroptosis in cancer therapy

Cancer is the second leading cause of death worldwide, behind cardiovascular disease (H. Wang et al., 2016). However, improvements in cancer survival rates due to the use of precision medicine or immunotherapy

There are a number of ways to counteract this resistance, including inhibition of glutathione synthesis, overexpression of transferrin receptors, underexpression of or defective ferritin (including the chaperone proteins that fill the ferritin) and/or ferroportin proteins (which export labile iron bound to chaperones outside the cell) (Zhang et al., 2022b).

2.3 Oxford Nanopore Episequencing

Nanopore sequencing is a next-generation sequencing technology that enables the real-time analysis of DNA or RNA molecules as they pass through a nanoscale pore. This technique is based on the principle that as a DNA strand translocates through a nanopore, changes in the electrical current can be measured and used to deduce the sequence of the DNA molecule Y. Wang et al., 2021. The raw electrical signal is translated into DNA sequence information through a process called base-calling. DNA methylation involves the addition of a methyl group to a cytosine base (5-methylcytosine). Nanopore sequencing can detect these modifications as changes in the electrical current during translocation using specialized algorithms and basecallers. One of the key advantages of nanopore sequencing is its ability to generate long reads, providing information about entire genomic regions in a single continuous sequence.

Adaptive sampling is a feature associated with PromethION that helps optimize the sequencing process by adjusting data collection parameters in real-time based on the characteristics of the DNA being sequenced. The system monitors various parameters such as the quality of the signal, the speed at which the DNA is translocating through the nanopore, and other relevant metrics (Martin et al., 2022). The software requires a user to upload a file of whitelisted reference sequences, and

the system can be set to either deplete or enrich for these on a specified set of channels.

In order to achieve this, the software basecalls the first few hundred bases of each read and compares it with the target reference sequences. Matching or unmatching sequences are rejected, depending on whether the software is set to enrich or deplete. By using this, coverage of our genes of interest is enriched, greatly reducing noise.

Chapter 3

Aims

Chapter 4

Methods

4.1 Introduction

Firstly, differential gene expression analysis was performed. Then, feature reduction of the base-level modification information obtained from ONT episequencing was carried out. Finally, we trained a predictive model using the reduced feature matrix.




Figure 4.1: Overview workflow.

4.2 Samples

To start off, expression data from 4 cell lines was gathered, namely 2 resistant and 2 sensitive neuroblastoma cell lines in quadruplicate (16 samples in total). These cell lines were characterized in terms of sensitivity towards ferroptosis beforehand experimentally. Furthermore, neuroblastoma is highly iron addicted which makes it exceptionally vulnerable to ferroptosis (Floros et al., 2021). Metadata for each cell line can be found in Table 4.1.

Cell Line	Tumor Type	Ferroptosis Sensitivity	Gender	Age (months)
IMR-32	Neuroblastoma	Sensitive	Male	13
SH-EP	Neuroblastoma	Sensitive	Female	48
SH-SY5Y	Neuroblastoma	Resistant	Female	48
SK-N-BE(2)-C	Neuroblastoma	Resistant	Male	26

Table 4.1: Cell Line Metadata

4.3 Gene Whitelist

Genes thought to be associated with ferroptosis were compiled into a custom gene whitelist. The full list can be found in Appendix X.

4.4 RNA-seq

4.5 ONT-seq

For the episequencing, a PromethION nanopore sequencing device based at OHMX.bio was used, containing 24 channels per flow cell. PromethION offers a significantly higher number of channels compared to other ONT platforms like MinION and GridION. This high channel count enhances the platform's capacity for parallel sequencing.

From the whitelist, a BED file was constructed. Additionally, each gene was extended by 15kB, after which overlapping genes were joined together.

Library prep involved following protocol SQK-NBD114.24, describing how to carry out native barcoding of genomic DNA (gDNA) using the Native Barcoding Kit 24 V14. Raw signal data in pod5 format was basecalled with a SUP model and 5(h)mC calling in CG context using GPU hardware on a GridION unit installed at OHMX.bio, and then subsequently demultiplexed. The basecalling model was set to fast, with the intent of making the decision of whether to reverse the voltage of the pore as fast as possible. The resulting unmapped bams were then aligned against a reference genome, and finally sorted and indexed using SAMtools (Danecek et al., 2021).

4.6 Differential Gene Expression Analysis

4.7 Feature Reduction

Since the episequencing returns many thousands of individual detected CpGs by locus, the number of features in our model will be staggering if used in its raw form. The first reduction of features will be actualized by bundling CpG sites into regions (e.g. CpG islands). A second dimensionality reduction will also be performed based on drivers derived from expression data, to remove sites or regions without genes that are differentially expressed by sensitivity.

Chapter 5

Results

Chapter 6

Discussion

Appendix A

Appendix

Bibliography

- Zhang, Chen et al. (Feb. 2022a). "Ferroptosis in cancer therapy: a novel approach to reversing drug resistance". en. In: *Mol. Cancer* 21.1, p. 47.
- Tait, Stephen W. G., Gabriel Ichim, and Douglas R. Green (May 2014). "Die another way – non-apoptotic mechanisms of cell death". In: *Journal of Cell Science* 127.10, pp. 2135–2144. ISSN: 0021-9533. DOI: 10.1242/jcs.093575. URL: <http://dx.doi.org/10.1242/jcs.093575>.
- Ye, Jing et al. (2018). "Non-apoptotic cell death in malignant tumor cells and natural compounds". In: *Cancer Letters* 420, pp. 210–227. ISSN: 0304-3835. DOI: <https://doi.org/10.1016/j.canlet.2018.01.061>. URL: <https://www.sciencedirect.com/science/article/pii/S0304383518300946>.
- Zou, Yilong and Stuart L. Schreiber (2020). "Progress in Understanding Ferroptosis and Challenges in Its Targeting for Therapeutic Benefit". In: *Cell Chemical Biology* 27.4. Special Issue on Chemical Biology of Ferroptosis, pp. 463–471. ISSN: 2451-9456. DOI: <https://doi.org/10.1016/j.chembiol.2020.03.015>. URL: <https://www.sciencedirect.com/science/article/pii/S2451945620301124>.
- Volm, Manfred and Thomas Efferth (Dec. 2015). "Prediction of Cancer Drug Resistance and Implications for Personalized Medicine". In: *Frontiers in Oncology* 5. ISSN: 2234-943X. DOI: 10.3389/fonc.2015.00282. URL: <http://dx.doi.org/10.3389/fonc.2015.00282>.
- Wang, Xiancan et al. (2023). "Effects of DNA, RNA, and Protein Methylation on the Regulation of Ferroptosis". In: *International Journal of Biological Sciences* 19.11, pp. 3558–3575. ISSN: 1449-2288. DOI: 10.7150/ijbs.85454. URL: <http://dx.doi.org/10.7150/ijbs.85454>.
- Li, Yuanyuan and Trygve O. Tollefsbol (2011). "DNA Methylation Detection: Bisulfite Genomic Sequencing Analysis". In: *Epigenetics Protocols*. Humana Press, pp. 11–

21. ISBN: 9781617793165. DOI: 10.1007/978-1-61779-316-5_2. URL: http://dx.doi.org/10.1007/978-1-61779-316-5_2.
- Heiss, Jonathan A. and Allan C. Just (June 2018). "Identifying mislabeled and contaminated DNA methylation microarray data: an extended quality control toolset with examples from GEO". In: *Clinical Epigenetics* 10.1. ISSN: 1868-7083. DOI: 10.1186/s13148-018-0504-1. URL: <http://dx.doi.org/10.1186/s13148-018-0504-1>.
- Gombert, Sara et al. (Oct. 2023). "Comparison of methylation estimates obtained via MinION nanopore sequencing and sanger bisulfite sequencing in the TRPA1 promoter region". In: *BMC Medical Genomics* 16.1. ISSN: 1755-8794. DOI: 10.1186/s12920-023-01694-6. URL: <http://dx.doi.org/10.1186/s12920-023-01694-6>.
- Flynn, Robert et al. (May 2022). "Evaluation of nanopore sequencing for epigenetic epidemiology: a comparison with DNA methylation microarrays". In: *Human Molecular Genetics* 31.18, pp. 3181–3190. ISSN: 1460-2083. DOI: 10.1093/hmg/ddac112. URL: <http://dx.doi.org/10.1093/hmg/ddac112>.
- Yan, Hong-fa et al. (Feb. 2021). "Ferroptosis: mechanisms and links with diseases". In: *Signal Transduction and Targeted Therapy* 6.1. ISSN: 2059-3635. DOI: 10.1038/s41392-020-00428-9. URL: <http://dx.doi.org/10.1038/s41392-020-00428-9>.
- Wang, Shufen et al. (Oct. 2019). "Transferrin receptor 1-mediated iron uptake plays an essential role in hematopoiesis". In: *Haematologica* 105.8, pp. 2071–2082. ISSN: 1592-8721. DOI: 10.3324/haematol.2019.224899. URL: <http://dx.doi.org/10.3324/haematol.2019.224899>.
- Knovich, Mary Ann et al. (May 2009). "Ferritin for the clinician". In: *Blood Reviews* 23.3, pp. 95–104. ISSN: 0268-960X. DOI: 10.1016/j.blre.2008.08.001. URL: <http://dx.doi.org/10.1016/j.blre.2008.08.001>.
- Cabantchik, Zvi loav (Mar. 2014). "Labile iron in cells and body fluids: physiology, pathology, and pharmacology". In: *Frontiers in Pharmacology* 5. ISSN: 1663-9812. DOI: 10.3389/fphar.2014.00045. URL: <http://dx.doi.org/10.3389/fphar.2014.00045>.
- Ayala, Antonio, Mario F. Muñoz, and Sandro Argüelles (2014). "Lipid Peroxidation: Production, Metabolism, and Signaling Mechanisms of Malondialdehyde and 4-

- Hydroxy-2-Nonenal". In: *Oxidative Medicine and Cellular Longevity* 2014, pp. 1–31. ISSN: 1942-0994. DOI: 10.1155/2014/360438. URL: <http://dx.doi.org/10.1155/2014/360438>.
- Gaschler, Michael M. and Brent R. Stockwell (Jan. 2017). "Lipid peroxidation in cell death". In: *Biochemical and Biophysical Research Communications* 482.3, pp. 419–425. ISSN: 0006-291X. DOI: 10.1016/j.bbrc.2016.10.086. URL: <http://dx.doi.org/10.1016/j.bbrc.2016.10.086>.
- Ademowo, O. S. et al. (May 2017). "Lipid (per) oxidation in mitochondria: an emerging target in the ageing process?" In: *Biogerontology* 18.6, pp. 859–879. ISSN: 1573-6768. DOI: 10.1007/s10522-017-9710-z. URL: <http://dx.doi.org/10.1007/s10522-017-9710-z>.
- Xie, Yangchun et al. (June 2023). "GPX4 in cell death, autophagy, and disease". In: *Autophagy* 19.10, pp. 2621–2638. ISSN: 1554-8635. DOI: 10.1080/15548627.2023.2218764. URL: <http://dx.doi.org/10.1080/15548627.2023.2218764>.
- Averill-Bates, Diana A. (2023). "The antioxidant glutathione". In: *Antioxidants*. Elsevier, pp. 109–141. ISBN: 9780443157684. DOI: 10.1016/bs.vh.2022.09.002. URL: <http://dx.doi.org/10.1016/bs.vh.2022.09.002>.
- Liu, Yu'e et al. (Aug. 2023). "The diversified role of mitochondria in ferroptosis in cancer". In: *Cell Death & Disease* 14.8. ISSN: 2041-4889. DOI: 10.1038/s41419-023-06045-y. URL: <http://dx.doi.org/10.1038/s41419-023-06045-y>.
- Gao, Minghui et al. (Jan. 2019). "Role of Mitochondria in Ferroptosis". In: *Molecular Cell* 73.2, 354–363.e3. ISSN: 1097-2765. DOI: 10.1016/j.molcel.2018.10.042. URL: <http://dx.doi.org/10.1016/j.molcel.2018.10.042>.
- Wang, Haidong et al. (Oct. 2016). "Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015". In: *The Lancet* 388.10053, pp. 1459–1544. ISSN: 0140-6736. DOI: 10.1016/S0140-6736(16)31012-1. URL: [http://dx.doi.org/10.1016/S0140-6736\(16\)31012-1](http://dx.doi.org/10.1016/S0140-6736(16)31012-1).
- Zhang, Chen et al. (Feb. 2022b). "Ferroptosis in cancer therapy: a novel approach to reversing drug resistance". In: *Molecular Cancer* 21.1. ISSN: 1476-4598. DOI: 10.1186/s12943-022-01530-y. URL: <http://dx.doi.org/10.1186/s12943-022-01530-y>.

- Wang, Yunhao et al. (Nov. 2021). "Nanopore sequencing technology, bioinformatics and applications". In: *Nature Biotechnology* 39.11, pp. 1348–1365. ISSN: 1546-1696. DOI: 10.1038/s41587-021-01108-x. URL: <http://dx.doi.org/10.1038/s41587-021-01108-x>.
- Martin, Samuel et al. (Jan. 2022). "Nanopore adaptive sampling: a tool for enrichment of low abundance species in metagenomic samples". In: *Genome Biology* 23.1. ISSN: 1474-760X. DOI: 10.1186/s13059-021-02582-x. URL: <http://dx.doi.org/10.1186/s13059-021-02582-x>.
- Floros, Konstantinos V. et al. (Jan. 2021). "MYCN-Amplified Neuroblastoma Is Addicted to Iron and Vulnerable to Inhibition of the System Xc-/Glutathione Axis". In: *Cancer Research* 81.7, pp. 1896–1908. ISSN: 1538-7445. DOI: 10.1158/0008-5472.can-20-1641. URL: <http://dx.doi.org/10.1158/0008-5472.CAN-20-1641>.
- Danecek, Petr et al. (Feb. 2021). "Twelve years of SAMtools and BCFtools". In: *GigaScience* 10.2. giab008. ISSN: 2047-217X. DOI: 10.1093/gigascience/giab008. eprint: <https://academic.oup.com/gigascience/article-pdf/10/2/giab008/36332246/giab008.pdf>. URL: <https://doi.org/10.1093/gigascience/giab008>.

