

Your Title

Your Subtitle

Your name

Email: XXX@xxx.cuhk.edu.hk

Office: Pavilion of Harmony, CUHK

The Chinese University of Hong Kong

March 27, 2024





- 1 Cite and Footnote
- 2 Text, Lists, Tables and Figures
- 3 Columns, Code, Links and Footnote
- 4 Equations and Blocks
- 5 References



- 1 Cite and Footnote
- 2 Text, Lists, Tables and Figures
- 3 Columns, Code, Links and Footnote
- 4 Equations and Blocks
- 5 References



Attention Is All You Need^[1]

[1] Vaswani et al., “Attention is All you Need”, 2017.



Long Short-Term Memory^[2]

[2] Hochreiter and Schmidhuber, “Long Short-Term Memory”, 1997.



- 1 Cite and Footnote
- 2 Text, Lists, Tables and Figures
- 3 Columns, Code, Links and Footnote
- 4 Equations and Blocks
- 5 References



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.

- Time-dependent Schrödinger's equation:

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle$$



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.

- Time-dependent Schrödinger's equation:

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle$$

- Mathematically, a function $f(x)$ is linear iff $f(u + v) = f(u) + f(v)$ and $f(cu) = cf(u)$.



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.

- Time-dependent Schrödinger's equation:

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle$$

- Mathematically, a function $f(x)$ is linear iff $f(u + v) = f(u) + f(v)$ and $f(cu) = cf(u)$.



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.

- Time-dependent Schrödinger's equation:

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle$$

- Mathematically, a function $f(x)$ is linear iff $f(u + v) = f(u) + f(v)$ and $f(cu) = cf(u)$.

Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur.



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.

- Time-dependent Schrödinger's equation:

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle$$

- Mathematically, a function $f(x)$ is linear iff $f(u + v) = f(u) + f(v)$ and $f(cu) = cf(u)$.

Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur.

Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.




Figure 1: Convex Surface



Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat.

Index	Areas (m^2)	Rent (HKD)
1	40	134072
2	92	182241
3	37	134731
4	124	204325
5	88	187375
...



- 1 Cite and Footnote
- 2 Text, Lists, Tables and Figures
- 3 Columns, Code, Links and Footnote
- 4 Equations and Blocks
- 5 References



- ① Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium.
- ② Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit.
- ③ Totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo.

- ① Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium.
- ② Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit.
- ③ Totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo.



Figure 2: Lenna



Algorithm 1 An algorithm with caption

Require: $n \geq 0$ **Ensure:** $y = x^n$ $y \leftarrow 1$ $X \leftarrow x$ $N \leftarrow n$ **while** $N \neq 0$ **do** **if** N is even **then** $X \leftarrow X \times X$ $N \leftarrow \frac{N}{2}$ **else if** N is odd **then** $y \leftarrow y \times X$ $N \leftarrow N - 1$ **end if****end while**

▷ This is a comment



- Beamer (LaTeX) - Wikipedia
- Please refer to page 2.
- [https://en.wikipedia.org/wiki/Beamer_\(LaTeX\)](https://en.wikipedia.org/wiki/Beamer_(LaTeX))



- **Beamer** is a \LaTeX document class for creating presentation slides, with a wide range of templates and a set of features for making slideshow effects. It supports pdf \LaTeX , \LaTeX + dvips, Lua \LaTeX and Xe \LaTeX . The name is taken from the German word “Beamer” as a pseudo-anglicism for “video projector”.¹

¹[https://en.wikipedia.org/wiki/Beamer_\(LaTeX\)](https://en.wikipedia.org/wiki/Beamer_(LaTeX))



- 1 Cite and Footnote
- 2 Text, Lists, Tables and Figures
- 3 Columns, Code, Links and Footnote
- 4 Equations and Blocks**
- 5 References



Example

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat.



Theorem

$\mathbf{X}^T \mathbf{X}$ is invertible $\iff \mathbf{X}$ has linearly independent columns.



Theorem

$\mathbf{X}^T \mathbf{X}$ is invertible $\iff \mathbf{X}$ has linearly independent columns.

Proof.

Firstly, note that $\mathbf{X}^T \mathbf{X} \in \mathbf{R}^{n \times n}$. We denote $N(\mathbf{X})$ as the kernel (nullspace) of \mathbf{X} , and $R(\mathbf{X})$ as the range (column space) of \mathbf{X} . We prove $\mathbf{X}^T \mathbf{X}$ and \mathbf{X} share the same kernel such that once $N(\mathbf{X}) = 0$, $N(\mathbf{X}^T \mathbf{X}) = 0$ and vice versa. □



Proof.

1) Prove $N(\mathbf{X}) \subset N(\mathbf{X}^T \mathbf{X})$

$\forall v \in N(\mathbf{X}), \mathbf{X}^T \mathbf{X}v = \mathbf{X}^T \mathbf{0} = \mathbf{0}$

$\implies v \in N(\mathbf{X}^T \mathbf{X}) \implies N(\mathbf{X}) \subset N(\mathbf{X}^T \mathbf{X}).$



Proof.

1) Prove $N(\mathbf{X}) \subset N(\mathbf{X}^T \mathbf{X})$

$$\forall v \in N(\mathbf{X}), \mathbf{X}^T \mathbf{X}v = \mathbf{X}^T \mathbf{0} = \mathbf{0}$$

$$\implies v \in N(\mathbf{X}^T \mathbf{X}) \implies N(\mathbf{X}) \subset N(\mathbf{X}^T \mathbf{X}).$$

2) Prove $N(\mathbf{X}^T \mathbf{X}) \subset N(\mathbf{X})$

$$\forall v \neq \mathbf{0} \in N(\mathbf{X}^T \mathbf{X}), \mathbf{X}^T \mathbf{X}v = \mathbf{0} \implies v \in N(\mathbf{X}^T) \text{ or } \mathbf{X}v \in N(\mathbf{X}^T).$$

However, we have $R(\mathbf{X}) \perp N(\mathbf{X}^T)$ and $\mathbf{X}v \in R(\mathbf{X})$,

$$\implies \mathbf{X}v \perp N(\mathbf{X}^T) \implies \mathbf{X}v \notin N(\mathbf{X}^T) \implies v \in N(\mathbf{X}^T)$$

$$\implies N(\mathbf{X}^T \mathbf{X}) \subset N(\mathbf{X})$$



Proof.

1) Prove $N(\mathbf{X}) \subset N(\mathbf{X}^T \mathbf{X})$

$$\forall v \in N(\mathbf{X}), \mathbf{X}^T \mathbf{X}v = \mathbf{X}^T \mathbf{0} = \mathbf{0}$$

$$\implies v \in N(\mathbf{X}^T \mathbf{X}) \implies N(\mathbf{X}) \subset N(\mathbf{X}^T \mathbf{X}).$$

2) Prove $N(\mathbf{X}^T \mathbf{X}) \subset N(\mathbf{X})$

$$\forall v \neq \mathbf{0} \in N(\mathbf{X}^T \mathbf{X}), \mathbf{X}^T \mathbf{X}v = \mathbf{0} \implies v \in N(\mathbf{X}^T) \text{ or } \mathbf{X}v \in N(\mathbf{X}^T).$$

However, we have $R(\mathbf{X}) \perp N(\mathbf{X}^T)$ and $\mathbf{X}v \in R(\mathbf{X})$,

$$\implies \mathbf{X}v \perp N(\mathbf{X}^T) \implies \mathbf{X}v \notin N(\mathbf{X}^T) \implies v \in N(\mathbf{X}^T)$$

$$\implies N(\mathbf{X}^T \mathbf{X}) \subset N(\mathbf{X})$$

$$1), 2) \implies N(\mathbf{X}^T \mathbf{X}) = N(\mathbf{X})$$





- 1 Cite and Footnote
- 2 Text, Lists, Tables and Figures
- 3 Columns, Code, Links and Footnote
- 4 Equations and Blocks
- 5 References**



- [1] Ashish Vaswani et al. “Attention is All you Need”. In: *Advances in Neural Information Processing Systems*. Ed. by I. Guyon et al. Vol. 30. Curran Associates, Inc., 2017.
- [2] Sepp Hochreiter and Jürgen Schmidhuber. “Long Short-Term Memory”. In: *Neural Computation* 9.8 (Nov. 1997), pp. 1735–1780. ISSN: 0899-7667.



- [Overleaf Documentation](#)
- [Learn LaTeX in 30 Minutes](#)
- [LaTeX Beamer - Overleaf](#)
- [Beamer Presentations: A Tutorial for Beginners](#)