

JDemetra+ online documentation

Anna Smyk, Alain Quartier-la-Tente, Tanguy Barthelemy, Karsten Webel

12/23/2022

Table of contents

Preface	8
JDemetra+	9
Introduction	9
Version 2.2.3 and version 3	9
Main functions	9
Seasonal adjustment algorithms	9
Temporal Disaggregation and benchmarking	10
Trend-cycle estimation	10
Nowcasting	10
Structure of this book	10
Algorithms	11
Tools	11
Underlying Statistical Methods	11
How to use this book	11
Audience	11
 I Algorithms	 12
Seasonal Adjustment	13
Motivation	13
Data frequencies : high and low	13
Unobserved Components (UC)	13
Detecting seasonal patterns	14
Available algorithms	14
Pre-treatment : Reg-ARIMA (or TRAMO)	15
why ?	15
Global outline	15
Default Specifications	16
Spans	17
Decomposition Scheme	20
Calendar correction	22
Outliers and intervention variables	23
Reg-Arima Model	24

Modelling	25
X-11 Decomposition	25
Default specifications	25
Quick Launch	26
Output 1: series	26
Output 2: final parameters	27
Output 3: diagnostics	27
User-defined parameters	29
SEATS Decomposition	33
Default specifications	33
Quick Launch	33
Output 1: series	33
Output2: Final Params	37
Output3: Diagnostics	37
Fine-tuning : user defined parameters	38
STL	39
SSF	39
Quality assessment	39
Residual seasonality	39
Residual calendar effects	39
Matrix	39
Add other diagnostics node	39
Seasonal adjustment of high-frequency data	40
Motivation: data specificities	40
Available algorithms	40
Unobserved Components	41
Identifying seasonal patterns	41
Spectral analysis	41
Seasonality tests	41
Pre-adjustment	41
Calendar correction	41
Outliers and intervention variables	41
Linearization	41
Decomposition	41
Extended X-11	41
Arima Model Based (AMB) Decomposition	41
STL decomposition	41
State Space framework	41
Quality assessment	42
Residual seasonality	42
Residual Calendar effects	42

Outlier detection	43
Motivation	43
With Reg Arima models	43
Specific TERROR tool	43
With structural models (BSM)	43
Calendar and user-defined corrections	44
Overview of Calendar effects in JDemetra	44
Summary of the method used in JDemetra+ to compute trading day and work- ing day effects	45
Mean and seasonal effects of calendar variables	49
Impact of the mean effects on the decomposition	51
Linear transformations of the calendar variables	54
Handling of specific holidays	55
Holidays with a validity period	57
Different Kinds of calendars	58
Tests for residual trading days	58
Benchmarking and temporal disaggregation	60
Benchmarking overview	60
Tools	61
Benchmarking with GUI	61
Benchmarking in R	64
References	64
Trend-cycle estimation	66
Motivation	66
Estimation Methods	66
Tools	66
rjd3 highfreq package	66
rjdfilters package	66
Nowcasting	67
Motivation	67
Tools	67
II Tools	68
Graphical User Interface	69
Overview	69
Installation Procedure	69
Running JDemetra+	70
Closing JDemetra+	71

R packages	72
Available algorithms	72
Organisation overview	72
RJDemetra and JD+ version 2.2.3	72
Installation procedure	73
Utility packages	73
rjd3toolkit	73
rjd3modelling	74
rjd3sa package	75
Seasonal adjustment packages	77
rjd3arima	77
rjd3x13 and rjd3tramoseats	77
Common functions	77
rjd3x13	77
rjd3tramoseats	78
Example	78
rjdemetra3	78
rjd3highfreq	78
rjd3stl	79
Add-in package	79
ggdemetra	79
ggdemetra3	79
rjdfilters	80
rjdfilters (1)	80
Create moving average <code>moving_average()</code>	80
Apply a moving average	81
rjd3sts	82
rjd3sts	82
rjd3bench	82
rjd3bench	82
Interaction with workspaces and GUI	82
Cruncher	83
Overview	83
Installation procedure	83
Running the cruncher in R	83
Revision Policies and Production	84
Revision Policies	84
Plug-ins for JDemetra+	85
Main functions	85

Tool selection issues and heuristics	86
III Methods	87
Spectral Analysis Principles and Tools	88
Spectral analysis concepts	88
Theoretical spectral density of an ARIMA model	89
Spectral density estimation	89
Method 1: The periodogram	89
Method 2: Autoregressive spectrum estimation	94
Method 3: Tukey spectrum	96
Identification of spectral peaks	96
Spectral graphs	105
Reg-Arima models	114
Overview	114
X-11 decomposition	115
Introduction	115
M-stats	131
Detailed tables	133
STL: Local regression decomposition	139
SEATS decomposition	140
Introduction	140
ARIMA modelling of the input series	140
Derivation of the models for the components	141
Estimation of the components with the Wiener-Kolmogorow filter	151
PsiE-weights	162
Local Polynomials Methods for Trend Estimation	165
Chapter building process	165
Tests	166
Introduction	166
Tests on residuals	166
Ljung-Box	166
Box-Pierce	167
Dornik-Hansen	167
Seasonality tests	169
F-test on seasonal dummies	171
Identification of spectral peaks	173

Friedman test for stable seasonality test	173
Moving seasonality test	176
Combined seasonality test	177
State space modelling	179
Introduction	179
State space forms (SSF)	179
General form	179
Temporal disaggregation and benchmarking	181
Benchmarking Underlying Theory	181
Temporal Disaggregation underlying Theory	183
References	184

Preface

Welcome to the JDemetra+ online documentation.

JDemetra+ is a software for seasonal adjustment and other time series functions, developed in the framework of Eurostat’s “Centre of Excellence on Statistical Methods and Tools” by the National Bank of Belgium with the support of the Bundesbank.

To learn more about this project <https://ec.europa.eu/eurostat/cros/content/centre-excellence-statistical-methods-and-tools>.

JDemetra+

Introduction

JD+ is a library of algorithms for seasonal adjustment and time series econometrics. You can learn more about the history of the project here ([link to below](#))

[link to key references](#) - [handbooks](#) (3) - [sets of guidelines](#) (2 or 3 ?)

Version 2.2.3 and version 3

[approach](#)

as v2 still widely used ...

Main functions

Seasonal adjustment algorithms

All are available for low and high frequency data.

Algorithms	Access	Key features
X13-Arima		
Tramo-Seats		
STL		
State Space Models (STS)		

Algorithms	Access	Key features
------------	--------	--------------

Temporal Disaggregation and benchmarking

Algorithms	Access	Key features
Chow-lin		
Fernandez		

Trend-cycle estimation

Algorithms	Access	Key features
------------	--------	--------------

Nowcasting

Algorithms	Access	Key features
------------	--------	--------------

Structure of this book

This book is divided in four parts, allowing the user to access the resources from different perspectives.

Algorithms

This part provides a step by step description of all the algorithms featured in JD+, grouped by purpose - seasonal adjustment - benchmarking - temporal disaggregation - ... links

Tools

JDemetra+ offers 3 kinds of tools - A Graphical User Interface (GUI) which can be enhanced with plug-ins - A set of R packages - A Cruncher for mass production in seasonal adjustment

Underlying Statistical Methods

This part gives details about the underlying statistical methods to foster a more in-depth understanding of the algorithms. Those methods are described in the light and spirit of their use as building blocks of the algorithms presented above, not aiming at all at their comprehensive coverage.

How to use this book

Audience

This book targets the beginner as well as seasoned methodologist interested in using JDemetra+ software for any the purposes listed below.

The documentation is built in layers allowing to skip details and complexity in the first steps

Part I

Algorithms

Seasonal Adjustment

Hint at GUI chap for global principles - ws structure (new SAP) - global specification handling
- file structure explanation

Hint at tool selection (link to chapter)

Motivation

The primary aim of the seasonal adjustment process is to remove seasonal fluctuations from the time series. Seasonal fluctuations are quasi-periodic infra-annual movements. They can mask...

[insert def SA from b_ov]

- infra annual periodic numbers
- no unambiguous definition
- depends on its practical estimation

Data frequencies : high and low

Until recently seasonal were designed to tackle monthly frequencies or lower in JD+, X13-
Seats Arima : 2,4,12 Seats : 2,3,4,6,12

For infra monthly data

Unobserved Components (UC)

The main components, each representing the impact of certain types of phenomena on the time series (X_t) , are:

- The trend (T_t) that captures long-term and medium-term behaviour;
- The seasonal component (S_t) representing intra-year fluctuations, monthly or quarterly, that are repeated more or less regularly year after year;

- The irregular component (I_t) combining all the other more or less erratic fluctuations not covered by the previous components.

In general, the trend consists of 2 sub-components:

- The long-term evolution of the series;
- The cycle, that represents the smooth, almost periodic movement around the long-term evolution of the series. It reveals a succession of phases of growth and recession. Trend and cycle are not separated in SA algorithms.

Two decomposition models:

- The additive model: $X_t = T_t + S_t + I_t$;
- The multiplicative model: $X_t = T_t \times S_t \times I_t$.

In the Reg-Arima (link) or Tramo(link) modelling the the multiplicative model is estimated in logs; $\log(X_t) = \log(T_t) + \log(S_t) + \log(I_t)$; Seats decomposition also relies on logs whereas X-11 decomp operates...

Detecting seasonal patterns

Seasonality tests (chap M_Tests)

Available algorithms

(link to table)

The two most popular are..

X-13ARIMA is a seasonal adjustment program developed and supported by the U.S. Census Bureau. It is based on the U.S. Census Bureau's earlier X-11 program, the X-11-ARIMA program developed at Statistics Canada, the X-12-ARIMA program developed by the U.S. Census Bureau.

Tramo-Seats

Those will be the most detailed in this doc, but there alternative solutions.

STL

BSM

X13 and TS are two step algos, pretreatment +dcomp (no pre ptreatment in STL + all at once in BSM)

- pre-treatment objectives and methods
- decomposition objective and methods

Pre-treatment : Reg-ARIMA (or TRAMO)

why ?

Global outline

let's call it regArima modelling (and tramo only when TRamo differences are outlined)

GOAL : remove deterministic effects to improve decomposition (details on modelling: [link](#))

global structure regression with ARIMA structure for residuals

$$Y_t = \sum \alpha_i O_{it} + \sum \beta_j C_{jt} + \sum \gamma_i U_{it} + Y_{lin,t}$$

where - O_{it} are the i final outliers (Ao, LS, TC) from an automatic detection procedure ([link to outliers chap](#))

- C_{it} are the calendar regressors (automatic or user-defined) ([link to calendar chap](#))
- U_{it} are all the other user-defined regressors ([link to outliers and regressors chap](#))
- $Y_{lin,t} \sim ARIMA(p, d, q)(P, D, Q)$

In the following sections we cover in more details: not the tight structure - Default params: (quick launch) INPUT (?) - Setting User-defined parameters (and not at the end post diagnostics ?) - Output1: series - Output2: Final Params - Output3: Diagnostics for each of these steps - calendar (auto and user) - outliers (auto and user) - adding intervention variables (here: just how to feed the model, details on regressor and calendar specif chapter) - aima models parameters and diagno

RegArima modelling part can be of an SA process or run on its own - GUI - R (slightly different specs,...)

WARNING : set params + retrieve their values: global principles

in GUI

Retrieve: in the specification window and panels for main (to detail ?)

in R

Default Specifications

here general explanations

default specification are set for the whole sa procedure: pre-treatment end decomposition

they are slightly different for X13-ARIMA and tramo seats

the user chooses the algo + one default spec

this default spec can be modified with user defined params for most of its parts, this is tackled in details for every sub part of the reg arima (tramo) modelling and corresponding decomposition (x11 and seats)

Starting point for X13-ARIMA

Spec identifier	Log/level detection	Outliers detection	Calendar effects	ARIMA
RSA0	NA	NA	NA	Airline(+mean)
RSA1	automatic	AO/LS/TC	NA	Airline(+mean)
RSA2c	automatic	AO/LS/TC	2 TD vars+Easter	Airline(+mean)
RSA3	automatic	AO/LS/TC	NA	automatic
RSA4c	automatic	AO/LS/TC	2 TD vars+Easter	automatic
RSA5	automatic	AO/LS/TC	7 TD vars+Easter	automatic
X-11	NA	NA	NA	NA

explanations :

NA : non applied

X11

RSA0

RSA3: no calendar effect correction

automatic: test is performed,

outliers : AO/LS/TC type of outliers automatically detected under a critical T-Stat value (default value=4)

calendar

- 2 regressors: weekdays vs week-ends + LY

- 7 regressors: each week day vs sunadys + LY
- always tested
- easter tested (length = 6 days in tramo, 8 days in X13 arima)

arima model automatic detection

TEST : RSA0 : mult or add

TD vars include LY (link to calendar)

Strating point for Tramo-Seats

Spec identifier	Log/level detection	Outliers detection	Calendar effects	ARIMA
RSA0	<i>NA</i>	<i>NA</i>	<i>NA</i>	Airline(+mean)
RSA1	automatic	AO/LS/TC	<i>NA</i>	Airline(+mean)
RSA2	automatic	AO/LS/TC	2 TD vars+Easter	Airline(+mean)
RSA3	automatic	AO/LS/TC	<i>NA</i>	automatic
RSA5	automatic	AO/LS/TC	6 TD vars+Easter	automatic
RSAfull	automatic	AO/LS/TC	automatic	automatic

Principle of user setting parameters: can be done from one of the default specs or any spec in “save” as mode very similar inGUI and R

Spans

Estimation span

principle + some differences between tramo and reg-arima

Specifies the span (data interval) of the time series to be used in the seasonal adjustment process (no computations outside) The user can restrcit the span

Table Common settings

Option	Description (expected format)
All	default
From	first observation included (yyyy-mm-dd)
To	last observation included (yyyy-mm-dd)
Between	interval [from ; to] inculded (yyyy-mm-dd to yyyy-mm-dd)

Option	Description (expected format)
First	number of obs from the begining of the series included (dynamic) (integer)
Last	number of obs from the end of the series (dynamic)(integer)
Excluding	excluding N first obs and P last obs from the computation,dynamic) (integer)
Preliminarycheck	to exclude highly problematic series e.g. the series with a number of identical observations and/or missing values above pre-specified threshold values. (True/False)

Setting in GUI image1: specification window and expanded nodes

Setting in R, part of modifying a specification ([link](#))

```
library(RJDemetra)
# estimation interval: option with static dates
user_spec_1<-x13_spec(spec = c("RSA5c", "RSA0", "RSA1", "RSA2c", "RSA3", "RSA4c", "X11"),
preliminary.check = TRUE,
estimate.from = "2012-06-01",
estimate.to = "2019-12-01")

# estimation interval: option with dynamic numbers of observations

#
# spec can be applied on different series and therefore exclude different dates
user_spec_2<-x13_spec(spec = c("RSA5c", "RSA0", "RSA1", "RSA2c", "RSA3", "RSA4c", "X11"),
estimate.first = 12)

# eestimation on the last 120 obs
user_spec_3<-x13_spec(spec = c("RSA5c", "RSA0", "RSA1", "RSA2c", "RSA3", "RSA4c", "X11"),
estimate.last = 120)

#excluding first 24 and last 36 observations
user_spec_4<-x13_spec(spec = c("RSA5c", "RSA0", "RSA1", "RSA2c", "RSA3", "RSA4c", "X11"),
estimate.exclFirst = 24,
estimate.exclLast = 36)

# Retrieve settings
```

For comprehensive details about `x13_spec` function see {RJDemetra} documentation for `tramo`
`seats` : the same ..one example outlining differences

[V3(RJDemetra+ V3) Example in X13]

Tramo-seats

(JD+ version 2) With RJDemetra, example in Tramo-Seats:

```
#excluding first 24 and last 36 observations
user_spec_1<-tramoseats_spec( spec = c("RSAfull", "RSA0", "RSA1", "RSA2", "RSA3", "RSA4",
estimate.exclFirst = 24,
estimate.exclLast = 36)
```

Model span

Principle: same as estimation span ([link](#)) above

Specifies the span (data interval) of the time series to be used for the estimation of the RegARIMA model coefficients. The RegARIMA model is then applied to the whole series. With this argument the data early in the series can be prevented from affecting the forecasts, or, alternatively, data later in the series are excluded from the modelling span, so that they do not influence the regression estimates. The available parameter values are

Additionnal parameters

Model span specific parameters

Tolerance	Convergence tolerance for the nonlinear estimation. The absolute changes in the log-likelihood are compared to Tolerance to check the convergence of the estimation iterations. The default setting is 0.0000001.
Tramo specific parameters	
Exact ML	When this option is marked, an exact maximum likelihood estimation is performed. Alternatively, the Unconditional Least Squares method is used. However, in the current version of RJDemetra+ it is not recommended to change this parameter's value
Unit Root Limit	Limit for the autoregressive roots. If the inverse of a real root of the autoregressive polynomial of the ARIMA model is higher than this limit, the root is set equal to 1. The default parameter value is 0.96.

Retrieve in GUI:

image specification window and nodes

Retrieve in R: not possible (to confirm), estimation span and model span must be equal

so additional parameters for model span are set on the estimation span

Tramo seats exemple

```
#excluding first 24 and last 36 observations
user_spec_1<-tramoseats_spec( spec = c("RSAfull", "RSA0", "RSA1", "RSA2", "RSA3", "RSA4",
estimate.tol = 0.0000001,
estimate.eml = FALSE,
estimate.urfinal = 0.98)
```

[V3 model span specif parameters]

Decomposition Scheme

Principle

écriture add

écriture multi

if multi : log

écriture log

consequences on X11 decomp and seats decomp

Transformation test – a test is performed to choose between an additive decomposition (no transformation) (link to reg A chap to detail this)

Settings

Function

transform {function=}

Transformation of data. 2 The user can choose between:

None – no transformation of the data;

Log – takes logs of the data;

Auto – the program tests for the log-level specification. This option is recommended for automatic modelling of many series.

The default setting is Auto.

Reg-Arima specific settings

AIC difference

`transform {aicdiff=}`

Defines the difference in AICC needed to accept no transformation over a log transformation when the automatic transformation

selection option is invoked. The option is disabled when `Function` is not set to `Auto`. The de

Adjust

`transform {adjust=}`

Options for proportional adjustment for the leap year effect. The option is available when `Function` is set to `Log`. `Adjust` can be set to:

`LeapYear` – performs a leap year adjustment of monthly or quarterly data;

`LengthOfPeriod` – performs a length-of-month adjustment on monthly data or length-of-quarter adjustment on quarterly data;

`None` – does not include a correction for the length of the period.

The default setting is `None`

Tramo specific settings

Fct

Transformation; fct

Controls the bias in the log/level pre-test (the function is active when **Function** is set to *Auto*); **Fct** > 1 favours levels, **Fct** < 1 favours logs. The default setting is 0.95.

Set and Retrieve in GUI

Image specification window + expanded nodes : RegA+ Tramo

Retrieve: in the specification window and panels for main (to detail ?)

Set and in R

X13

```
#excluding first 24 and last 36 observations
user_spec <-x13_spec(spec = c("RSA5c", "RSA0", "RSA1", "RSA2c", "RSA3", "RSA4c", "X11"),
transform.function ="Log", # c(NA, "Auto", "None", "Log"),
transform.adjust = "LeapYear", #c(NA, "None", "LeapYear", "LengthOfPeriod"),
transform.aicdiff = -3)
```

```
#Retrieve settings
```

Tramo seats settings

```
#transfo
user_spec_1<-tramoseats_spec( spec = c("RSAfull", "RSA0", "RSA1", "RSA2", "RSA3", "RSA4",
transform.function = "Auto", #c(NA, "Auto", "None", "Log"),
transform.fct = 0.5)

# Retrieve settings
```

Calendar correction

Rationale : why correct (in a nutshell)

Method regression as deterministic effects, details of regressor building in calendar chapter

Default specifications for calendar correction

Options related to calendar regressors choice are embedded into default specifications ([link](#))

National calendars not taken into account.

- working days
- trading days
- Leap year regressor
- Test: Remove/ Add / None
- Easter
- Test add / remove / non
- duration (enabled when testing removed)
- Julian
- pre-test

0.0.0.0.1 * GUI: Setting this options and retrieveing parameters which are in use

0.0.0.0.2 * R: Setting this options and retrieveing parameters which are in use

Customizing calendars

How to correct this ?

solution 1: if working with GUI build a new calendar in GUI (building details in chap C) set this option import it : [here](#)

GUI : how to use it or customize HTML file structure [explanation](#)

Question: can rjd3modelling generate an HTML calendar for GUI

set this option in GUI

set this option in R

solution 2: import external regressors can be built with rjd3modelling ([link](#)) which can then be used in via are or imported via GUI

set this option in GUI

set this option in R

(generic sentence to quasi repeat at the end of each part) Once regressors set, the reg arima (tramo) model will be estimated globally with all the other regression variable and taking into account arima model specificities as well. That is why diagnostics are all jointly displayed at the end of the process. ([link](#))

[link to: worked example: french calendar in R](#)

Outliers and intervention variables

rationale

improve decomposition ([cf guidelines](#))

method

regression

Outlier types

Automatic detection

for BSM cf below modifying specifics - type - critical value - span

([link to revision policies](#))

Pre-specified Outliers

Other User-defined input : building regressors

out of calendars and outliers warning : calendars won't be allocated to calendars, if not...

rjd3modelling

Outliers vs intervention variables

assigning to componenets

Allocation of effects

(in teh decomp part for x11 and seats, table with allocation post decomp)

0.0.0.0.1 * output

Reg-Arima Model

Tramo and Reg-Arima are very similar...details in M chapter, parameters and output are the same, treated as one here

Context : here in put is spcified (auto or user) focus on global model estimation

Quick launch (= automatic run)

Output1: series (if relevant) Retrieve regressors GUI (cruncher) R

Retrieve Pre-adjustment series Table

in GUI (cruncher)

in R

Output2: Final Params what is selected and estimated by the software (not in the specifs above)

arima orders (if relevant) and coefficients

GUI

R

Output3: Diagnostics

Regression Results - tables with results for all variables (auto and user-defined) student, F test - tests

Arima model results

Goodness of fit Test on Arima model residuals list retrieve in GUI retrieve in R

Residual calendar effects what kind of test is performed ? spectral peaks - residual effect in SA or I series

Goodness of fit (principle : first diagnostics then what can be fine tuned)

User defined parameters ARIMA model parameters what can be modified

Modifying specifications for Arima model GUI

R

Modelling

(small addendum) the reg arima part can be run without decomposition slight differences in parameters

ref to additional chapter ?

X-11 Decomposition

this part should allow to use x-11 via R as well as via GUI

Default specifications

link to table from reg-arima model

if identifiers slightly different in GUI, make it clear + pb of names in V3

in calendar effects : what is tested, link to calendar chapter

typos : RJDemetra doc (pull req)

- 7 TD vars

Quick Launch

(linked from intro JDemetra+ intro chapter)

Output 1: series

Series	Final X11 name	Final Results	Reallocation of pre-adjustment effects
Raw series (forecasts)		y (y_f)	
Final seasonal component	D16	s (s_f)	
Final trend	D12	t (t_f)	
Final irregular Calendar component	D13	i (i_f)	
Seasonal without calendar	D10		

make clear reallocation of outliers effects (frequent questions)

reallocation of intervention variables effects

att : diff D10 and D16 ?

X-11 gives access to a great part of it's intermediate computations (link to M chapter)

Here we focus on the final components main series (= 2nd part of D tables, outliers effects added)

List of series (edit : table with name and meaning)

Retrieve in GUI (forecasts glued)

link to chap GUI : generate output in GUI (generic) link to chap cruncher : generate output with cruncher

Retrieve in R (forecast in a specific variable)

Link to appendix: final output overview

Output 2: final parameters

Relevant if parameters not set manually, or any parameters automatically selected by the software without having a fixed default value. (The rest of the parameters is set in the spec) To manually set those parameters and see all the fixed default values see Specifications / parameters section

Final trend filter : length of Henderson filter applied for final estimation (in the second part of the D step).

Final seasonal filter: length of Henderson filter applied for final estimation (in the second part of the D step).

Retrieve via GUI image ? Node Decomposition(X11) > Final Filters (make node clickable, link to GUI chapter, handling of nodes)

Retrieve in R From the `model_sa` object (link to R package chap global explanation)

```
model_sa$decomposition$s_filter
model_sa$decomposition$t_filter

# add version 3 equivalent
```

Output 3: diagnostics

X11 produces the following type diagnostics or quality measures Table with link to detail - SI ratios

SI-ratios

It is a plot (paste example)

Description, and computation details

Retrieve in GUI NODE Main Results > SI-Ratios

in GUI values cannot be retrieved Retrieve in R

```
# data frame with values
model_sa$decomposition$si_ratio
# customizable plot
plot(model_sa, type= "cal-seas-irr",first_date = c(2015, 1))

# add version 3 equivalent
```

M-statistics

At the end of the decomposition, X-11 algorithm provides quality measures of the decomposition called “M statistics”: 11 statistics (M1 to M11) and 2 summary indicators (Q et Q-M2). By design (clickable) $0 < M_x < 3$ and acceptance region is $M_x \leq 1$ Formula for Q (They are based on F-tables)

- **M1** The relative contribution of the irregular over three months span
- **M2** The relative contribution of the irregular component to the stationary portion of the variance
- **M3** The amount of month to month change in the irregular component as compared to the amount of month to month change in the trend-cycle (I/C-ratio)
- **M5** MCD (Months for Cyclical Dominance): The number of months it takes the change in the trend-cycle to surpass the amount of change in the irregular
- **M6** The amount of year to year change in the irregular as compared to the amount of year to year change in the seasonal (only valid for 3x5 seasonal filter)
- **M7** The amount of moving seasonality present relative to the amount of stable seasonality
- **M8** The size of the fluctuations in the seasonal component throughout the whole series
- **M9** The average linear movement in the seasonal component throughout the whole series
- **M10** Same as 8, calculated for recent years only (4 years, N-2 to N-5)
- **M11** Same as 9, calculated for recent years only

For further details link to Lothian-Mory (quid in M_meth X11)

Retrieve in GUI NODE Decomposition(X-11) > Quality Measures > Summary

Retrieve in R

```
model_sa$decomposition$mstats  
# add version 3 equivalent
```

Detailed Quality measures

List (+ link to chap M: add desc from Lothian Mory/ current doc) make clear: on which x11 series is the measure based

in GUI all the diagnostics below can be retrieved in NODE Decomposition(X-11) > Quality Measures > Details

If a stat can be retrieved in R, the code example is in the corresponding paragraph, otherwise it is not possible

- 0.0.0.0.1 *** Average differences without regard to sign over the indicated span
- 0.0.0.0.2 *** Relative contribution to the variance of the differences in the components of the original series
- 0.0.0.0.3 *** Average differences with regard to sign and standard deviation over indicated span
- 0.0.0.0.4 *** Average duration of run
- 0.0.0.0.5 *** I/C ratio over indicated span
- 0.0.0.0.6 *** I/C ratio (global)
- 0.0.0.0.7 *** Relative contribution to the stationary part of the variance in the original series
- 0.0.0.0.8 *** Autocorrelations in the irregular
- 0.0.0.0.9 *** Heteroskedasticity
Cochran test on equal variances within each period
- 0.0.0.0.10 *** Moving seasonality ratios (MSR)

User-defined parameters

Here is described how to change default values or automatic choices.

0.0.0.0.1 * General settings

- **Mode**

- check if this option still works, if so add and edit instructions from old page)
- if not but button present : explain that the mode is determined in pre-adjustment (function)

- **Seasonal component**

- check if still relevant, idem as above
- in v.2.3 if not ticked, S estimated but options on seasonal filter not available

- **Forecasts horizon**

Length of the forecasts generated by the RegARIMA model - in months (positive values) - years (negative values) - if set to is set to 0, the X-11 procedure does not use any model-based forecasts but the original X-11 type forecasts for one year. - default value: -1, thus one year from the Arima model

- **Backcasts horizon**

Length of the backcasts generated by the RegARIMA model - in months (positive values) - years (negative values) - default value: 0

0.0.0.0.2 * Irregular correction

- **LSigma**

- sets lower sigma (standard deviation) limit used to down-weight the extreme irregular values in the internal seasonal adjustment iterations, learn more here (LINK to M_ chapter)
- values in $[0, U\sigma]$
- default value is 1.5

- **USigma**

- sets upper sigma (standard deviation)
- values in $[L\sigma, +\infty]$
- default value is 2.5

- **Calendarsigma**

- allows to set different **LSigma** and **USigma** for each period
- values
 - * None (default)

- * All: standard errors used for the extreme values detection and adjustment computed separately for each calendar month/quarter
- * Signif: groups determined by cochrane test (check)
- * Sigmavec: set two customized groups of periods

- **Excludeforecasts**

- ticked : forecasts and backcasts from the RegARIMA model not used in Irregular Correction
- unticked (default): forecasts and backcasts used

0.0.0.0.3 * Seasonality extraction filters choice

Specifies which be used to estimate the seasonal factors for the entire series (link to relevant part in M chapter):

- **Seasonal filter**

- default value: *MSR* (Moving seasonality ratio), automatic choice of final seasonal filter (cf msr definition and decision table: link), initial filters are computed with 3×3
- choices : 3×1 , 3×3 , 3×5 , 3×9 , 3×15 or Stable
- “Stable” : constat factor for each calendar period (simple moving average of all $S + I$ values for each period)

User choices will be applied to all steps or only to final phase D step.

The seasonal filters can be selected for the entire series, or for a particular month or quarter.

- **Details on seasonal filters**

Sets different seasonal filters by period in order to account for seasonal heteroskedasticity (link to M chapter)

- default value: empty

0.0.0.0.4 * Trend estimation filters

- **Automatic Henderson filter** our user-defined

- default: length 13
- unticked: user defined length choice

- **Henderson filter** length choice

- values: odd number in $[3, 101]$

– default value: 13

Check: will user choice be applied to all steps or only to final phase D step

Parameter setting in GUI

Specification (general or particular to one series) image

All the parameters above can be set with the specification ([link to general explanations](#)) box

Parameter setting in R packages

extensive help on functions available in package help pages Rcode snippets

In R, to implement any param change, it is required to retrieve current spec, modify it and apply it again (see T R packages chapter for details). ([specific link](#))

here example changing all the settings (just remove irrelevant changes)

Rjdemetra (v2) Edit : here static R code link to a “worked example” with dynamic code ticked box in GUI corresponds to ...? in R

```
#Creating a modified specification, customizing all available X11 parameters
modified_spec<- x13_spec(current_sa_model,
  #x11.mode="?",
  #x11.seasonalComp = "?",
  x11.fcasts = -2,
  x11.bcasts = -1,
  x11.lsigma = 1.2,
  x11.usigma = 2.8,
  x11.calendarSigma = NA, # EDIT with example
  x11.sigmaVector = NA,
  x11.excludeFcasts = NA
  # filters
  x11.trendAuto = NA, # needed inf value ?
  x11.trendma = 23,
  x11.seasonalma = "S3X9
  # details on seasonal filters ?)

#New SA estimation : apply modified_spec
modified_sa_model<-x13(raw_series,modified_spec)
```

EDIT : [link to package help page v2 +v3](#)

SEATS Decomposition

policy: what is readable in GUI should be explained (even if shortly with link to details here)
intermediate series ?

decomp of linearized series in level or log

Default specifications

link to Tramo-Seats part

if identifiers slightly different in GUI, make it clear + pb of names in V3

in calendar effects : what is tested, link to calendar chapter

typos : RJDemetra doc (pull req)

- 7 TD vars

Quick Launch

overall procedure

in GUI

in R

Output 1: series

global indications: __f: forecasts - __e - -ef

Stochastic series

decomposition of the linearized series or of its log if multiplicative model

components: t_lin, s_lin, i_lin

if log, everything is displayed in log

Series Name	Meaning
decomposition.y_lin	
decomposition.y_lin_f	
decomposition.y_lin_ef	

Series Name	Meaning
decomposition.t_lin	
decomposition.t_lin_f	
decomposition.t_lin_e	
decomposition.t_lin_ef	
decomposition.sa_lin	
decomposition.sa_lin_f	
decomposition.sa_lin_e	
decomposition.sa_lin_ef	
decomposition.s_lin	
decomposition.s_lin_f	
decomposition.s_lin_e	
decomposition.s_lin_ef	
decomposition.i_lin	
decomposition.i_lin_f	
decomposition.i_lin_e	
decomposition.i_lin_ef	

Retrieve in GUI

NODE : Decompostion > Stocastic series

forecast f series are pasted at the end of the correspoding series

- __e series are labelled with (stde)

__ef appear in graphs (but values cannot be retrieved)

IMAGE1 GUI

- image 1 : decompostion node and series
- image 2 trend graph
- image 3 seasonal graph

Retrieve in R

Components (Level)

if additive model : same as

decompostion of the linearisezed series or of its log if multiplicative model

components: t_lin, s_lin, i_lin

if log, everything is displayed in log

Series Name	Meaning
decomposition.y_lin	
decomposition.y_lin_f	
decomposition.y_lin_ef	
decomposition.t_lin	
decomposition.t_lin_f	
decomposition.t_lin_e	
decomposition.t_lin_f	
decomposition.sa_lin	
decomposition.sa_lin_f	
decomposition.sa_lin_e	
decomposition.sa_lin_ef	
decomposition.s_lin	
decomposition.s_lin_f	
decomposition.s_lin_e	
decomposition.s_lin_ef	
decomposition.i_lin	
decomposition.i_lin_f	
decomposition.i_lin_e	
decomposition.i_lin_ef	

Retrieve in GUI

NODE : Decompostion > Stocastic series

forecast f series are pasted at the end of the correspodng series

- __e series are labelled with (stde)

__ef appear in graphs (but values cannot be retrieved)

IMAGE1 GUI

- image 1 : decompostion node and series
- image 2 trend graph
- image 3 seasonal graph

Retrieve in R

Biais correction voir JP (done between lin and and comp ?)

Final series

decompostion of the linearisezed series or of its log if multiplicative model

components: t_lin, s_lin, i_lin

if log, everything is dispalayed in log

Series Name	Meaning
decomposition.y_lin	
decomposition.y_lin_f	
decomposition.y_lin_ef	
decomposition.t_lin	
decomposition.t_lin_f	
decomposition.t_lin_e	
decomposition.t_lin_ef	
decomposition.sa_lin	
decomposition.sa_lin_f	
decomposition.sa_lin_e	
decomposition.sa_lin_ef	
decomposition.s_lin	
decomposition.s_lin_f	
decomposition.s_lin_e	
decomposition.s_lin_ef	
decomposition.i_lin	
decomposition.i_lin_f	
decomposition.i_lin_e	
decomposition.i_lin_ef	

Retrieve in GUI

NODE : Decompostion > Stocastic series

forecast f series are pasted at the end of the corresponding series

- __e series are labelled with (stde)

__ef appear in graphs (but values cannot be retrieved)

IMAGE1 GUI

- image 1 : decompostion node and series
- image 2 trend graph
- image 3 seasonal graph

Retrieve in R

make clear reallocation of outliers effects (frequent questions)

Table

GUI

R

Output2: Final Params

what is chosen automatically or default (cf estp) Relevant if parameters not set manually, or any parameters automatically selected by the software without having a fixed default value. (The rest of the parameters is set in the spec) To manually set those parameters and see all the fixed default values see Specifications / parameters section

here all have a default value, nothing to be automatically selected

Output3: Diagnostics

SEATS (model based) specific diagnostics check GUI (more plots, values) vs R links to methods chapter, here just be able to “read” (knowledge assumed) (here direct base on gui plots, and the add what can be retrieved in R, if cannot be retrieved in R : could it be rebuild from available elements) Obj : simple description res : current doc, jpslides, doc esp, Q JP

- WK analysis

components final estimators

- Error analysis autocorrelation of the errors (sa, trend) revisions of the errors
- Growth rates
- Model based tests
- Significant seasonality
- Stationnary variance decomposition

Fine-tuning : user defined parameters

- Prediction length

Forecast span used in the decomposition default: one year (-1) (years are set in negative values, positive values indicate number of periods)

- Approximation Mode

Modification type for inadmissible models None (default) Legacy Noisy

- MA unit root boundary

Modulus threshold for resetting MA “near-unit” roots [0,1] default (0.95)

- Trend Boundary Modulus threshold for assigning positive real AR Roots [0,1] default (0.5)
- Seasonal Tolerance Degree threshold for assigning complex AR roots [0,10] default (2)
- Seasonal Boundary (unique) Modulus threshold for assigning negative real AR roots [0,1] default (0.8)
- Seasonal Boundary (unique) Same modulus threshold unique seasonal AR roots [0,1] default (0.8)
- Method

Algorithm used for estimation of unobserved components

Burman (default)

KalmanSmoother

McElroyMatrix

Further explanation (cf slides JP)

link to M chapter for details on root allocation

0.0.0.0.1 * Set in GUI

In specification window (link to GUI global exp)

IMAGE

0.0.0.0.2 * Set in R

version 2 (RJDemetra)

```

tramoseats_spec(
  spec = c("RSAfull", "RSA0", "RSA1", "RSA2", "RSA3", "RSA4", "RSA5"),
  fcst.horizon = NA_integer_,
  seats.predictionLength = NA_integer_,
  seats.approx = c(NA, "None", "Legacy", "Noisy"),
  seats.trendBoundary = NA_integer_,
  seats.seasBoundary = NA_integer_,
  seats.seasBoundary1 = NA_integer_,
  seats.seasTol = NA_integer_,
  seats.maBoundary = NA_integer_,
  seats.method = c(NA, "Burman", "KalmanSmoother", "McElroyMatrix")
)

```

in version 3 with {rjd3tramoseats}

STL

use on linearized data data, no pre-adjustment (2nd layer)

SSF

(2nd layer)

Quality assessment

Residual seasonality

Residual calendar effects

(all GUI based analysis) NODES below bechamrking

Matrix

Add other diagnostics node

Seasonal adjustment of high-frequency data

Motivation: data specificities

Infra monthly data displays multiple and non integer periodicities which cannot be dealt with classical versions of SA algorithms. JD+ provides tailored versions of these algorithms.

[Link to worked example \(Daily French births\)](#)

Available algorithms

code here and/or link to R packages chapter

Col1	Algorithm	GUI v 3.0	R package / function
Pre-treatment	Fractionnal Airline Model	yes	
Decompsition	Extended SEATS Fractionnal Airline Model	yes	
	Extended X-11	yes	
	Extended STL	no	
One-Step	SSF Framework	no	

Unobserved Components

Identifying seasonal patterns

Spectral analysis

Seasonality tests

Pre-adjustment

Calendar correction

generate

Outliers and intervention variables

add

Linearization

Decomposition

Extended X-11

Arima Model Based (AMB) Decomposition

STL decomposition

State Space framework

one step for pre-treatment and decomp

Quality assessment

Residual seasonality

Residual Calendar effects

Outlier detection

(in or outside a seasonal adjustment process)

Motivation

With Reg Arima models

Specific TERROR tool

With structural models (BSM)

Calendar and user-defined corrections

This chapter describes the generating process of calendar regressors, outliers and other input variables. The use of this variables inside a seasonal adjustment process is described in the relevant chapter on SA or on SA of HF data.

Overview of Calendar effects in JDemetra

edit : this has evolved a lot with v3 definition possibilities via GUI and R have to be re described

The following description of the calendar effects in JDemetra+ is strictly based on PALATE, J. (2014).

A natural way for modelling calendar effects consists of distributing the days of each period into different groups. The regression variable corresponding to a type of day (a group) is simply defined by the number of days it contains for each period. Usual classifications are:

- Trading days (7 groups): each day of the week defines a group (Mondays,...,Sundays);
- Working days (2 groups): week days and weekends.

The definition of a group could involve partial days. For instance, we could consider that one half of Saturdays belong to week days and the second half to weekends.

Usually, specific holidays are handled as Sundays and they are included in the group corresponding to "non-working days". This approach assumes that the economic activity on national holidays is the same (or very close to) the level of activity that is typical for Sundays. Alternatively, specific holidays can be considered separately, e.g. by the specification that divided days into three groups:

- Working days (Mondays to Fridays, except for specific holidays),
- Non-working days (Saturdays and Sundays, except for specific holidays),
- Specific holidays.

Summary of the method used in JDemetra+ to compute trading day and working day effects

The computation of trading day and working days effects is performed in four steps:

1. Computation of the number of each weekday performed for all periods.
2. Calculation of the usual contrast variables for trading day and working day.
3. Correction of the contrast variables with specific holidays (for each holiday add +1 to the number of Sundays and subtract 1 from the number of days of the holiday). The correction is not performed if the holiday falls on a Sunday, taking into account the validity period of the holiday.
4. Correction of the constant variables for long term mean effects, > taking into account the validity period of the holiday; see below > for the different cases.

The corrections of the constant variables may receive a weight corresponding to the part of the holiday considered as a Sunday.

An example below illustrates the application of the above algorithm for the hypothetical country in which three holidays are celebrated:

- New Year (a fixed holiday, celebrated on 01 January);
- Shrove Tuesday (a moving holiday, which falls 47 days before Easter Sunday, celebrated until the end of 2012);
- Freedom day (a fixed holiday, celebrated on 25 April).

The consecutive steps in calculation of the calendar for 2012 and 2013 years are explained below.

First, the number of each day of the week in the given month is calculated as it is shown in table below.

Number of each weekday in different months

Month	Mon	Tue	Wed	Thu	Fri	Sat	Sun
Jan-12	5	5	4	4	4	4	5
Feb-12	4	4	5	4	4	4	4
Mar-12	4	4	4	5	5	5	4
Apr-12	5	4	4	4	4	4	5
May-12	4	5	5	5	4	4	4
Jun-12	4	4	4	4	5	5	4
Jul-12	5	5	4	4	4	4	5
Aug-12	4	4	5	5	5	4	4

Month	Mon	Tue	Wed	Thu	Fri	Sat	Sun
Sep-12	4	4	4	4	4	5	5
Oct-12	5	5	5	4	4	4	4
Nov-12	4	4	4	5	5	4	4
Dec-12	5	4	4	4	4	5	5
Jan-13	4	5	5	5	4	4	4
Feb-13	4	4	4	4	4	4	4
Mar-13	4	4	4	4	5	5	5
Apr-13	5	5	4	4	4	4	4
May-13	4	4	5	5	5	4	4
Jun-13	4	4	4	4	4	5	5
Jul-13	5	5	5	4	4	4	4
Aug-13	4	4	4	5	5	5	4
Sep-13	5	4	4	4	4	4	5
Oct-13	4	5	5	5	4	4	4
Nov-13	4	4	4	4	5	5	4
Dec-13	5	5	4	4	4	4	5

Next, the contrast variables are calculated (table below) as a result of the linear transformation applied to the variables presented in table below.

Contrast variables (series corrected for leap year effects)

Month	Mon	Tue	Wed	Thu	Fri	Sat	Length
Jan-12	0	0	-1	-1	-1	-1	0
Feb-12	0	0	1	0	0	0	0.75
Mar-12	0	0	0	1	1	1	0
Apr-12	0	-1	-1	-1	-1	-1	0
May-12	0	1	1	1	0	0	0
Jun-12	0	0	0	0	1	1	0
Jul-12	0	0	-1	-1	-1	-1	0
Aug-12	0	0	1	1	1	0	0
Sep-12	-1	-1	-1	-1	-1	0	0
Oct-12	1	1	1	0	0	0	0
Nov-12	0	0	0	1	1	0	0
Dec-12	0	-1	-1	-1	-1	0	0
Jan-13	0	1	1	1	0	0	0
Feb-13	0	0	0	0	0	0	-0.25
Mar-13	-1	-1	-1	-1	0	0	0
Apr-13	1	1	0	0	0	0	0
May-13	0	0	1	1	1	0	0

Month	Mon	Tue	Wed	Thu	Fri	Sat	Length
Jun-13	-1	-1	-1	-1	-1	0	0
Jul-13	1	1	1	0	0	0	0
Aug-13	0	0	0	1	1	1	0
Sep-13	0	-1	-1	-1	-1	-1	0
Oct-13	0	1	1	1	0	0	0
Nov-13	0	0	0	0	1	1	0
Dec-13	5	5	4	4	4	4	0

In the next step the corrections for holidays is done in the following way:

- New Year: In 2012 it falls on a Sunday. Therefore no correction is applied. In 2013 it falls on a Tuesday. Consequently, the following corrections are applied to the number of each weekday in January: Tuesday -1, Sunday +1, so the following corrections are applied to the contrast variables: -2 for Tuesday and -1 for the other contrast variables.
- Shrove Tuesday: It is a fixed day of the week holiday that always falls on Tuesday. For this reason in 2012 the following corrections are applied to the number of each weekday in February: Tuesday -1, Sunday +1, so the following corrections are applied to the contrast variables: -2 for the contrast variable associated with Tuesday, and -1 for the other contrast variables. The holiday expires at the end of 2012. Therefore no corrections are made for 2013.
- Freedom Day: In 2012 it falls on a Wednesday. Consequently, the following corrections are applied to the number of each weekday in April: Wednesday -1, Sunday +1, so the following corrections are applied to the contrast variables: -2 for Wednesday and -1 for the other contrast variables. In 2013 it falls on Thursday. Therefore, the following corrections are applied to the number of each weekday in April: Thursday -1, Sunday +1, so the following corrections are applied to the contrast variables: -2 for Thursday, and -1 for the other contrast variables.

The result of these corrections is presented in table below.

Contrast variables corrected for holidays

Month	Mon	Tue	Wed	Thu	Fri	Sat	Length
Jan-12	0	0	-1	-1	-1	-1	0
Feb-12	-1	-2	0	-1	-1	-1	0.75
Mar-12	0	0	0	1	1	1	0
Apr-12	-1	-2	-3	-2	-2	-2	0
May-12	0	1	1	1	0	0	0
Jun-12	0	0	0	0	1	1	0
Jul-12	0	0	-1	-1	-1	-1	0

Month	Mon	Tue	Wed	Thu	Fri	Sat	Length
Aug-12	0	0	1	1	1	0	0
Sep-12	-1	-1	-1	-1	-1	0	0
Oct-12	1	1	1	0	0	0	0
Nov-12	0	0	0	1	1	0	0
Dec-12	0	-1	-1	-1	-1	0	0
Jan-13	-1	-1	0	0	-1	-1	0
Feb-13	0	0	0	0	0	0	-0.25
Mar-13	-1	-1	-1	-1	0	0	0
Apr-13	0	0	-1	-2	-1	-1	0
May-13	0	0	1	1	1	0	0
Jun-13	-1	-1	-1	-1	-1	0	0
Jul-13	1	1	1	0	0	0	0
Aug-13	0	0	0	1	1	1	0
Sep-13	0	-1	-1	-1	-1	-1	0
Oct-13	0	1	1	1	0	0	0
Nov-13	0	0	0	0	1	1	0
Dec-13	0	0	-1	-1	-1	-1	0

Finally, the long term corrections are applied on each year of the validity period of the holiday.

- New Year: Correction on the contrasts: +1, to be applied to January of 2012 and 2013.
- Shrove Tuesday: It may fall either in February or in March. It will fall in March if Easter is on or after 17 April. Taking into account the theoretical distribution of Easter, it gives: $\text{prob}(\text{March}) = +0.22147$, $\text{prob}(\text{February}) = +0.77853$. The correction of the contrasts will be +1.55707 for Tuesday in February 2012 and +0.77853 for the other contrast variables. The correction of the contrasts will be +0.44293 for Tuesday in March 2012, +0.22147 for the other contrast variables.
- Freedom Day: Correction on the contrasts: +1, to be applied to April of 2012 and 2013.

The modifications due to the corrections described above are presented in table below.

Trading day variables corrected for the long term effects

Month	Mon	Tue	Wed	Thu	Fri	Sat	Length
Jan-12	1	1	0	0	0	0	0
Feb-12	-0.22115	-0.44229	0.778853	-0.22115	-0.22115	-0.22115	0.75
Mar-12	0.221147	0.442293	0.221147	1.221147	1.221147	1.221147	0
Apr-12	0	-1	-2	-1	-1	-1	0
May-12	0	1	1	1	0	0	0

Month	Mon	Tue	Wed	Thu	Fri	Sat	Length
Jun-12	0	0	0	0	1	1	0
Jul-12	0	0	-1	-1	-1	-1	0
Aug-12	0	0	1	1	1	0	0
Sep-12	-1	-1	-1	-1	-1	0	0
Oct-12	1	1	1	0	0	0	0
Nov-12	0	0	0	1	1	0	0
Dec-12	0	-1	-1	-1	-1	0	0
Jan-13	0	0	1	1	0	0	0
Feb-13	0	0	0	0	0	0	-0.25
Mar-13	-1	-1	-1	-1	0	0	0
Apr-13	1	1	0	-1	0	0	0
May-13	0	0	1	1	1	0	0
Jun-13	-1	-1	-1	-1	-1	0	0
Jul-13	1	1	1	0	0	0	0
Aug-13	0	0	0	1	1	1	0
Sep-13	0	-1	-1	-1	-1	-1	0
Oct-13	0	1	1	1	0	0	0
Nov-13	0	0	0	0	1	1	0
Dec-13	0	0	-1	-1	-1	-1	0

Mean and seasonal effects of calendar variables

The calendar effects produced by the regression variables that fulfil the definition presented above include a mean effect (i.e. an effect that is independent of the period) and a seasonal effect (i.e. an effect that is dependent of the period and on average it is equal to 0). Such an outcome is inappropriate, as in the usual decomposition of a series the mean effect should be allocated to the trend component and the fixed seasonal effect should be affected to the corresponding component. Therefore, the actual calendar effect should only contain effects that don't belong to the other components.

In the context of JDemetra+ the mean effect and the seasonal effect are long term theoretical effects rather than the effects computed on the time span of the considered series (which should be continuously revised).

The mean effect of a calendar variable is the average number of days in its group. Taking into account that one year has on average 365.25 days, the monthly mean effects for a working days are, as shown in the table below, 21.7411 for week days and 8.696 for weekends.

Monthly mean effects for the Working day variable

Groups of Working day effect	Mean effect
Week days	$365.25/12*5/7 = \mathbf{21.7411}$
Weekends	$365.25/12*2/7 = \mathbf{8.696}$
Total	$365.25/12 = \mathbf{30.4375}$

The number of days by period is highly seasonal, as apart from February, the length of each month is the same every year. For this reason, any set of calendar variables will contain, at least in some variables, a significant seasonal effect, which is defined as the average number of days by period (Januaries..., first quarters...) outside the mean effect. Removing that fixed seasonal effects consists of removing for each period the long term average of days that belong to it. The calculation of a seasonal effect for the working days classification is presented in the table below.

The mean effect and the seasonal effect for the calendar periods

Period	Average number of days	Average number of week days	Mean effect	Seasonal effect
January	31	$31*5/7=22.1429$	21.7411	0.4018
February	28.25	$28.25*5/7=20.1786$	21.7411	-1.5625
March	31	$31*5/7=22.1429$	21.7411	0.4018
April	30	$30*5/7=21.4286$	21.7411	-0.3125
May	31	$31*5/7=22.1429$	21.7411	0.4018
June	30	$30*5/7=21.4286$	21.7411	-0.3125
July	31	$31*5/7=22.1429$	21.7411	0.4018
August	31	$31*5/7=22.1429$	21.7411	0.4018
September	30	$30*5/7=21.4286$	21.7411	-0.3125
October	31	$31*5/7=22.1429$	21.7411	0.4018
November	30	$30*5/7=21.4286$	21.7411	-0.3125
December	31	$31*5/7=22.1429$	21.7411	0.4018
Total	365.25	260.8929	260.8929	0

For a given time span, the actual calendar effect for week days can be easily calculated as the difference between the number of week days in a specific period and the sum of the mean effect and the seasonal effect assigned to this period, as it is shown in the table below for the period 01.2013 – 06.2013.

The calendar effect for the period 01.2013 - 06.2013

Time period (t)	Week days	Mean effect	Seasonal effect	Calendar effect
Jan-2013	23	21.7411	0.4018	0.8571

Time period (t)	Week days	Mean effect	Seasonal effect	Calendar effect
Feb-2013	20	21.7411	-1.5625	-0.1786
Mar-2013	21	21.7411	0.4018	-1.1429
Apr-2013	22	21.7411	-0.3125	0.5714
May-2013	23	21.7411	0.4018	0.8571
Jun-2013	20	21.7411	-0.3125	-1.4286
Jul-2013	23	21.7411	0.4018	0.8571

The distinction between the mean effect and the seasonal effect is usually unnecessary. Those effects can be considered together (simply called mean effects) and be computed by removing from each calendar variable its average number of days by period. These global means effect are considered in the next section.

Impact of the mean effects on the decomposition

When the ARIMA model contains a seasonal difference – something that should always happen with calendar variables – the mean effects contained in the calendar variables are automatically eliminated, so that they don't modify the estimation. The model is indeed estimated on the series/regression variables after differencing. However, they lead to a different linearised series (y_{lin}). The impact of other corrections (mean and/or fixed seasonal) on the decomposition is presented in the next paragraph. Such corrections could be obtained, for instance, by applying other solutions for the long term corrections or by computing them on the time span of the series.

Now the model with "correct" calendar effects (denoted as C), i.e. effects without mean and fixed seasonal effects, can be considered. To simplify the problem, the model has no other regression effects.

For such a model the following relations hold:

$$y_{lin} = y - C$$

$$T = F_T(y_{lin})$$

$$S = F_S(y_{lin}) + C$$

$$I = F_I(y_{lin})$$

where:

T - the trend;

S - the seasonal component;

I - the irregular component;

F_X - the linear filter for the component X.

Consider next other calendar effects (\widetilde{C}) that contain some mean (cm, integrated to the final trend) and fixed seasonal effects (cs, integrated to the final seasonal). The modified equations are now:

$$\widetilde{C} = C + cm + cs$$

$$\widetilde{y}_{\text{lin}} = y - \widetilde{C} = y_{\text{lin}} - cm - cs$$

$$\widetilde{T} = F_T(\widetilde{y}_{\text{lin}}) + cm$$

$$\widetilde{S} = F_S(\widetilde{y}_{\text{lin}}) + C + cs$$

$$\widetilde{I} = F_I(\widetilde{y}_{\text{lin}})$$

Taking into account that F_X is a linear transformation and that¹

$$F_T(cm) = cm$$

$$F_T(cs) = 0$$

$$F_S(cm) = 0$$

¹In case of SEATS the properties can be trivially derived from the matrix formulation of signal extraction. They are also valid for X-11 (additive).

$$F_S(\text{cs}) = cs$$

$$F_I(\text{cm}) = 0$$

$$F_I(\text{cs}) = 0$$

The following relationships hold:

$$\tilde{T} = F_T(\tilde{y}_{\text{lin}}) + cm = F_T(y_{\text{lin}}) - cm + cm = T$$

$$\tilde{S} = F_S(\tilde{y}_{\text{lin}}) + C + cs = F_S(y_{\text{lin}}) - cs + C + cs = S$$

$$\tilde{I} = I$$

If we don't take into account the effects and apply the same approach as in the “correct” calendar effects, we will get:

$$\check{T} = F_T(\tilde{y}_{\text{lin}}) = T - cm$$

$$\check{S} = F_S(\tilde{y}_{\text{lin}}) + \check{C} = S + cm$$

$$\check{I} = F_I(\tilde{y}_{\text{lin}}) = I$$

The trend, seasonal and seasonally adjusted series will only differ by a (usually small) constant.

In summary, the decomposition does not depend on the mean and fixed seasonal effects used for the calendar effects, provided that those effects are integrated in the corresponding final components. If these corrections are not taken into account, the main series of the decomposition will only differ by a constant.

Linear transformations of the calendar variables

As far as the RegARIMA and the TRAMO models are considered, any non-degenerated linear transformation of the calendar variables can be used. It will produce the same results (likelihood, residuals, parameters, joint effect of the calendar variables, joint F-test on the coefficients of the calendar variables...). The linearised series that will be further decomposed is invariant to any linear transformation of the calendar variables.

However, it should be mentioned that choices of calendar corrections based on the tests on the individual t statistics are dependent on the transformation, which is rather arbitrary. This is the case in old versions of TRAMO-SEATS. That is why the joint F-test (as in the version of TRAMO-SEATS implemented in TSW+) should be preferred.

An example of a linear transformation is the calculation of the contrast variables. In the case of the usual trading day variables, they are defined by the following transformation: the 6 contrast variables ($No. (Mondays) - No. (Sundays)$, ... $No. (Saturdays) - No. (Sundays)$) used with the length of period.

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} \text{Mon} \\ \text{Tue} \\ \text{Wed} \\ \text{Thu} \\ \text{Fri} \\ \text{Sat} \\ \text{Sun} \end{bmatrix} = \begin{bmatrix} \text{Mon} - \text{Sun} \\ \text{Tue} - \text{Sun} \\ \text{Wed} - \text{Sun} \\ \text{Thu} - \text{Sun} \\ \text{Fri} - \text{Sun} \\ \text{Sat} - \text{Sun} \\ \text{Length of period} \end{bmatrix}$$

For the usual working day variables, two variables are used: one contrast variable and the length of period

$$\begin{bmatrix} 1 & -\frac{5}{2} \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \text{Week} \\ \text{Weekend} \end{bmatrix} = \begin{bmatrix} \text{Contrast week} \\ \text{Length of period} \end{bmatrix}$$

The Length of period variable is defined as a deviation from the length of the month (in days) and the average month length, which is equal to 30.4375. Instead, the leap-year variable can be used here (see Regression sections in [RegARIMA](#) or [Tramo](#))².

Such transformations have several advantages. They suppress from the contrast variables the mean and the seasonal effects, which are concentrated in the last variable. So, they lead to fewer correlated variables, which are more appropriate to be included in the regression model. The sum of the effects of each day of the week estimated with the trading (working) day contrast variables cancel out.

²GÓMEZ, V., and MARAVALL, A (2001b).

Handling of specific holidays

check vs GUI (v3) and rjd3 modelling

Three types of holidays are implemented in JDemetra+:

- Fixed days, corresponding to the fixed dates in the year (e.g. New Year, Christmas).
- Easter related days, corresponding to the days that are defined in relation to Easter (e.g. Easter +/- n days; example: Ascension, Pentecost).
- Fixed week days, corresponding to the fixed days in a given week of a given month (e.g. Labor Day celebrated in the USA on the first Monday of September).

From a conceptual point of view, specific holidays are handled in exactly the same way as the other days. It should be decided, however, to which group of days they belong. Usually they are handled as Sundays. This convention is also used in JDemetra+. Therefore, except if the holiday falls on a Sunday, the appearance of a holiday leads to correction in two groups, i.e. in the group that contains the weekday, in which holiday falls, and the group that contains the Sundays.

Country specific holidays have an impact on the mean and the seasonal effects of calendar effects. Therefore, the appropriate corrections to the number of particular days (which are usually the basis for the definition of other calendar variables) should be applied, following the kind of holidays. These corrections are applied to the period(s) that may contain the holiday. The long term corrections in JDemetra+ don't take into account the fact that some moving holidays could fall on the same day (for instance the May Day and the Ascension). However, those events are exceptional, and their impact on the final result is usually not significant.

Fixed day

The probability that the holiday falls on a given day of the week is $1/7$. Therefore, the probability to have 1 day more that is treated like Sunday is $6/7$. The effect on the means for the period that contains the fixed day is presented in the table below (the correction on the calendar effect has the opposite sign).

The effect of the fixed holiday on the period, in which it occurred

Sundays	Others days	Contrast variables
+ $6/7$	- $1/7$	$1/7 - (+ 6/7) = -1$

Easter related days

Easter related days always fall the same week day (denoted as Y in the table below: The effects of the Easter Sunday on the seasonal means). However, they can fall during different periods (months or quarters). Suppose that, taking into account the distribution of the dates for Easter and the fact that this holiday falls in one of two periods, the probability that Easter falls during the period m is p , which implies that the probability that it falls in the period $m + 1$ is $1 - p$. The effects of Easter on the seasonal means are presented in the table below.

The effects of the Easter Sunday on the seasonal means

Period	Sundays	Days X Others	days Contrast Y	Other contrasts
m	p	$0 - 2p - p$	$m + 1$	$1 - p$

The distribution of the dates for Easter may be approximated in different ways. One of the solutions consists of using some well-known algorithms for computing Easter on a very long period. JDemetra+ provides the Meeus/Jones/Butcher's and the Ron Mallen's algorithms (they are identical till year 4100, but they slightly differ after that date). Another approach consists in deriving a raw theoretical distribution based on the definition of Easter. It is the solution used for Easter related days. It is shortly explained below.

The date of Easter in the given year is the first Sunday after the full moon (the Paschal Full Moon) following the northern hemisphere's vernal equinox. The definition is influenced by the Christian tradition, according to which the equinox is reckoned to be on 21 March³ and the full moon is not necessarily the astronomically correct date. However, when the full moon falls on Sunday, then Easter is delayed by one week. With this definition, the date of Easter Sunday varies between 22 March and 25 April. Taking into account that an average lunar month is 29.530595 days the approximated distribution of Easter can be derived. These calculations do not take into account the actual ecclesiastical moon calendar.

For example, the probability that Easter Sunday falls on 25 March is 0.004838 and results from the facts that the probability that 25 March falls on a Sunday is $1/7$ and the probability that the full moon is on 21 March, 22 March, 23 March or 24 March is $5/29.53059$. The probability that Easter falls on 24 April is 0.01708 and results from the fact that the probability that 24 April is Sunday is $1/7$ and takes into account that 18 April is the last acceptable date for the full moon. Therefore the probability that the full moon is on 16 April or 17 April is $1/29.53059$ and the probability that the full moon is on 18 April is $1.53059/29.53059$.

The approximated distribution of Easter dates

Day	Probability
22 March	$1/7 * 1/29.53059$
23 March	$1/7 * 2/29.53059$

³In fact, astronomical observations show that the equinox occurs on 20 March in most years.

Day	Probability
24 March	$1/7 * 3/29.53059$
25 March	$1/7 * 4/29.53059$
26 March	$1/7 * 5/29.53059$
27 March	$1/7 * 6/29.53059$
28 March	$1/29.53059$
29 March	$1/29.53059$
...	...
18 April	$1/29.53059$
19 April	$1/7 * (6 + 1.53059)/29.53059$
20 April	$1/7 * (5 + 1.53059)/29.53059$
21 April	$1/7 * (4 + 1.53059)/29.53059$
22 April	$1/7 * (3 + 1.53059)/29.53059$
23 April	$1/7 * (2 + 1.53059)/29.53059$
24 April	$1/7 * (1 + 1.53059)/29.53059$
25 April	$1/7 * 1.53059/29.53059$

Fixed week days

Fixed week days always fall on the same week day (denoted as Y in the table below) and in the same period. Their effect on the seasonal means is presented in the table below.

The effect of the fixed week holiday on the period, in which it occurred

Sundays	Day Y	Others days
+ 1	- 1	0

The impact of fixed week days on the regression variables is zero because the effect itself is compensated by the correction for the mean effect.

Holidays with a validity period

When a holiday is valid only for a given time span, JDemetra+ applies the long term mean corrections only on the corresponding period. However, those corrections are computed in the same way as in the general case.

It is important to note that using or not using mean corrections will impact in the estimation of the RegARIMA and TRAMO models. Indeed, the mean corrections do not disappear after differencing. The differences between the SA series computed with or without mean corrections will no longer be constant.

Different Kinds of calendars

see link with GUI

This scenario presents how to define different kinds of calendars. These calendars can be applied to the specifications that take into account country-specific holidays and can be used for detecting and estimating the calendar effects.

The calendar effects are those parts of the movements in the time series that are caused by different number of weekdays in calendar months (or quarters). They arise as the number of occurrences of each day of the week in a month (or a quarter) differs from year to year. These differences cause regular effects in some series. In particular, such variation is caused by a leap year effect because of an extra day inserted into February every four years. As with seasonal effects, it is desirable to estimate and remove calendar effects from the time series.

The calendar effects can be divided into a mean effect, a seasonal part and a structural part. The mean effect is independent from the period and therefore should be allocated to the trend-cycle. The seasonal part arises from the properties of the calendar that recur each year. For one thing, the number of working days of months with 31 calendar days is on average larger than that of months with 30 calendar days. This effect is part of the seasonal pattern captured by the seasonal component (with the exception of leap year effects). The structural part of the calendar effect remains to be determined by the calendar adjustment. For example, the number of working days of the same month in different years varies from year to year.

Both X-12-ARIMA/X-13ARIMA-SEATS and TRAMO/SEATS estimate calendar effects by adding some regressors to the equation estimated in the pre-processing part (RegARIMA or TRAMO, respectively). Regressors mentioned above are generated from the default calendar or the user defined calendar.

The calendars of JDemetra+ simply correspond to the usual trading days contrast variables based on the Gregorian calendar, modified to take into account some specific holidays. Those holidays are handled as "Sundays" and the variables are properly adjusted to take into account the long term mean effects.

Tests for residual trading days

We consider below tests on the seasonally adjusted series (sa_t) or on the irregular component (irr_t). When the reasoning applies on both components, we will use y_t . The functions *stdev* stands for "standard deviation" and *rms* for "root mean squares"

The tests are computed on the log-transformed components in the case of multiplicative decomposition.

TD are the usual contrasts of trading days, 6 variables (no specific calendar).

Non significant irregular

When irr_t is not significant, we don't compute the test on it, to avoid irrelevant results. We consider that irr_t is significant if $stdev(irr_t) > 0.01$ (multiplicative case) or if $stdev(irr_t)/rms(sa_t) > 0.01$ (additive case).

F test

The test is the usual joint F-test on the TD coefficients, computed on the following models:

0.0.0.0.1 * Autoregressive model (AR modelling option)

We compute by OLS:

$$y_t = \mu + \alpha y_{t-1} + \beta T D_t + \epsilon_t$$

0.0.0.0.2 * Difference model

We compute by OLS:

$$\Delta y_t - \overline{\Delta y_t} = \beta T D_t + \epsilon_t$$

So, the latter model is a restriction of the first one ($\alpha = 1, \mu = \mu = \overline{\Delta y_t}$)

The tests are the usual joint F-tests on β ($H_0 : \beta = 0$).

By default, we compute the tests on the 8 last years of the components, so that they might highlight moving calendar effects.

Remark:

In Tramo, a similar test is computed on the residuals of the Arima model. More exactly, the F-test is computed on $e_t = \beta T D_t + \epsilon_t$, where e_t are the one-step-ahead forecast errors.

Benchmarking and temporal disaggregation

In this chapter we describe the practical implementation, the underlying theory in a dedicated chapter.[\(link\)](#)

Benchmarking overview

Often one has two (or multiple) datasets of different frequency for the same target variable. Sometimes, however, these data sets are not coherent in the sense that they don't match up. Benchmarking^[1] is a method to deal with this situation. An aggregate of a higher-frequency measurement variables is not necessarily equal to the corresponding lower-frequency less-aggregated measurement. Moreover, the sources of data may have different reliability levels. Usually, less frequent data are considered more trustworthy as they are based on larger samples and compiled more precisely. The more reliable measurements, hence often the less frequent, will serve as benchmark.

In seasonal adjustment methods benchmarking is the procedure that ensures the consistency over the year between adjusted and non-seasonally adjusted data. It should be noted that the [ESS Guidelines on Seasonal Adjustment (2015)] (<https://ec.europa.eu/eurostat/documents/3859598/6830795/KGQ-15-001-EN-N.pdf/d8f1e5f5-251b-4a69-93e3-079031b74bd3>), do not recommend benchmarking as it introduces a bias in the seasonally adjusted data. The U.S. Census Bureau also points out that “*forcing the seasonal adjustment totals to be the same as the original series annual totals can degrade the quality of the seasonal adjustment, especially when the seasonal pattern is undergoing change. It is not natural if trading day adjustment is performed because the aggregate trading day effect over a year is variable and moderately different from zero*”^[2]. Nevertheless, some users may need that the annual totals of the seasonally adjusted series match the annual totals of the original, non-seasonally adjusted series^[3].

According to the [ESS Guidelines on Seasonal Adjustment (2015)] (<https://ec.europa.eu/eurostat/documents/3859598/6830795/KGQ-15-001-EN-N.pdf/d8f1e5f5-251b-4a69-93e3-079031b74bd3>), the only benefit of this approach is that there is consistency over the year between adjusted and the non-seasonally adjusted data; this can be of particular interest when low-frequency (e.g. annual) benchmarking figures officially exist (e.g. National Accounts, Balance of Payments, External Trade, etc.) and where users' needs for time consistency are stronger.

Tools

Benchmarking with GUI

1. With the [pre-defined specifications](#) the benchmarking functionality is not applied by default following the *ESS Guidelines on Seasonal Adjustment* (2015) recommendations. It means that once the user has seasonally adjusted the series with a pre-defined specification the *Benchmarking* node is empty. To execute benchmarking click on the *Specifications* button and activate the checkbox in the *Benchmarking* section.




Figure 1: Text

Benchmarking option – a default view

2. Three parameters can be set here. *Target* specifies the target variable for the benchmarking procedure. It can be either the *Original* (the raw time series) or the *Calendar Adjusted* (the time series adjusted for calendar effects). *Rho* is a value of the AR(1) parameter (set between 0 and 1). By default it is set to 1. Finally, *Lambda* is a parameter that relates to the weights in the regression equation. It is typically equal to 0 (for an additive decomposition), 0.5 (for a proportional decomposition) or 1 (for a multiplicative decomposition). The default value is 1.
3. To launch the benchmarking procedure click on the **Apply** button. The results are displayed in four panels. The top-left one compares the original output from the seasonal adjustment procedure with the result from applying a benchmarking to the seasonal

adjustment. The bottom-left panel highlights the differences between these two results. The outcomes are also presented in a table in the top-right panel. The relevant statistics concerning relative differences are presented in the bottom-right panel.




Figure 2: Text

The results of the benchmarking procedure

- Both pictures and the table can be copied the usual way (see the *Simple seasonal adjustment of a single time series* scenario).

Options for benchmarking results

- To export the result of the benchmarking procedure (*benchmarking.result*) and the target data (*benchmarking.target*) one needs to once execute the seasonal adjustment with benchmarking using the multi-processing option (see the *Simple seasonal adjustment of multiple time series* scenario). Once the multi-processing is executed, select the *Output* item from the *SAProcessing* menu.

The *SAProcessing* menu

- Expand the "+" menu and choose an appropriate data format (here Excel has been chosen). It is possible to save the results in TXT, XLS, CSV, and CSV matrix formats. Note that the *available content of the output depends on the output type*.

Exporting data to an Excel file

- Chose the output items that refer to the results from the benchmarking procedure, move them to the window on the right and click **OK**.

Exporting the results of the benchmarking procedure



Figure 3: Text




Figure 4: Text




Figure 5: Text

Benchmarking in R

package rjd3bench orga doc - here - in package - example

References




Figure 6: Text

Trend-cycle estimation

Motivation

Estimation Methods

Tools

rjd3 highseq package

rjdfilters package

Nowcasting

Motivation

Underlying Theory: references ?

Tools

- plug in ?
- R package ?

Part II

Tools

Graphical User Interface

Overview

Installation Procedure

JDemetra+ is a stand-alone application packed in a zip package. To run JDemetra+ the Java RE 8 or higher is needed. Java RE can be downloaded from [Oracle website](#).

The official release of JDemetra+ is accessible at [a dedicated Github page](#). The site presents all available releases - both official releases (labelled in green as latest releases) and pre-releases (labelled in red) - packed in zip packages. From the *Latest release* section either choose the installer appropriate for your operating system (Windows, Linux, Mac OS, Solaris) or take the portable zip-file. The installation process is straightforward and intuitive. For example, when the zip-file is chosen and downloaded, then under Windows OS the application can be found in the “bin”-folder of the installation/unpacked zip. To open an application, double click on nbdemetra.exe or nbdemetra64.exe depending on the system version (nbdemetra.exe for the 32-bit system version and nbdemetra64.exe for the 64-bit system version).




Figure 7: Text

Launching JDemetra+

If the launching of JDemetra+ fails, you can try the following operations:

- Check if Java SE Runtime Environment (JRE) is properly installed by typing in the following command in a terminal: `java -version`
- Check the logs in your home directory:
 - %appdata%/.nbdemetra/dev/var/log/ for Windows;
 - ~/.nbdemetra/dev/var/log/ for Linux and Solaris;
 - ~/Library/Application Support/.nbdemetra/dev/var/log/ for Mac OS X.

In order to remove a previously installed JDemetra+ version, the user should delete an appropriate JDemetra+ folder.

Running JDemetra+

To open an application, navigate to the destination folder and double click on *nbdemetra.exe* or *nbdemetra64.exe* depending on the system version (*nbdemetra.exe* for the 32-bit system version and *nbdemetra64.exe* for the 64-bit system version).




Figure 8: Text

Running JDemetra+

Closing JDemetra+

To close the application, select *File* → *Exit* from the [File menu](#).




Figure 9: Text

Closing JDemetra+

The other way is to click on the close box in the upper right-hand corner of the JDemetra+ window. If there is any unsaved work, JDemetra+ will display a warning and provide the user with the opportunity to save it. The message box is shown below.




Figure 10: Text

The warning from leaving JDemetra+ without saving the workspace

R packages

Available algorithms

Table 24: Main Algorithms accessible via R packages

Domain	Algorithm	Package	Comments
Seasonal Adjustment	X13-Arima		
	Tramo-Seats		
	STL()	rjd3highfreq	
	Basic	rjd3stl	
Trend estimation	Structural Models	rjd3sts	
	Moving Averages, Local		
	Polynomial		
	idem	rjd3highfreq	
Benchmarking and TD	rjd3bench		

Organisation overview

RJDemetra and JD+ version 2.2.3

Note on RJDemetra

examples in A_sa : v2 and v3 rest of the chapter on version 3

(écriture standard des noms des packages..cf K) a suite (order)

dependencies

utility packages

general output organisation

Installation procedure

```
#install.packages("remotes")
remotes::install_github("palatej/rjd3toolkit")
remotes::install_github("palatej/rjd3modelling")
remotes::install_github("palatej/rjd3sa")
remotes::install_github("palatej/rjd3arima")
remotes::install_github("palatej/rjd3x13")
remotes::install_github("palatej/rjd3tramoseats")
remotes::install_github("palatej/rjdemetra3")
remotes::install_github("palatej/rjdfilters")
remotes::install_github("palatej/rjd3sts")
remotes::install_github("palatej/rjd3highfreq")
remotes::install_github("palatej/rjd3stl")
remotes::install_github("palatej/rjd3bench")
remotes::install_github("AQLT/ggdemetra3")
```

Below you will find a comprehensive list and main functions by categories
for detailed function, you can refer to each package's own R documentation

Utility packages

rjd3toolkit

Contains several utility functions used in other `rjd` packages and several functions to perform test.

Utility

Tests - Normality tests: Bowman-Shenton (`bowmanshenton()`), Doornik-Hansen (`doornikhansen()`), Jarque-Bera (`jarquebera()`)

- Runs tests (randomness of data): mean or the median (`testofruns()`) or up and down runs test (`testofupdownruns()`)
- autocorrelation functions (usual, inverse, partial)
- `aggregate()` to aggregate a time series to a higher frequency

Example

```
library(rjd3toolkit)
set.seed(100)
x = rnorm(1000); y = rlnorm(1000)
bowmanshenton(x) # normal distribution
bowmanshenton(y) # log-normal distribution
testofruns(x) # random data
testofruns(y) # random data
testofruns(1:1000) # non-random data
autocorrelations(x)
autocorrelations.inverse(x)
autocorrelations.partial(x)
```

rjd3modelling

Purpose : creating input variables (regressors) for to be used in Reg-Arima

- create user-defined calendar and trading-days regressors: `calendar.new()` (create a new calendar), `calendar.holiday()` (add a specific holiday, e.g. christmas), `calendar.easter()` (easter related day) and `calendar.fixedday()`
- outliers regressors (AO, LS, TC, SO, Ramp, intervention variables), calendar related regressors (stock, leap year, periodic dummies and contrasts, trigonometric variables) -> to be added quadratic ramps
- Range-mean regression test (to choose log transformation), Canova-Hansen (`td.ch()`) and trading-days f-test (`td.f()`)
- specification functions for `rjd3x13` and `rjd3tramoseats`

more explanations and examples needed here

Example of calendar specification

```
library(rjd3modelling)
fr_cal <- calendar.new()
calendar.holiday(fr_cal, "NEWYEAR")
calendar.holiday(fr_cal, "EASTERMONDAY")
calendar.holiday(fr_cal, "MAYDAY")
calendar.fixedday(fr_cal, month = 5, day = 8,
                  start = "1953-03-20")
```

```
# calendar.holiday(fr_cal, "WHITMONDAY") # Equivalent to:
calendar.easter(fr_cal, offset = 61)

calendar.fixedday(fr_cal, month = 7, day = 14)
# calendar.holiday(fr_cal, "ASSUMPTION")
calendar.easter(fr_cal, offset = 61)
calendar.holiday(fr_cal, "ALLSAINTSDAY")
calendar.holiday(fr_cal, "ARMISTICE")
calendar.holiday(fr_cal, "CHRISTMAS")
```

Use `holidays()` to get the days of the holidays and `htd()` to get the trading days regressors

```
holidays(fr_cal, "2020-12-24", 10, single = T)
s = ts(0, start = 2020, end = c(2020, 11), frequency = 12)
# Trading-days regressors (each day has a different effect, sunday as contrasts)
td_reg <- htd(fr_cal, s = s, groups = c(1, 2, 3, 4, 5, 6, 0))
# Working-days regressors (Monday = ... = Friday; Saturday = Sunday = contrasts)
wd_reg <- htd(fr_cal, s = s, groups = c(1, 1, 1, 1, 1, 0, 0))
# Monday = ... = Friday; Saturday; Sunday = contrasts
wd_reg <- htd(fr_cal, s = s, groups = c(1, 1, 1, 1, 1, 2, 0))
wd_reg
```

more explanations on contrasts and references links to calendar correction chapters

Example of outliers

```
s = ts(0, start = 2000, end = 2005, frequency = 12)
ao = ao.variable(s = s, date = "2001-03-01")
ls = ls.variable(s = s, date = "2001-01-01")
tc = tc.variable(s = s, date = "2001-01-01", rate = 0.7)
so = so.variable(s = s, date = "2003-05-01")
ramp = ramp.variable(s = s, range = c("2001-01-01", "2001-12-01"))
plot(ts.union(ao, ls, tc, so, ramp), plot.type = "single",
     col = c("red", "lightgreen", "orange", "blue", "black"))
```

rjd3sa package

Seasonality tests:

(for each code snippet example and link to `M_Test` chapter)

- Canova-Hansen (`seasonality.canovahansen()`)
- X-12 combined test (`seasonality.combined()`)
- F-test on seasonal dummies (`seasonality.f()`)
- Friedman Seasonality Test (`seasonality.friedman()`)
- Kruskal-Wallis Seasonality Test (`seasonality.kruskalwallis()`)
- Periodogram Seasonality Test (`seasonality.periodogram()`)
- QS Seasonality Test (`seasonality.qs()`)

EXP : Always correct the trend and remove the mean before seasonality tests:

```
library(rjd3sa)
y = diff(rjd3toolkit::ABS$X0.2.09.10.M, 1); y = y - mean(y)
seasonality.f(y, 12)
seasonality.friedman(y, 12)
seasonality.kruskalwallis(y, 12)
seasonality.combined(y, 12)
```

Seasonal adjustment packages

rjd3arima

`rjd3arima` is devoted to formatting the output of Arima related results

rjd3x13 and rjd3tramoseats

attention : splitting between package description here and practical usage in `A_sa` chapter, here general philosophy of the package

Common functions

In `RJDemetra` you have one function to set the specification (`regarima_spec_x13()`, `regarima_spec_tramo()`, `x13_spec()` and `tramoseats_spec()`) now one function for each part of the specification

Common functions (defined in `rjd3modelling`) to set the specification of the preprocessing:

`set_arima()`, `set_automodel()`, `set_basic()`, `set_easter()`, `set_estimate()`, `set_outlier()`, `set_tradingdays()`, `set_transform()`, `add_outlier()` and `remove_outlier()`, `add_ramp()` and `remove_ramp()`

- warning `add_usrdefvar()` not yet available

rjd3x13

Main functions:

- Specification: created with `spec_x11_default()`, `spec_x13_default()`, `spec_regarima_default()` and customized with `rjd3arima` functions + `set_x11()`
- Apply model with `x11()`, `x13()`, `fast.x13()`, `regarima()`, `fast.regarima()`
- Refresh policies: `regarima.refresh()` and `x13.refresh()`

rjd3tramoseats

Main functions:

- Specification: created with `spec_tramoseats_default()`, `spec_tramo_default()` and customized with `rjd3arima` functions + `set_seats()`
- Apply model with `tramoseats()`, `fast.tramoseats()`, `tramo()`, `fast.tramo()`
- Refresh policies: `tramo.refresh()` and `tramoseats.refresh()`

Example

```
# {r tramoseats, out.height="70%"}
# spec = spec_tramoseats_default("rsafull") |>
#   set_easter(type = "IncludeEasterMonday") |>
#   set_tradingdays(test = "Separate_T") |>
#   set_seats(algorithm = "KalmanSmoother")
# m = rjd3tramoseats::tramoseats(y, spec)
# # More informations:
# names(m)
# m$result
```

rjdemetra3

Functions to manipulate JDemetra+ workspaces:

- Still in construction: you can load an existing workspace but not create a new one (use `jws.load()` for example)
- Will contain all the functionalities of `rjdworkspace`

rjd3highfreq

Seasonal adjustment of high frequency data:

- fractional and multi airline decomposition
- Extension of X-11 decomposition with non integer periodicity

`rjd3stl` : STL, MSTL, ISTL, loess

rjd3stl

Add-in package

table ggdemetra rjworkspace

attention version 2 and version: adaptation under construction version 2, retrieve R tools info here

ggdemetra

ggdemetra3

Like ggdemetra but compatible with rjdemetra3: ggplot2 to add seasonal adjustment statistics to your plot. Also compatible with high-frequency methods (WIP):

```
library(ggdemetra3)
spec <- spec_x13_default("rsa3") |> set_tradingdays(option = "WorkingDays")
p_ipi_fr <- ggplot(data = ipi_c_eu_df, mapping = aes(x = date, y = FR)) +
  geom_line() +
  labs(title = "SA - IPI-FR",
        x = NULL, y = NULL)
p_sa <- p_ipi_fr +
  geom_sa(component = "y_f(12)", linetype = 2,
          spec = spec) +
  geom_sa(component = "sa", color = "red") +
  geom_sa(component = "sa_f", color = "red", linetype = 2)
p_sa
p_sa +
  geom_outlier(geom = "label_repel",
               coefficients = TRUE,
               ylim = c(NA, 65), force = 10,
               arrow = arrow(length = unit(0.03, "npc"),
                             type = "closed", ends = "last"),
               digits = 2)
```

rjdfilters

rjdfilters (1)

- easily create/combine/apply moving averages `moving_average()` (much more general than `stats::filter()`) and study their properties: plot coefficients (`plot_coef()`), gain (`plot_gain()`), phase-shift (`plot_phase()`) and different statics (`diagnostic_matrix()`)
- trend-cycle extraction with different methods to treat endpoints:
- `lp_filter()` local polynomial filters of Proietti and Luati (2008) (including Musgrave): Henderson, Uniform, biweight, Trapezoidal, Triweight, Tricube, “Gaussian”, Triangular, Parabolic (= Epanechnikov)
- `rkhs_filter()` Reproducing Kernel Hilbert Space (RKHS) of Dagum and Bianconcini (2008) with same kernels
- `fst_filter()` FST approach of Grun-Rehomme, Guggemos, and Ladiray (2018)
- `dfa_filter()` derivation of AST approach of Wildi and McElroy (2019)
- change the filter used in X-11 for TC extraction

Create moving average `moving_average()`

```
library(rjdfilters)
m1 = moving_average(rep(1,3), lags = 1); m1 # Forward MA
m2 = moving_average(rep(1,3), lags = -1); m2 # centered MA
m1 + m2
m1 - m2
m1 * m2
```

Can be used to create all the MA of X-11:

```
e1 <- moving_average(rep(1,12), lags = -6)
e1 <- e1/sum(e1)
e2 <- moving_average(rep(1/12, 12), lags = -5)
# used to have the 1rst estimate of the trend
tc_1 <- M2X12 <- (e1 + e2)/2
coef(M2X12) |> round(3)
```



```

si_1 <- 1 - tc_1
M3 <- moving_average(rep(1/3, 3), lags = -1)
M3X3 <- M3 * M3
# M3X3 moving average applied to each month
coef(M3X3) |> round(3)
M3X3_seasonal <- to_seasonal(M3X3, 12)
coef(M3X3_seasonal) |> round(3)
s_1 = M3X3_seasonal * si_1
s_1_norm = (1 - M2X12) * s_1
sa_1 <- 1 - s_1_norm
henderson_mm = moving_average(lp_filter(horizon = 6)$
                              filters.coef[, "q=6"],
                              lags = -6)

tc_2 <- henderson_mm * sa_1
si_2 <- 1 - tc_2
M5 <- moving_average(rep(1/5, 5), lags = -2)
M5X5_seasonal <- to_seasonal(M5 * M5, 12)
s_2 = M5X5_seasonal * si_2
s_2_norm = (1 - M2X12) * s_2
sa_2 <- 1 - s_2_norm
tc_f <- henderson_mm * sa_2

par(mai = c(0.3, 0.3, 0.2, 0))
layout(matrix(c(1,1,2,3), 2, 2, byrow = TRUE))

plot_coef(tc_f);plot_coef(sa_2, col = "orange", add = TRUE)
legend("topleft",
      legend = c("Final TC filter", "Final SA filter"),
      col= c("black", "orange"), lty = 1)
plot_gain(tc_f);plot_gain(sa_2, col = "orange", add = TRUE)
plot_phase(tc_f);plot_phase(sa_2, col = "orange", add = TRUE)

```

Apply a moving average

```

y <- retailsa$AllOtherGenMerchandiseStores
trend <- y * tc_1
sa <- y * sa_1
plot(window(ts.union(y, trend, sa), start = 2000),
      plot.type = "single",

```

```
col = c("black", "orange", "lightblue"))
```

rjd3sts

rjd3sts

Interface to structural time series and state space models

Several examples available here https://github.com/palatej/test_rjd3sts

rjd3bench

rjd3bench

Benchmarking and temporal disaggregation

Several examples here: https://github.com/palatej/test_rjd3bench

Interaction with workspaces and GUI

Cruncher

Overview

Installation procedure

Running the cruncher in R

Revision Policies and Production

Revision Policies

Plug-ins for JDemetra+

Main functions

table

Tool selection issues and heuristics

- ws structure
- stand alone

Part III

Methods

Spectral Analysis Principles and Tools

Spectral analysis concepts

A time series x_t with stationary covariance, mean μ and k^{th} autocovariance $E(x_t - \mu)(x_{t-k} - \mu) = \gamma(k)$ can be described as a weighted sum of periodic trigonometric functions: $\sin(\omega t)$ and $\cos(\omega t)$, where $\omega = \frac{2\pi}{T}$ denotes frequency. Spectral analysis investigates this frequency domain representation of x_t to determine how important cycles of different frequencies are in accounting for the behavior of x_t .

Assuming that the autocovariances $\gamma(k)$ are absolutely summable ($\sum_{k=-\infty}^{\infty} |\gamma(k)| < \infty$), the autocovariance generating function, which summarizes these autocovariances through a scalar valued function, is given by equation [1]⁴.

$$acgf(z) = \sum_{k=-\infty}^{\infty} z^k \gamma(k),$$

where z denotes complex scalar.

Once the equation [1] is divided by π and evaluated at some $z = e^{-i\omega} = \cos\omega - i\sin\omega$, where $i = \sqrt{-1}$ and ω is a real scalar, $-\infty < \omega < \infty$, the result of this transformation is called a population spectrum $f(\omega)$ for x_t , given in equation [2]⁵.

$$f(\omega) = \frac{1}{\pi} \sum_{k=-\infty}^{\infty} e^{-ik\omega} \gamma(k)$$

Therefore, the analysis of the population spectrum in the frequency domain is equivalent to the examination of the autocovariance function in the time domain analysis; however it provides an alternative way of inspecting the process. Because $f(\omega)d$ is interpreted as a contribution to the variance of components with frequencies in the range $(\omega, \omega + d\omega)$, a peak in the spectrum indicates an important contribution to the variance at frequencies near the value that corresponds to this peak.

As $e^{-i\omega} = \cos\omega - i\sin\omega$, the spectrum can be also expressed as in equation [3].

$$f(\omega) = \frac{1}{\pi} \sum_{k=-\infty}^{\infty} (\cos\omega k - i\sin\omega k) \gamma(k)$$

⁴HAMILTON, J.D. (1994).

⁵HAMILTON, J.D. (1994).

Since $\gamma(k) = \gamma(-k)$ (i.e. $\gamma(k)$ is an even function of k) and $\sin(-x) = -\sin x$, [3] can be presented as equation

$$f(\omega) = \frac{1}{\pi} [\gamma(0) + 2 \sum_{k=1}^{\infty} \gamma(k) \cos k\omega]$$

,

This implies that if autocovariances are absolutely summable the population spectrum exists and is a continuous, real-valued function of ω . Due to the properties of trigonometric functions ($\cos(-\omega k) = \cos(\omega k)$ and $\cos(\omega + 2\pi j)k = \cos(\omega k)$) the spectrum is a periodic, even function of ω , symmetric around $\omega = 0$. Therefore, the analysis of the spectrum can be reduced to the interval $(-\pi, \pi)$. The spectrum is non-negative for all $\omega \in (-\pi, \pi)$.

The shortest cycle that can be distinguished in a time series lasts two periods. The frequency which corresponds to this cycle is $\omega = \pi$ and is called the Nyquist frequency. The frequency of the longest cycles that can be observed in the time series with n observations is $\omega = \frac{2\pi}{n}$ and is called the fundamental (Fourier) frequency.

Note that if x_t is a white noise process with zero mean and variance σ^2 , then for all $|k| > 0$ $\gamma(k) = 0$ and the spectrum of x_t is constant ($f(\omega) = \frac{\sigma^2}{\pi}$) since each frequency in the spectrum contributes equally to the variance of the process⁶.

The aim of spectral analysis is to determine how important cycles of different frequencies are in accounting for the behaviour of a time series⁷. Since spectral analysis can be used to detect the presence of periodic components, it is a natural diagnostic tool for detecting trading day effects as well as seasonal effects⁸. Among the tools used for spectral analysis are the autoregressive spectrum and the periodogram.

The explanations given in the subsections of this node derive mainly from DE ANTONIO, D., and PALATE, J. (2015) and BROCKWELL, P.J., and DAVIS, R.A. (2006).

comment1: end old intro: ok

Theoretical spectral density of an ARIMA model

Spectral density estimation

Method 1: The periodogram

For any given frequency ω the sample periodogram is the sample analog of the sample spectrum. In general, the periodogram is used to identify the periodic components of unknown

⁶BROCKWELL, P.J., and DAVIS, R.A. (2002).

⁷HAMILTON, J.D. (1994).

⁸SOKUP, R.J., and FINDLEY, D. F. (1999).

frequency in the time series. X-13ARIMA-SEATS and TRAMO-SEATS use this tool for detecting seasonality in raw time series and seasonally adjusted series. Apart from this it is applied for checking randomness of the residuals from the ARIMA model.

To define the periodogram, first consider the vector of complex numbers⁹:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \cdot \\ \cdot \\ \cdot \\ x_n \end{bmatrix} \in \mathbb{C}^n$$

where \mathbb{C}^n is the set of all column vectors with complex-valued components.

The Fourier frequencies associated with the sample size n are defined as a set of values $\omega_j = \frac{2\pi j}{n}$, $j = -[\frac{n-1}{2}], \dots, [\frac{n}{2}]$, $-\pi < \omega_j \leq \pi$, $j \in F_n$, where $[n]$ denotes the largest integer less than or equal to n . The Fourier frequencies, which are called harmonics, are given by integer multiples of the fundamental frequency $\frac{2\pi}{n}$.

Now the n vectors $e_j = n^{-\frac{1}{2}}(e^{-i\omega_j}, e^{-i2\omega_j}, \dots, e^{-in\omega_j})'$ can be defined. Vectors e_1, \dots, e_n are orthonormal in the sense that:

$$\mathbf{e}_j^* \mathbf{e}_k = n^{-1} \sum_{r=1}^n e^{ir(\omega_j - \omega_k)} = \begin{cases} 1, & \text{if } j = k \\ 0, & \text{if } j \neq k \end{cases}$$

where \mathbf{e}_j^* denotes the row vector, which k^{th} component is the complex conjugate of the k^{th} component of \mathbf{e}_j .¹⁰ These vectors are a basis of F_n , so that any $\mathbf{x} \in \mathbb{C}^n$ can be expressed as a sum of n components:

$$\mathbf{x} = \sum_{j=-[\frac{n-1}{2}]}^{[\frac{n}{2}]} a_j \mathbf{e}_j$$

where the coefficients $a_j = \mathbf{e}_j^* \mathbf{x} = n^{-\frac{1}{2}} \sum_{t=1}^n x_t e^{-it\omega_j}$ are derived from [3] by multiplying the equation on the left by \mathbf{e}_j^* and using [1].

The sequence of $\{a_j, j \in F_n\}$ is referred as a discrete Fourier transform of $\mathbf{x} \in \mathbb{C}^n$ and the periodogram $I(\omega_j)$ of \mathbf{x} at Fourier frequency $\omega_j = \frac{2\pi j}{n}$ is defined as the square of the Fourier transform $\{a_j\}$ of \mathbf{x} :

⁹BROCKWELL, P.J., and DAVIS, R.A. (2002).

¹⁰For details see BROCKWELL, P.J., and DAVIS, R.A. (2006).

$$I(\omega_j) = |a_j|^2 = n^{-1} \left| \sum_{t=1}^n x_t e^{-it\omega_j} \right|^2$$

From [2] and [3] it can be shown that in fact the periodogram decomposes the total sum of squares $\sum_{t=1}^n |x_t|^2$ into a sums of components associated with the Fourier frequencies

$$\omega_j$$

:

$$\sum_{t=1}^n |x_t|^2 = \sum_{j=-\lfloor \frac{n-1}{2} \rfloor}^{\lfloor \frac{n}{2} \rfloor} |a_j|^2 = \sum_{j=-\lfloor \frac{n-1}{2} \rfloor}^{\lfloor \frac{n}{2} \rfloor} I(\omega_j)$$

If $\mathbf{x} \in R^n$, ω_j and $-\omega_j$ are both in $[-\pi, \pi]$ and a_j is presented in its polar form (i.e. $a_j = r_j \exp(i\theta_j)$), where r_j is the modulus of a_j , then [3] can be rewritten in the form:

$$\mathbf{x} = a_0 \mathbf{e}_0 + \sum_{j=1}^{\lfloor \frac{n-1}{2} \rfloor} 2^{1/2} r_j (\mathbf{c}_j \cos \theta_j - \mathbf{s}_j \sin \theta_j) + a_{n/2} \mathbf{e}_{n/2}$$

The orthonormal basis for R^n is $\{\mathbf{e}_0, \mathbf{c}_1, \mathbf{s}_1, \dots, \mathbf{c}_{\lfloor \frac{n-1}{2} \rfloor}, \mathbf{s}_{\lfloor \frac{n-1}{2} \rfloor}, \mathbf{e}_{\frac{n}{2}} (\text{excluded if } n \text{ is odd})\}$, where:

\mathbf{e}_0 is a vector composed of n elements equal to $n^{-1/2}$, which implies that $\mathbf{a}_0 \mathbf{e}_0 = (n^{-1} \sum_{t=1}^n x_t, \dots, n^{-1} \sum_{t=1}^n x_t)'$;

$$\mathbf{c}_j = \left(\frac{n}{2}\right)^{-1/2} (\cos \omega_j, \cos 2\omega_j, \dots, \cos n\omega_j)', \text{ for } 1 \leq j \leq \lfloor \frac{(n-1)}{2} \rfloor$$

;

$$\mathbf{s}_j = \left(\frac{n}{2}\right)^{-1/2} (\sin \omega_j, \sin 2\omega_j, \dots, \sin n\omega_j)', \text{ for } 1 \leq j \leq \lfloor \frac{(n-1)}{2} \rfloor$$

;

$$\mathbf{e}_{n/2} = (-(n^{-\frac{1}{2}}), n^{-\frac{1}{2}}, \dots, -(n^{-\frac{1}{2}}), n^{-\frac{1}{2}})'$$

.

Equation [5] can be seen as an OLS regression of x_t on a constant and the trigonometric terms. As the vector of explanatory variables includes n elements, the number of explanatory variables in [5] is equal to the number of observations. HAMILTON, J.D. (1994) shows that the explanatory variables are linearly independent, which implies that an OLS regression yields

[7]

[8]

[9]

[10]

$$[11]$$

With [5] the total sum of squares $\sum_{t=1}^n |x_t|^2$ can be decomposed into $2 \times \lfloor \frac{n-1}{2} \rfloor$ components corresponding to \mathbf{c}_j and \mathbf{s}_j , which are grouped to produce the “frequency ω_j ” component for $1 \geq j \geq \lfloor \frac{n-1}{2} \rfloor$. As it is shown in the table below, the value of the periodogram at the frequency ω_j is the contribution of the j th harmonic to the total sum of squares $\sum_{t=1}^n |x_t|^2$.

Decomposition of sum of squares into components corresponding to the harmonics

	Frequency	Degrees of freedom	Sum of squares decomposition
$\omega_0(\text{mean})$	$ 1 $	$a_0^2 = n^{-1}(\sum_{t=1}^n x_t)^2 = I(0)$	$ \omega_1 2 2r_1^2 = 2\ a_1\ ^2 = 2I(\omega_1) : : :$
ω_k	$ 2 $	$2r_k^2 = 2\ a_k\ ^2 = 2I(\omega_k)$	$:$
		$:$	
$\omega_{n/2}$	$= \pi$ (excluded if n is odd)	$ 1 $	$a_{n/2}^2 = I(\pi) \textbf{Total } \mathbf{n} \sum_{t=1}^{\mathbf{n}} \mathbf{x}_t^2 $

Source: DE ANTONIO, D., and PALATE, J. (2015).

Obviously, if series were random then each component $I(\omega_j)$ would have the same expectation. On the contrary, when the series contains a systematic sine component having a frequency j and amplitude A then the sum of squares $I(\omega_j)$ increases with A . In practice, it is unlikely that the frequency j of an unknown systematic sine component would exactly match any of the frequencies, for which periodogram have been calculated. Therefore, the periodogram would show an increase in intensities in the immediate vicinity of j .¹¹

Note that in JDemetra+ the periodogram object corresponds exactly to the contribution to the sum of squares of the standardised data, since the series are divided by their standard deviation for computational reasons.

Using the decomposition presented in table above the periodogram can be expressed as:

$$I(\omega_j) = r_j^2 = \frac{1}{2}(\alpha_j^2 + \beta_j^2) = \frac{1}{n} \left(\sum_{t=1}^n x_t \cos\left(t \frac{2\pi j}{n}\right) \right)^2 + \frac{1}{n} \left(\sum_{t=1}^n x_t \sin\left(t \frac{2\pi j}{n}\right) \right)^2$$

[12]

where $j = 0, \dots, \lfloor \frac{n}{2} \rfloor$.

Since $\mathbf{x} - \bar{\mathbf{x}}$ are generated by an orthonormal basis, and $\bar{\mathbf{x}} = a_0 \mathbf{e}_0$ [5] can be rearranged to show that the sum of squares is equal to the sum of the squared coefficients:

$$\mathbf{x} - a_0 \mathbf{e}_0 = \sum_{j=1}^{\lfloor (n-1)/2 \rfloor} (\alpha_j \mathbf{c}_j + \beta_j \mathbf{s}_j) + a_{n/2} \mathbf{e}_{n/2}$$

[13]

Thus the sample variance of

$$x_t$$

can be expressed as:

$$n^{-1} \sum_{t=1}^n (x_t - \bar{x})^2 = n^{-1} \left(\sum_{k=1}^{\lfloor (n-1)/2 \rfloor} 2r_j^2 + a_{n/2}^2 \right)$$

[14]

where $a_{n/2}^2$ is excluded if n is odd.

The term $2r_j^2$ in [14] is then the contribution of the j^{th} harmonic to the variance and [14] shows then how the total variance is partitioned.

The periodogram ordinate $I(\omega_j)$ and the autocovariance coefficient $\gamma(k)$ are both quadratic forms of x_t . It can be shown that the periodogram and autocovariance function are related

¹¹BOX, G.E.P., JENKINS, G.M., and REINSEL, G.C. (2007).

and the periodogram can be written in terms of the sample autocovariance function for any non-zero Fourier frequency ω_j :¹²

$$I(\omega_j) = \sum_{|k| < n} \hat{\gamma}(k) e^{-ik\omega_j} = \hat{\gamma}(0) + 2 \sum_{k=1}^{n-1} \hat{\gamma}(k) \cos(k\omega_j)$$

and for the zero frequency $I(0) = n|\bar{x}|^2$.

Once comparing [15] with an expression for the spectral density of a stationary process:

$$f(\omega) = \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma(k) e^{-ik\omega} = \frac{1}{2\pi} \left(\gamma(0) + 2 \sum_{k=1}^{\infty} \gamma(k) \cos(k\omega) \right)$$

It can be noticed that the periodogram is a sample analog of the population spectrum. In fact, it can be shown that the periodogram is asymptotically unbiased but inconsistent estimator of the population spectrum $f(\omega)$.^[75] Therefore, the periodogram is a wildly fluctuating, with high variance, estimate of the spectrum. However, the consistent estimator can be achieved by applying the different linear smoothing filters to the periodogram, called lag-window estimators. The lag-window estimators implemented in JDemetra+ includes square, Welch, Tukey, Barlett, Hanning and Parzen. They are described in DE ANTONIO, D., and PALATE, J. (2015). Alternatively, the model-based consistent estimation procedure, resulting in autoregressive spectrum estimator, can be applied.

comment2: end part theory>spectral analysis>periodogram

Method 2: Autoregressive spectrum estimation

BROCKWELL, P.J., and DAVIS, R.A. (2006) point out that for any real-valued stationary process (x_t) with continuous spectral density $f(\omega)$ it is possible to find both $AR(p)$ and $MA(q)$ processes which spectral densities are arbitrarily close to $f(\omega)$. For this reason, in some sense, (x_t) can be approximated by either $AR(p)$ or $MA(q)$ process. This fact is a basis of one of the methods of achieving a consistent estimator of the spectrum, which is called an autoregressive spectrum estimation. It is based on the approximation of the stochastic process (x_t) by an autoregressive process of sufficiently high order p :

$$x_t = \mu + (\phi_1 B + \dots + \phi_p B^p)x_t + \varepsilon_t$$

where ε_t is a white-noise variable with mean zero and a constant variance.

¹²The proof is given in BROCKWELL, P.J., and DAVIS, R.A. (2006).

The autoregressive spectrum estimator for the series x_t is defined as: ¹³

$$\hat{s}(\omega) = 10 \times \log_{10} \frac{\sigma_x^2}{2\pi |1 - \sum_{k=1}^p \hat{\phi}_k e^{-ik\omega}|^2}$$

where:

ω – frequency, $0 \leq \omega \leq \pi$;

σ_x^2 – the innovation variance of the sample residuals;

$\hat{\phi}_k$ – AR(k) coefficient estimates of the linear regression of $x_t - \bar{x}$ on $x_{t-k} - \bar{x}$, $1 \leq k \leq p$.

The autoregressive spectrum estimator is used in the visual spectral analysis tool for detecting significant peaks in the spectrum. The criterion of *visual significance*, implemented in JDemetra+, is based on the range $\hat{s}^{\max} - \hat{s}^{\min}$ of the $\hat{s}(\omega)$ values, where $\hat{s}^{\max} = \max_k \hat{s}(\omega_k)$; $\hat{s}^{\min} = \min_k \hat{s}(\omega_k)$; and $\hat{s}(\omega_k)$ is k^{th} value of autoregressive spectrum estimator.

The particular value is considered to be visually significant if, at a trading day or at a seasonal frequency ω_k (other than the seasonal frequency $\omega_{60} = \pi$), $\hat{s}(\omega_k)$ is above the median of the plotted values of $\hat{s}(\omega_k)$ and is larger than both neighbouring values $\hat{s}(\omega_{k-1})$ and $\hat{s}(\omega_{k+1})$ by at least $\frac{6}{52}$ times the range $\hat{s}^{\max} - \hat{s}^{\min}$.

Following the suggestion of SOUKUP, R.J., and FINDLEY, D.F. (1999), JDemetra+ uses an autoregressive model spectral estimator of model order 30. This order yields high resolution of strong components, meaning peaks that are sharply defined in the plot of $\hat{s}(\omega)$ with 61 frequencies. The minimum number of observations needed to compute the spectrum is set to $n = 80$ for monthly data and to $n = 60$ for quarterly series while the maximum number of observations considered for the estimation is 121. Consequently, with these settings it is possible to identify up to 30 peaks in the plot of 61 frequencies. By choosing $\omega_k = \frac{k}{60}$ for $k = 0, 1, \dots, 60$ the density estimates are calculated at exact seasonal frequencies (1, 2, 3, 4, 5 and 6 cycles per year).

The model order can also be selected based on the AIC criterion (in practice it is much lower than 30). A lower order produces the smoother spectrum, but the contrast between the spectral amplitudes at the trading day frequencies and neighbouring frequencies is weaker, and therefore not as suitable for automatic detection.

SOUKUP, R.J., and FINDLEY, D.F. (1999) also explain that the periodogram can be used in the *visual significance* test as it has as good as those of the AR(30) spectrum abilities to detect trading day effect, but also has a greater false alarm rate¹⁴.

comment2: end part theory>spectral analysis>auto-regressive spectrum

¹³Definition from ‘X-12-ARIMA Reference Manual’ (2011).

¹⁴The false alarm rate is defined as the fraction of the 50 replicates for which a visually significant spectral peak occurred at one of the trading day frequencies being considered in the designated output spectra (SOUKUP, R.J., and FINDLEY, D.F. (1999)).

Method 3: Tukey spectrum

Identification of spectral peaks

comment3: start part theory>spectral analysis>identification of spectral peaks

Identification of seasonal peaks in a Tukey periodogram and in an autoregressive spectrum

In order to decide whether a series has a seasonal component that is predictable (stable) enough, these tests use visual criteria and formal tests for the periodogram. The periodogram is calculated using complete years, so that the set of Fourier frequencies contains exactly all seasonal frequencies¹⁵.

The tests rely on two basic principles:

- The peaks associated with seasonal frequencies should be larger than the median spectrum for all frequencies and;
- The peaks should exceed the spectrum of the two adjacent values by more than a critical value.

JDemetra+ performs this test on the original series. If these two requirements are met, the test results are displayed in green. The statistical significance of each of the seasonal peaks (i.e. frequencies $\frac{\pi}{6}$, $\frac{\pi}{3}$, $\frac{\pi}{2}$, $\frac{2\pi}{3}$ and $\frac{5\pi}{6}$ corresponding to 1, 2, 3, 4 and 5 cycles per year) is also displayed. The seasonal and trading days frequencies depends on the frequency of time series. They are shown in the table below. The symbol d denotes a default frequency and is described below the table.

The seasonal and trading day frequencies by time series frequency

{: .table .table-style}	Number of months per full period	Seasonal frequency	Trading day frequency (radians)
2.714	6	$\frac{\pi}{6}, \frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}, \frac{5\pi}{6}, \pi$	d
4	3	π	d
1.292, 1.850, 2.128	3	π	d
2	2	π	d

The calendar (trading day or working day) effects, related to the variation in the number of different days of the week per period, can induce periodic patterns in the data that can be similar to those resulting from pure seasonal effects. From the theoretical point of view, trading day variability is mainly due to the fact that the average number of days in the months or quarters is not equal to a multiple of 7 (the average number of days of a month in the year of 365.25 days is equal to $\frac{365.25}{12} = 30.4375days$). This effect occurs $\frac{365.25}{12} \times \frac{1}{7} = 4.3482timespermonth$: one time for each one of the four complete weeks of each month, and a residual of 0.3482 cycles per month, i.e. $0.3482 \times 2\pi = 2.1878radians$. This turns out to be a fundamental frequency for the effects associated with monthly data. In JDemetra+

¹⁵For definition of the periodogram and Fourier frequencies see section [Spectral Analysis](#)

the fundamental frequency corresponding to 0.3482 cycles per month is used in place of the closest frequency $\frac{k}{60}$. Thus, the quantity $\frac{\pi \times 42}{60}$ is replaced by $\omega_{42} = 0.3482 \times 2\pi = 2.1878$. The frequencies neighbouring ω_{42} , i.e. ω_{41} and ω_{43} are set to, respectively, $2.1865 - \frac{1}{60}$ and $2.1865 + \frac{1}{60}$.

The default frequencies (d) for calendar effect are: 2.188 (monthly series) and 0.280 (quarterly series). They are computed as:

$$\omega_{ce} = \frac{2\pi}{7} \left(n - 7 \times \left\lfloor \frac{n}{7} \right\rfloor \right)$$

, [1]

where:

$n = \frac{365.25}{s}$, $s = 4$ for quarterly series and $s = 12$ for monthly series.

Other frequencies that correspond to trading day frequencies are: 2.714 (monthly series) and 1.292, 1.850, 2.128 (quarterly series).

In particular, the calendar frequency in monthly data (marked in red on the figure below) is very close to the seasonal frequency corresponding to 4 cycles per year $\omega_{40} = \frac{2}{3}\pi = 2.0944$.




Figure 11: Text

Periodogram with seasonal (grey) and calendar (red) frequencies highlighted

This implies that it may be hard to disentangle both effects using the frequency domain techniques.

comment3: end part theory>spectral analysis>identification of spectral peaks

in Tukey spectrum

comes from Identification of seasonal peaks in a Tukey spectrum

Tukey Spectrum definition

The Tukey spectrum belongs to the class of lag-window estimators. A lag window estimator of the spectral density $f(\omega) = \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma(k) e^{ik\omega}$ is defined as follows:

$$\hat{f}_L(\omega) = \frac{1}{2\pi} \sum_{|h| \leq r} w(h/r) \hat{\gamma}(h) e^{ih\omega}$$

where $\hat{\gamma}(\cdot)$ is the sample autocovariance function, $w(\cdot)$ is the lag window, and r is the truncation lag. $|w(x)|$ is always less than or equal to one, $w(0) = 1$ and $w(x) = 0$ for $|x| > 1$. The simple idea behind this formula is to down-weight the autocovariance function for high lags where $\hat{\gamma}(h)$ is more unreliable. This estimator requires choosing r as a function of the sample size such that $r/n \rightarrow 0$ and $r \rightarrow \infty$ when $n \rightarrow \infty$. These conditions guarantee that the estimator converges to the true density.

JDemetra+ implements the so-called Blackman-Tukey (or Tukey-Hanning) estimator, which is given by $w(h/r) = 0.5(1 + \cos(\pi h/r))$ if $|h/r| \leq 1$ and 0 otherwise.

The choice of large truncation lags r decreases the bias, of course, but it also increases the variance of the spectral estimate and decreases the bandwidth.

JDemetra+ allows the user to modify all the parameters of this estimator, including the window function.

Graphical Test

The current JDemetra+ implementation of the seasonality test is based on a $F(d_1, d_2)$ approximation that has been originally proposed by Maravall (2012) for TRAMO-SEATS. This test is has been designed for a Blackman-Tukey window based on a particular choices of the truncation lag r and sample size. Following this approach, we determine visually significant peaks for a frequency ω_j when

$$\frac{2f_x(\omega_j)}{[f_x(\omega_{j+1}) + f_x(\omega_{j-1})]} \geq CV(\omega_j)$$

where $CV(\omega_j)$ is the critical value of a $F(d_1, d_2)$ distribution, where the degrees of freedom are determined using simulations. For $\omega_j = \pi$, we have a significant peak when $\frac{f_x(\omega_{[n/2]})}{[f_x(\omega_{[(n-1)/2])}] \geq CV(\omega_j)$

Two significant levels for this test are considered: $\alpha = 0.05$ (code “t”) and $\alpha = 0.01$ (code “T”).

As opposed to the [AR spectrum](#), which is computed on the basis of the last 120 data points, we will use here all available observations. Those critical values have been calculated given the recommended truncation lag $r = 79$ for a sample size within the interval $\in [80, 119]$ and $r = 112$ for $n \in [120, 300]$. The F approximation is less accurate for sample sizes larger than 300. For quarterly data, $r = 44$, but there are no recommendations regarding the required sample size.

Use

The test can be applied directly to any series by selecting the option *Statistical Methods » Seasonal Adjustment » Tools » Seasonality Tests*. This is an example of how results are displayed for the case of a monthly series:

4. Identification of seasonal peaks in a Tukey periodogram and in an auto-regressive spectrum

Seasonality present

T or t for Tukey periodogram, A or a for auto-regressive spectrum; 'T' or 'A' for very significant peaks, 't' or 'a' for significant peaks, '_' otherwise

AT.AT.AT.AT.A-

Figure 12: tktest

JDemetra+ considers critical values for $\alpha = 1\%$ (code “T”) and $\alpha = 5\%$ (code “t”) at each one of the seasonal frequencies represented in the table below, e.g. frequencies $\frac{\pi}{6}$, $\frac{\pi}{3}$, $\frac{\pi}{2}$, $\frac{2\pi}{3}$ and $\frac{5\pi}{6}$ corresponding to 1, 2, 3, 4, 5 and 6 cycles per year in this example, since we are dealing with monthly data. The codes “a” and “A” correspond to the so-called [AR spectrum](#), so ignore them for the moment.

The seasonal and trading day frequencies by time series frequency

Number of months per full period	Seasonal frequency	Trading day frequency (radians)
12	$\frac{\pi}{6}, \frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}, \frac{5\pi}{6}, \pi$	$d, 2.714$
6	$\frac{\pi}{3}, \frac{2\pi}{3}, \pi$	d
4	$\frac{\pi}{2}, \pi$	$d, 1.292, 1.850, 2.128$
3	π	d
2	π	d

Currently, only seasonal frequencies are tested, but the program allows you to manually plot the Tukey spectrum and focus your attention on both seasonal and trading day frequencies.

References

- Tukey, J. (1949). The sampling theory of power spectrum estimates., Proceedings Symposium on Applications of Autocorrelation Analysis to Physical Problems, NAVEXOS-P-735, Office of Naval Research, Washington, 47-69

in AR Spectrum definition

comes from: “Identification of seasonal peaks in autoregressive spectrum”

The estimator of the spectral density at frequency $\lambda \in [0, \pi]$ will be given by the assumption that the series will follow an AR(p) process with large p . The spectral density of such model, with an innovation variance $var(x_t) = \sigma_x^2$, is expressed as follows:

$$10 \times \log_{10} f_x(\lambda) = 10 \times \log_{10} \frac{\sigma_x^2}{2\pi |\phi(e^{i\lambda})|^2} = 10 \times \log_{10} \frac{\sigma_x^2}{2\pi \left| 1 - \sum_{k=1}^p \phi_k e^{ik\lambda} \right|^2}$$

where ϕ_k denotes the AR(k) coefficient, and $e^{-ik\lambda} = \cos(-ik\lambda) + i \sin(-ik\lambda)$.

Soukup and Findely (1999) suggest the use of $p=30$, which in practice much larger than the order that would result from the AIC criterion. The minimum number of observations needed to compute the spectrum is set to $n=80$ for monthly data (or $n=60$) for quarterly series. In turn, the maximum number of observations considered for the estimation is $n=121$. This choice offers enough resolution, being able to identify a maximum of 30 peaks in a plot of 61 frequencies: by choosing $\lambda_j = \pi j/60$, for $j = 0, 1, \dots, 60$, we are able to calculate our density estimates at exact seasonal frequencies (1, 2, 3, 4, 5 and 6 cycles per year). Note that x cycles per year can be converted into cycles per month by simply dividing by twelve, $x/12$, and to radians by applying the transformation $2\pi(x/12)$.

The traditional trading day frequency corresponding to 0.348 cycles per month is used in place of the closest frequency $\pi j/60$. Thus, we replace $\pi 42/60$ by $\lambda_{42} = 0.348 \times 2\pi = 2.1865$. The frequencies neighbouring λ_{42} are set to $\lambda_{41} = 2.1865 - 1/60$ and $\lambda_{43} = 2.1865 + 1/60$. The periodogram below illustrates the proximity of this trading day frequency λ_{42} (red shade) and the frequency corresponding to 4 cycles per year $\lambda_{40} = 2.0944$. This proximity is precisely what poses the identification problems: the AR spectrum boils down to a smoothed version of the periodogram and the contribution of the trading day frequency may be obscured by the leakage resulting from the potential seasonal peak at λ_{40} , and vice-versa.

Periodogram with seasonal (grey) and calendar (red) frequencies highlighted

JDemetra+ allows the user to modify the number of lags of this estimator and to change the number of observations used to determine the AR parameters. These two options can improve the resolution of this estimator.




Figure 13: Text

Graphical Test

The statistical significance of the peaks associated to a given frequency can be informally tested using a visual criterion, which has proved to perform well in simulation experiments. Visually significant peaks for a frequency λ_j satisfy both conditions:

- $\frac{f_x(\lambda_j) - \max\{f_x(\lambda_{j+1}), f_x(\lambda_{j-1})\}}{[\max_k f_x(\lambda_k) - \min_i f_x(\lambda_i)]} \geq CV(\lambda_j)$, where $CV(\lambda_j)$ can be set equal to $6/52$ for all j
- $f_x(\lambda_j) > \text{median}_j \{f_x(\lambda_j)\}$, which guarantees $f_x(\lambda_j)$ it is not a local peak.

The first condition implies that if we divide the range $\max_k f_x(\lambda_k) - \min_i f_x(\lambda_i)$ in 52 parts (traditionally represented by stars) the height of each pick should be at least 6 stars.

Use

The test can be applied directly to any series by selecting the option *Statistical Methods » Seasonal Adjustment » Tools » Seasonality Tests*. This is an example of how results are displayed for the case of a monthly series:

4. Identification of seasonal peaks in a Tukey periodogram and in an auto-regressive spectrum

Seasonality present

T or t for Tukey periodogram, A or a for auto-regressive spectrum; 'T' or 'A' for very significant peaks, 't' or 'a' for significant peaks, '_' otherwise

AT.AT.AT.AT.A-

Figure 14: artest

JDemetra+ considers critical values for $\alpha = 1\%$ (code "A") and $\alpha = 5\%$ (code "a") at each one of the seasonal frequencies represented in the table below, e.g. frequencies $\frac{\pi}{6}$, $\frac{\pi}{3}$, $\frac{\pi}{2}$, $\frac{2\pi}{3}$ and $\frac{5\pi}{6}$

corresponding to 1, 2, 3, 4, 5 and 6 cycles per year in this example, since we are dealing with monthly data. The codes “t” and “T” correpond to the so-called [Tukey spectrum](#), so ignore them for the moment.

The seasonal and trading day frequencies by time series frequency

{: .table .table-style} Number of months per full period Seasonal frequency Trading day frequency (radians)		
2.714	6 $\frac{\pi}{3}, \frac{2\pi}{3}, \pi$ d	4 $\frac{\pi}{2}, \pi$ d , 1.292, 1.850, 2.128
		3 π d
		2 π d
		12 $\frac{\pi}{6}, \frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}, \frac{5\pi}{6}, \pi$ d ,

Currently, only seasonal frequencies are tested, but the program allows you to manually plot the AR spectrum and focus your attention on both seasonal and trading day frequencies. Agustin Maravall has conducted a simulation experiment to calculate $CV(\lambda_{42})$ (trading day frequency) and proposes to set for all j equal to the critical value associated to the trading frequency, but this is currently not part of the current automatic testing procedure of JDemetra+.

References

- Soukup, R.J., and D.F. Findley (1999) On the Spectrum Diagnosis used by X12-ARIMA to Indicate the Presence of Trading Day Effects After Modeling or Adjustment. In Proceedengs of the American Statistical Association. Business and Economic Statistics Section, 144-149, Alexandria, VA.

in a Periodogram

comes from: Identification of seasonal peaks in periodogram

The periodogram $I(\omega_j)$ of $\mathbf{X} \in \mathbb{C}^n$ is defined as the squared of the Fourier transform

$$I(\omega_j) = a_j^2 = n^{-1} \left| \sum_{t=1}^n \mathbf{X}_t e^{-it\omega_j} \right|^2,$$

where the Fourier frequencies ω_j are given by multiples of the fundamental frequency $\frac{2\pi}{n}$:

$$\omega_j = \frac{2\pi j}{n}, -\pi < \omega_j \leq \pi$$

An orthonormal basis in \mathbb{R}^n :

$$\left\{ e_0, \quad c_1, s_1, \quad \dots \quad, \quad c_{[(n-1)/2]}, s_{[(n-1)/2]} \quad, \quad e_{n/2} \right\},$$

where $e_{n/2}$ is excluded if n is odd,
can be used to project the data and obtain the spectral decomposition

Thus, the periodogram is given by the projection coefficients and represents the contribution of the j th harmonic to the total sum of squares, as illustrated by Brockwell and Davis (1991):

Source	Degrees of freedom
Frequency ω_0	1
Frequency ω_1	2
\vdots	\vdots
Frequency ω_k	2
\vdots	\vdots
Frequency $\omega_{n/2} = \pi$ (excluded if n is odd)	1
=====	=====
Total	n

In JDemetra+, the periodogram of $\mathbf{X} \in \mathbb{R}^n$ is computed for the standardized time series.

Defining a F-test

Brockwell and Davis (1991, section 10.2) exploit the fact that the periodogram can be expressed as the projection on the orthonormal basis defined above to derive a test. Thus, under the null hypothesis:

- $2I(\omega_k) = \|P_{\bar{s}p_{\{c_k, s_k\}}} \mathbf{X}\|^2 \sim \sigma^2 \chi^2(2)$, for Fourier frequencies $0 < \omega_k = 2\pi k/n < \pi$
- $I(\pi) = \|P_{\bar{s}p_{\{e_{n/2}\}}} \mathbf{X}\|^2 \sim \sigma^2 \chi^2(1)$, for π

Because $I(\omega_k)$ is independent from the projection error sum of squares, we can define our F-test statistic as follows:

- $\frac{2I(\omega_k)}{\|\mathbf{X} - P_{\bar{s}p_{\{e_0, c_k, s_k\}}}\mathbf{X}\|^2} \frac{n-3}{2} \sim F(2, n-3)$, for Fourier frequencies $0 < \omega_k = 2\pi k/n < \pi$
- $\frac{I(\pi)}{\|\mathbf{X} - P_{\bar{s}p_{\{e_0, e_{n/2}\}}}\mathbf{X}\|^2} \frac{n-2}{1} \sim F(1, n-2)$, for π

where - $\|\mathbf{X} - P_{\bar{s}p_{\{e_0, c_k, s_k\}}} \mathbf{X}\|^2 = \sum_{i=1}^n \mathbf{X}_i^2 - I(0) - 2I(\omega_k) \sim \sigma^2 \chi^2(n-3)$ for Fourier frequencies $0 < \omega_k = 2\pi k/n < \pi$ - $\|\mathbf{X} - P_{\bar{s}p_{\{e_0, e_{n/2}\}}} \mathbf{X}\|^2 = \sum_{i=1}^n \mathbf{X}_i^2 - I(0) - I(\pi) \sim \sigma^2 \chi^2(n-2)$ for π

Thus, we reject the null if our F-test statistic computed at a given seasonal frequency (different from π) is larger than $F_{1-\alpha}(2, n-3)$. If we consider π , our test statistic follows a $F_{1-\alpha}(1, n-2)$ distribution.

Seasonality test

The implementation of JDemetra+ considers simultaneously the whole set of seasonal frequencies (1, 2, 3, 4, 5 and 6 cycles per year). Thus, the resulting test-statistic is:

$$\frac{2I(\pi/6) + 2I(\pi/3) + 2I(2\pi/3) + 2I(5\pi/6) + \delta I(\pi)}{\left\| \mathbf{X} - P_{\bar{s}p_{\{e_0, c_1, s_1, c_2, s_2, c_3, s_3, c_4, s_4, c_5, s_5, \delta e_{n/2}\}}} \mathbf{X} \right\|^2} \frac{n-12}{11} \sim F(11-\delta, n-12+\delta)$$

where $\delta = 1$ if n is even and 0 otherwise.

In small samples, the test performs better when the periodogram is evaluated as the exact seasonal frequencies. JDemetra+ modifies the sample size to ensure the seasonal frequencies belong to the set of Fourier frequencies. This strategy provides a very simple and effective way to eliminate the leakage problem.

Example of how results are displayed:

5. Periodogram

Test on the sum of the values of a periodogram at seasonal frequencies

Seasonality present

Distribution: F with 11 degrees of freedom in the nominator and 180 degrees of freedom in the denominator

Value: 45.1387

PValue: 0.0000

Figure 15: periodtest

References

Brockwell, P.J., and R.A. Davis (1991). Times Series: Theory and Methods. Springer Series in Statistics.

Spectral graphs

probably move this part to GUI (Tools), just leave a link

comment3: start part case studies > spectral graphs

This scenario is designed for advanced users interested in an in-depth analysis of time series in the frequency domain using three spectral graphs. Those graphs can also be used as a complementary analysis for a better understanding of the results obtained with some of the tests described above.

Economic time series are usually presented in a time domain (X-axis). However, for analytical purposes it is convenient to convert the series to a frequency domain due to the fact that any stationary time series can be expressed as a combination of cosine (or sine) functions. These functions are characterized with different periods (amount of time to complete a full cycle) and amplitudes (maximum/minimum value during the cycle).

The tool used for the analysis of a time series in a frequency domain is called a spectrum. The peaks in the spectrum indicate the presence of cyclical movements with periodicity between two months and one year. A seasonal series should have peaks at the seasonal frequencies. Calendar adjusted data are not expected to have peak at with a calendar frequency.

The periodicity of the phenomenon at frequency f is $\frac{2\pi}{f}$. It means that for a monthly time series the seasonal frequencies $\frac{\pi}{6}$, $\frac{\pi}{3}$, $\frac{\pi}{2}$, $\frac{2\pi}{3}$, $\frac{5\pi}{6}$ and π correspond to 1, 2, 3, 4, 5 and 6 cycles per year. For example, the frequency $\frac{\pi}{3}$ corresponds to a periodicity of 6 months (2 cycles per year are completed). For the quarterly series there are two seasonal frequencies: $\frac{\pi}{2}$ (one cycle per year) and π (two cycles per year). A peak at the zero frequency always corresponds to the trend component of the series. Seasonal frequencies are marked as grey vertical lines, while violet vertical lines represent the trading-days frequencies. The trading day frequency is 0.348 and derives from the fact that a daily component which repeats every seven days goes through 4.348 cycles in a month of average length 30.4375 days. It is therefore seen to advance 0.348 cycles per month when the data are obtained at twelve equally spaced times in 365.25 days (the average length of a year).

The interpretation of the spectral graph is rather straightforward. When the values of a spectral graph for low frequencies (i.e. one year and more) are large in relation to its other values it means that the long-term movements dominate in the series. When the values of a spectral graph for high frequencies (i.e. below one year) are large in relation to its other values it means that the series are rather trendless and contains a lot of noise. When the values of a spectral graph are distributed randomly around a constant without any visible peaks, then it is highly probable that the series is a random process. The presence of seasonality in a time series is manifested in a spectral graph by the peaks on the seasonal frequencies.

Spectral graphs in GUI

Auto-regressive spectrum's properties




Figure 16: Text

1. The spectral graphs are available from: *Tools* → *Spectral analysis*.




Figure 17: Text

Tools for spectral analysis

2. When the first option is chosen JDemetra+ displays an empty *Auto-regressive spectrum* window. To start an analysis drag a single time series from the *Providers* window and drop it into the *Drop data here* area.

Launching an auto-regressive spectrum

3. An auto-regressive spectrum graph available in JDemetra+ is based on the relevant tool from the X-13ARIMA-SEATS program. It shows the spectral density (spectrum) function, which reformulates the content of the stationary time series' autocovariances in terms of amplitudes at frequencies of half a cycle per month or less. The number of observations, data transformations and other options such as the specification of the frequency grid and the order of the autoregressive polynomial (30 by default) can be specified by opening the *Window* → *Properties* from the main menu.

The *Auto-regressive - Properties* window contains the following options:

- **Log** - a log transformation of a time series;
- **Differencing** - transforms a data by calculating a regular (order 1,2..) or seasonal (order 4, 12, depending on the time series frequency) differences;




Figure 18: Text

- **Differencing lag** - the number of lags that the program will use to take differences. For example, if *Differencing lag* = 3 then the differencing filter does not apply to the first lag (default) but to the third lag.
 - **Last years** - a number of years at the end of the time series taken to produce autoregressive spectrum. By default, it is 0, which means that the whole time series is considered.
 - **Auto-regressive polynomial order** - the number of lags in the AR model that is used to estimate the spectral density. By default, the order of the autoregressive polynomial is set to 30 lags.
 - **Resolution** - the value 1 plots the spectral density estimate for the frequencies $\omega_j = \frac{2\pi j}{n}$, where $n \in (-\pi; \pi)$ is the size of the sample used to estimate the AR model. Increasing this value, which is set to 5 by default, will increase the precision of this grid.
4. The seasonality test described above uses an empirical criterion to check whether the series has a seasonal component that is predictable (stable) enough that it can be estimated with reasonable success. The peak in the [auto-regressive spectrum](#) has to be greater than the median of the 61 spectrum ordinates and has to exceed the two adjacent spectral values by more than a critical value. When such a case is detected, the test results are displayed in green.

An example of an-auto-regressive spectrum




Figure 19: Text

5. The second spectral graph is a periodogram. To perform the analysis of a single time series using this tool, choose *Tools* → *Spectral analysis* → *Periodogram* and drag and drop a series from the *Providers* window to the empty *Periodogram* window.




Figure 20: Text

Launching a periodogram

6. The sample size and data transformations can be specified by opening the *Window* → *Properties*, in the main menu. The *Periodogram - Properties* window contains the following options:
 - **Log** - a log transformation of a time series;

- **Differencing** - transforms the data by calculating regular (order 1,2..) or seasonal (order 4, 12, depending on the time series frequency) differences;
- **Differencing lag** - the number of lags that you will use to take differences. For example, if *Differencing lag* = 3 then the differencing filter does not apply to the first lag (default) but to the third lag.
- **Last years** - the number of years at the end of the time series taken to produce periodogram. By default it is 0, which means that the whole time series is considered.




Figure 21: Text

Periodogram's properties

7. The [periodogram](#) was one of the earliest tools used for the analysis of time series in the frequency domain. It enables the user to identify the dominant periods (or frequencies) of a time series. In general, the periodogram is a wildly fluctuating estimate of the spectrum with a high variance and is less stable than an auto-regressive spectrum.

An example of a periodogram




Figure 22: Text

8. The third spectral graph is the Tukey spectrum. To perform the analysis of time series using this tool, choose *Tools* → *Spectral analysis* → *Tukey spectrum* and drag and drop a single series from the *Providers* window to the empty *Periodogram* window.




Figure 23: Text

Launching a Tukey spectrum

9. The Tukey spectrum estimates the spectral density by smoothing the periodogram.

An example of a Tukey spectrum

10. The options for the Tuckey window can be specified by opening the *Window* → *Properties*




Figure 24: Text

from the main menu. The *Periodogram - Properties* window contains the following options:

- **Log** - a log transformation of a time series.
- **Differencing** - transforms the data by calculating regular (order 1, 2..) or seasonal (order 4, 12, depending on the time series frequency) differences.
- **Differencing lag** - the number of lags that you will use to take differences. For example, if *Differencing lag* = 3 then the differencing filter does not apply to the first lag (default) but to the third lag.
- **Taper part** – parameter larger than 0 and smaller or equal to one that shapes the curvature of the smoothing function that is applied to the auto-covariance function.
- **Window length** – the size of the window that is used to smooth the auto-covariance function. A value of zero includes the whole series.
- **Window type** – it refers to the weighting scheme that it is used to smooth the auto-covariance function. The available windows types (*Square*, *Welch*, *Tukey*, *Barlett*, *Hamming*, *Parzen*) are suitable to estimate the spectral density.

Tukey spectrum's properties

comment3: end part case studies > spectral graphs




Figure 25: Text

Reg-Arima models

Overview

The primary aim of seasonal adjustment is to remove the unobservable seasonal component from the observed series. The decomposition routines implemented in the seasonal adjustment methods make specific assumptions concerning the input series. One of the crucial assumptions is that the input series is stochastic, i.e. it is clean of deterministic effects. Another important limitation derives from the symmetric linear filter used in TRAMO-SEATS and X-13ARIMA-SEATS. A symmetric linear filter cannot be applied to the first and last observations with the same set of weights as for the central observations^[1]. Therefore, for the most recent observations these filters provide estimates that are subject to revisions.

To overcome these constraints both seasonal adjustment methods discussed here include a modelling step that aims to analyse the time series development and provide a better input for decomposition purposes. The tool that is frequently used for this purpose is the ARIMA model, as discussed by BOX, G.E.P., and JENKINS, G.M. (1970). However, time series are often affected by the outliers, other deterministic effects and missing observations. The presence of these effects is not in line with the ARIMA model assumptions. The presence of outliers and other deterministic effects impede the identification of an optimal ARIMA model due to the important bias in the estimation of parameters of [sample autocorrelation functions](#) (both global and partial)^[3]. Therefore, the original series need to be corrected for any deterministic effects and missing observations. This process is called linearisation and results in the stochastic series that can be modelled by ARIMA.

For this purpose both TRAMO and RegARIMA use regression models with ARIMA errors. With these models TRAMO and RegARIMA also produce forecasts.

X-11 decomposition

Introduction

A complete documentation of the X-11 method is available in LADIRAY, D., and QUENNEVILLE, B. (2001). The X-11 program is the result of a long tradition of non-parametric smoothing based on moving averages, which are weighted averages of a moving span of a time series (see hereafter). Moving averages have two important drawbacks:

- They are not resistant and might be deeply impacted by outliers;
- The smoothing of the ends of the series cannot be done except with asymmetric moving averages which introduce phase-shifts and delays in the detection of turning points.

These drawbacks adversely affect the X-11 output and stimulate the development of this method. To overcome these flaws first the series are modelled with a RegARIMA model that calculates forecasts and estimates the regression effects. Therefore, the seasonal adjustment process is divided into two parts.

- In a first step, the RegARIMA model is used to clean the series from non-linearities, mainly outliers and calendar effects. A global ARIMA model is adjusted to the series in order to compute the forecasts.
- In a second step, an enhanced version of the X-11 algorithm is used to compute the trend, the seasonal component and the irregular component.

The flow diagram for seasonal adjustment with X-13ARIMA-SEATS using the X-11 algorithm.

Moving averages

The moving average of coefficient θ_i is defined as:

$$M(X_t) = \sum_{k=-p}^{+f} \theta_k X_{t+k}$$




Figure 26: Text

The value at time t of the series is therefore replaced by a weighted average of p “past” values of the series, the current value, and f “future” values of the series. The quantity $p + f + 1$ is called the moving average order. When p is equal to f , that is, when the number of points in the past is the same as the number of points in the future, the moving average is said to be centred. If, in addition, $\theta_{-k} = \theta_k$ for any k , the moving average M is said to be symmetric. One of the simplest moving averages is the symmetric moving average of order $P = 2p + 1$ where all the weights are equal to $\frac{1}{P}$.

This moving average formula works well for all time series observations, except for the first p values and last f values. Generally, with a moving average of order $p + f + 1$ calculated for instant t with points p in the past and points f in the future, it will be impossible to smooth out the first p values and the last f values of the series because of lack of input to the moving average formula.

In the X-11 method, symmetric moving averages play an important role as they do not introduce any phase-shift in the smoothed series. But, to avoid losing information at the series ends, they are either supplemented by *ad hoc* asymmetric moving averages or applied on the series extended by forecasts.

For the estimation of the seasonal component, X-13ARIMA-SEATS uses $P \times Q$ composite moving averages, obtained by composing a simple moving average of order P , which coefficients are all equal to $\frac{1}{P}$, and a simple moving average of order Q , which coefficients are all equal to $\frac{1}{Q}$.

The composite moving averages are widely used by the X-11 method. For an initial estimation of trend X-11 method uses a 2×4 moving average in case of a quarterly time series while for a monthly time series a 2×12 moving average is applied. The 2×4 moving average is an average of order 5 with coefficients $\frac{1}{8} \{1, 2, 2, 2, 1\}$. It eliminates frequency $\frac{\pi}{2}$ corresponding to period 4 and therefore it is suitable for seasonal adjustment of the quarterly series with a constant seasonality. The 2×12 moving average, with coefficients $\frac{1}{24} \{1, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 1\}$ that retains linear trends, eliminates order-12 constant seasonality and minimises the variance of the irregular component. The 2×4 and 2×12 moving averages are also used in the X-11 method to normalise the seasonal factors. The composite moving averages are also used to extract the seasonal component. These, which are used in the purely automatic run of the X-11 method (without any intervention from the user) are 3×3 , 3×5 and 3×9 .

In the estimation of the trend also Henderson moving averages are used. These filters have been chosen for their smoothing properties. The coefficients of a Henderson moving average of order $2p + 1$ may be calculated using the formula:

$$\theta_i = \frac{315 \left[(n-1)^2 - i^2 \right] \left[n^2 - i^2 \right] \left[(n+1)^2 - i^2 \right] \left[3n^2 - 16 - 11i^2 \right]}{8n(n^2 - 1)(4n^2 - 1)(4n^2 - 9)(4n^2 - 25)}$$

,

2

where: $n = p + 2n = p + 2$.

The basic algorithm of the X-11 method

The X-11 method is based on an iterative principle of estimation of the different components using appropriate moving averages at each step of the algorithm. The successive results are saved in tables. The list of the X-11 tables displayed in JDemetra+ is included at the end of this section.

The basic algorithm of the X-11 method will be presented for a monthly time series X_t that is assumed to be decomposable into trend, seasonality and irregular component according to an additive model $X_t = TC_t + S_t + I_t$.

A simple seasonal adjustment algorithm can be thought of in eight steps. The steps presented below are designed for the monthly time series. In the algorithm that is run for the quarterly time series the 2×4 moving average instead of the 2×12 moving average is used.

Step 1: Estimation of Trend by 2×12 moving average:

$$TC_t^{(1)} = M_{2 \times 12}(X_t)$$

3

Step 2: Estimation of the Seasonal-Irregular component:

$$(S_t + I_t)^{(1)} = X_t - TC_t^{(1)}$$

4

Step 3: Estimation of the Seasonal component by

$$\mathbf{3 \times 3}$$

moving average over each month:

$$S_t^{(1)} - M_{3 \times 3} \left[(S_t + I_t)^{(1)} \right]$$

5

The moving average used here is a 3×3 moving average over 5 terms, with coefficients $\frac{1}{9} \{1, 2, 3, 2, 1\}$. The seasonal component is then centred using a 2×12 moving average.

$$\tilde{S}_t^{(1)} = S_t^{(1)} - M_{2 \times 12} (S_t^{(1)})$$

6

Step 4: Estimation of the seasonally adjusted series:

$$SA_t^{(1)} = (TC_t + I_t)^{(1)} = X_t - \tilde{S}_t^{(1)}$$

7

This first estimation of the seasonally adjusted series must, by construction, contain less seasonality. The X-11 method again executes the algorithm presented above, changing the moving averages to take this property into account.

Step 5: Estimation of Trend by 13-term Henderson moving average:

$$TC_t^{(2)} = H_{13} (SA_t^{(1)})$$

8

Henderson moving averages, while they do not have special properties in terms of eliminating seasonality (limited or none at this stage), have a very good smoothing power and retain a local polynomial trend of degree 2 and preserve a local polynomial trend of degree 3.

Step 6: Estimation of the Seasonal-Irregular component:

$$(S_t + I_t)^{(2)} = X_t - TC_t^{(2)}$$

Step 7: Estimation of the Seasonal component by 3×5 moving average over each month:

$$S_t^{(2)} - M_{3 \times 3} \left[(S_t + I_t)^{(2)} \right]$$

The moving average used here is a 3×5 moving average over 7 terms, of coefficients

$$\frac{1}{15} \{1, 2, 3, 3, 3, 2, 1\}$$

and retains linear trends. The coefficients are then normalised such that their sum over the whole 12-month period is approximately cancelled out:

$$\widetilde{S}_t^{(2)} = S_t^{(2)} - M_{2 \times 12} (S_t^{(2)})$$

Step 8: Estimation of the seasonally adjusted series:

$$SA_t^{(2)} = (TC_t + I_t)^{(2)} = X_t - \widetilde{S}_t^{(2)}$$

The whole difficulty lies, then, in the choice of the moving averages used for the estimation of the trend in steps 1 and 5 on the one hand, and for the estimation of the seasonal component in steps 3 and 5. The course of the algorithm in the form that is implemented in JDemetra+ is presented in the figure below. The adjustment for trading day effects, which is present in the original X-11 program, is omitted here, as since calendar correction is performed by the RegARIMA model, JDemetra+ does not perform further adjustment for these effects in the decomposition step.

A workflow diagram for the X-11 algorithm based upon training material from the Deutsche Bundesbank

0.0.0.0.1 * The iterative principle of X-11

To evaluate the different components of a series, while taking into account the possible presence of extreme observations, X-11 will proceed iteratively: estimation of components, search for disruptive effects in the irregular component, estimation of components over a corrected series, search for disruptive effects in the irregular component, and so on.

The Census X-11 program presents four processing stages (A, B, C, and D), plus 3 stages, E, F, and G, that propose statistics and charts and are not part of the decomposition per se. In stages B, C and D the basic algorithm is used as is indicated in the figure below.

A workflow diagram for the X-11 algorithm implemented in JDemetra+. Source: Based upon training material from the Deutsche Bundesbank

- **Part A: Pre-adjustments**

This part, which is not obligatory, corresponds in X-13ARIMA-SEATS to the first cleaning of the series done using the RegARIMA facilities: detection and estimation of outliers and calendar effects (trading day and Easter), forecasts and backcasts^[61] of the series. Based on these results, the program calculates prior adjustment factors that are applied to the raw series. The series thus corrected, Table B1 of the printouts, then proceeds to part B.

- **Part B: First automatic correction of the series**

This stage consists of a first estimation and down-weighting of the extreme observations and, if requested, a first estimation of the calendar effects. This stage is performed by applying the basic algorithm detailed earlier. These operations lead to Table B20, adjustment values for extreme observations, used to correct the unadjusted series and result in the series from Table C1.

- **Part C: Second automatic correction of the series**

Applying the basic algorithm once again, this part leads to a more precise estimation of replacement values of the extreme observations (Table C20). The series, finally “cleaned up”, is shown in Table D1 of the printouts.

- **Part D: Seasonal adjustment**

This part, at which our basic algorithm is applied for the last time, is that of the seasonal adjustment, as it leads to final estimates:

- of the seasonal component (Table D10);
- of the seasonally adjusted series (Table D11);
- of the trend component (Table D12);
- of the irregular component (Table D13).

- **Part E: Components modified for large extreme values**

Parts E includes:

- Components modified for large extreme values;
- Comparison the annual totals of the raw time series and seasonally adjusted time series;
- Changes in the final seasonally adjusted series;
- Changes in the final trend;
- Robust estimation of the final seasonally adjusted series.

The results from part E are used in part F to calculate the quality measures.

- **Part F: Seasonal adjustment quality measures**

Part F contains statistics for judging the quality of the seasonal adjustment. JDemetra+ presents selected output for part F, i.e.:

- M and Q statistics;
- Tables.

- **Part G: Graphics**

Part G presents spectra estimated for:

- Raw time series adjusted a priori (Table B1);
- Seasonally adjusted time series modified for large extreme values (Table E2);
- Final irregular component adjusted for large extreme values (Table E3).

Originally, graphics were displayed in character mode. In JDemetra+, these graphics are replaced favourably by the usual graphics software.

The Henderson moving average and the trend estimation

In iteration B (Table B7), iteration C (Table C7) and iteration D (Table D7 and Table D12) the trend component is extracted from an estimate of the seasonally adjusted series using Henderson moving averages. The length of the Henderson filter is chosen automatically by X-13ARIMA-SEATS in a two-step procedure.

It is possible to specify the length of the Henderson moving average to be used. X-13ARIMA-SEATS provides an automatic choice between a 9-term, a 13-term or a 23-term moving average. The automatic choice of the order of the moving average is based on the value of an indicator called $\frac{T}{C}$ ratio which compares the magnitude of period-on-period movements in the irregular component with those in the trend. The larger the ratio, the higher the order of the moving

average selected. Moreover, X-13ARIMA-SEATS allows the user to choose manually any odd-numbered Henderson moving average. The procedure used in each part is very similar; the only differences are the number of options available and the treatment of the observations in the both ends of the series. The procedure below is applied for a monthly time series.

In order to calculate $\frac{\bar{I}}{\bar{C}}$ ratio a first decomposition of the SA series (seasonally adjusted) is computed using a 13-term Henderson moving average.

For both the trend (C) and irregular (I) components, the average of the absolute values for monthly growth rates (multiplicative model) or for monthly growth (additive model) are computed. They are denoted as \bar{C} and \bar{I} , receptively, where $\bar{C} = \frac{1}{n-1} \sum_{t=2}^n |C_t - C_{t-1}|$ and $\bar{I} = \frac{1}{n-1} \sum_{t=2}^n |I_t - I_{t-1}|$.

Then the value of $\frac{\bar{I}}{\bar{C}}$ ratio is checked and in iteration B:

- If the ratio is smaller than 1, a 9-term Henderson moving average is selected;
- Otherwise, a 13-term Henderson moving average is selected.

Then the trend is computed by applying the selected Henderson filter to the seasonally adjusted series from Table B6. The observations at the beginning and at the end of the time series that cannot be computed by means of symmetric Henderson filters are estimated by ad hoc asymmetric moving averages.

In iterations C and D:

- If the ratio is smaller than 1, a 9-term Henderson moving average is selected;
- If the ratio is greater than 3.5, a 23-term Henderson moving average is selected.
- Otherwise, a 13-term Henderson moving average is selected.

The trend is computed by applying selected Henderson filter to the seasonally adjusted series from Table C6, Table D7 or Table D12, accordingly. At the both ends of the series, where a central Henderson filter cannot be applied, the asymmetric ends weights for the 7 term Henderson filter are used.

0.0.0.0.2 * Choosing the composite moving averages when estimating the seasonal component

In iteration D, Table D10 shows an estimate of the seasonal factors implemented on the basis of the modified SI (Seasonal – Irregular) factors estimated in Tables D4 and D9bis. This component will have to be smoothed to estimate the seasonal component; depending on the importance of the irregular in the SI component, we will have to use moving averages of varying length as in the estimate of the Trend/Cycle where the $\frac{\bar{I}}{\bar{C}}$ ratio was used to select the length of the Henderson moving average. The estimation includes several steps.

Step 1: Estimating the irregular and seasonal components

An estimate of the seasonal component is obtained by smoothing, month by month and therefore column by column, Table D9bis using a simple 7-term moving average, i.e. of coefficients $\frac{1}{7} \{1, 1, 1, 1, 1, 1, 1\}$. In order not to lose three points at the beginning and end of each column, all columns are completed as follows. Let us assume that the column that corresponds to the month is composed of N values $\{x_1, x_2, x_3, \dots, x_{N-1}, x_N\}$. It will be transformed into a series $\{x_{-2}, x_{-1}, x_0, x_1, x_2, x_3, \dots, x_{N-1}, x_N, x_{N+1}, x_{N+2}, x_{N+3}\}$ with $x_{-2} = x_{-1} = x_0 = \frac{x_1 + x_2 + x_3}{3}$ and $x_{N+1} = x_{N+2} = x_{N+3} = \frac{x_N + x_{N-1} + x_{N-2}}{3}$. We then have the required estimates: $S = M_7(D9bis)$ and $I = D9bis - S$.

Step 2: Calculating the Moving Seasonality Ratios

For each i^{th} month the mean annual changes for each component is obtained by calculating

$$\bar{S}_i = \frac{1}{N_i - 1} \sum_{t=2}^{N_i} |S_{i,t} - S_{i,t-1}|$$

and

$$\bar{I}_i = \frac{1}{N_i - 1} \sum_{t=2}^{N_i} |I_{i,t} - I_{i,t-1}|$$

, where N_i refers to the number of months in the data, and the moving seasonality ratio of month i :

$$MSR_i = \frac{\bar{I}_i}{\bar{S}_i}$$

. These ratios are presented in *Details* of the *Quality Measures* node under the *Decomposition (X11)* section. These ratios are used to compare the year-on-year changes in the irregular component with those in the seasonal component. The idea is to obtain, for each month, an indicator capable of selecting the appropriate moving average for the removal of any noise and providing a good estimate of the seasonal factor. The higher the ratio, the more erratic the series, and the greater the order of the moving average should be used. As for the rest, by default the program selects the same moving average for each month, but the user can select different moving averages for each month.

Step 3: Calculating the overall Moving Seasonality Ratio

The overall Moving Seasonality Ratio is calculated as follows:

$$MSR_i = \frac{\sum_i N_i \bar{I}_i}{\sum_i N_i \bar{S}_i}$$

Step 4: Selecting a moving average and estimating the seasonal component

Depending on the value of the ratio, the program automatically selects a moving average that is applied, column by column (i.e. month by month) to the Seasonal/Irregular component in Table D8 modified, for extreme values, using values in Table D9.

The default selection procedure of a moving average is based on the Moving Seasonality Ratio in the following way:

- If this ratio occurs within zone A ($MSR < 2.5$), a 3×3 moving average is used; if it occurs within zone C ($3.5 < MSR < 5.5$), a 3×5 moving average is selected; if it occurs within zone E ($MSR \geq 6.5$), a 3×9 moving average is used;
- If the MSR occurs within zone B or D, one year of observations is removed from the end of the series, and the MSR is recalculated. If the ratio again occurs within zones B or D, we start over again, removing a maximum of five years of observations. If this does not work, i.e. if we are again within zones B or D, a 3×5 moving average is selected.

The chosen symmetric moving average corresponds, as the case may be 5 (3×3), 7 (3×5) or 11 (3×9) terms, and therefore does not provide an estimate for the values of seasonal factors in the first 2 (or 3 or 5) and the last 2 (or 3 or 5) years. These are then calculated using associated asymmetric moving averages.

Moving average selection procedure, source: DAGUM, E. B.(1999)

0.0.0.0.3 * Identification and replacement of extreme values

X-13ARIMA-SEATS detects and removes outliers in the RegARIMA part. However, if there is a seasonal heteroscedasticity in a time series i.e. the variance of the irregular component is different in different calendar months. Examples for this effect could be the weather and snow-dependent output of the construction sector in Germany during winter, or changes in Christmas allowances in Germany and resulting from this a transformation in retail trade turnover before Christmas. The ARIMA model is not on its own able to cope with this characteristic. The practical consequence is given by the detection of additional extreme values by X-11. This may not be appropriate if the seasonal heteroscedasticity is produced by political interventions or other influences. The ARIMA models assume a constant variance and are therefore not by themselves able to cope with this problem. Choosing longer (in the case of diverging weather conditions in the winter time for the construction sector) or shorter filters (in the case of a changing pattern of retail trade turnover in the Christmas time) may be reasonable in such cases. It may even be sensible to take into account the possibility of period-specific (e.g. month-specific) standard deviations, which can be done by changing the default settings of the **calendarsigma** parameter (see [Specifications-X13](#) section). The value of the **calendarsigma** parameter will have an impact on the method of calculation of the moving standard deviation in the procedure for extreme values detection presented below.

Step 1: Estimating the seasonal component

The seasonal component is estimated by smoothing the SI component separately for each period using a 3×3 moving average, i.e.:

$$\frac{1}{9} \times \begin{pmatrix} 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, \\ 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, \\ 3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, \\ 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, \\ 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, \end{pmatrix}$$

14

Step 2: Normalizing the seasonal factors

The preliminary seasonal factors are normalized in such a way that for one year their average is equal to zero (additive model) or to unity (multiplicative model).

Step 3: Estimating the irregular component

The initial normalized seasonal factors are removed from the Seasonal-Irregular component to provide an estimate of the irregular component.

Step 4: Calculating a moving standard deviation

By default, a moving standard deviation of the irregular component is calculated at five-year intervals. Each standard deviation is associated with the central year used to calculate it. The values in the central year, which in the absolute terms deviate from average by more than the **Usigma** parameter are marked as extreme values and assigned a zero weight. After excluding the extreme values the moving standard deviation is calculated once again.

Step 5: Detecting extreme values and weighting the irregular

The default settings for assigning a weight to each value of irregular component are:

- Values which are more than **Usigma** (2.5, by default) standard deviations away (in the absolute terms) from the 0 (additive) or 1 (multiplicative) are assigned a zero weight;
- Values which are less than 1.5 standard deviations away (in the absolute terms) from the 0 (additive) or 1 (multiplicative) are assigned a full weight (equal to one);
- Values which lie between 1.5 and 2.5 standard deviations away (in the absolute terms) from the 0 (additive) or 1 (multiplicative) are assigned a weight that varies linearly between 0 and 1 depending on their position.

The default boundaries for the detection of the extreme values can be changed with **LSigma** and **USigma** parameters

Step 6: Adjusting extreme values of the seasonal-irregular component

Values of the SI component are considered extreme when a weight less than 1 is assigned to their irregular. Those values are replaced by a weighted average of five values:

- The value itself with its weight;
- The two preceding values, for the same period, having a full weight(if available);
- The next two values, for the same period, having full a weight (if available).

When the four full-weight values are not available, then a simple average of all the values available for the given period is taken.

This general algorithm is used with some modification in parts B and C for detection and replacement of extreme values.

0.0.0.0.4 * X-11 tables

The list of tables produced by JDemetra+ is presented below. It is not identical to the output produced by the original X-11 program.

Part A – Preliminary Estimation of Outliers and Calendar Effects.

This part includes prior modifications to the original data made in the RegARIMA part:

- Table A1 – Original series;
- Table A1a – Forecast of Original Series;
- Table A2 – Leap year effect;
- Table A6 – Trading Day effect (1 or 6 variables);
- Table A7 – The Easter effect;
- Table A8 – Total Outlier Effect;
- Table A8i – Additive outlier effect;
- Table A8t – Level shift effect;
- Table A8s – Transitory effect;
- Table A9 – Effect of user-defined regression variables assigned to the seasonally adjusted series or for which the component has not been defined;
- Table 9sa – Effect of user-defined regression variables assigned to the seasonally adjusted series;

- Table9u – Effect of user-defined regression variables for which the component has not been defined.

Part B – Preliminary Estimation of the Time Series Components:

- Table B1 – Original series after adjustment by the RegARIMA model;
- Table B2 – Unmodified Trend (preliminary estimation using composite moving average);
- Table B3 – Unmodified Seasonal – Irregular Component (preliminary estimation);
- Table B4 – Replacement Values for Extreme SI Values;
- Table B5 – Seasonal Component;
- Table B6 – Seasonally Adjusted Series;
- Table B7 – Trend (estimation using Henderson moving average);
- Table B8 – Unmodified Seasonal – Irregular Component;
- Table B9 – Replacement Values for Extreme SI Values;
- Table B10 – Seasonal Component;
- Table B11 – Seasonally Adjusted Series;
- Table B13 – Irregular Component;
- Table B17 – Preliminary Weights for the Irregular;
- Table B20 – Adjustment Values for Extreme Irregulars.

Part C – Final Estimation of Extreme Values and Calendar Effects:

- Table C1 – Modified Raw Series;
- Table C2 – Trend (preliminary estimation using composite moving average);
- Table C4 – Modified Seasonal – Irregular Component;
- Table C5 – Seasonal Component;
- Table C6 – Seasonally Adjusted Series;
- Table C7 – Trend (estimation using Henderson moving average);
- Table C9 – Seasonal – Irregular Component;
- Table C10 – Seasonal Component;
- Table C11 – Seasonally Adjusted Series;
- Table C13 – Irregular Component;

- Table C20 – Adjustment Values for Extreme Irregulars.

Part D – Final Estimation of the Different Components:

- Table D1 – Modified Raw Series;
- Table D2 – Trend (preliminary estimation using composite moving average);
- Table D4 – Modified Seasonal – Irregular Component;
- Table D5 – Seasonal Component;
- Table D6 – Seasonally Adjusted Series;
- Table D7 – Trend (estimation using Henderson moving average);
- Table D8 – Unmodified Seasonal – Irregular Component;
- Table D9 – Replacement Values for Extreme SI Values;
- Table D10 – Final Seasonal Factors;
- Table D10A – Forecast of Final Seasonal Factors;
- Table D11 – Final Seasonally Adjusted Series;
- Table D11A – Forecast of Final Seasonally Adjusted Series;
- Table D12 – Final Trend (estimation using Henderson moving average);
- Table D12A – Forecast of Final Trend Component;
- Table D13 – Final Irregular Component;
- Table D16 – Seasonal and Calendar Effects;
- Table D16A – Forecast of Seasonal and Calendar Component;
- Table D18 – Combined Calendar Effects Factors.

Part E – Components Modified for Large Extreme Values:

- Table E1 – Raw Series Modified for Large Extreme Values;
- Table E2 – SA Series Modified for Large Extreme Values;
- Table E3 – Final Irregular Component Adjusted for Large Extreme Values;
- Table E11 – Robust Estimation of the Final SA Series.

Part F – Quality indicators:

- Table F2A – Changes, in the absolute values, of the principal components;
- Table F2B – Relative contribution of components to changes in the raw series;

- Table F2C – Averages and standard deviations of changes as a function of the time lag;
- Table F2D – Average duration of run;
- Table F2E – I/C ratio for periods span;
- Table F2F – Relative contribution of components to the variance of the stationary part of the original series;
- Table F2G – Autocorrelogram of the irregular component.

Filter length choice

A seasonal filter is a weighted average of a moving span of fixed length within a time series that can be used to remove a fixed seasonal pattern. X-13ARIMA-SEATS uses several of these filters, according to the needs of the different stages of the program. As only X-13ARIMA-SEATS allows the user to manually select seasonal filters, this case study can be applied only to the X-13ARIMA-SEATS specifications.

The automatic seasonal adjustment procedure uses the default options to select the most appropriate moving average. However there are occasions when the user will need to specify a different seasonal moving average to that identified by the program. For example, if the SI values do not closely follow the seasonal component, it may be appropriate to use a shorter moving average. Also the presence of sudden breaks in the seasonal pattern – e.g. due to changes in the methodology – can negatively impact on the automatic selection of the most appropriate seasonal filter. In such cases the usage of short seasonal filters in the selected months or quarters can be considered. Usually, a shorter seasonal filter (3×1) allows seasonality to change very rapidly over time. However, a very short seasonal filter should not normally be used, as it might often lead to large revisions as new data becomes available. If a short filter is to be used it will usually be limited to one month/quarter with a known reason for wanting to capture a rapidly changing seasonality.

In the standard situation one seasonal filter is applied to all individual months/quarters. The estimation of seasonal movements is therefore based on the sample windows of equal lengths for each individual month/quarter (i.e. for each month/quarter the seasonal filter length or the number of years representing the major part of the seasonal filter weights is identical). This approach relies on the assumption that the number of past periods in which the conditions causing seasonal behaviour are sufficiently homogenous is the same in all months/quarters. However, this assumption does not always hold. Seasonal causes may change in one month, while staying the same in others¹⁶. For instance, seasonal heteroskedasticity might require different filter lengths in different months or quarters.

¹⁶When the series are non-stationary differentiation is performed before the seasonality tests.

Another interesting example is industrial production in Germany. It can be influenced by school holidays, since many employees have school-age children, which interrupt their working pattern during these school holidays. Consequently, businesses may temporarily suspend or lower production during these periods. Since school holidays do not occur at the same time throughout Germany and their timing varies from year to year in the individual federal states, the effect is not completely captured by seasonal adjustment. And since school holidays are treated as usual working days, these effects are not captured by calendar adjustment either. The majority of school holidays in Germany can take place either in July or in August. This yields higher variances in the irregular component for these months compared to the rest of the year. Therefore, in this case a longer seasonal filter is used for these months to account for this.

Another example might be given by German retail trade. Due to changes in the consumers' behaviour around Christmas – possibly more gifts of money – the seasonal peak in December has become steadily less pronounced. To account for this moving seasonality, shorter seasonal filters in December than during the rest of the year need to be applied.

JDemetra+ offers the options to assign a different seasonal filter length to each period (month or quarter). The program offers these options in the *single spec* mode as well as in the *multispec* mode, albeit they are available only in the *Specifications* window, after a document is created.

M-stats

The details about the measures are given below.

- $M1$ measures the contribution of the irregular component to the total variance. When it is above 1 some changes in outlier correction should be considered.
- $M2$, which is a very similar to $M1$, is calculated on the basis of the contribution of the irregular component to the stationary portion of the variance. When it is above 1, some changes in an outlier correction should be considered.
- $M3$ compares the irregular to the trend taken from a preliminary estimate of the seasonally adjusted series. If this ratio is too large, it is difficult to separate the two components from each other. When it is above 1 some changes in outlier correction should be considered.
- $M4$ tests the randomness of the irregular component. A value above 1 denotes a correlation in the irregular component. In such case a shorter seasonal moving average filter should be considered.
- $M5$ is used to compare the significance of changes in the trend with that in the irregular. When it is above 1 some changes in outlier correction should be considered.

- $M6$ checks the SI (seasonal – irregular components ratio). If annual changes in the irregular component are too small in relation to the annual changes in the seasonal component, the 3×5 seasonal filter used for the estimation of the seasonal component is not flexible enough to follow the seasonal movement. In such case a longer seasonal moving average filter should be considered. It should be stressed that $M6$ is calculated only if the 3×5 filter has been applied in the model.
- $M7$ is the combined test for the presence of an identifiable seasonality. The test compares the relative contribution of stable and moving seasonality¹⁷.
- $M8$ to $M11$ measure if the movements due to the short-term quasi-random variations and movements due to the long-term changes are not changing too much over the years. If the changes are too strong then the seasonal factors could be erroneous. In such case a correction for a seasonal break or the change of the seasonal filter should be considered.

The Q statistic is a composite indicator calculated from the M statistics.

Edit : problem with tables display

$$Q = \frac{10M1 + 11M2 + 10M3 + 8M4 + 11M5 + 10M6 + 18M7 + 7M8 + 7M9 + 4M10 + 4M11}{100}$$

[1]

$Q - M2$ (also called $Q2$) is the Q statistic for which the $M2$ statistics was excluded from the formula, i.e.:

$$Q - M2 = \frac{10M1 + 10M3 + 8M4 + 11M5 + 10M6 + 18M7 + 7M8 + 7M9 + 4M10 + 4M11}{89}$$

[2]

If a time series does not cover at least 6 years, the $M8$, $M9$, $M10$ and $M11$ statistics cannot be calculated. In this case the Q statistic is computed as:

$$Q = \frac{14M1 + 15M2 + 10M3 + 8M4 + 11M5 + 10M6 + 32M7}{100}$$

The model has a satisfactory quality if the Q statistic is lower than 1.

The tables displayed in the *Quality measures* → *Details* node correspond to the F-set of tables produced by the original X-11 algorithm. To facilitate the analysis of the results, the numbers and the names of the tables are given under each table following the convention used in LADIRAY, D., and QUENNEVILLE, B. (1999).

¹⁷See section [Combined seasonality tests](#).

Detailed tables

The first table presents the average percent change without regard to sign of the percent changes (multiplicative model) or average differences (additive model) over several periods (from 1 to 12 for a monthly series, from 1 to 4 for a quarterly series) for the following series:

- O – Original series (Table A1);
- CI – Final seasonally adjusted series (Table D11);
- I – Final irregular component (Table D13);
- C – Final trend (Table D12);
- S – Final seasonal factors (Table D10);
- P – Preliminary adjustment coefficients, i.e. regressors estimated by the RegARIMA model (Table A2);
- $TD\&H$ – Final calendar component (Tables A6 and A7);
- $Mod.O$ – Original series adjusted for extreme values (Table E1);
- $Mod.CI$ – Final seasonally adjusted series corrected for extreme values (Table E2);
- $Mod.I$ – Final irregular component adjusted for extreme values (Table E3).

In the case of an additive decomposition, for each component the average absolute changes over several periods are calculated as¹⁸:

$$\text{Component}_d = \frac{1}{n-d} \sum_{t=d+1}^n |Table_t - Table_{t-d}|$$

[4]

where:

d – time lag in periods (from a monthly time series d varies from 4 or from 1 to 12);

n – total number of observations per period;

Component – the name of the component;

Table – the name of the table that corresponds to the component.

Table F2A – changes, in the absolute values, of the principal components

¹⁸For the multiplicative decomposition the following formula is used:

$$\text{Component}_d = \frac{1}{n-d} \sum_{t=d+1}^n \left| \frac{Table_t}{Table_{t-d}} - 1 \right|$$

Average percent change without regard to sign over the indicated span

Span	O	CI	I	C	S	P	TD&H	Mod.O	Mod.CI	Mod.I
1	7,50	3,81	3,49	1,42	6,99	0,00	0,00	7,75	3,57	3,29
2	5,33	4,88	3,90	2,88	3,57	0,00	0,00	5,40	4,61	3,55
3	8,23	5,75	3,74	4,39	7,16	0,00	0,00	8,53	5,50	3,39
4	6,36	6,75	3,76	5,94	0,00	0,00	0,00	6,74	6,74	3,56

Figure 27: Text

Next, Table F2B of relative contributions of the different components to the differences (additive model) or percent changes (multiplicative model) in the original series is displayed. They express the relative importance of the changes in each component. Assuming that the components are independent, the following relation is valid:

$$O_d^2 \approx C_d^2 + S_d^2 + I_d^2 + P_d^2 + TD\&H_d^2$$

. [5]

In order to simplify the analysis, the approximation can be replaced by the following equation:

$$O_d^{*2} = C_d^2 + S_d^2 + I_d^2 + P_d^2 + TD\&H_d^2$$

. [6]

The notation is the same as for Table F2A. The column Total denotes total changes in the raw time series.

Data presented in Table F2B indicate the relative contribution of each component to the percent changes (differences) in the original series over each span, and are calculated as:

$$\frac{I_d^2}{O_d^{*2}},$$

$$\frac{C_d^2}{O_d^{*2}},$$

$$\frac{S_d^2}{O_d^{*2}},$$

$$\frac{P_d^2}{O_d^{*2}}$$

and $\frac{TD\&H_d^2}{O_d^{*2}}$

where: $O_d^{*2} = I_d^2 + C_d^2 + S_d^2 + P_d^2 + TD\&H_d^2$.

The last column presents the *Ratio* calculated as:

$$100 \times \frac{O_d^{*2}}{O_d^2},$$

which is an indicator of how well the approximation

$$(O_d^*)^2 \approx O_d^2$$

holds.

Relative contributions to the variance of the percent change in the components of the original series

Span	I	C	S	P	TD&H	Total	Ratio
1	17,53	3,27	79,20	0,00	0,00	100,00	102,79
2	37,38	24,71	37,91	0,00	0,00	100,00	115,35
3	13,97	23,47	62,56	0,00	0,00	100,00	112,79
4	26,47	73,53	0,00	0,00	0,00	100,00	105,49

Figure 28: Text

Table F2B – relative contribution of components to changes in the raw series

When an additive decomposition is used, Table F2C presents the average and standard deviation of changes calculated for each time lag d , taking into consideration the sign of the changes of the raw series and its components. In case of a multiplicative decomposition the respective table shows the average percent differences and related standard deviations.

Average percent change with regard to sign and standard deviation over indicated span

Span	O		I		C		S		CI	
	Avg	S.D.	Avg	S.D.	Avg	S.D.	Avg	S.D.	Avg	S.D.
1	1,97	8,67	0,05	3,73	1,41	0,48	0,53	8,01	1,46	3,81
2	3,19	5,72	0,15	4,48	2,86	0,97	0,24	4,42	3,02	4,72
3	4,97	9,47	0,09	4,52	4,36	1,44	0,55	8,30	4,46	4,97
4	5,93	3,81	0,10	4,32	5,90	1,90	0,00	0,00	6,01	5,06

Figure 29: Text

Table F2C – Averages and standard deviations of changes as a function of the time lag

Average duration of run is an average number of consecutive monthly (or quarterly) changes in the same direction (no change is counted as a change in the same direction as the preceding change). JDemetra+ displays this indicator for the seasonally adjusted series, for the trend and for the irregular component.

Average duration of run.

CI	8.44
I	1.31
C	15.20

Figure 30: Text

Table F2D – Average duration of run

The $\frac{I}{C}$ ratios for each value of time lag d , presented in Table F2E, are computed on a basis of the data in Table F2A.

<u>I/C Ratio for indicated span.</u>	
1	0.150
2	0.052
3	0.039
4	0.031

I/C Ratio: 0.314

Figure 31: Text

Table F2E – $\frac{I}{C}$ ratio for periods span

The relative contribution of components to the variance of the stationary part of the original series is calculated for the irregular component (I), trend made stationary¹⁹ (C), seasonal component (S) and calendar effects (TD&H). The short description of the calculation method is given in LADIRAY, D., and QUENNEVILLE, B. (1999).

<u>Relative contribution of the components to the stationary portion of the variance in the original series.</u>	
I	0.01
C	99.56
S	0.15
P	0.00
TD&H	0.00
Total	99.72

Figure 32: Text

Table F2F – Relative contribution of components to the variance of the stationary part of the original series

The last table shows the autocorrelogram of the irregular component from Table D13. In the case of multiplicative decomposition it is calculated for time lags between 1 and the number of periods per year +2 using the formula²⁰:

¹⁹The component is estimated by extracting a linear trend from the trend component presented in Table D12.

²⁰For the additive decomposition the formula is:

$$Corr_k I_t = \frac{\sum_{t=k+1}^N (I_t \times I_{t-k})}{\sum_{t=1}^N (I_t)^2}$$

$$\text{Corr}_k I = \frac{\sum_{t=k+1}^N (I_t - 1)(I_{t-k} - 1)}{\sum_{t=1}^N (I_t - 1)^2}$$

[7]

where N is number of observations in the time series and k the lag.

Autocorrelation of the irregular:

1	-0.601
2	0.200
3	0.019
4	-0.147
5	0.187
6	-0.138

Figure 33: Text

Table F2G – Autocorrelation of the irregular component

The Cochran test is design to identify the heterogeneity of a series of variances. X-13-ARIMA-SEATS uses this test in the extreme value detection procedure to check if the irregular component is heteroskedastic. In this procedure the standard errors of the irregular component are used for an identification of extreme values. If the null hypothesis that for all the periods (months, quarters) the variances of the irregular component are identical is rejected, the standard errors will be computed separately for each period (in case the option *Calendar-sigma=signif* has been selected).

Heteroskedasticity (Cochran test on equal variances within each period)

Test statistic	Critical value (5% level)	Decision
0.1303	0.15	Null hypothesis is not rejected.

Figure 34: Text

Cochran test

For each i^{th} month we will be looking at the mean annual changes for each component by calculating:

$$\bar{S}_i = \frac{1}{N_i - 1} \sum_{t=2}^{N_i} |S_{i,t} - S_{i,t-1}|$$

and

$$\bar{I}_i = \frac{1}{N_i - 1} \sum_{t=2}^{N_i} |I_{i,t} - I_{i,t-1}|$$

,

where N_i refers to the number of months i in the data, and the moving seasonality ratio of month i :

$$MSR_i = \frac{\bar{I}_i}{\bar{S}_i}$$

These ratios are published in Table D9A in X13ARIMA-SEATS software. In JDemetra+ they are presented in the details of the quality measures.

The [Moving Seasonality Ratio \(MSR\)](#) is used to measure the amount of noise in the Seasonal-Irregular component. By studying these values, the user can [select for each period the seasonal filter](#) that is the most suitable given the noisiness of the series.

Moving Seasonality Ratios (MSR):

Period	I	S	MSR
1	0.0597	0.0211	2.8292
2	0.0808	0.0135	5.9850
3	0.0767	0.0139	5.5038
4	0.0777	0.0262	2.9640

Figure 35: Text

Table D9a – Moving seasonality ratios

STL: Local regression decomposition

SEATS decomposition

Introduction

SEATS is a program for estimating unobserved components in a time series. It follows the ARIMA-model-based (AMB) method, developed from the work of CLEVELAND, W.P., and TIAO, G.C. (1976), BURMAN, J.P. (1980), HILLMER, S.C., and TIAO, G.C. (1982), BELL, W.R., and HILLMER, S.C. (1984) and MARAVALL, A., and PIERCE, D.A. (1987).

In JDemetra+ the input for the model based signal extraction procedure is always provided by TRAMO and includes the original series y_t , the linearized series x_t (i.e. the original series y_t with the deterministic effects removed), the ARIMA model for the stochastic (linearized) time series x_t and the deterministic effects (calendar effects, outliers and other regression variable effects)²¹. SEATS decomposes the linearized series (and the ARIMA model) into trend, seasonal, transitory and irregular components, provides forecasts for these components, together with the associated standard errors, and finally assign the deterministic effects to each component yielding the *final* components²². The Minimum Mean Square Error (MMSE) estimators of the components are computed with a Wiener-Kolmogorov filter applied to the finite series extended with forecasts and backcasts²³.

ARIMA modelling of the input series

One of the fundamental assumptions made by SEATS is that the linearized time series x_t follows the ARIMA model

$$\phi(B)\delta(B)x_t = \theta(B)a_t$$

[1]

where:

B – the backshift operator ($Bx_t = x_{t-1}$);

²¹In the original software SEATS can be used either with TRAMO, operating on the input received from the latter, or alone, fitting an ARIMA model to the series.

²²GÓMEZ, V., and MARAVALL, A. (1998).

²³GÓMEZ, V., and MARAVALL, A. (1997).

$\delta(B)$ – a non-stationary autoregressive (AR) polynomial in B (unit roots);

$\theta(B)$ – an invertible moving average (MA) polynomial in B and in B^S , which can be expressed in the multiplicative form

$$(1 + \vartheta_1 B + \dots + \vartheta_q B^q) (1 + \Theta_1 B^s + \dots + \Theta_Q B^{sQ}) ;$$

$\phi(B)$ – a stationary autoregressive (AR) polynomial in B and in B^S containing regular and seasonal unit roots, with s representing the number of observations per year;

a_t – a white-noise variable with the variance $V(a)$.

It should be noted that the stochastic time series can be predicted using its past observations and making an error. The variable a_t , which is assumed to be white noise, is the fundamental *innovation* to the series at time t , that is the part that cannot be predicted based on the past history of the series.

Denoting $\varphi(B) = \phi(B)\delta(B)$, [1] can be written in a more concise form as

$$\varphi(B) x_t = \theta(B) a_t$$

, [2]

where $\varphi(B)$ contains both the stationary and the nonstationary roots.

Derivation of the models for the components

Let us consider the additive decomposition model

$$x_t = \sum_{i=1}^k x_{it}$$

, [3]

where i refers to the orthogonal components: trend, seasonal, transitory or irregular. Apart from the irregular component, supposed to be a white noise, it is assumed that each component follows the ARIMA model which can be represented, using the notation of [2], as:

$$\varphi_i(B) x_{it} = \theta_i(B) a_{it}$$

, [4]

where $\varphi_i(B) = \phi_i(B)\delta_i(B)$, x_{it} is the i -th unobserved component, $\varphi_i(B)$ and $\theta_i(B)$ are finite polynomials of order p_i and q_i , respectively, and a_{it} , the disturbance associated with such component, is a white noise process with zero mean and constant variance $V(a_i)$ and a_{it}

and a_{jt} are not correlated for $i \neq j$ and for any t . These disturbances are functions of the innovations in the series and are called "pseudo-innovations" in the literature concerning the AMB decomposition as they refer to the components that are never observed ²⁴. In the JDemetra+ documentation the term "innovations" is used to refer to the "pseudo-innovations".

The following assumptions hold for [4]. For each i the polynomials $\phi_i(B)$, $\delta_i(B)$ and $\theta_i(B)$ are prime and of finite order. The roots of $\delta_i(B)$ lie on the unit circle; those of $\phi_i(B)$ lie outside, while all the roots of $\theta_i(B)$ are on or outside the unit circle. This means that nonstationary and noninvertible components are allowed. Since different roots of the AR polynomial induce peaks in the spectrum²⁵ of the series at different frequencies, and given that different components are associated with the spectral peaks for different frequencies, it is assumed that for $i \neq j$ the polynomials $\phi_i(B)$ and $\phi_j(B)$ do not share any common root (they are coprime). Finally, it is assumed that the polynomials $\theta_i(B)$, $i = 1, \dots, k$ share no unit root in common, guaranteeing the invertibility of the overall series. In fact, since the unit root of $\theta_i(B)$ induce a spectral zero, when the polynomials $\theta_i(B)$, $i = 1, \dots, k$ share no unit root in common, there is no frequency for which all component spectra become zero²⁶.

Since aggregation of ARIMA models yields ARIMA models, the series x_t will also follow an ARIMA model, as in [2], and consequently the following identity can be derived:

$$\frac{\theta(B)}{\varphi(B)} a_t = \sum_{i=1}^k \frac{\theta_i(B)}{\varphi_i(B)} a_{it}$$

. [5]

In the ARIMA model based approach implemented in SEATS, the ARIMA model identified and estimated for the observed series x_t is decomposed to derive the models for the components. In particular, the AR polynomials for the components, $\varphi_i(B)$, are easily derived through the factorization of the AR polynomial $\varphi(B)$:

$$\varphi(B) = \prod_{i=1}^k \varphi_i(B)$$

, [6]

while the MA polynomials for the components, together with the innovation variances $V(a_i)$, cannot simply be obtained through the relationship:

$$\theta(B) a_t = \sum_{i=1}^k \varphi_{ni}(B) \theta_i(B) a_{it}$$

²⁴GÓMEZ, V., and MARAVALL, A. (2001a).

²⁵For description of the spectrum see section [Spectral Analysis](#).

²⁶MARAVALL, A. (1995).

, [7]

where $\varphi_{ni}(B)$ is the product of all $\varphi_j(B)$, $j = 1, \dots, k$, except from $\varphi_i(B)$. Further assumptions are therefore needed to cope with the underidentification problem: i) $p_i \geq q_i$ and ii) the canonical decomposition, i.e. the decomposition that allocate all additive white noise to the irregular component (yielding noninvertible components except the irregular).

To understand how SEATS factorizes the AR polynomials, first a concept of a root will be explored²⁷.

The equation [2] can be expressed as:

$$\psi^{-1}(B)x_t = a_t(1 + \varphi_1 B + \dots \varphi_p B^p)x_t = (1 + \theta_1 B + \dots \theta_q B^q)a_t$$

, [8]

Let us now consider [2] in the inverted form:

$$\theta(B)y_t = \varphi(B)a_t$$

, [9]

If both sides of [8] are multiplied by x_{t-k} with $k > q$, and expectations are taken, the right hand side of the equation vanishes and the left hand side becomes:

$$\varphi(B)\gamma_k = \gamma_k + \varphi_1\gamma_{k-1} + \dots \varphi_p\gamma_{k-p} = 0$$

, [10]

where B operates on the subindex k .

The autocorrelation function γ_k is a solution of [10] with the characteristic equation:

$$z^p + \varphi_1 z^{p-1} + \dots \varphi_{p-1} z + \varphi_p = 0$$

. [11]

If z_1, \dots, z_p are the roots of [11], the solutions of [10] can be expressed as:

$$\gamma_k = \sum_{i=1}^p z_i^k, [12]$$

and will converge to zero as $k \rightarrow \infty$ when $|r_i| < 1$, $i = 1, \dots, p$. From [10] and [12] it can be noticed that $z_1 = B_1^{-1}$, meaning that z_1, \dots, z_p are the inverses of the roots B_1, \dots, B_p of the polynomial $\varphi(B)$. The convergence of γ_k implies that the roots of the $\varphi(B)$ are larger than 1 in modulus (lie outside the unit circle). Therefore, from the equation

²⁷Description based on KAISER, R., and MARAVALL, A. (2000) and MARAVALL, A. (2008c).

$$\varphi(B)^{-1} = \frac{1}{(1 - z_1) \dots (1 - z_1)}$$

[13]

it can be derived that $\varphi(B)^{-1}$ is convergent and all its inverse roots are less than 1 in modulus.

Equation [11] has real and complex roots (solutions). Complex number $x = a + bi$, with a and b both real numbers, can be represented as $x = r(\cos(\omega) + i \sin(\omega))$, where i is the imaginary unit ($i^2 = -1$), r is the modulus of x , that is $r = |x| = \sqrt{a^2 + b^2}$ and ω is the argument (frequency). When roots are complex, they are always in pairs of complex conjugates. The representation of the complex number $x = a + bi$ has a geometric interpretation in the complex plane established by the real axis and the orthogonal imaginary axis.




Figure 36: Text

Geometric representation of a complex number and of its conjugate

Representing the roots of the characteristic equation [11] in the complex plane enhances understanding how they are allocated to the components. When the modulus r of the roots in z are greater than 1 (i.e. modulus of the roots in $\varphi(B) < 1$), the solution of the characteristic equation has a systematic explosive process, which means that the impact of the given impulse on the time series is more and more pronounced in time. This behaviour is not in line with the

developments that can be identified in actual economic series. Therefore, the models estimated by TRAMO-SEATS (and X-13ARIMA-SEATS) have never inverse roots in B with modulus greater than 1.

The characteristic equations associated with the regular and the seasonal differences have roots in $\varphi(B)$ with modulus $r = 1$. They are called non-stationary roots and can be represented on the unit circle. Let us consider the seasonal differencing operator applied to a quarterly time series $(1 - B^4)$. Its characteristic equation is $(z^4 - 1) = 0$ with solutions given by $z = \sqrt[4]{1}$, i.e. $z_{1,2} = \pm 1$ and $z_{3,4} = \pm i$. The first two solutions are real and the last two are complex conjugates. They are represented by the black points on the unit circle on the figure below.




Figure 37: Text

Unit roots on the unit circle

For the seasonal differencing operator $(1 - B^{12})$ applied to the monthly time series the characteristic equation $(z^{12} - 1) = 0$ has twelve non-stationary solutions given by $z = \sqrt[12]{1}$: two real and ten complex conjugates, represented by the white circles in unit roots figure above.

$$z_t = A^t \cos(\omega t + W).$$

[14]

A – amplitude;

 W - phase (angle at $t = 0$).

For quarterly series the seasonal movements are produced by complex conjugates roots with angular frequencies at $\frac{\pi}{2}$ (one cycle per year) and π (two cycles per year). The corresponding number of cycles per year and the length of the movements are presented in the table below.

	Angular frequency (ω)	Frequency (cycles per unit time) (f)	Cycles per year	Length of the movement measured in quarters (τ)
$\frac{\pi}{2}$	0.25	1	4	π
0.5	2	2		

Seasonal frequencies for a monthly time series

Angular frequency (ω)										Frequency (cycles per unit time) (f)									
Cycles per year										Length of the movement measured in months (τ)									
4 $\frac{2\pi}{3}$ 0.333 4 3 $\frac{5\pi}{6}$ 0.417 5 2.4										$\frac{\pi}{6}$ 0.083 1 12 $\frac{\pi}{3}$ 0.167 2 6 $\frac{\pi}{2}$ 0.250 3									

0.500	6	2
-------	---	---

²⁸For details see MARAVALL, A., CAPORELLO, G., PÉREZ, D., and LÓPEZ, R. (2014).

- Roots of $(1 - B)^d$ are assigned to trend component.
- Roots of $(1 - B^s)^{d_s} = ((1 - B)(1 + B + \dots + B^{s-1}))^{d_s}$ are assigned to the trend component (root of $(1 - B)^{d_s}$) and to the seasonal component (roots of $(1 + B + \dots + B^{s-1})^{d_s}$).
- When the modulus of the inverse of a real positive root of $\varphi(B)$ is greater than k or equal to k , where k is the threshold value controlled by the *Trend boundary* parameter (in the original SEATS it is controlled by *rmod*)²⁹, then the root is assigned to the trend component. Otherwise it is assigned to the transitory component.
- Real negative inverse roots of $\varphi_p(B)$ associated with the seasonal two-period cycle are assigned to the seasonal component if their modulus is greater than k , where k is the threshold value controlled by the *Seasonal boundary* and the *Seas. boundary (unique)* parameters. Otherwise they are assigned to the transitory component.
- Complex roots, for which the argument (angular frequency) is close enough to the seasonal frequency are assigned to the seasonal component. Closeness is controlled by the *Seasonal tolerance* and *Seasonal tolerance (unique)* parameters (in the original SEATS it is controlled by *epsphi*). Otherwise they are assigned to the transitory component.
- If d_s (seasonal differencing order) is present and $Bphi < 0$ ($Bphi$ is the estimate of the seasonal autoregressive parameter), the real positive inverse root is assigned to the trend component and the other $(s - 1)$ inverse roots are assigned to the seasonal component. When $d_s = 0$, the root is assigned to the seasonal when $Bphi < -0.2$ and/or the overall test for seasonality indicates presence of seasonality. Otherwise it goes to the transitory component. Also, when $Bphi > 0$, roots are assigned to the transitory component.

For further details about JDemetra+ parameters see section [TramoSeats](#).

It should be highlighted that when $Q > P$, where Q and P denote the orders of the polynomials $\varphi(B)$ and $\theta(B)$, the SEATS decomposition yields a pure MA $(Q - P)$ component (hence transitory). In this case the transitory component will appear even when there is no AR factor allocated to it.

Once these rules are applied, the factorization of the AR polynomial presented by [2] yields to the identification of the AR polynomials for the components which contain, respectively, the AR roots associated with the trend component, the seasonal component and the transitory component.³⁰

Then with the partial fraction expansion the spectrum of the final components are obtained.

²⁹In JDemetra+ this argument is called *Trend boundary*.

³⁰The AR roots close to or at the trading day frequency generates a stochastic trading day component. A stochastic trading day component is always modelled as a stationary ARMA(2,2), where the AR part contains the roots close to the TD frequency, and the MA(2) is obtained from the model decomposition (MARAVALL, A., and PÉREZ, D. (2011)). This component, estimated by SEATS, is not implemented by the current version of JDemetra+.

For example, the Airline model for a monthly time series:

$$(1 - B)(1 - B^{12})x_t = (1 + \theta_1 B)(1 + \Theta_1 B^{12}) a_t$$

, [15]

is decomposed by SEATS into the model for the trend component:

$$(1 - B)(1 - B)c_t = (1 + \theta_{c,1}B + \theta_{c,2}B^2)a_{c,t}$$

, [16]

and the model for the seasonal component:

$$(1 + B + \dots + B^{11})s_t = (1 + \theta_{s,1}B + \dots + \theta_{s,11}B^{11})a_{s,t},$$

[17]

As a result, the Airline model is decomposed as follows:

$$\frac{(1 + \theta_1 B)(1 + \Theta_1 B^{12})}{(1 - B)(1 - B)}a_t = \frac{(1 + \theta_{s,1}B + \dots + \theta_{s,11}B^{11})}{(1 + B + \dots + B^{11})}a_{s,t} + \frac{(1 + \theta_{c,1}B + \theta_{c,2}B^2)}{(1 - B)(1 - B)}a_{c,t} + u_t$$

. [18]

The transitory component is not present in this case and the irregular component is the white noise.

The partial fractions decomposition is performed in a frequency domain. In essence, it consists in portioning of the pseudo-spectrum³¹ of x_t into additive spectra of the components. When the AMB decomposition of the ARIMA model results in the non-negative spectra for all components, the decomposition is called admissible³². In such case an infinite number of admissible decompositions exists, i.e. decompositions that yield the non-negative spectra of all components. Therefore, the MA polynomials and the innovation variances cannot be yet identified from the model of x_t . As sketched above, to solve this underidentification problem and identify a unique decomposition, it is assumed that for each component the order of the MA polynomial is no greater than the order of the AR polynomial and the canonical solution

³¹The term pseudo-spectrum is used for a non-stationary time series, while the term spectrum is used for a stationary time series.

³²If the ARIMA model estimated in TRAMO does not accept an admissible decomposition, SEATS replaces it with a decomposable approximation. The modified model is therefore used to decompose the series. There are also other rare situations when the ARIMA model chosen by TRAMO is changed by SEATS. It happens when, for example, the ARIMA models generate unstable seasonality or produce a senseless decomposition. Such examples are discussed by MARAVALL, A. (2009).

of S.C. Hillmer and G.C. Tiao is applied³³, i.e. all additive white noise is added to the irregular component. As a consequence all components derived from the canonical decomposition, except from the irregular, have a spectral minimum of zero and are thus noninvertible³⁴. Given the stochastic features of the series, it can be shown by that the canonical decomposition produces as stable as possible trend and seasonal components since it maximizes the variance of the irregular and minimizes the variance of the other components³⁵. However, there is a price to be paid as canonical components can produce larger revisions in the preliminary estimators of the component³⁶ than any other admissible decomposition.

The figure below represents the pseudo-spectrum for the canonical trend and an admissible trend.




Figure 38: Text

A comparison of canonical trend and admissible trend

A pseudo-spectrum is denoted by $g_i(\omega)$, where ω represents the angular frequency. The pseudo-spectrum of x_{it} is defined as the Fourier transform of ACGF of x_t which is expressed as:

$$\frac{\psi_i(B) \psi_i(F)}{\delta_i(B) \delta_i(F)} V(a_i)$$

³³HILLMER, S.C., and TIAO, G.C. (1982).

³⁴GÓMEZ, V., and MARAVALL, A. (2001a).

³⁵HILLMER, S.C., and TIAO, G.C. (1982).

³⁶MARAVALL, A. (1986).

, [19]

where:

$$\psi_i(F) = \frac{\theta_i(F)}{\phi_i(F)}$$

$$\psi_i(B) = \frac{\theta_i(B)}{\phi_i(B)}$$

B is the backward operator,

F is the forward operator.

A pseudo-spectrum for a monthly time series x_t is presented in the figure below: The pseudo-spectrum for a monthly series. The frequency $\omega = 0$ is associated with the trend, frequencies in the range $[0 + \epsilon_1, \frac{\pi}{6} - \epsilon_2]$ with $[0 + \epsilon_1, \frac{\pi}{6} - \epsilon_2]$ $\epsilon_1, \epsilon_2 > 0$ and $\epsilon_1 < \frac{\pi}{6} - \epsilon_2$ are usually associated with the business-cycle and correspond to a period longer than a year and bounded³⁷. The frequencies in the range $[\frac{\pi}{6}, \pi]$ are associated with the short term movements, whose cycle is completed in less than a year. If a series contains an important periodic component, its spectrum reveals a peak around the corresponding frequency and in the ARIMA model it is captured by an AR root. In the example below spectral peaks occur at the frequency $\omega = 0$ and at the seasonal frequencies $(\frac{\pi}{6}, \frac{2\pi}{6}, \frac{3\pi}{6}, \frac{4\pi}{6}, \frac{5\pi}{6}, \pi)$.³⁸




Figure 39: Text

The pseudo-spectrum for a monthly series

³⁷Ibid.

³⁸KAISER, R., and MARAVALL, A. (2000).

In the decomposition procedure, the pseudo-spectrum of the time series x_t is divided into the spectra of its components (in the example figure below, four components were obtained).




Figure 40: Text

The pseudo-spectra for the components

Estimation of the components with the Wiener-Kolmogorow filter

The various components are estimated using Wiener-Kolmogorow (WK) filters. JDemetra+ includes three options to estimate the WK filter, namely *Burman*, *KalmanSmoother* and *MCElroyMatrix*³⁹. Here the first of abovementioned options, proposed by BURMAN, J.P. (1980) will be explained.

The estimation procedure and the properties of the WK filter are easier to explain with a two-component model. Let the seasonally adjusted series (s_t) be the signal of interest and the seasonal component (n_t) be the remainder, "the noise". The series is given by the model [2] and from [4] the models for theoretical components are:

$$\varphi_s(B)s_t = \theta_s(B)a_{st}$$

³⁹The choice of the estimation method is controlled by the *Method* parameter, explained in the [SEATS specification](#) section.

[20]

and

$$\varphi_n(B)n_t = \theta_n(B)a_{nt}$$

. [21]

From [6] and [7] it is clear that $\varphi(B) = \varphi_s(B)\varphi_n(B)$ and $\theta(B)a_t = \theta_s(B)a_{st} + \theta_n(B)a_{nt}$.

As the time series components are never observed, their estimators have to be used. Let us note X_T an infinite realization of the time series x_t . SEATS computes the Minimum Mean Square Error (MMSE) estimator of s_t , e.g. the estimator

$$\hat{s}_t$$

that minimizes

$$E[(s_t - \hat{s}_t)^2 | X_T]$$

. Under the normality assumption

$$\hat{s}_{t|T}$$

is also equal to the conditional expectation

$$E(s_t | X_T)$$

, so it can be presented as a linear function of the elements in

$$X_T$$

.⁴⁰ WHITTLE (1963) shows that the MMSE estimator of

$$\hat{s}_t$$

is:

$$\hat{s}_t = k_s \frac{\psi_s(B)\psi_s(F)}{\psi(B)\psi(F)} x_t$$

, [22]

where

$$\psi(B) = \frac{\theta(B)}{\phi(B)}$$

,

$$F = B^{-1}$$

⁴⁰MARAVALL, A. (2008c).

and

$$k_s = \frac{V(a_s)}{V(a)}$$

,

$$V(a_s)$$

is the variance of

$$a_{st}$$

and

$$V(a)$$

is the variance of

$$a_t$$

.

Expressing the

$$\psi(B)$$

polynomials as functions of the AR and MA polynomials, after cancelation of roots, the estimator of

$$s_t$$

can be expressed as:

$$\hat{s}_t = k_s \frac{\theta_s(B) \theta_s(F) \varphi_n(B) \delta_n(B) \varphi_n(F) \delta_n(F)}{\theta(B) \theta(F)} x_t$$

, [23]

where:

$$\nu_s(B, F) = k_s \frac{\theta_s(B) \theta_s(F) \varphi_n(B) \delta_n(B) \varphi_n(F) \delta_n(F)}{\theta(B) \theta(F)}$$

[24]

is a WK filter.

Equation [24] shows that the WK filter is two-sided (uses observations both from the past and from the future), centered (the number of points in the past is the same as in the future) and symmetric (for any k the weight applied to x_{t-k} and x_{t+k} is the same), which allows the phase effect to be avoided. Due to invertibility of $\theta(B)$ (and $\theta(F)$) the filter is convergent in the past and in the future.

The estimator can be presented as

$$\hat{s}_t = \nu_i(B, F) x_t$$

, [25]

where

$$\nu_i(B, F) = \nu_0 + \sum_{j=1}^{\infty} \nu_{ij}(B^j + F^j)$$

is the WK filter.

The example of the WK filters obtained for the pseudo-spectra of the series illustrated above is shown on the figure below: WK filters for components.




Figure 41: Text

WK filters for components

The WK filter from [24] can also be expressed as a ratio of two pseudo-autocovariance generating functions (p-ACGF). The p-ACGF function summarizes the sequence of absolutely summable autocovariances of a stationary process x_t (see section [Spectral Analysis](#)).

The ACGF function of an ARIMA process is expressed as:

$$acgf(B) = \frac{\theta(B) \theta(F)}{\phi(B) \delta(B) \phi(F) \delta(F)} V(a)$$

[26]

And, the WK filter can be rewritten as:

$$\nu_s(B, F) = \frac{\gamma_s(B, F)}{\gamma(B, F)}$$

, [27]

where:

$$\gamma_s(B, F) = \frac{\theta_s(B) \theta_s(F)}{\phi_s(B) \delta_s(B) \phi_s(F) \delta_s(F)} V(a_s)$$

is the p-ACGF of

$$s_t$$

;

$\gamma(B, F) = \frac{\theta(B)\theta(F)}{\phi(B)\delta(B)\phi(F)\delta(F)} V(a)$ is the p-ACGF of x_t .

From [24] it can be seen that the WK filter depends on both the component and the series models. Consequently, the estimator of the component and the WK filter reflect the characteristic of data and by construction, the WK filter adapts itself to the series under consideration. Therefore, the ARIMA model is of particular importance for the SEATS method. Its misspecification results in an incorrect decomposition.

This adaptability, if the model has been correctly determined, avoids the dangers of under and overestimation with an ad-hoc filtering. For example, for the series with a highly stochastic seasonal component the filter adapts to the width of the seasonal peaks and the seasonally adjusted series does not display any spurious seasonality⁴¹. Examples of WK filters for stochastic and stable seasonal components are presented on the figure below.

WK filters for stable and stochastic seasonal components

The derivation of the components requires an infinite realization of x_t in the direction of the past and of the future. However, the convergence of the WK filter guarantees that, in practice, it could be approximated by a truncated (finite) filter and, in most applications, for large

$$k$$

the estimator for the central periods of the series can be safely seen as generated by the WK filter⁴²:

⁴¹MARAVALL, A. (1995).

⁴²MARAVALL, A., and PLANAS, C. (1999).




Figure 42: Text

$$\hat{s}_t = \nu_k x_{t-k} + \dots + \nu_0 x_t + \dots + \nu_k x_{t+k}$$

. [28]

When $T > 2L + 1$, where T is the last observed period, and L is an a priori number that typically expands between 3 and 5 years, the estimator expressed by [23] can be assumed as the final (historical) estimator for the central observations of the series⁴³. In practice, the Wiener-Kolmogorov filter is applied to x_t extended with forecasts and backcasts from the ARIMA model. The final or historical estimator of

$$\hat{s}_t$$

, is obtained with a doubly infinite filter, and therefore contains an error

$$e_{st}$$

called final estimation error, which is equal

$$e_{st} = s_t - \hat{s}_t$$

.

⁴³MARAVALL, A. (1998).

In the frequency domain, the Wiener-Kolmogorov filter $\nu(B, F)$ that provides the final estimator of s_t is expressed as the ratio of the s_t and x_t pseudo-spectra:

$$\tilde{\nu}(\omega) = \frac{g_s(\omega)}{g_x(\omega)}$$

. [29]

The function $\tilde{\nu}(\omega)$ is also referred as the gain of the filter.⁴⁴ GÓMEZ, V., and MARAVALL, A. (2001a) show that when for some frequency the signal (the seasonally adjusted series) dominates the noise (seasonal fluctuations) the gain $\tilde{\nu}(\omega)$ approaches 1. On the contrary, when for some frequency the noise dominates the gain $\tilde{\nu}(\omega)$ approaches 0.

The spectrum of the estimator of the seasonal component is expressed as:

$$g_{\hat{s}}(\omega) = \left[\frac{g_s(\omega)}{g_x(\omega)} \right]^2 g_x(\omega)$$

, [30]

where $[\tilde{\nu}(\omega)]^2 = \left[\frac{g_s(\omega)}{g_x(\omega)} \right]^2 = \left[\frac{g_s(\omega)}{g_s(\omega) + g_n(\omega)} \right]^2 = \left[\frac{1}{1 + \frac{1}{r(\omega)}} \right]^2$ is the squared gain of the filter and $r(\omega) = \frac{g_s(\omega)}{g_n(\omega)}$ represents the signal-to-noise ratio.

For each ω , the MMSE estimation gives the signal-to-noise ratio. If this ratio is high, then the contribution of that frequency to the estimation of the signal will be also high. Assume that the trend is a signal that needs to be extracted from a seasonal time series. Then $R(0) = 1$ and the frequency $\omega = 0$ will only be used for trend estimations. For seasonal frequencies $R(\omega) = 0$, so that these frequencies are ignored in computing the trend resulting in spectral zeros in $g_{\hat{s}}(\omega)$. For this reason, unlike the spectrum of the component, the component spectrum contains dips as it can be seen on the figure below: Component spectrum and estimator spectrum for trend.

Component spectrum and estimator spectrum for trend

From the equation [29] it is clear that the squared gain of the filter determines how the variance of the series contributes to the variance of the seasonal component for the different frequencies. When $\tilde{\nu}(\omega) = 1$, the full variation of x_t for that frequency is passed to

$$\hat{s}_t$$

, while if

$$\tilde{\nu}(\omega) = 0$$

⁴⁴GÓMEZ, V., and MARAVALL, A. (2001a).




Figure 43: Text

the variation of x_t for that frequency is fully ignored in the computation of

$$\hat{s}_t$$

. These two cases are well illustrated by the figure below that shows the square gain of the WK filter for two series already analysed in the figure above (Figure: WK filters for stable and stochastic seasonal components).




Figure 44: Text

The squared gain of the WK filter for stable and stochastic seasonal components.

Since $r(\omega) \geq 0$, then $\tilde{\nu}(\omega) \leq 1$ and from [29] it can be derived that $g_{\hat{s}}(\omega) = \tilde{\nu}(\omega) g_s(\omega)$. As a result, the estimator will always underestimate the component, i.e. it will be always more stable than the component.⁴⁵

Since $g_{\hat{n}}(\omega) < g_n(\omega)$ and $g_{\hat{s}}(\omega) < g_s(\omega)$ the expression: $g_x(\omega) - [g_{\hat{n}}(\omega) + g_{\hat{s}}(\omega)] \geq 0$ is the cross-spectrum. As it is positive, the MMSE yields correlated estimators. This effect emerges

⁴⁵Ibid.

since variance of estimator is smaller than the variance of component. Nevertheless, if at least one non-stationary component exists, cross-correlations estimated by TRAMO-SEATS will tend to zero as cross-covariances between estimators of the components are finite. In practice, the inconvenience caused by this property will likely be of little relevance.

Preliminary estimators for the components

GÓMEZ, V., and MARAVALL, A. (2001a) point out that *the properties of the estimators have been derived for the final (or historical) estimators. For a finite (long enough) realization, they can be assumed to characterize the estimators for the central observations of the series, but for periods close to the beginning of the end the filter cannot be completed and some preliminary estimator has to be used.* Indeed, the historical estimator shown in [28] is obtained for the central periods of the series. However, when t approaches T (last observation), the WK filter requires observations, which are not available yet. For this reason a preliminary estimator needs to be used.

To introduce preliminary estimators let us consider a semi-finite realization $[x_{-\infty}, \dots, x_T]$, where T is the last observed period. The preliminary estimator of

$$x_{it}$$

obtained at T

$$(T - t = k \geq 0)$$

can be expressed as

$$\hat{x}_{it|t+k} = \nu_i(B, F) x_{t|T}^e$$

, [31]

where

$$\nu_i(B, F)$$

is the WK filter and

$$x_{t|T}^e$$

is the extended series, such that $x_{t|T}^e = x_t$ for $t \leq T$ and

$$x_{t|T}^e = \hat{x}_{t|T}$$

for

$$t > T$$

, where

$$\hat{x}_{t|T}$$

denotes the forecast of x_t obtained at period T .

The future k values necessary to apply the filter are not yet available and are replaced by their optimal forecasts from the ARIMA model on

$$x_t$$

. When

$$k = 0$$

the preliminary estimator becomes the concurrent estimator. As the forecasts are linear functions of present and past observations of

$$x_t$$

, the preliminary estimator

$$\hat{x}_{it}$$

will be a truncated asymmetric filter applied to

$$x_t$$

that generates a phase effect⁴⁶.

When a new observation

$$x_{T+1}$$

becomes available the forecast

$$\hat{x}_{T+1|T}$$

is replaced by the observation and the forecast

$$\hat{x}_{iT+j|T}$$

,

$$j > 1$$

are updated to

$$x_{T+j|T+1}$$

resulting in the revision error⁴⁷. The total error in the preliminary estimator

$$d_{it|t+k}$$

is expressed as a sum of the final estimation error (

$$e_{it}$$

) and the revision error (

$$r_{it|t+k}$$

⁴⁶KAISER, R., and MARAVALL, A. (2000).

⁴⁷MARAVALL, A. (1995).

), i.e.:

$$d_{it|t+k} = x_{it} - \hat{x}_{it|t+k} = (x_{it} - \hat{x}_{it}) + (\hat{x}_{it} - \hat{x}_{it|t+k}) = e_{it} + r_{it|t+k}$$

, [32]

where:

$$x_{it} - i^{th}$$

component;

$$\hat{x}_{it|t+k}$$

- the estimator of

$$x_{it}$$

when the last observation is

$$x_{t+k}$$

.

Therefore the preliminary estimator is subject not only to the final error but also to a revision error, which are orthogonal to each other⁴⁸. The revision error decreases as

$$k$$

increases, until it can be assumed equal to 0 for large enough

$$k$$

.

It's worth remembering that SEATS estimates the unobservable components of the time series so the "true" components are never observed. Therefore, MARAVALL, A. (2009) stresses that *the error in the historical estimator is more of academic rather than practical interest. In practice, interest centres on revisions. (...) the revision standard deviation will be an indicator of how far we can expect to be from the optimal estimator that will be eventually attained, and the speed of convergence of $\theta(B)^{-1}$ will dictate the speed of convergence of the preliminary estimator to the historical one.* The analysis of an error is therefore useful for making decision concerning the revision policy, including the policy for revisions and horizon of revisions.

⁴⁸MARAVALL, A. (2009).

PsiE-weights

The estimator of the component is calculated as

$$\hat{x}_{it} = \nu_s(B, F) x_t$$

. By replacing

$$x_{it} = \frac{\theta(B)}{\gamma(B)\delta(B)} a_t$$

, the component estimator can be expressed as⁴⁹:

$$\hat{x}_{it} = \xi_s(B, F) a_t$$

, [33]

where $\xi_s(B, F) = \dots + \xi_j B^j + \dots + \xi_1 B + \xi_0 + \xi_{-1} F \dots \xi_{-j} F^j + \dots$

This representation shows the estimator as a filter applied to the innovation

$$a_t$$

, rather than on the series

$$x_t$$

⁵⁰. Hence, the filter from [32] can be divided into two components: the first one, i.e.

$$\dots + \xi_j B^j + \dots + \xi_1 B + \xi_0$$

, applies to prior and concurrent innovations, the second one, i.e.

$$\xi_{-1} F + \dots + \xi_{-j} F^j$$

applies to future (i.e. posterior to

$$t$$

) innovations. Consequently,

$$\xi_j$$

determines the contribution of

$$a_{t-j}$$

to

$$\hat{s}_t$$

while

$$\xi_{-j}$$

⁴⁹The section is based on KAISER, R., and MARAVALL, A. (2000).

⁵⁰See section PsiE-weights. For further details see MARAVALL, A. (2008).

determines the contribution of

$$a_{t+j}$$

to

$$\hat{s}_t$$

. Finally, the estimator of the component can be expressed as:

$$\hat{x}_{it} = \xi_i(B)^- a_t + \xi_i(F)^+ a_{t+1}$$

, [34]

where:

$\xi_i(B)^- a_t$ is an effect of starting conditions, present and past innovations in series;

$\xi_i(F)^+ a_{t+1}$ is an effect of future innovations.

For the two cases already presented in figure *WK filters for stable and stochastic seasonal components* and figure *The squared gain of the WK filter for stable and stochastic seasonal components* above, the psi-weights are shown in the figure below.




Figure 45: Text

It can be shown that

$$\xi_{-1}, \dots, \xi_{-j}$$

are convergent and

$$\xi_j, \dots, \xi_1, \xi_0$$

are divergent. From [33] , the concurrent estimator is equal to

$$\hat{x}_{it|t} = E_t x_{it} = E_t \hat{x}_{it} = \xi_i(B)^- a_t$$

, [35]

so that the revision

$$r_{it} = \hat{x}_{it} - \hat{x}_{it|t} = \xi_i(F)^+ a_{t+1}$$

[36]

is a zero-mean stationary MA process. As a result, historical and preliminary estimators are cointegrated. From expression [25] the relative size of the full revision and the speed of convergence can be obtained.

Local Polynomials Methods for Trend Estimation

Chapter building process

This chapter provides details on the the statistical methods whereas practical estimation steps are described “here” (Chap On trend estimation) and are run with the `rjfilters` or `rjd3highfreq` packages.

Tests

Introduction

This chapter describes all the tests available in JDemetra+, via Graphical User interface and/or R packages. An outline of the underlying theoretical principles of each test is provided.

The procedure to apply these tests in context is described when their use is relevant in the chapters dedicated to algorithms description, mainly on seasonal adjustment.(not sure) Links to use in context are available below.

Tests on residuals

table with all tests by purpose and accessibility

Table 27: Tests on Residuals

Test	Purpose	GUI	R package
Ljung-Box	autocorrelation		
Box-Pierce	autocorrelation		
Doornik-Hansen	normality		

Ljung-Box

The Ljung-Box Q-statistics are given by:

$$LB(k) = n \times (n + 2) \times \sum_{k=1}^K \frac{\rho_{a,k}^2}{n - k}$$

, [1]

where $\rho_{a,k}^2$ is the autocorrelation coefficient at lag k of the residuals \hat{a}_t , n is the number of terms in differenced (? differenciated ?) series, K is the maximum lag being considered, set in JDemetra+ to 24 (monthly series) or 8 (quarterly series).

If the residuals are random (which is the case for residuals from a well specified model), they will be distributed as $\chi^2_{(K-m)}$, where m is the number of parameters in the model which has been fitted to the data. (edit: not the residuals, but $\hat{\rho}$)

The Ljung-Box and Box-Pierce tests sometimes fail to reject a poorly fitting model. Therefore, care should be taken not to accept a model on a basis of their results. For the description of autocorrelation concept see section [Autocorrelation function and partial autocorrelation function](#).

Box-Pierce

The Box-Pierce Q-statistics are given by:

$$BP(k) = n \sum_{k=1}^K \rho_{a,k}^2$$

where:

$\rho_{a,k}^2$ is the autocorrelation coefficient at lag k of the residuals \hat{a}_t .

n is the number of terms in differenced (differenciaded?) series;

K is the maximum lag being considered, set in JDemetra+ to 24 (monthly series) or 8 (quarterly series).

If the residuals are random (which is the case for residuals from a well specified model), they will be distributed as $\chi^2_{(K-m)}$ degrees of freedom, where m is the number of parameters in the model which has been fitted to the data.(edit: same as above)

Dornik-Hansen

The Doornik-Hansen test for multivariate normality (DOORNIK, J.A., and HANSEN, H. (2008)) is based on the skewness and kurtosis of multivariate data that is transformed to ensure independence. It is more powerful than the Shapiro-Wilk test for most tested multivariate distributions⁵¹.

The skewness and kurtosis are defined, respectively, as:

$$s = \frac{m_3}{\sqrt{m_2}^3}$$

and $k = \frac{m_4}{m_2^2} - 3$, where: $m_i = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^i$, $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ and n is a number of (non-missing) residuals.

⁵¹The description of the test derives from DOORNIK, J.A., and HANSEN, H. (2008).

The Doornik-Hansen test statistic derives from SHENTON, L.R., and BOWMAN, K.O. (1977) and uses transformed versions of skewness and kurtosis.

The transformation for the skewness s into z_1 is as in D'AGOSTINO, R.B. (1970):

$$\beta = \frac{3(n^2 + 27n - 70)(n + 1)(n + 3)}{(n - 2)(n + 5)(n + 7)(n + 9)}$$

$$\omega^2 = -1 + \sqrt{2(\beta - 1)}$$

$$\delta = \frac{1}{\sqrt{\log(\omega^2)}}$$

$$y = s \sqrt{\frac{(\omega^2 - 1)(n + 1)(n + 3)}{12(n - 2)}}$$

$$z_1 = \delta \log(y + \sqrt{y^2 - 1})$$

The kurtosis k is transformed from a gamma distribution to χ^2 , which is then transformed into standard normal z_2 using the Wilson-Hilferty cubed root transformation:

$$\delta = (n - 3)(n + 1)(n^2 + 15n - 4)$$

$$a = \frac{(n - 2)(n + 5)(n + 7)(n^2 + 27n - 70)}{6\delta}$$

$$c = \frac{(n - 7)(n + 5)(n + 7)(n^2 + 2n - 5)}{6\delta}$$

$$l = \frac{(n + 5)(n + 7)(n^3 + 37n^2 + 11n - 313)}{12\delta}$$

$$\alpha = a + c \times s^2$$

$$\chi = 2l(k - 1 - s^2)$$

$$z_2 = \sqrt{9\alpha} \left(\frac{1}{9\alpha} - 1 + \sqrt[3]{\frac{\chi}{2\alpha}} \right)$$

Finally, the Doornik-Hansen test statistic is defined as the sum of squared transformations of the skewness and kurtosis. Approximately, the test statistic follows a χ^2 distribution, i.e.:

$$DH = z_1^2 + z_2^2 \sim \chi^2(2)$$

Seasonality tests

table with all tests by purpose and accessibility

Table 28: Seasonality tests

Test	Purpose	GUI	R package
QS test	Autocorrelation at seasonal lags		
F-test with seasonal dummies	Stable seasonality		
Identification of spectral peaks	Seasonal frequencies		
Friedman test	Stable seasonality		
Two-way variance analysis	Moving seasonality		

QS Test on autocorrelation at seasonal lags

The QS test is a variant of the [Ljung-Box](#) test computed on seasonal lags, where we only consider positive auto-correlations

More exactly,

$$QS = n(n+2) \sum_{i=1}^k \frac{[\max(0, \hat{\gamma}_{i \cdot l})]^2}{n - i \cdot l}$$

where

$$k = 2$$

, so only the first and second seasonal lags are considered. Thus, the test would check the correlation between the actual observation and the observations lagged by one and two years. Note that

$$l = 12$$

when dealing with monthly observations, so we consider the autocovariances

$$\hat{\gamma}_{12}$$

and

$$\hat{\gamma}_{24}$$

alone. In turn,

$$k = 4$$

in the case of quarterly data.

Under H_0 , which states that the data are independently distributed, the statistic follows a

$$\chi(k)$$

distribution. However, the elimination of negative correlations makes it a bad approximation. The p-values would be given by $P(\chi^2(k) > Q)$ for $k = 2$. As $P(\chi^2(2)) > 0.05 = 5.99146$ and $P(\chi^2(2)) > 0.01 = 9.21034$, $QS > 5.99146$ and $QS > 9.21034$ would suggest rejecting the null hypothesis at 95% and 99% significance levels, respectively.

Modification

Maravall (2012) proposes approximating the correct distribution (p-values) of the QS statistic using simulation techniques. Using 1000K replications of sample size 240, the correct critical values would be 3.83 and 7.09 with confidence levels of 95% and 99%, respectively (lower than the 5.99146 and 9.21034 shown above). For each of the simulated series, he obtains the distribution by assuming $QS = 0$ when

$$\hat{\gamma}_{12}$$

, so in practice this test will detect seasonality only when any of these conditions hold: - Statistically significant positive autocorrelation at lag 12 - Nonnegative sample autocorrelation at lag 12 and statistically significant positive autocorrelation at lag 24

Use

The test can be applied directly to any series by selecting the option *Statistical Methods » Seasonal Adjustment » Tools » Seasonality Tests*. This is an example of how results are displayed for the case of a monthly series:

The test can be applied to the input series before any seasonal adjustment method has been applied. It can also be applied to the seasonally adjusted series or to the irregular component.

1. Tests on autocorrelations at seasonal lags

Seasonality present

ac(12)=0.8238

ac(24)=0.7006

Distribution: Chi2 with 2 degrees of freedom

Value: 258.5028

PValue: 0.0000

Figure 46: qs

References

- LJUNG G. M. and G. E. P. BOX (1978). “On a Measure of a Lack of Fit in Time Series Models”. *Biometrika* 65 (2): 297–303. doi:10.1093/biomet/65.2.297
- MARAVALL, A. (2011). “Seasonality Tests and Automatic Model Identification in Tramo-Seats”. Manuscript
- MARAVALL, A. (2012). “Update of Seasonality Tests and Automatic Model Identification in TRAMO-SEATS”. Bank of Spain (November 2012)

F-test on seasonal dummies

The F-test on seasonal dummies checks for the presence of deterministic seasonality. The model used here uses seasonal dummies (mean effect and 11 seasonal dummies for monthly data, mean effect and 3 for quarterly data) to describe the (possibly transformed) time series behaviour. The test statistic checks if the seasonal dummies are jointly statistically not significant. When this hypothesis is rejected, it is assumed that the deterministic seasonality is present and the test results are displayed in green.

This test refers to Model-Based χ^2 and F-tests for Fixed Seasonal Effects proposed by LYTRAS, D.P., FELDPASCH, R.M., and BELL, W.R. (2007) that is based on the estimates of the regression dummy variables and the corresponding t-statistics of the RegARIMA model, in which the ARIMA part of the model has a form (0,1,1)(0,0,0). The consequences of a misspecification of a model are discussed in LYTRAS, D.P., FELDPASCH, R.M., and BELL, W.R. (2007).

For a monthly time series the RegARIMA model structure is as follows:

$$(1 - B) (y_t - \beta_1 M_{1,t} - \dots - \beta_{11} M_{11,t} - \gamma X_t) = \mu + (1 - B)a_t$$

, [1]

where:

$$M_{j,t} = \begin{cases} 1 & \text{in month } j = 1, \dots, 11 \\ -1 & \text{in December} \\ 0 & \text{otherwise} \end{cases} \quad \text{- dummy variables;}$$

y_t – the original time series;

B – a backshift operator;

X_t – other regression variables used in the model (e.g. outliers, calendar effects, user-defined regression variables, intervention variables);

μ – a mean effect;

a_t – a white-noise variable with mean zero and a constant variance.

In the case of a quarterly series the estimated model has a form:

$$(1 - B)(y_t - \beta_1 M_{1,t} - \dots - \beta_3 M_{3,t} - \gamma X_t) = \mu + (1 - B)a_t$$

, [2]

where:

$$M_{j,t} = \begin{cases} 1 & \text{in quarter } j = 1, \dots, 3 \\ -1 & \text{in the fourth quarter} \\ 0 & \text{otherwise} \end{cases} \quad \text{- dummy variables;}$$

One can use the individual t-statistics to assess whether seasonality for a given month is significant, or a chi-squared test statistic if the null hypothesis is that the parameters are collectively all zero. The chi-squared test statistic is $\hat{\chi}^2 = \hat{\beta}' [Var(\hat{\beta})]^{-1} \hat{\beta}$ in this case compared to critical values from a $\chi^2(df)$ -distribution, with degrees of freedom $df = 11 \setminus$ (monthly series) or $df = 3$ (quarterly series). Since the $Var(\hat{\beta})$ computed using the estimated variance of α_t may be very different from the actual variance in small samples, this test is corrected using the proposed F statistic:

$$F = \frac{\hat{\chi}^2}{s-1} \times \frac{n-d-k}{n-d}$$

where n is the sample size, d is the degree of differencing, s is time series frequency (12 for a monthly series, 4 for a quarterly series) and k is the total number of regressors in the RegARIMA model (including the seasonal dummies $M_{j,t}$ and the intercept).

This statistic follows a

$$F_{s-1, n-d-k}$$

distribution under the null hypothesis.

Identification of spectral peaks

link to relevant part in spectral analysis chapter ?

Friedman test for stable seasonality test

The Friedman test is a non-parametric method for testing that samples are drawn from the same population or from populations with equal medians. The significance of the month (or quarter) effect is tested. The Friedman test requires no distributional assumptions. It uses the rankings of the observations. If the null hypothesis of no stable seasonality is rejected at the 0.10% significance level then the series is considered to be seasonal and the test's outcome is displayed in green.

The test statistic is constructed as follows. Consider first the matrix of data

$$\{x_{ij}\}_{n \times k}$$

with

$$n$$

rows (the blocks, i.e. number of years in the sample),

$$k$$

columns (the treatments, i.e. either 12 months or 4 quarters, depending on the frequency of the data).

The data matrix needs to be replaced by a new matrix

$$\{r_{ij}\}_{n \times k}$$

, where the entry

$$r_{ij}$$

is the rank of

$$x_{ij}$$

within block

$$i$$

.

The test statistic is given by

$$Q = \frac{SS_t}{SS_e}$$

where

$$SS_t = n \sum_{j=1}^k (\bar{r}_{.j} - \bar{r})^2$$

and

$$SS_e = \frac{1}{n(k-1)} \sum_{i=1}^n \sum_{j=1}^k (r_{ij} - \bar{r})^2$$

It represents the variance of the average ranking across treatments j relative to the total.

Under the hypothesis of no seasonality, all months can be equally treated. For the sake of completeness: -

$$\bar{r}_{.j}$$

is the average ranks of each treatment (month) j within each block (year) - The average rank is given by

$$\bar{r} = \frac{1}{nk} \sum_{i=1}^n \sum_{j=1}^k (r_{ij})$$

For large

$$n$$

or

$$k$$

, i.e. $n > 15$ or $k > 4$, the probability distribution of

$$Q$$

can be approximated by that of a chi-squared distribution. Thus, the p-value is given by

$$P(\chi_{k-1}^2 > Q)$$

.

Use

The test can be applied directly to any series by selecting the option *Statistical Methods » Seasonal Adjustment » Tools » Seasonality Tests*. This is an example of how results are displayed for the case of a monthly series:

If the null hypothesis of no stable seasonality is rejected at the 1% significance level, then the series is considered to be seasonal and the outcome of the test is displayed in green.

The test can be applied to the input series before any seasonal adjustment method has been applied. It can also be applied to the seasonally adjusted series or to the irregular component.

2. Non parametric (Friedman) test
Based on the rank of the observations in each year

Seasonality present
Distribution: Chi2 with 11 degrees of freedom
Value: 142.8654
PValue: 0.0000

Figure 47: friedman

In the case of X-13ARIMA-SEATS, the test is applied to the preliminary estimate of the unmodified Seasonal-Irregular component⁵² (time series shown in Table B3). In this estimate, the number of observations is lower than in the final estimate of the unmodified Seasonal-Irregular component. Thus, the number of degrees of freedom in the stable seasonality test is lower than the number of degrees of freedom in the test for the [presence of seasonality assuming stability](#). For example, X-13ARIMA-SEATS uses a centred moving average of order 12 to calculate the preliminary estimation of trend. Consequently, the first six and last six points in the series are not computed at this stage of calculation. The preliminary estimation of the trend is then used for the calculation of the preliminary estimation of the unmodified Seasonal-Irregular.

Related tests

- When using this kind of design for a binary response, one instead uses the Cochran's Q test.
- Kendall's W is a normalization of the Friedman statistic between 0 and 1.
- The Wilcoxon signed-rank test is a nonparametric test of non-independent data from only two groups.

References

- Friedman, Milton (December 1937). "The use of ranks to avoid the assumption of normality implicit in the analysis of variance". *Journal of the American Statistical Association* (American Statistical Association) 32 (200): 675–701. doi:10.2307/2279372. JSTOR 2279372.
- Friedman, Milton (March 1939). "A correction: The use of ranks to avoid the assumption of normality implicit in the analysis of variance". *Journal of the American Statistical Association* (American Statistical Association) 34 (205): 109. doi:10.2307/2279169. JSTOR 2279169.

⁵²The unmodified Seasonal-Irregular component corresponds to the Seasonal-Irregular factors with the extreme values.

- Friedman, Milton (March 1940). “A comparison of alternative tests of significance for the problem of m rankings”. The Annals of Mathematical Statistics 11 (1): 86–92. doi:10.1214/aoms/1177731944. JSTOR 2235971.

Moving seasonality test

The evolutive seasonality test is based on a two-way analysis of variance model. The model uses the values from complete years only. Depending on the decomposition type for the Seasonal – Irregular component it uses [1] (in the case of a multiplicative model) or [2] (in the case of an additive model):

$$|SI_{ij} - 1| = X_{ij} = b_i + m_j + e_{ij}$$

, [1]

$$|SI_{ij}| = X_{ij} = b_i + m_j + e_{ij}$$

, [2]

where:

m_j – the monthly or quarterly effect for j -th period, $j = (1, \dots, k)$, where $k = 12$ for a monthly series and $k = 4$ for a quarterly series;

b_i – the annual effect i , ($i = 1, \dots, N$) where N is the number of complete years;

e_{ij} – the residual effect.

The test is based on the following decomposition:

$$S^2 = S_A^2 + S_B^2 + S_R^2,$$

[3]

where:

$$S^2 = \sum_{j=1}^k \sum_{i=1}^N (\bar{X}_{ij} - \bar{X}_{..})^2$$

–the total sum of squares;

$$S_A^2 = N \sum_{j=1}^k (\bar{X}_{.j} - \bar{X}_{..})^2$$

– the inter-month (inter-quarter, respectively) sum of squares, which mainly measures the magnitude of the seasonality;

$$S_B^2 = k \sum_{i=1}^N (\bar{X}_{i\bullet} - \bar{X}_{\bullet\bullet})^2$$

– the inter-year sum of squares, which mainly measures the year-to-year movement of seasonality;

$$S_R^2 = \sum_{i=1}^N \sum_{j=1}^k (\bar{X}_{ij} - \bar{X}_{i\bullet} - \bar{X}_{\bullet j} + \bar{X}_{\bullet\bullet})^2$$

– the residual sum of squares.

The null hypothesis H_0 is that $b_1 = b_2 = \dots = b_N$ which means that there is no change in seasonality over the years. This hypothesis is verified by the following test statistic:

$$F_M = \frac{\frac{S_B^2}{(n-1)}}{\frac{S_R^2}{(n-1)(k-1)}}$$

which follows an F -distribution with $k - 1$ and $n - k$ degrees of freedom.

Combined seasonality test

This test combines the Kruskal-Wallis test along with test for the presence of seasonality assuming stability (F_S), and evaluative seasonality test for detecting the presence of identifiable seasonality (F_M). Those three tests are calculated using the final unmodified SI component. The main purpose of the combined seasonality test is to check whether the seasonality of the series is identifiable. For example, the identification of the seasonal pattern is problematic if the process is dominated by highly moving seasonality⁵³. The testing procedure is shown in the figure below.

Combined seasonality test, source: LADIRAY, D., QUENNEVILLE, B. (2001)

⁵³DAGUM, E.B. (1987).




Figure 48: Text

State space modelling

Introduction

State space forms play a central role in JD+. In a first point, we present the model used throughout the library. In a second point, we give an overview of the current implementation. The specificity of the state space framework of JD+ is stressed and the main modules are shortly presented. We finish the document with two examples based on the state space framework of JD+. The first one explains the basic structural models implemented in the “nbdemetra-sa-advanced” plug-in of JD+. The second one shows how the framework can be used to implement time-varying trading days models.

State space forms (SSF)

General form

The general linear gaussian state-space model can be written in many different ways. The measurement equation and the state equation considered in JD+ 3.0 are presented below.

$$\begin{aligned} y_t &= Z_t \alpha_t + \epsilon_t, & \epsilon_t &\sim \mathcal{N}(0, \sigma^2 H_t) \\ \alpha_{t+1} &= T_t \alpha_t + \mu_t, & \mu_t &\sim \mathcal{N}(0, \sigma^2 V_t) \end{aligned}$$

y_t is the observation at period t , α_t is the state vector. ϵ_t, μ_t are assumed to be serially independent at all time points and independent between them at all time points.

The residuals of the state equation will be modelled as $\mu_t = S_t \xi_t$, $\xi_t \sim \mathcal{N}(0, \sigma^2 I)$.

In other words, $V_t = S_t S_t'$.

The initial conditions of the filter are defined as follows:

$$\alpha_{-1} = a_{-1} + A_{-1} \delta + \mu_{-1}$$

$$\delta \sim \mathcal{N}(0, \sigma^2 \kappa I)$$

$$\mu_{-1} \sim \mathcal{N}(0, \sigma^2 P_*)$$

where κ is arbitrary large.

P_* is the variance of the stationary part of the initial state vector and models the diffuse part.

The model used in JD+ is quasi-identical to the model discussed in Durbin-Koopman.

Temporal disaggregation and benchmarking

Benchmarking Underlying Theory

Benchmarking⁵⁴ is a procedure widely used when for the same target variable the two or more sources of data with different frequency are available. Generally, the two sources of data rarely agree, as an aggregate of higher-frequency measurements is not necessarily equal to the less-aggregated measurement. Moreover, the sources of data may have different reliability. Usually it is thought that less frequent data are more trustworthy as they are based on larger samples and compiled more precisely. The more reliable measurement is considered as a benchmark.

Benchmarking also occurs in the context of seasonal adjustment. Seasonal adjustment causes discrepancies between the annual totals of the seasonally unadjusted (raw) and the corresponding annual totals of the seasonally adjusted series. Therefore, seasonally adjusted series are benchmarked to the annual totals of the raw time series⁵⁵. Therefore, in such a case benchmarking means the procedure that ensures the consistency over the year between adjusted and non-seasonally adjusted data. It should be noted that the ‘*ESS Guidelines on Seasonal Adjustment*’ (2015) do not recommend benchmarking as it introduces a bias in the seasonally adjusted data. Also the U.S. Census Bureau points out that: *Forcing the seasonal adjustment totals to be the same as the original series annual totals can degrade the quality of the seasonal adjustment, especially when the seasonal pattern is undergoing change. It is not natural if trading day adjustment is performed because the aggregate trading day effect over a year is variable and moderately different from zero.*⁵⁶ Nevertheless, some users may prefer the annual totals for the seasonally adjusted series to match the annual totals for the original, non-seasonally adjusted series⁵⁷. According to the ‘*ESS Guidelines on Seasonal Adjustment*’ (2015), the only benefit of this approach is that there is consistency over the year between adjusted and non-seasonally adjusted data; this can be of particular interest when low-frequency (e.g. annual) benchmarking figures officially exist (e.g. National Accounts, Balance of Payments, External Trade, etc.) where user needs for time consistency are stronger.

The benchmarking procedure in JDemetra+ is available for a single seasonally adjusted series and for an indirect seasonal adjustment of an aggregated series. In the first case, univariate

⁵⁴Description of the idea of benchmarking is based on DAGUM, B.E., and CHOLETTE, P.A. (1994) and QUENNEVILLE, B. et al (2003). Detailed information can be found in: DAGUM, B.E., and CHOLETTE, P.A. (2006).

⁵⁵DAGUM, B.E., and CHOLETTE, P.A. (2006).

⁵⁶‘*X-12-ARIMA Reference Manual*’ (2011).

⁵⁷HOOD, C.C.H. (2005).

benchmarking ensures consistency between the raw and seasonally adjusted series. In the second case, the multivariate benchmarking aims for consistency between the seasonally adjusted aggregate and its seasonally adjusted components.

Given a set of initial time series

$$\{z_{i,t}\}_{i \in I}$$

, the aim of the benchmarking procedure is to find the corresponding

$$\{x_{i,t}\}_{i \in I}$$

that respect temporal aggregation constraints, represented by

$$X_{i,T} = \sum_{t \in T} x_{i,t}$$

and contemporaneous constraints given by

$$q_{k,t} = \sum_{j \in J_k} w_{kj} x_{j,t}$$

or, in matrix form:

$$q_{k,t} = w_k x_t$$

.

The underlying benchmarking method implemented in JDemetra+ is an extension of Cholette's⁵⁸ method, which generalises, amongst others, the additive and the multiplicative Denton procedure as well as simple proportional benchmarking.

The JDemetra+ solution uses the following routines that are described in DURBIN, J., and KOOPMAN, S.J. (2001):

- The multivariate model is handled through its univariate transformation,
- The smoothed states are computed by means of the disturbance smoother.

The performance of the resulting algorithm is highly dependent on the number of variables involved in the model ($\propto n^3$). The other components of the problem (number of constraints, frequency of the series, and length of the series) are much less important ($\propto n$).

From a theoretical point of view, it should be noted that this approach may handle any set of linear restrictions (equalities), endogenous (between variables) or exogenous (related to external values), provided that they don't contain incompatible equations. The restrictions can also be relaxed for any period by considering their "observation" as missing. However, in practice, it appears that several kinds of contemporaneous constraints yield unstable results. This is more especially true for constraints that contain differences (which is the case for non-binding constraints). The use of a special square root initializer improves in a significant way the stability of the algorithm.

⁵⁸CHOLETTE, P.A. (1979).

Temporal Disaggregation underlying Theory

References