

Quantum Gravity and Quantum Computing

(State of the art)

G. Valdivia

University of Houston

gvaldivi@cougar.net.uh.edu

$$|\psi(t=0)\rangle = |\psi_1(t=0)\rangle \otimes |\psi_2(t=0)\rangle$$

$$= \frac{1}{\sqrt{2}}(|L\rangle_1 + |R\rangle_1) \frac{1}{\sqrt{2}}(|L\rangle_2 + |R\rangle_2)$$

$$= \frac{1}{2} [|LL\rangle_{12} + |LR\rangle_{12} + |RL\rangle_{12} + |RR\rangle_{12}]$$

$$V(t=0, t>0) = e^{-\frac{i}{\hbar} H t} \quad ; \quad H = \phi(x) = \frac{G m_1 m_2}{x_{12}}$$

$$* U|LL\rangle_{12} = e^{\frac{-i}{\hbar} \phi(x)t} |LL\rangle_{12} = e^{\frac{-i}{\hbar} \frac{Gm_1 m_2}{x_{12}} t} |LL\rangle_{12} \longrightarrow x_{12} = d$$


$$U|LR\rangle_{12} = \dots$$

$$U |R\rangle_{12} = e^{\frac{-i}{\hbar} \phi(x) t} |R\rangle_{12} = e^{\frac{-i}{\hbar} \frac{G m_1 m_2}{x_{12}} t} |R\rangle_{12} \longrightarrow x_{12} = d - \Delta x$$


$$\bigcup |RR\rangle_{12} = \dots$$

$$\begin{aligned}
 |\Psi(t>0)\rangle &= \frac{1}{2} \left[\underbrace{e^{\frac{-iG_{m_1 m_2}}{\hbar} \frac{\Delta}{2}}}_{e^{i\phi}} |LL\rangle_{12} + \underbrace{e^{\frac{-iG_{m_1 m_2}}{\hbar} \frac{\Delta+\Delta x}{2}}}_{e^{i\phi_{LR}}} |LR\rangle_{12} + \underbrace{e^{\frac{-iG_{m_1 m_2}}{\hbar} \frac{\Delta-\Delta x}{2}}}_{e^{i\phi_{RL}}} |RL\rangle_{12} + \underbrace{e^{\frac{-iG_{m_1 m_2}}{\hbar} \frac{\Delta}{2}}}_{e^{i\phi}} |RR\rangle_{12} \right] \\
 &= \frac{1}{\sqrt{2}} e^{i\phi} \left[\frac{1}{\sqrt{2}} |L\rangle_1 \left(|L\rangle_2 + e^{i\Delta\phi_{LR}} |R\rangle_2 \right) + \frac{1}{\sqrt{2}} |R\rangle_1 \left(e^{i\Delta\phi_{RL}} |L\rangle_2 + |R\rangle_2 \right) \right]
 \end{aligned}$$

Entangled state \longrightarrow We need to measure the entanglement!



Gravitational Entanglement Witnessing in SG interferometry



Step 1: Spin dependent spatial splitting of the center of mass state of the test mass m_j :

$$|C_1\rangle \xrightarrow{\frac{1}{\sqrt{2}}} \left(|\uparrow\rangle_1 + |\downarrow\rangle_1 \right) \longrightarrow \frac{1}{\sqrt{2}} \left(|L\uparrow\rangle_1 + |R\downarrow\rangle_1 \right)$$

$$|C_2\rangle \xrightarrow{\frac{1}{\sqrt{2}}} \left(|\uparrow\rangle_2 + |\downarrow\rangle_2 \right) \longrightarrow \frac{1}{\sqrt{2}} \left(|L\uparrow\rangle_2 + |R\downarrow\rangle_2 \right)$$

$$|C_i\rangle = \frac{1}{\sqrt{2}} (|L\rangle_i + |R\rangle_i)$$

Step 2:

$$|\Psi(t=0)\rangle = \frac{1}{\sqrt{2}} \left(|L\uparrow\rangle_1 + |R\downarrow\rangle_1 \right) \frac{1}{\sqrt{2}} \left(|L\uparrow\rangle_2 + |R\downarrow\rangle_2 \right)$$

$$|\Psi(t>0)\rangle = \frac{1}{\sqrt{2}} e^{i\phi} \left[\frac{1}{\sqrt{2}} |\uparrow\rangle_1 \left(|\uparrow\rangle_2 + e^{i\Delta\phi_{LR}} |\downarrow\rangle_2 \right) + \frac{1}{\sqrt{2}} |\downarrow\rangle_1 \left(e^{i\Delta\phi_{RL}} |\uparrow\rangle_2 + |\downarrow\rangle_2 \right) \right]$$

Step 3: Brings back the superposition through the unitary transformation

$$|L\uparrow\rangle_{1,2} \rightarrow |C,\uparrow\rangle$$

$$|R\downarrow\rangle_{1,2} \rightarrow |C,\downarrow\rangle$$

$$|\Psi(t_f)\rangle = \frac{1}{\sqrt{2}} e^{i\phi} \left[\frac{1}{\sqrt{2}} |\uparrow\rangle_1 \left(|\uparrow\rangle_2 + e^{i\Delta\phi_{LR}} |\downarrow\rangle_2 \right) + \frac{1}{\sqrt{2}} |\downarrow\rangle_1 \left(e^{i\Delta\phi_{RL}} |\uparrow\rangle_2 + |\downarrow\rangle_2 \right) \right] |C_1 C_2\rangle$$

Entangled state of the spins of the 2 test masses!

Entangled State:

$$|\uparrow\rangle_1 \left(|\uparrow\rangle_2 + e^{i\Delta\phi_{LR}} |\downarrow\rangle_2 \right) + |\downarrow\rangle_1 \left(e^{i\Delta\phi_{RL}} |\uparrow\rangle_2 + |\downarrow\rangle_2 \right) \neq |\Omega\rangle \otimes |\chi\rangle$$

Product State:

$$|\uparrow\rangle_1 \left(|\uparrow\rangle_2 + e^{i\Delta\phi_{LR}} |\downarrow\rangle_2 \right) + |\downarrow\rangle_1 \left(e^{i\Delta\phi_{RL}} |\uparrow\rangle_2 + |\downarrow\rangle_2 \right) \propto |\Omega\rangle \otimes |\chi\rangle$$

$$\text{If } -\Delta\phi_{LR} = 2\pi n + \Delta\phi_{RL}:$$

$$= \left[e^{i\Delta\phi_{LR}} |\uparrow\rangle_1 + |\downarrow\rangle_1 \right] \left[e^{i\Delta\phi_{RL}} |\uparrow\rangle_2 + |\downarrow\rangle_2 \right]$$

Let us define:

$$W = \sigma_x^{(1)} \otimes \sigma_z^{(2)} + \sigma_y^{(1)} \otimes \sigma_y^{(2)}$$

$$W = \begin{pmatrix} 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & -1 \\ 1 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{pmatrix}$$

$$|\psi(t_f)\rangle = \frac{1}{2} e^{i\phi} \begin{pmatrix} 1 \\ e^{i\Delta\phi_{LR}} \\ e^{i\Delta\phi_{RL}} \\ 1 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} i\phi \\ e^{i\phi} \\ e^{i\Delta\phi_{LR}} \\ e^{i\Delta\phi_{RL}} \\ e^{i\phi} \end{pmatrix}$$

Therefore:

$$|\langle W \rangle| = |\cos(\Delta\phi_{RL}) - \cos(\Delta\phi_{LR}) + \cos(\Delta\phi_{RL} - \Delta\phi_{LR}) - 1|$$

If $-\Delta\phi_{LR} = 2\pi n + \Delta\phi_{RL}$:


$$|\langle W \rangle| = \left| \frac{\cos(2\Delta\phi_{LR}) - 1}{2} \right| \in [0; 1] \rightarrow \text{Product State}$$

Thus:


$$|\langle W \rangle| > 1 \rightarrow \text{Entangled State}$$

Quantum Circuit:

$$\begin{aligned} * \quad |\Psi(t=0)\rangle &= \frac{1}{\sqrt{2}} (|L\uparrow\rangle_1 + |R\downarrow\rangle_1) \frac{1}{\sqrt{2}} (|L\uparrow\rangle_2 + |R\downarrow\rangle_2) \\ &= \frac{1}{2} [|L\uparrow\rangle_1 |L\uparrow\rangle_2 + |L\uparrow\rangle_1 |R\downarrow\rangle_2 + |R\downarrow\rangle_1 |L\uparrow\rangle_2 + |R\downarrow\rangle_1 |R\downarrow\rangle_2] \\ &= \frac{1}{2} [\underset{0000}{|\uparrow\uparrow LL\rangle}_{12} + \underset{0101}{|\uparrow\downarrow LR\rangle}_{12} + \underset{1010}{|\downarrow\uparrow RL\rangle}_{12} + \underset{1111}{|\downarrow\downarrow RR\rangle}_{12}] \end{aligned}$$



$$|L\rangle \rightarrow H|L\rangle = \frac{1}{\sqrt{2}} (|L\rangle + |R\rangle) = |C\rangle$$



$$\rightarrow \underset{1111}{|\downarrow\downarrow RR\rangle} + \underset{1010}{|\downarrow\downarrow RL\rangle} + \underset{0101}{|\uparrow\uparrow LR\rangle} + \underset{0000}{|\uparrow\uparrow LL\rangle}$$

$$\begin{aligned}
 |\Psi(t>0)\rangle &= \frac{1}{\sqrt{2}} e^{i\phi} \left[\frac{1}{\sqrt{2}} |\uparrow\uparrow\rangle_1 \left(|\uparrow\uparrow\rangle_2 + e^{i\Delta\phi_{LR}} |\downarrow\downarrow\rangle_2 \right) + \frac{1}{\sqrt{2}} |\downarrow\downarrow\rangle_1 \left(e^{i\Delta\phi_{RL}} |\uparrow\uparrow\rangle_2 + |\downarrow\downarrow\rangle_2 \right) \right] \\
 &= \frac{1}{2} \left[|\uparrow\uparrow LL\rangle e^{i\phi} + |\uparrow\downarrow LR\rangle e^{i\phi_{LR}} + |\downarrow\uparrow RL\rangle e^{i\phi_{RL}} + |\downarrow\downarrow RR\rangle e^{i\phi} \right]
 \end{aligned}$$


From U_1^+
From U_2^+
From U_3^+
From U_4^+

$$\begin{aligned}
 U_1^+ |\uparrow\uparrow\rangle &= U_1^+ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
 &= U_1^+ \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} e^{i\phi} \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad ; \quad U_1^+ = \begin{pmatrix} e^{i\phi} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}
 \end{aligned}$$

In what follows we will work only over the position states. Let us summarize:

$$|\uparrow\uparrow LL\rangle = |\uparrow\uparrow\rangle \quad ; \quad |\uparrow\downarrow LR\rangle = |\uparrow\downarrow\rangle \quad ; \quad |\downarrow\uparrow RL\rangle = |\downarrow\uparrow\rangle \quad ; \quad |\downarrow\downarrow RR\rangle = |\downarrow\downarrow\rangle$$

$$i) U_1^+ :$$




$$t=t_0: |\uparrow\uparrow\rangle_{12} + |\uparrow\downarrow\rangle_{12} + |\downarrow\uparrow\rangle_{12} + |\downarrow\downarrow\rangle_{12}$$

$$t=t_1: |\downarrow\downarrow\rangle_{12} + |\downarrow\uparrow\rangle_{12} + |\uparrow\downarrow\rangle_{12} + |\uparrow\uparrow\rangle_{12}$$

$$\begin{aligned}
 t=t_2: & \underbrace{|\downarrow\downarrow\rangle R_z(\frac{\phi}{2}) |\downarrow\downarrow\rangle}_{\downarrow \begin{pmatrix} 1 & 0 \\ 0 & e^{i\phi/2} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}} + \underbrace{|\downarrow\downarrow\rangle R_z(\frac{\phi}{2}) |\uparrow\uparrow\rangle}_{\downarrow \begin{pmatrix} 1 & 0 \\ 0 & e^{i\phi/2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}} + \underbrace{|\uparrow\downarrow\rangle R_z(\frac{\phi}{2}) |\downarrow\downarrow\rangle}_{\downarrow \begin{pmatrix} 1 & 0 \\ 0 & e^{i\phi/2} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}} + \underbrace{|\uparrow\downarrow\rangle R_z(\frac{\phi}{2}) |\uparrow\uparrow\rangle}_{\downarrow \begin{pmatrix} 1 & 0 \\ 0 & e^{i\phi/2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}} \\
 & \downarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 0 \\ e^{i\phi/2} \end{pmatrix} \quad \downarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 0 \\ e^{i\phi/2} \end{pmatrix} \quad \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
 & = e^{i\phi/2} |\downarrow\downarrow\rangle_{12} + |\uparrow\uparrow\rangle_{12} + e^{i\phi/2} |\uparrow\downarrow\rangle_{12} + |\uparrow\uparrow\rangle_{12}
 \end{aligned}$$

Convention:



$$t=t_3: e^{i\phi/2} |\downarrow\uparrow\rangle + |\downarrow\downarrow\rangle + e^{i\phi/2} |\uparrow\downarrow\rangle + |\uparrow\uparrow\rangle$$

$$t=t_4: e^{i\phi/2} |\downarrow\rangle \underbrace{R_z(-\phi/2)|\uparrow\rangle}_{|\uparrow\rangle} + |\downarrow\rangle \underbrace{R_z(-\phi/2)|\downarrow\rangle}_{e^{-i\phi/2}|\downarrow\rangle} + e^{i\phi/2} |\uparrow\rangle \underbrace{R_z(-\phi/2)|\downarrow\rangle}_{e^{-i\phi/2}|\downarrow\rangle} + |\uparrow\rangle \underbrace{R_z(-\phi/2)|\uparrow\rangle}_{|\uparrow\rangle}$$


$$= e^{i\phi/2} |\downarrow\uparrow\rangle_{12} + e^{-i\phi/2} |\downarrow\downarrow\rangle_{12} + |\uparrow\downarrow\rangle_{12} + |\uparrow\uparrow\rangle_{12}$$

$$t=t_5: e^{i\phi/2} |\downarrow\downarrow\rangle_{12} + e^{-i\phi/2} |\downarrow\uparrow\rangle_{12} + |\uparrow\downarrow\rangle_{12} + |\uparrow\uparrow\rangle_{12}$$

$$t=t_6: e^{i\phi} |\downarrow\downarrow\rangle + |\downarrow\uparrow\rangle + |\uparrow\downarrow\rangle + |\uparrow\uparrow\rangle$$

$$t=t_7: e^{i\phi} |\uparrow\uparrow\rangle + |\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle + |\downarrow\downarrow\rangle$$

ii) U_2^\dagger :




$$t=t_8: e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi\frac{LR}{2}} |\uparrow\downarrow\rangle + e^{-i\phi\frac{LR}{2}} |\downarrow\uparrow\rangle + |\downarrow\downarrow\rangle$$

$$t=t_9: e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi\frac{LR}{2}} |\uparrow\downarrow\rangle + e^{-i\phi\frac{LR}{2}} |\downarrow\downarrow\rangle + |\downarrow\uparrow\rangle$$

$$t=t_{10}: e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi\frac{LR}{2}} |\uparrow\downarrow\rangle + |\downarrow\downarrow\rangle + |\downarrow\uparrow\rangle$$

$$t=t_{11}: e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi\frac{LR}{2}} |\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle + |\downarrow\downarrow\rangle$$


iii) U_3^\dagger :



$t = t_{17}$:

$$e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi_{LR}} |\uparrow\downarrow\rangle + e^{i\phi_{RL}} |\downarrow\uparrow\rangle + |\downarrow\downarrow\rangle$$

* U_4^\dagger :




$t = t_{22}$:

$$e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi_{LR}} |\uparrow\downarrow\rangle + e^{i\phi_{RL}} |\downarrow\uparrow\rangle + e^{i\phi} |\downarrow\downarrow\rangle$$

LL
 LR
 RL
 RR

Thus, up to now:


$$|\Psi_{(t>0)}\rangle = \frac{1}{2} \left[|\uparrow\uparrow LL\rangle e^{i\phi} + |\uparrow\downarrow LR\rangle e^{i\phi_{LR}} + |\downarrow\uparrow RL\rangle e^{i\phi_{RL}} + |\downarrow\downarrow RR\rangle e^{i\phi} \right]$$



$t = t_{25}$:

$$\frac{1}{2} \left[e^{i\phi} |\uparrow\uparrow\rangle + e^{i\phi_{LR}} |\uparrow\downarrow\rangle + e^{i\phi_{RL}} |\downarrow\uparrow\rangle + e^{i\phi} |\downarrow\downarrow\rangle \right] |c_1\rangle |c_2\rangle$$

Convention:



$$|\Psi(t_f)\rangle = \frac{1}{\sqrt{2}} e^{i\phi} \left[\frac{1}{\sqrt{2}} |\uparrow\rangle_1 \left(|\uparrow\rangle_2 + e^{i\Delta\phi_{LR}} |\downarrow\rangle_2 \right) + \frac{1}{\sqrt{2}} |\downarrow\rangle_1 \left(e^{i\Delta\phi_{RL}} |\uparrow\rangle_2 + |\downarrow\rangle_2 \right) \right] |e_1 e_2\rangle$$





Fig. 5 Entanglement witness (W) vs $\Delta\phi_{LR} + \Delta\phi_{RL}$ plot. Data points in the upper half of the horizontal red line denote entangled states. The section of the X-axis in between two vertical green lines are the corresponding $\Delta\phi_{LR} + \Delta\phi_{RL}$ values for which the state is entangled

$$2.9113 < \Delta\phi_{LR} + \Delta\phi_{RL} < 4.2647$$




$$46.7439 \mu < \tau < 60.4742 \mu$$

Millisecond Coherence in a Superconducting Qubit

Aaron Somoroff,¹ Quentin Ficheux,¹ Raymond A. Mencia,¹ Haonan Xiong¹,¹
 Roman Kuzmin,¹ and Vladimir E. Manucharyan^{1,2}


¹*Department of Physics, Joint Quantum Institute, and Quantum Materials Center,
 University of Maryland, College Park, Maryland 20742, USA*

²*École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland*

 (Received 21 April 2021; revised 24 March 2023; accepted 10 May 2023; published 29 June 2023)

Improving control over physical qubits is a crucial component of quantum computing research. Here we report a superconducting fluxonium qubit with *uncorrected* coherence time $T_2^* = 1.48 \pm 0.13$ ms, exceeding the state of the art for transmons by an order of magnitude. The average gate fidelity was benchmarked at 0.99991(1). Notably, even in the millisecond range, the coherence time is limited by material absorption and could be further improved with a more rigorous fabrication. Our demonstration may be useful for suppressing errors in the next generation quantum processors.

DOI: 10.1103/PhysRevLett.130.267001

[4] On the possibility of laboratory evidence for quantum superposition of geometries 

Marios Christodoulou^{a,*}, Carlo Rovelli^b

^a Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China
^b CPT, Aix-Marseille Université, Université de Toulon, CNRS, F-13288 Marseille, France

References :

- [1] Bose, S., Mazumdar, A., Morley, G. W., Ulbricht, H., Toroš, M., Paternostro, M., ... & Milburn, G. (2017). Spin entanglement witness for quantum gravity. *Physical review letters*, 119(24), 240401.
- [2] Manabputra, Behera, B. K., & Panigrahi, P. K. (2020). A simulational model for witnessing quantum effects of gravity using IBM quantum computer. *Quantum Information Processing*, 19, 1-12.
- [3] Somoroff, A., Ficheux, Q., Mencia, R. A., Xiong, H., Kuzmin, R., & Manucharyan, V. E. (2023). Millisecond coherence in a superconducting qubit. *Physical Review Letters*, 130(26), 267001.

Our Team !!

