

Scalar Field Dark Matter Spectator During Inflation: The Effect of Self-interaction

Luis E. Padilla* and Tonatiuh Matos

Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, A.P. 14-740, 07000 México D.F., México.

J. Alberto Vázquez† and Gabriel Germán

Instituto de Ciencias Físicas, Universidad Nacional Autónoma de México,

Apdo. Postal 48-3, 62251 Cuernavaca, Morelos, México

(Dated: December 17, 2018)

Nowadays cosmological inflation is the most accepted mechanism to explain the primordial seeds that led to the structure formation observed in the Universe. Current observations are in well agreement to initial adiabatic conditions, which imply that single-scalar-field inflation may be enough to describe the early Universe. However, there are several scenarios where the existence of more than a single field could be relevant during this period, for instance, the situation where the so-called spectator is present. In this paper we apply the formalism of two-field inflationary models to two different spectator scenarios (jav: dos escenarios?). Firstly, to the possibility that an ultra-light scalar field dark matter coexist with the inflaton, where such ultra-light field could be auto-interacting or massive. Secondly, to the curvaton scenario (jav: ya no hay, o si?). For both cases (jav: both?) we compare the results to current cosmological data and provide constraints on the parameters describing the scalar field potential. (jav: 25 veces 'consider' se repite! – y ya cambie algunas)

PACS numbers: ????

I. INTRODUCTION

It is well accepted that the primordial seeds of the structure formation in the Universe were generated by quantum fluctuations given a scalar field (SF) during an inflationary epoch. The simplest scenario where density perturbations are carried out by a single inflaton is preferred since the initial perturbations are nearly adiabatic [1–3]. However the presence of any light field, other than the inflaton, could also fluctuate during inflation and contribute to the primordial density perturbations. A particular interest is the possibility that this extra SF can be used as the dark matter component (DM). This scenario is usually known as scalar field dark matter (SFDM) and assumes that the DM is made of bosonic excitations of an ultra-light SF. The typical mass of this field that satisfies astrophysical observations is around $m \sim 10^{-22} \text{eV}/c^2$, which might include self-interaction.

The idea of scalar fields as DM was initially introduced by [4] and since then rediscovered with different names, for instance Scalar Field Dark Matter (SFDM) [5], Fuzzy DM [6], Wave DM [7, 8], Bose-Einstein Condensate DM [9] or Ultra-light Axion DM [10, 11], amongst many others. The purpose of this SFDM is to resolve the apparent conflict, with observations, that exhibits the cold dark matter (CDM) formed of weakly-interacting massive particles (WIMPs) [12, 13]. Some of the CDM weaknesses may appear at small-scales within galaxies, e.g. cuspy

halo density profiles, overproduction of satellite dwarfs within the Local Group and many others, see for example [14–18]). Since then, the SFDM model has been successfully tested by different observations; for a review on ultra-light SFDM see [19–22].

The paper is organized as follows: First, in section II we review the basics about the inflationary scenario. Then in section III we present the basic observables used to constraint our models. Once the mathematical background and the observational constrictions are given, in section IV we analyze two different models (jav: cuales?). First, we assume a massive SFDM spectator during inflation. We provide some limits for the mass of the scalar field using isocurvature constrictions and compare them with the actual constraints given by astrophysical and cosmological observations. Then we consider a self-interacting term for the SFDM. We constrain their parameters using isocurvature perturbations and compare our results with cosmological and astrophysical observations. Finally in section V our conclusions are given.

II. INFLATIONARY SCENARIO: SFDM AS AN SPECTATOR

In this section we briefly review the inflationary process for spectator-like SFs during inflation following [26] (see also (lp: Referenciar nuestro paper) for a more recent review). Throughout this paper we use natural units ($\hbar = c = 1$).

A SF ϕ_i living during the inflationary era is thought to acquire quantum fluctuations with a primordial power

* epadilla@fis.cinvestav.mx

† javazquez@icf.unam.mx

spectrum measured at Hubble exit as

$$P_{\phi_i} \simeq \left(\frac{H_*}{2\pi} \right)^2. \quad (1)$$

Here and after quantities with subscript $*$ are evaluated at the Hubble exit. Assuming that within the inflationary era there exists several scalar fields, then it is convenient to work on a rotating basis by defining the *adiabatic field* σ parallel to the trajectory of the field-space and the *entropy fields* s_i perpendicular to the same trajectory. If the background trajectory is a straight-line that evolves in the direction of the field – causing the inflationary process –, the scenario is therefore called the *inflaton scenario* (IS). In this case quantum fluctuations $\delta\sigma$ of the adiabatic field and δs_i – also called *spectator fields* –, are frozen at Hubble exit¹ and start evolving until they re-enter the horizon. Considering the case with one extra SF, additional to the inflaton, then the primordial power spectrum is given by

$$P_{\sigma^*}(k) = P_s^*(k) \simeq \left(\frac{H_*}{2\pi} \right)^2. \quad (2)$$

On the other hand, the curvature and isocurvature perturbations are defined as

$$R \equiv \frac{H}{\dot{\sigma}} \delta\sigma, \quad S \equiv \frac{H}{\dot{\sigma}} \delta s. \quad (3)$$

Then the final power spectra, evaluated at the beginning of the radiation-domination era, are thus

$$P_R = P_S \simeq P|_*, \quad (4)$$

where at linear order in slow-roll parameters

$$P|_* = \frac{1}{2\epsilon} \left(\frac{H_*}{2\pi M_{pl}} \right)^2, \quad (5)$$

with $M_{pl} = 1.221 \times 10^{19} GeV$ the Planck mass.

Gravitational waves.– Given the fact that scalar and tensor perturbations are decoupled at linear order, the amplitude of the gravitational waves spectrum and the tensor-to-scalar ratio r , in the IS, remain the same similarly to the case with no extra spectator fields during the inflationary process.

We can see then that the incorporation of these new fields during the IS will produce only isocurvature perturbations. Such perturbations can be used to constrain the free parameters of our SFDM models as we will see later.

III. CONSTRAINTS ON INFLATIONARY PARAMETERS

In the standard approximation the inflationary observables are given by the tensor-to-scalar ratio r , the spectral index for adiabatic perturbations n_R and the amplitude for adiabatic perturbations A_r^2 . The constraints of these parameters are quoted at the pivot scale $k_0 = 0.05 Mpc^{-1}$ by [1–3, 27–29] (lp: [Agregar referencia del nuevo Planck](#))

$$A_r^2(k_0) = (2.215^{+0.032}_{-0.079}) \times 10^{-9}, \quad \text{at 68\% CL}, \quad (6a)$$

$$r_{k_0} < 0.064 \quad \text{at 95\% CL}, \quad (6b)$$

$$n_R(k_0) = 0.968 \pm 0.006. \quad (6c)$$

Using these measurements we are able to constrain the value of the Hubble expansion rate during inflation H_* as [30, 31]

$$r = 1.6 \times 10^{-5} \left(\frac{H_*}{10^{12} GeV} \right)^2. \quad (7)$$

As we have noted before, if more than one SF is present during inflation we will obtain isocurvature perturbations generated by extra scalar fields perpendicular to the trajectory on field space. Parameterizing the isocurvature power spectrum for dark matter in terms of the curvature power, we have

$$P_{DM}(k) = \frac{\beta_{iso}(k)}{1 - \beta_{iso}(k)} P_R(k), \quad (8)$$

where $P_{DM} = \langle \delta\rho_{DM^*}/\rho_{DM} \rangle$, $\delta\rho_{DM}$ are the isocurvature perturbations for the dark matter (DM) generated by extra scalar fields during inflation and ρ_{DM} is the initial condition of DM. The uncorrelated scale-invariant DM isocurvature is constrained by Planck data [1, 2] at pivot scale k_0 as

$$\beta_{iso}(k_0) < 0.038 \quad \text{at 95\% CL}. \quad (9)$$

Notice that isocurvature perturbations can be used to constrain the inflationary scale, just by combining equations (7), (8) and (9).

IV. CONSTRAINING MASSIVE AND SELF-INTERACTING ULTRA-LIGHT SFDM MODELS

In this section we assume the possibility that an ultra-light SFDM candidate coexists with the inflaton during the inflationary epoch. We require that the SFDM candidate be a stable spectator field with negligible classical dynamics and energy density. Such scenario is reached by considering the inflaton scenario, i.e., within the field

¹ If the trajectory is curved in the field-space, the entropy and adiabatic perturbations are correlated at Hubble exit and the perturbations continue evolving until inflation ends up [26]

space, the evolution of the system is on the inflaton direction ϕ whereas the direction perpendicular to the trajectory corresponds to the SFDM ψ . Notice this requirement implies that our dark matter candidate evolves much slower than the inflaton and its density is smaller than the associated to the inflaton.

As we mentioned before it is possible to constrain the free parameters of our model when isocurvature perturbations are taken into account. For this reason in the next subsections we review the cosmological history that a massive and a self-interacting scalar field should have gone through the evolution of the Universe, to then match the present values of the field with those during the inflationary era and therefore take into account isocurvature constrictions. We also consider different cosmological and astrophysical constraints for each SFDM model in order to compare our results.

First of all let us remember the basics about SFs. The dynamical evolution of any SF is governed by the Klein-Gordon (KG) equation

$$\square\Psi - 2\frac{dV}{d|\Psi|^2}\Psi = 0. \quad (10)$$

When a SF is complex it is convenient to use a Madelung transformation [34]

$$\Psi = \eta \exp[i\theta], \quad (11)$$

where $\eta \equiv |\Psi|$ is the magnitude of field Ψ and θ its phase. With this new decomposition the KG equation is separated in its real and imaginary components as

$$\ddot{\eta} + 3H\dot{\eta} + 2\frac{dV}{d|\psi|^2}\eta - \omega^2\eta = 0, \quad (12a)$$

$$\dot{\omega}\eta + (2\dot{\eta} + 3H\eta)\omega = 0, \quad (12b)$$

where $\omega = \dot{\theta}$. Equation (12b) can be exactly integrated obtaining

$$a^3\eta^2\omega = Q, \quad (13)$$

where Q is a charge of the SF related to the total number of particles [37–41]. Plugging this last equation into (12a) we obtain the radial component of the scalar field follows

$$\ddot{\eta} + 3H\dot{\eta} + M^2\eta - \frac{Q^2}{\eta^3} = 0. \quad (14)$$

The term containing Q is given by the complex nature of the SF [38] and can be interpreted as a “centrifugal force” [41]; $M^2 \equiv 2(dV/d|\psi|^2)$ is seen as an effective mass term of the scalar field. Notice that if we assume Ψ is the SFDM and we take $Q^2/\eta^3 \ll 1$, by assuming the SFDM candidate fulfills the slow-roll condition during inflation, then the field η will remain frozen at value η_i until $H \sim M$. Here and for the rest of this work subindex i means values right after inflation ends. Then, when

$H \sim M$ will start evolving depending on the effective mass term. In order to get the slow-roll behavior of the SFDM it is necessary that $Q \simeq 0$, as explained in (lp: Referencia paper Abril)². On the other hand if Ψ is the inflaton, ϕ it is usually considered as a real field, then in this case $\theta = Q = 0$.

In this work we consider only situations where the general potential can be decomposed as $V(\phi, \psi) = V_{inf}(\phi) + V_{SFDM}(|\psi|^2)$.

A. Real ultra-light SFDM candidate

1. Cosmological history

The possibility that an ultra-light SFDM candidate could coexist with the inflaton has been recently studied in [44] considering a potential of the form

$$V(\phi, |\psi|^2) = V(\phi) + \frac{1}{2}m^2\psi^2. \quad (15)$$

and by fixing $Q = 0$ for the SFDM in Eq. (14). Such study was performed by assuming the SFDM is an axion-like particle. However their results can be easily extrapolated for mostly any massive SFDM candidate. For this particular case we observe that $M^2 = m^2$. As mentioned above when $H \gg m$ the term with m^2 in equation (12a) can be neglected. Bering in mind the field is slowly rolling during the inflationary era, we can neglect the second derivatives in (12a), and then the field ψ remains frozen at its initial value by Hubble dragging during the early universe [42]. When the $m \sim H$ condition is approached the SFDM starts evolving and oscillates as a massive field. During its oscillation phase the dependence of ψ respect to a is $\psi \sim 1/a^{3/2}$, while its density behaves as $\rho_\psi \sim 1/a^3$ [37, 38]. So the scalar density of the SFDM can be written as

$$\rho_\psi = \begin{cases} \frac{1}{2}m^2\psi_i^2 & \text{when } H \gg m, \\ \frac{1}{2}m^2\psi_i^2 \left(\frac{a_{osc}}{a}\right)^3 & \text{when } H \ll m. \end{cases} \quad (16)$$

For typical masses of a SFDM candidate, $m \sim 10^{-22}eV$, the field started oscillating during the radiation-dominated Universe. During this period the Hubble parameter evolves in terms of the scale factor as $H \propto a^{-2}$, and the KG equation (12a) can be solved exactly in terms of a . In Ref. [44] the initial conditions for the massive case were obtained (jav: they) by using the evolution of the form (16) and taking into account that the entropy of the Universe is conserved. If the total

² In fact the inflationary behavior is an attractor solution of the KG equation for a real field in the limit when $M^2 \ll H^2$ [35, 36]. In this limit the typical dynamics of a real SF is a stiff-like epoch, followed by an inflationary-like era.

amount of dark matter is composed of SFDM particles they obtain (jav: they?)

$$\psi_i^2 \simeq \frac{10^{34} \text{GeV}^2}{0.6} \left(\frac{g_{*osc}}{3.36} \right)^{-3/4} \left(\frac{g_{s*osc}}{3.91} \right) \left(\frac{m}{10^{-22} \text{eV}} \right)^{-1/2}. \quad (17)$$

Here g_{*osc} and g_{s*osc} are the effective degrees of freedom associated to the total particles and to the entropy of the SFDM oscillations. In particular, for the case of an ultra-light SFDM that started its oscillations during the radiation-dominated Universe $g_{osc} = 3.36$ and $g_{sosc} = 3.91$.

2. Constraints from isocurvature perturbations

For this case we demand the energy density contribution of the SFDM being small during inflation (DM dominates right after radiation-matter equality) and hence it is necessary that

$$\frac{m^2}{2} < \frac{V(\phi)}{\psi_i^2} \simeq \frac{H_*^2 M_p^2}{\psi_i^2}, \quad (18)$$

where during inflation our field remains frozen at value ψ_i . Notice that for an ultra-light SFDM candidate ($m \sim 10^{-22} \text{eV}$) the above expression is fulfilled for most of the initial conditions given by ψ_i . On the other hand we can constraint isocurvature perturbations generated by a SFDM using Eqn. (8) or equivalently (9). The way it is done was studied in ref [44] by noticing that we can re-express the primordial spectrum as $\delta\rho_\psi/\rho_\psi = 2\delta\psi/\psi_i$ (since from Eq. (16) we have $\rho_{SFDM} \propto \psi^2$) which implies that $P_{SFDM} = 4P_\psi/\psi_i^2$, where P_ψ is given by equation (1) or (2) and ψ_i by equation (17), and then compare it with P_{DM} from equation (8). When such comparison is done they finally obtain the result

$$\frac{m}{10^{-22} \text{eV}} < \left(\frac{2 \times 10^{-4}}{r} \right)^2. \quad (19)$$

Then the isocurvature constrictions allow us to constraint the mass parameter of the SFDM with the measurement of the tensor-to-scalar ratio.

The above relation for the mass parameter must be in agreement with the cosmological and astrophysical constrictions obtained by this model. We need to stress out that we cannot use all the constrictions in the literature since some of them consider different cosmological evolutions for the SFDM. For example in [37] it was studied CMB and Big Bang Nucleosynthesis (BBN) by understanding the SFDM was generated right after inflation with a stiff-like equation of state ($p \simeq \rho$). Then, these kind of restrictions are not applicable to our model.

Other constraints. - We start by considering CMB constrictions. In reference [61] the CMB was studied in the form of Planck temperature power spectrum, here they

obtained $m \gtrsim 10^{-24} \text{eV}$. Considering the hydrodynamical representation of the SFDM model, Ref. [62] suggests the SFDM's quantum pressure as the origin of the offset between dark matter and ordinary matter in Abel 3827. For this purpose they required a mass $m \simeq 2 \times 10^{-24} \text{eV}$. When the model is tested with the dynamics of dwarf spheroidal galaxies (dSphs) – Fornax and Sculpture–, in reference [63] was obtained a mass constriction of $m < 0.4 \times 10^{-22} \text{eV}$ at 97.5% The constriction obtained when the survival of the cold clump in Ursa and the distribution of globular clusters in Fornax is considered requires a mass $m \sim 0.3 - 1 \times 10^{-22} \text{eV}$ [64]. Explaining the half-light mass in the ultra-faint dwarfs fits the mass term to be $m \sim 3.7 - 5.6 \times 10^{-22} \text{eV}$ [65]. The model has also been constrained by considering the process of reionization. In [66] it was obtained, using N-body simulations and demanding an ionized fraction of HI of 50% by $z = 8$, the result of $m > 2.6 \times 10^{-23} \text{eV}$. Finally, using the Lyman- α forest flux power spectrum demands that the mass parameter fulfills $m \gtrsim 20 - 30 \times 10^{-22} \text{eV}$ [67, 68].

Figure 1 displays the aforementioned constraints on a m vs r plane. In order to simplify the lecture of the figure we have only plotted the upper(lower) value for the constrictions that fit the mass of the SFDM with an upper(lower) limmit. We have also added arrows that points out the region that remains valid for such constrictions. Firstly, the gray region is fulfilled by isocurvature observations (19). The dot-dashed black line corresponds to the equality values in Eq. (19). Then, the figure must be interpreted as follows: Suppose we have measured a value for r . Notice that such value will cross with the dot-dashed black line for a given mass m_{max} . Then, the masses that will continue allowed by the model must be the ones that are lower than m_{max} .

The region that fulfil observations obtained by CMB is specified in green, while the value provided by Abel 3827 is given by the dot-dashed red line. The region for dwarf spheroidal galaxies is indicated in blue, Ursa with Fornax in light blue, ultra-faint dwarfs in teal, reionization in purple and Liman- α in orange. We notice that isocurvature perturbations cannot constrain observations of the dynamics of dSphs galaxies given that both provide an upper limit for the mass of the SFDM. However, the detectability of gravitational waves and the different constrictions by cosmological and astrophysical observations can be used to test the massive model. For example, if we ignore by the moment the dynamics of dSphs galaxies and we would like to fulfill at least observations provided by Abel 3827, we should not detect gravitational waves until $r \simeq 1.3 \times 10^{-3}$ (fuchsia stright-line), while if we were interested in fulfilled all the other observations we should not detect gravitational waves until $r \lesssim 2.33 \times 10^{-5}$ (gold stright-line).

These results are important given that [46] demonstrated that an ultra-light axion-like dark matter candidate must be presented during inflation. Then, if r is detected in the near future, it could represent a strong

constraint for the axion-like particle model. Notice that if we relax the mechanism under this particle is created or if we add an auto-interacting component, we should expect these restrictions be less affective to the model.

We also plotted the actual upper limit for r in a navy blue dashed-line. By the moment this value is not so restrictive for the model since it represents an upper value for r . The only information it can provide us is that the masses smaller than the mass m_{upper} where the blue dashed-line and the black dot-dashed-line are crossed, are allowed by the data. However, masses bigger than m_{upper} can not be discarded since the only way it could be possible to do it is if r is detected.




FIG. 1: Isocurvature constraints for the SFDM candidate.

B. Real Self-interacting SFDM Candidate

1. Cosmological history

In this section a self-interacting SFDM with a positive interaction is considered. This scenario is described by the general potential

$$V(\phi, |\psi|^2) = V(\phi) + \frac{1}{2}m^2\psi^2 + \frac{1}{4}\lambda\psi^4, \quad (20)$$

and fixing again $Q = 0$ for the SFDM in equation [14].

Notice that for this case $M^2 = m^2 + \lambda\psi^2$. As we have previously discussed the effective mass of the field after inflation remains constant at $M^2 = m^2 + \lambda\psi_i^2$ until $M \sim H$. Then, depending of each contribution to M^2 , we can have two different dynamics.

Weakly self-interacting regime.- This limit is obtained when the constant term m^2 dominates, that is when

$$m^2 \gg \lambda\psi_i^2. \quad (21)$$

In this regime it is possible to ignore the autointeracting term in equation (14) when the oscillations of the

scalar field begins. However, by ignoring this term the field behaves as a massive field and from (16) the field value always decreases. Therefore the autointeracting term never dominates and all the cosmological history remains the same as in the pure massive SFDM scenario. In fact, thanks to the decreasing behavior of this scenario we can consider that this regime is fulfilled always that $m^2 \geq \lambda|\psi_i|^2/2$ or equivalently when $\lambda \leq 2m^2/|\psi_i|^2$. If the SFDM oscillations start at the same time than the massive case (which is a good approximation since the effective mass of the SFDM is $M^2 = m^2 + \lambda\psi_i^2 \leq 2m^2$), we observe from (17) that it must be fulfilled that

$$\left(\frac{\lambda}{10^{-96}}\right) \leq 1.2 \left(\frac{m}{10^{-22}eV}\right)^{5/2}. \quad (22)$$

We plot in figure 2 the weak limit obtained by our approximation. However this overestimates the maximum value of λ since the dust-like behavior is obtained when the λ term is completely negligible.




FIG. 2: Weakly self-interacting regime

Strong self-interacting regime.- This scenario is obtained when

$$m^2 \ll \lambda\psi_i^2. \quad (23)$$

Here the SFDM follows an attractor solution during the inflationary era.

Attractor behavior of the SF during inflation.- In the strong self-interacting regime the SFDM follows the attractor solution [42]³

$$\psi_{att} = \left(2\lambda \int_{\phi}^{\phi_0} V_{,\phi}^{-1} d\phi\right)^{-1/2}, \quad (24)$$

³ The study done in reference [42] was for a curvaton-like scalar field in a chaotic-like inflationary scenario. However their results can be used as well in this context were the attractor behavior can be easily obtained for whichever inflationary potential.

where ϕ_0 is the value of the inflaton at the beginning of inflation. In the above expression we can identify two possible branches:

- $\psi_{att} < \sqrt{2}m/\sqrt{\lambda}$

The SFDM follows the attractor solution until $\psi \simeq \sqrt{2}m/\sqrt{\lambda}$. Then the field reaches $\psi_i = \sqrt{2}m/\sqrt{\lambda}$ for the rest of inflation. Notice that this value corresponds to the upper limit that the weakly self-interacting regime allows. Then the field starts evolving when $H \sim M \simeq m$ behaving as a massive SF. In this way the constrictions given in the non-interacting case apply and the initial conditions are also fixed by ψ_i . Using both relations the λ value is approximated

$$\left(\frac{\lambda}{10^{-96}}\right) \simeq 1.2 \left(\frac{g_{*osc}}{3.36}\right)^{3/4} \left(\frac{g_{s*osc}}{3.91}\right)^{-1} \left(\frac{m}{10^{-22}eV}\right)^{5/2}. \quad (25)$$

The mass term and the auto-interacting constant are rescaled by appropriate values, that is, the mass term is measured in units of $10^{-22}eV$ while the auto-interacting constant in terms of 10^{-96} (jav: unidades)(lp: Según yo es adimensional.).

- $\psi_{att} > \sqrt{2}m/\sqrt{\lambda}$

In this scenario the dynamics of the inflaton, given by (24), implies that the initial condition of the field after the inflationary period is

$$\psi_{att}^i = \left(2\lambda \int_{\phi_{end}}^{\phi_0} V_{,\phi}^{-1} d\phi\right)^{-1/2}, \quad (26)$$

where ϕ_{end} is the value of the inflaton at the end of inflation. We need to stress out that this is the value of the field until its oscillation period starts (i.e. when $M \sim H$).

In this scenario we observe that at the time the SFDM starts its oscillations its effective mass is quadratic in the field. In that regime the scalar field evolves as $\psi \sim 1/a$ and its energy density as $\rho_\psi \sim 1/a^4$, behaving as radiation. Then, when $m^2 \sim \lambda\psi_t^2$ the effective scalar field mass is now constant, obtaining the dust-like behavior already analyzed. Therefore, the history of the scalar field density is

$$\rho_\psi = \begin{cases} \frac{1}{4}\lambda^2\psi_i^4 & \text{when } H \gg \lambda\psi_i^4 \\ \frac{1}{4}\lambda\psi_i^4 \left(\frac{a_{osc}}{a}\right)^4 & \text{when } H_t \leq \lambda\psi_i^4 \leq H \\ \frac{1}{2}m^2\psi_t^2 \left(\frac{a_t}{a}\right)^3 & \text{when } H \leq m^2 \text{ and } \lambda\psi^2 < m^2 \end{cases} \quad (27)$$

Here sub-index t means quantities measured at transition between radiation-like to dust-like behavior of the SFDM and

$$\psi_i^2 = \left[\frac{2m^2}{\lambda}\psi_t^2\right]^{1/2} \left(\frac{a_t}{a_{osc}}\right)^2. \quad (28)$$

Notice that, for simplicity, we have taken an instantaneous transition between radiation-like to dust-like behaviors.

Since the auto-interacting KG equation cannot be solved exactly we work with approximated solutions. By using a pure approximated description of the system, [38] obtained the relation (see its equation 80 and 86 and also [37])⁴

$$\left(\frac{a_t}{a_{osc}}\right)^2 = \frac{3}{7^{1/3}f^2\left(\frac{a_s}{r_S}\right)}, \quad (29a)$$

where

$$f(\sigma) = \frac{1}{s^{1/3}(1+4s)^{1/6}}, \quad (29b)$$

with

$$s = \frac{4\sigma - 1 + \sqrt{(4\sigma - 1)^2 + 12\sigma}}{6}. \quad (29c)$$

Additionally $r_S = 2mG/c^2$ and $a_s = \hbar^2\lambda/4\pi m$. Then it follows that $a_s/r_S = \lambda M_p^2/m^2$. Rearranging the expression in a more convenient way we have

$$\sigma \simeq 5.93 \times 10^2 \left(\frac{m}{10^{-22}eV}\right)^{-2} \left(\frac{\lambda}{10^{-96}}\right). \quad (30)$$

Notice that when $a_t/a_{osc} \simeq 1$, i.e. $3/(7^{1/3}f^2(\sigma)) \sim 1$, there is no radiation-like epoch. This scenario should match with the non-interacting scenario that we present previously. Inserting equation (29a) into (28) yields to

$$\psi_i^2 = \frac{3}{7^{1/3}f^2(\sigma)} \left[\frac{2m^2}{\lambda}\psi_t^2\right]^{1/2}. \quad (31)$$

The relation (31) matches the field at ψ_t with its value right after inflation ends. Then if we obtain the value of ψ_t by comparing with quantities at present, with the above expressions we can also obtain the value of ψ_i . On the other hand, notice that at a_t the scalar field behaves as dust with an effective mass $M^2 = m^2 + \lambda\psi_t^2$. This implies that dust-like oscillations of the SF began a little before than in the non-interacting case. If we allow m to be ultra-light ($m \sim 10^{-22}eV$) and using the fact that m^2 is about the same order that $\lambda\psi_t^2$, we get that such oscillations start during the same epoch than in the non-interacting case. In fact because

⁴ The reference [38] obtained this relation by considering a Universe with only a SFDM content. However a similar analysis can be used in a Universe with several types of matter contents.

the decreasing behavior of the SF during the dust-like period ($\psi \sim 1/a^{3/2}$) the auto-interacting term contribution quickly vanishes and then the dynamics of the field is described only by the mass term m . Thus, once the dust-like behavior starts, the dynamics is described similarly to the non-interacting case, in such case the condition (17) is fulfilled by the SF as well, but interchanging subindex i with t^5 .

2. Constraints from isocurvature perturbations

As we have shown in the last section we have two different scenarios for this model: a weak self-interacting and a strong self-interacting. In the weak limit our SFDM behaves effectively as a massive field without auto-interaction, and in such case the constrictions for the massive field apply to this scenario as well. On the other hand when the auto-interacting term is big enough, the SFDM will have a new period with a behavior similar to a radiation-like fluid. In this way the constrictions we obtained before will not apply to this model anymore

In the strong self-interacting regime, during the inflationary era, the SFDM follows the attractor solution (24). The value that the homogeneous field acquired after inflation depends on the condition $\psi_{att} \leq \sqrt{2}m/\sqrt{\lambda} \equiv \psi_t$. For $\psi_{att} < \psi_t$ the field follows the attractor solution until $\psi \simeq \psi_t$. Then the SFDM is frozen at that value and starts oscillating as a massive field when $m \sim H$. We can constraint this scenario by noticing that it is the same than the massive case but with the initial condition $\psi_i = \psi_t$. Matching Eq. (17) with ψ_t and considering the constriction (19) we obtain

$$\left(\frac{\lambda}{10^{-96}} \right) \leq 1.2 \left(\frac{2 \times 10^{-4}}{r} \right)^5. \quad (32)$$

In figure 3 we have plotted the above condition that is valid during the strong self-interacting regime, when $\psi_{att} < \psi_t$. The pink region corresponds to the region allowed by isocurvature perturbations in this limit. As we observe the self-interacting term for this model can be constrained in a similar way than the mass parameter in the massive case. This scenario must fulfill the relation (19) as well, since its cosmological evolution after inflation is only like a massive SFDM.

Additionally, in this scenario, the inflationary potential fulfills the condition

$$\left(\int_{\phi}^{\phi_0} V_{,\phi}^{-1} d\phi \right)^{-1/2} < 2m. \quad (33)$$

⁵ In fact this is a lower limit for the strong auto-interacting case.




FIG. 3: Isocurvature constraints for the weakly self-interacting term. (jav: weak?)

We can see that it is very difficult to obtain this relation for an ultra-light SFDM candidate. For example, if we consider a chaotic-like inflationary potential, $V(\phi) = \frac{1}{2}M_{inf}^2\phi^2$, the above conditions implies that

$$\left(\log \frac{\phi_0}{\phi} \right)^{-1/2} < 2 \frac{m}{M_{inf}}. \quad (34)$$

However, for this potential the mass M_{inf} of the inflaton that best matches the observations⁶ is of order $M_{inf} \sim 10^{12} GeV$ [48]. If now we assume an ultra-light SFDM candidate with a mass $m \sim 10^{-22} eV$, the above conditions implies that the logarithmic part of the expression should be lower than $\sim 10^{-43}$. The inflationary behavior for a chaotic-like inflaton ends when $\phi_{end} \simeq 2M_{pl}$ [42, 48]. Moreover as it is explained in [42], the initial condition of the inflaton cannot be arbitrarily large since the stochastic behavior is significant for $\dot{\phi}H^{-1} < H/2\pi$. If the Universe starts when the inflaton escapes from this behavior we have that its initial condition should be

$$\phi_0 \sim 10^5 \times M_{pl} \left(\frac{10^{13} GeV}{M_{inf}} \right)^{1/2}. \quad (35)$$

where we can easily see that the condition given in (34) cannot be fulfilled. This implies that if we consider that an ultra-light self-interacting SFDM candidate coexists with the inflaton, it should be within the strong regime since their conditions are easier to satisfy.

When $\psi_{att} > \psi_t$ we have that the field follows the attractor solution during all the inflationary period. Hence the initial condition for the SFDM is given by (26)

$$\psi_{att}^i = \left(2\lambda \int_{\phi_{end}}^{\phi_0} V_{,\phi}^{-1} d\phi \right)^{-1/2}. \quad (36)$$

⁶ This chaotic-like inflationary potential is ruled-out now for observations, however we use it as an example in order to obtain general constraints for our models.

Then the SFDM remains frozen at value ψ_{att}^i until $M \sim H$ and starts oscillating with a quartic potential. In this scenario the SFDM density behaves as $\rho_{SFDM} \propto \psi^4$ and in such case we can write $\delta\rho_\psi/\rho_\psi = 4\delta\psi/\psi_i$. Therefore the primordial isocurvature perturbations for a strong self-interacting SFDM is given by

$$P_{SFDM}(k) = \left(\frac{2H_*}{\pi\psi_i} \right)^2. \quad (37)$$

In the last section we showed the relation of the initial condition with the value of the field today. Using eqs. (31) and (17), with $g_{*osc} = 3.36$ and $g_{s*osc} = 3.91$ and considering appropriate units we obtain

$$r < \frac{1.172 \times 10^{-4}}{7^{1/3} f^2(\sigma)} \left[\frac{2 \left(\frac{m}{10^{-22} \text{eV}} \right)^{3/2}}{\left(\frac{\lambda}{10^{-96}} \right)} \right]^{1/2}. \quad (38)$$

Similarly to the massive case, the above relation must be compared with observations in order to get constrictions for the strong self-interacting SFDM scenario.

Other constraints.- In [69] it was studied the possibility that a SFDM candidate could be self-interacting. In their work they constraint the ratio $\Lambda \equiv m/\lambda^{1/4}$ to be $\Lambda \sim 1\text{eV}$ by analyzing the line-of-sight velocity dispersion for the eight dSphs satellites of the Milky Way (MW). This study was complementary to the ones done in [70–73] where the SFDM model was studied by employing rotational curves of the most Dark Matter dominated galaxies from different surveys and where they obtained the constriction $\Lambda \sim 2.6 - 2.9\text{eV}$. On the other hand, in [74] the model was studied in a cosmological context by demanding that the SFDM candidate behaves as a dust-like component before the time of matter-radiation equality. In such work they obtain the result $\Lambda > 0.8\text{eV}$. Finally, it was also possible to test the self-interacting scenario by considering the number of extra relativistic species at BBN. When such observations are confronted with the SFDM it is obtained that it must be fulfilled that $\Lambda \gtrsim 4\text{eV}$ (see [71]).

In figure 4 we have plotted contour levels of the numerical value of the right side of the relation (38). The grey region corresponds to values larger than 0.064 which is the actual upper constraint on tensor-to-scalar ratio. This means that within that region we are certain that (38) is fulfilled and, by the moment, just this region is completely allowed by observations. With this in mind the plot should be understood as follows: let's suppose that in the near future we measure a value for r . Such value will coincide with a curve in figure 4. Then, the space of parameters that could continue be allowed by data should be the one where the contour levels are bigger than the detected value of r , while the one with a smaller value in the contour levels must be discarded. Similarly than in the massive case, if r is not detected but it continue being constrained with an upper limit,




FIG. 4: Isocurvature constraints for the strong auto-interacting scenario. (jav: explicar bien)

this implies that the regions with contour levels bigger than such upper value will be allowed by the data, but the regions with smaller values can not be discarded until r is detected.

The value of Λ that fulfil observations of dSph's Light-of-sight is given by the dot-dashed red line, while the region of parameters necessary for rotational curves is presented in blue. Similar than in the massive plot (fig. 1) we plotted the cosmological constriction in purple by drawing a curve that refers to the upper value of Λ and an arrow that point out the region that remains valid for the constriction. We did the same for the observations for BBN but in fuchsia. The white region corresponds to the weak limit. We can see from figure 4 that it is possible to fulfill observations for whichever value for the mass as long as the self-interacting constant is large enough. In other words, for a given mass, the measurement of r can only constraint the self-interacting constant with a lower limit. Isocurvature observations (or equivalently observations on r) can also help the model to constraint with an upper value other observations. As an example let's suppose that we want to be completely sure that the measurements obtained by rotational curves are fulfilled. Then, given the actual upper constraints in r , the only region of parameters that we can be sure that fulfil observations are the ones in the blue region that are also inside the gray region.

Remark: This scenario is of special interest given that the attractor solution justify the initial conditions for the SFDM model and because it is natural to avoid isocurvature perturbations when the auto-interacting term of the SF is big enough.

Similarly to the above description we can compute general constraints for the inflationary potential that should generate inflation on these kind of scenarios. First we

have that

$$\left(\int_{\phi}^{\phi_0} V_{,\phi}^{-1} d\phi \right)^{-1/2} > 2m, \quad (39)$$

which is very easy to fulfill as we saw in the chaotic-like example. Using isocurvature constricton we also have

$$r < \frac{0.6 \times 10^{40}}{\left(\frac{\lambda}{10^{-96}} \right)} \left(\frac{1}{\int_{\phi_{end}}^{\phi_0} V_{,\phi}^{-1} d\phi} \right), \quad (40)$$

that can also be satisfied as far as the auto-interacting term and the integral are small enough; for example in the chaotic-like scenario by using (34) and (35) and taking $M_{inf} \simeq 10^{-6} M_{pl}$, we can obtain the constricton

$$\left(\frac{\lambda}{10^{-96}} \right) < \frac{0.3288 \times 10^{84}}{r}. \quad (41)$$

that is easily satisfied for whichever value of λ of our interest. If now we compare (26) and (31) we have

$$\left(\int_{\phi_{end}}^{\phi_0} V_{,\phi}^{-1} d\phi \right)^{-1} = \frac{6}{7^{1/3} f^2(\sigma)} (2m^2 \lambda |\psi_t|^2)^{1/2}. \quad (42)$$

This relation is interpreted as follows: consider that the auto-interacting SFDM candidate coexists with the inflaton, and suppose there are several measurements constraining the mass parameter m as well as the auto-interacting parameter λ , therefore such constraints are translated into restrictions to the inflationary potential.

It is also necessary to be careful that the SFDM does not come to dominate the inflationary period. This is guarantee by demanding that

$$\lambda < \frac{H_*^2 M_p^2}{|\psi_i|^4}, \quad (43)$$

or in terms of (36)

$$\lambda > \left(4H_*^2 M_{pl}^2 \left(\int_{\phi_{end}}^{\phi_0} V_{,\phi}^{-1} d\phi \right)^2 \right)^{-1}. \quad (44)$$

Considering the chaotic-like example and using $H_* = 10^{14} GeV$, we obtain the constricton

$$\frac{\lambda}{10^{-96}} > 5.777 \times 10^{77}. \quad (45)$$

Notice that the above expresion fulfills astrophysical constrictions but it does not fulfills CMB and BBN. This may not be a problem for auto-interacting SFDM models since Eq. (45) is obtained for a chaotic-like inflationary potential and, as we already know, this inflationary model may be in problems from the actual constraints given by Planck ([lp: Referencia Planck 2018](#)).

C. Complex SFDM generalization

(jav: ————— aqui voy —————)

As we have seen at the beginning of this section, when we consider a complex scalar field its dynamics is modified only by the centrifugal term (see eq. (14)). However as it was mentioned in [38] such term does not affect the dynamics of the field at cosmological levels, obtaining then that a complex scalar field and a real scalar field have the same cosmological history in the Universe. In this way if we take that our complex SFDM fulfill slow-roll conditions during inflation then its constrictions for isocurvature perturbations must be the same than in the real field analogue.

V. CONCLUSIONS

-
- [1] P. A. R. Ade et al. (Planck), *Astron. Astrophys.* 571 , A16 (2014), arXiv:1303.5076 [astro- ph.CO]
 - [2]] P. A. R. Ade et al. (Planck), *Astron. Astrophys.* 571 , A22 (2014), arXiv:1303.5082 [astro- ph.CO]
 - [3] P. A. R. Ade et al. (Planck); DOI: 10.1051/0004-6361/201525898; arXiv:1502.02114 [astro-ph.CO]
 - [4] Baldeschi M., Gelmini G., Ruffini R., 1983, *Phys.Lett.B*, 12
 - [5] Matos T., Guzman F. S., 2000, *Class. Quant. Grav.*
 - [6] Hu W., Barkana R., Gruzinov A., 2000, *Physical Review Letters*, 85, 1158
 - [7] Bray H., 2010
 - [8] Schive H.-Y., Chiueh T., Broadhurst T., 2014a, *Nature Phys.*, 10, 496
 - [9] Böhmer C. G., Harko T., 2007, *J. Cosmology Astropart. Phys.*, 6, 025
 - [10] Marsh D. J. E., Ferreira P. G., 2010, *Phys. Rev. D*, 82, 103528
 - [11] Membrado M., Pacheco A. F., Sañudo J., 1989, *Phys.Rev.A*, 39, 4207
 - [12] Peebles P. J. E., 1982, *ApJ*, 263, L1.
 - [13] White S. D. M., Frenk C. S., Davis M. et al., 1987, *ApJ*, 313, 505.
 - [14] Bullock J. S., Boylan-Kolchin M., 2017, *ARA&A*, 55, 343
 - [15] Clowe D., Bradac M., Gonzalez A. H. et al., 2006, *ApJ*, 648, 2.
 - [16] Klypin A., Kravtsov A. V., Valenzuela O. et al., 1999, *ApJ*, 522, 82.
 - [17] Moore B., Ghigna S., Governato F. et al., 1999, *ApJ*, 524, L19.
 - [18] Penny S., Conselice C. J., De Rijcke S. et al., 2009, *MN-RAS*, 393, 1054.
 - [19] J. Magaña and T. Matos, *Journal of Physics: Conference Series*, Volume 378, conference 1

- [20] A. Suárez, V. H. Robles, and T. Matos, *Astrophys. Space Sci. Proc.* 38 , 107 (2014), 1302.0903.
- [21] Marsh D. J. E., 2016, *Phys. Rep.*, 643, 1
- [22] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, *Phys. Rev. D* 95 , 043541 (2017).
- [23] K. Enqvist and M. S. Sloth, *Nucl. Phys. B* 626 (2002) 395 doi:10.1016/S0550-3213(02)00043-3 [hep-ph/0109214]
- [24] D. H. Lyth and D. Wands, *Phys. Lett. B* 524 (2002) 5 doi:10.1016/S0370-2693(01)01366-1 [hep-ph/0110002]
- [25] T. Moroi and T. Takahashi, *Phys. Lett. B* 522 (2001) 215 [hep-ph/0110096]
- [26] Christian T. Byrnes, David Wands; *Phys. Rev. D* 74 (2006) 043529; DOI: 10.1103/PhysRevD.74.043529; arXiv:astro-ph/0605679v3
- [27] D. Barkats et al. (BICEP1), *Astrophys. J.* 783 , 67 (2014), arXiv:1310.1422 [astro-ph.C]
- [28] P. A. R. Ade et al. (BICEP2, Planck), *Phys. Rev. Lett.* 114 , 101301 (2015), arXiv:1502.00612 [astro-ph.CO]
- [29] P. A. R. Ade et al. (BICEP2, Keck Array), *Phys. Rev. Lett.* 116 , 031302 (2016), arXiv:1510.09217 [astro-ph.CO]
- [30] D. Lyth, *Physics Letters B* 147 , 403 (1984)
- [31] D. H. Lyth and E. D. Stewart, *Physics Letters B* 283 , 189 (1992).
- [32] V. Vennin, K. Koyama and D. Wands, *Inflation with an extra light scalar field after Planck* , *JCAP* 03 (2016) 024 [arXiv:1512.03403] [IN SPIRE].
- [33] K. Enqvist and T. Takahashi, *Journal of Cosmology and Astroparticle Physics* 2013 , 034 17 (2013), arXiv:1306.5958.
- [34] E. Madelung, *Zeit. F. Phys.* 40 , 322 (1927)
- [35] V.A. Belinsky, L.P. Grishchuk, I.M. Khalatnikov, and Ya.B. Zeldovich, *Phys. Lett. B* 155 , 232 (1985)
- [36] T. Piran and R.M. Williams, *Phys. Lett. B* 163 , 331 (1985)
- [37] B. Li, T. Rindler-Daller, and P.R. Shapiro, *Phys. Rev. D* 89 , 083536 (2014)
- [38] A. Suárez and P. H. Chavanis, *Phys. Rev. D* 95 , 063515 (2017) .
- [39] A. Suárez and P.H. Chavanis, *Phys. Rev. D* 92 , 023510 (2015)
- [40] A. Arbey, J. Lesgourgues, and P. Salati, *Phys. Rev. D* 65 , 083514 (2002)
- [41] J.-A. Gu and W.-Y.P. Hwang, *Phys. Lett. B* 517 , (2001)
- [42] K. Harigaya, M. Ibe, M. Kawasaki and T. T. Yanagida, *Phys. Rev. D* 87 , 063514 (2013) [arXiv:1211.3535 [hep-ph]].
- [43] P. A. R. Ade et al. [Planck Collaboration], *Astron. Astrophys.* 594 , A13 (2016) [arXiv:1502.01589 [astro-ph.CO]].
- [44] T. Kobayashi, R. Murgia, A. De Simone, V. Iršič, and M. Viel, *Phys. Rev. D* 96 , 123514 (2017) .
- [45] L. Husdal, (2016), 1609.04979.
- [46] L. Visinelli, *Phys. Rev. D* 96, 023013 (2017).
- [47] P.H. Chavanis, M. Lemou, and F. M e hats, *Phys. Rev. D* 12 , 123527 (2015)
- [48] D. H. Lyth and A. R. Liddle, *The primordial density perturbation; cosmology inflation and the origin of structure* , 2009, Cambridge University Press.
- [49] S.W. Randall, M. Markevitch, D. Clowe, A.H. Gonzalez, 42 and M. Bradac, *Astrophys. J.* 679 , 1173 (2008)
- [50] T. Moroi and T. Takahashi, *Phys. Rev. D* 66 , 063501 (2002) [arXiv:hep-ph/0206026].
- [51]] D. H. Lyth, C. Ungarelli and D. Wands, *Phys. Rev. D* 67 , 023503 (2003) [arXiv:astro-ph/0208055]
- [52] D. H. Lyth and D. Wands, *Phys. Rev. D* 68 , 103516 (2003) [arXiv:astro-ph/0306500].
- [53] C. He, D. Grin and W. Hu, *Phys. Rev. D* 92 , no. 6, 063018 (2015), arXiv:1505.00639.
- [54] D. Langlois and F. Vernizzi, *Mixed inflaton and curvaton perturbations* , *Phys. Rev. D* 70 (2004) 063522 [astro-ph/0403258] [IN SPIRE].
- [55] T. Moroi, T. Takahashi and Y. Toyoda, *Relaxing constraints on inflation models with curvaton* , *Phys. Rev. D* 72 (2005) 023502 [hep-ph/0501007] [IN SPIRE]
- [56] T. Moroi and T. Takahashi, *Implications of the curvaton on inflationary cosmology* , *Phys. Rev. D* 72 (2005) 023505 [astro-ph/0505339] [IN SPIRE].
- [57] K. Ichikawa, T. Suyama, T. Takahashi and M. Yamaguchi, *Non-Gaussianity, Spectral Index and Tensor Modes in Mixed Inflation and Curvaton Models* , *Phys. Rev. D* 78 (2008) 023513 [arXiv:0802.4138] [IN SPIRE].
- [58] T. Suyama, T. Takahashi, M. Yamaguchi and S. Yokoyama, *On Classification of Models of Large Local-Type Non-Gaussianity* , *JCAP* 12 (2010) 030 [arXiv:1009.1979] [IN SPIRE].
- [59] K. Enqvist and T. Takahashi, *Mixed inflaton and spectator field models after Planck* , *JCAP* 10 (2013) 034 [arXiv:1306.5958] [IN SPIRE].
- [60] A. D. Linde and V. F. Mukhanov, *Phys. Rev. D* 56 , 535 (1997) [astro-ph/9610219]
- [61] Hlozek R., Grin D., Marsh D. J. E., Ferreira P. G., 2015, *Phys. Rev. D*, 91, 103512
- [62] Paredes A., Michinel H., 2016, *Physics of the Dark Universe*, 12, 50
- [63] <https://arxiv.org/pdf/1609.05856.pdf>
- [64] Lora V., Magaña J., Bernal A., Sánchez-Salcedo F. J., Grebel E. K., 2012, *J. Cosmology Astropart. Phys.*, 2, 11
- [65] Calabrese E., Spergel D. N., 2016, *MNRAS*, 460, 4397
- [66] Sarkar A., Mondal R., Das S., Sethi S. K., Bharadwaj S., Marsh D. J. E., 2016, *J. Cosmology Astropart. Phys.*, 4, 012
- [67] Armengaud E., Palanque-Delabrouille N., Y'èche C., Marsh D. J. E., Baur J., 2017, preprint, (arXiv:1703.09126)
- [68] Iršič V., Viel M., Haehnelt M. G., Bolton J. S., Becker G. D., 2017, preprint, (arXiv:1703.04683)
- [69] <https://arxiv.org/pdf/1404.1054.pdf>
- [70] C.G. Boehmer and T. Harko, ?Can dark matter be a Bose-Einstein condensate?,? *JCAP* 0706 025 (2007) [arXiv:0705.4158[astro-ph]]
- [71] A. Arbey, J. Lesgourgues and P. Salati, ?Galactic halos of fluid dark matter,? *Phys. Rev. D* 68 023511 (2003) [astro-ph/0301533]
- [72] T. Harko, ?Bose-Einstein condensation of dark matter solves the core/cusp problem,? *JCAP* 1105 022 (2011) [arXiv:1105.2996[astro-ph.CO]]
- [73] V.H. Robles and T. Matos, ?Flat central density profile and constant DM surface density in galaxies from Scalar Field Dark Matter,? *Mon. Not. Roy. Astron. Soc.* 422 282-289 (2012) [arXiv:1201.3032[astro-ph.CO]]
- [74] A. Arbey, J. Lesgourgues and P. Salati, ?Cosmological constraints on quintessential halos,? *Phys. Rev. D* 65 083514 (2002) [astro-ph/0112324]