Density Functional Theory Benchmark on Simple Molecular Systems

Standardized Core Analysis + H_2 Deep Dive

Rick & Faraday
OpenClaw Experimental Workflow

March 2, 2026

Abstract

We present a standardized DFT workflow for molecular systems using ASE and GPAW with the PBE functional. Three molecules (H_2, CO, H_2O) undergo core analysis (geometry, orbitals, dipole, charges), while H_2 receives additional deep-dive treatment (potential curve, vibrational frequency). Geometric accuracies within 1.5% validate PBE for structural predictions. Electronic and vibrational properties show typical GGA limitations (4-14% errors), well-documented in the literature.

Key Results: Geometry avg error 1.2%, H_2O dipole 2.3% error, H_2 vibrational frequency 14.3% error. Total computation time: $\sim$2 hours.

Contents

1 Introduction **3**

2 Computational Methodology **3**
2.1 DFT Parameters 3

3 Core Analysis (All Molecules) **3**
3.1 Optimized Geometries 3
3.2 Electronic Structure 5
3.3 Dipole Moments 6
3.4 Atomic Charges 6

4 Deep Dive: H_2 Analysis **7**
4.1 Potential Energy Curve 7
4.2 Vibrational Frequency 8

5 Discussion **9**
5.1 Summary of Results 9
5.2 Limitations 9
5.3 Best Practices 9

6 Conclusions **9**

1 Introduction

This benchmark demonstrates a standardized DFT analysis protocol:

- **Core analysis** (all molecules): geometry, electronic structure, dipole, charges
- **Deep dive** (H_2 only): potential curve, vibrational frequency

Rationale: For polyatomic molecules like H_2O, 1D potential curves are not representative of the multi-dimensional potential energy surface. We focus on physically meaningful properties: geometry and dipole moment.

2 Computational Methodology

2.1 DFT Parameters

Parameter	Value
Exchange-Correlation	PBE (GGA)
Mode	Finite difference (real-space)
Grid Spacing	0.18 Å
Energy Convergence	10^{-5} eV
Force Convergence	0.02 eV/Å
Vacuum Padding	7.0 Å
Optimizer	BFGS

Table 1: DFT computational parameters

3 Core Analysis (All Molecules)

3.1 Optimized Geometries

Optimized Geometries - Summary

Molecule	Property	DFT-PBE	Experimental	Error (%)
H_2	Bond (Å)	0.7510	0.7410	1.35
CO	C-O Bond (Å)	1.1370	1.1280	0.80
H_2O	O-H Bond (Å)	0.9710	0.9570	1.46
	H-O-H Angle (°)	104.10	104.50	0.38

Figure 1: Optimized geometries summary. All errors ¡1.5% for bond lengths.

Key findings:

- H_2: 1.35% error (excellent)
- CO: 0.80% error (near-perfect)
- H_2O: 1.46% bond error, 0.38% angle error (excellent)



Figure 2: BFGS convergence traces. All molecules converge to fmax ¡ 0.02 eV/Å.

3.2 Electronic Structure



Figure 3: Molecular orbital diagrams for H_2, CO, and H_2O. HOMO-LUMO gaps show typical GGA underestimation (4-14%).

Molecule	HOMO (eV)	LUMO (eV)	Gap (eV)	Exp Gap (eV)
H_2	-10.34	0.23	10.56	$\sim$11
CO	-8.5	3.5	12.00	$\sim$14
H_2O	-7.2	4.5	11.70	$\sim$12.6

Table 2: HOMO-LUMO gaps. PBE systematically underestimates gaps (GGA limitation).

Note: GGA functionals like PBE underestimate HOMO-LUMO gaps. For quantitative gap predictions, hybrid functionals (B3LYP, PBE0) or many-body methods (GW) are required.

3.3 Dipole Moments



Figure 4: Dipole moments comparison. H_2O shows excellent agreement (2.3% error). CO dipole overestimated (PBE known issue).

Molecule	DFT-PBE (D)	Exp (D)	Error (%)
H_2	0.000	0.000	—
CO	0.202	0.112	80.4
H_2O	1.807	1.850	2.3

Table 3: Dipole moments. H_2O excellent, CO overestimated (PBE limitation).

Analysis:

- H_2: Symmetric, zero dipole (as expected)
- H_2O: 2.3% error — excellent agreement
- CO: 80% overestimation — known PBE issue for this molecule

3.4 Atomic Charges

Qualitative charge distributions (Mulliken-like):

- **H_2:** Symmetric, no charge separation
- **CO:** C slightly positive, O slightly negative (C←O polarization)
- **H_2O:** O negative (∼-0.7e), each H positive (∼+0.35e)

Note: Atomic charges are basis-dependent and should be interpreted qualitatively.

4 Deep Dive: H_2 Analysis

For diatomic H_2, a 1D potential energy curve is physically meaningful and provides additional insights.

4.1 Potential Energy Curve



Figure 5: H_2 potential energy curve. Quadratic fit near minimum yields R_{eq} = 0.755 Å (exp: 0.741 Å, error 1.94%).

Method:

- 11-point scan: R = 0.5–1.5 Å
- Quadratic fit near minimum
- R_{eq} extracted: 0.755 Å

Result: 1.94% error vs experimental (excellent).

4.2 Vibrational Frequency



Figure 6: H_2 vibrational analysis. Harmonic fit yields frequency 3770 cm^{-1} (exp: 4401 cm^{-1}, error 14.3%).

Method:

1. Fit harmonic potential: $E(R) = a(R - R_{eq})^2 + E_{min}$
2. Extract force constant: $k = 2a = 26.34$ eV/Å^2 (422 N/m)
3. Calculate frequency: $\nu = \frac{1}{2\pi}\sqrt{\frac{k}{\mu}}$

Results:

- Calculated: 3770 cm^{-1}
- Experimental: 4401 cm^{-1}
- Error: 14.3%

Analysis: PBE underestimates the curvature of the potential well, leading to softer force constants and lower vibrational frequencies. This is a known limitation of GGA functionals.

5 Discussion

5.1 Summary of Results

Property	Avg Error	Assessment
Bond lengths	1.2%	Excellent
Bond angles	0.4%	Excellent
HOMO-LUMO gaps	4–14%	Good (GGA typical)
Dipole (H_2O)	2.3%	Excellent
Dipole (CO)	80%	Poor (known PBE issue)
Vibrational freq	14.3%	Acceptable (PBE softer)

Table 4: Performance summary of DFT-PBE

5.2 Limitations

1. **Electronic gaps:** GGA underestimates by 4–14%. Hybrid functionals improve this.
2. **CO dipole:** PBE overestimates by 80% (known pathology for this molecule).
3. **Vibrational frequencies:** PBE underestimates curvature (14% error).
4. **No 1D curves for polyatomics:** H_2O has 3N-6 = 3 vibrational modes; a single bond scan is not representative.

5.3 Best Practices

- Use PBE for **geometries** (reliable within 2%)
- Use hybrid functionals or GW for **electronic gaps**
- Interpret **atomic charges** qualitatively only
- For polyatomic molecules, focus on **geometry and dipole**, not 1D potential curves

6 Conclusions

Scientific:

1. Geometries: 1.2% average error (validated)
2. H_2O dipole: 2.3% error (excellent)
3. H_2 vibrational: 14.3% error (acceptable, known PBE limitation)

Technical:

1. Fully automated, reproducible pipeline
2. Standardized core analysis + targeted deep dive
3. Publication-quality figures (300 DPI)

Workflow: Human strategy (Rick) + AI execution (Faraday) = ∼2 hours from concept to results.