

Lecture Notes on Exercise Course: Calculus A(I)

Wei Wang

Contents

1	Some rules	5
1.1	Submission rules	5
2	Preliminaries of calculus	7
2.1	Trigonometry	7
2.2	Some useful inequalities	7
2.3	Real numbers	9
3	Limits of sequences	11
3.1	Comparing the order	11
3.2	Basic properties on limits	12
3.3	Nonexistence of limits	12
3.4	Monotone sequence	12
3.5	Boundedness	12
3.6	Some important sequence and limits	13
3.7	Sandwich theorem	13
3.8	Cauchy's proposition	13
3.9	Limits with respect to the definition of e	14
3.10	Problems associated with fixed points	15
4	Limits of functions	17
4.1	Heine criterion	17
4.2	Riemannian function	17
4.3	Some basic calculations	18
4.4	Two important limits	18
4.5	Order analysis	18
5	Continuous function	21
5.1	Discontinuity	21
5.2	Uniqueness	21
5.3	Intermediate value theorem	21
5.4	Contraction maps	22
5.5	Boundedness, maximum, and minimum	23
5.6	Monotone functions	23
6	Differentiation	25
6.1	Definitions	25
6.2	Some tricks	26

7	Indefinite integral	27
7.1	Pairing method	27
7.2	By iterations	27
7.3	Some special integrals	28
8	Definite integral	31
8.1	Calculate limits	31
8.2	Divide into two parts	31
8.3	Approximation and Riemann-Lebesgue Lemma	32
8.4	Calculations by symmetry	32
8.5	Mean value theorem	32
8.6	Some inequalities	33
8.7	Combination of previous topics	34
9	Review of the midterm exam	35
9.1	Some questions about sequences	35
9.2	Some questions about integrals	35
10	Mean value theorem of derivatives	37
10.1	Constructing auxiliary functions	37
10.2	Related calculations	38
10.3	Comprehensive applications	38
11	L'Hospital's rule	41
12	Taylor expansion and further applications of MVT	43
12.1	The property of best approximation	43
12.2	Asymptotic behavior	43
12.3	Existences of particular points	44
12.4	Basic calculations	46
12.5	Domination and zero functions	47
13	Application of derivatives	49
13.1	Monotonicity and zero points	49
13.2	Uniqueness of ODE	50
13.3	Basic calculations	50
13.4	Proof of inequalities	50
14	Calculus with multi-variables I	51
15	Calculus with multi-variables II	55
15.1	Homogeneous functions	55
15.2	Nondegeneracy	55
15.3	Reverse problems	56
15.4	The implicit function theorem	56

Chapter 1

Some rules

1.1 Submission rules

The exercise course is on every Tuesday. In this course, you should submit the homework assigned last Tuesday and the Thursday before the last Thursday. For example, suppose that you submit the homework on September 10, then you have to submit those assigned on September 3 and August 29.

An exception is that on October 8, you must submit the homework assignment before September 24 (including this day). After that, you should follow the above rules.

You can send your homework to the email:2201110024@stu.pku.edu.cn, or write on paper to submit it to me on the exercise course. If you want to send it by email, you should name the title of this email with "name+student ID+the number of the homework." For example, "Zhang San 0000000000 Homework 1" in Chinese.

Grade of homework

We only care about whether you submit and complete the homework or not. In particular, if submit all the homework and complete all the problems assigned, then you will get full marks. Regular grades only come from the submission of homework.

Some useful links

We present some useful links associated with calculus.

Lecture notes by Yantong Xie:<https://darkoxie.github.io>

Mathstackexchange: <https://math.stackexchange.com>

Chapter 2

Preliminaries of calculus

2.1 Trigonometry

We introduce some results in trigonometry.

Some basic formulae

$$\begin{aligned}\sin a + \sin b &= 2 \sin \frac{a+b}{2} \cos \frac{a-b}{2}, \\ \sin a \cos b &= \frac{1}{2}(\sin(a+b) + \sin(a-b)).\end{aligned}$$

Exercise 2.1. What about $\sin a - \sin b$, $\cos a \pm \cos b$, $\sin a \sin b$, $\cos a \cos b$?

Reverse functions

Consider $f(x) = \sin x$. We see that it is increasing in $[-\pi/2, \pi/2]$ and we can define the reverse function of it in this interval by $f^{-1}(x) = \arcsin x$, where $x \in [-1, 1]$.

Exercise 2.2. Consider reverse functions of $\cos x$, $\tan x$.

Euler formula

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Exercise 2.3. Calculate $\sum_{k=1}^n \cos kx$.

2.2 Some useful inequalities

Cauchy's inequality

Theorem 2.4. For $\{a_i\}_{i=1}^n, \{b_i\}_{i=1}^n \subset \mathbb{R}$, we have

$$\left(\sum_{i=1}^n a_i b_i \right)^2 \leq \left(\sum_{i=1}^n a_i^2 \right) \left(\sum_{i=1}^n b_i^2 \right).$$

Proof. Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$. Consider $|a + tb|^2$ with $t \in \mathbb{R}$. □

Question 2.5. What is the condition such that the above inequality satisfies “=”?

Proposition 2.6. Let $a, b, c > 0$ be such that $a + b + c = 1$. We have

$$a^3 + b^3 + c^3 \geq \frac{a^2 + b^2 + c^2}{3}.$$

Proof. Applying Theorem 2.4, we have

$$(a^2 + b^2 + c^2)^2 \leq (a^3 + b^3 + c^3)(a + b + c) = a^3 + b^3 + c^3.$$

Additionally,

$$a^2 + b^2 + c^2 \geq ab + ac + bc.$$

This implies that $3(a^2 + b^2 + c^2) = (a + b + c)^2 = 1$. Combining all the above properties, the result follows directly. \square

Hölder's inequality

Theorem 2.7. For $\{a_i\}_{i=1}^n, \{b_i\}_{i=1}^n \subset \mathbb{R}_+$, we have

$$\left(\sum_{i=1}^n a_i b_i \right) \leq \left(\sum_{i=1}^n a_i^p \right)^{1/p} \left(\sum_{i=1}^n b_i^q \right)^{1/q},$$

where $1/p + 1/q = 1$ with $p, q > 0$.

Proof. It is a corollary of Yong's inequality, that is, for $a, b > 0$, $p, q > 0$ with $1/p + 1/q = 1$, we have $ab \leq a^p/p + b^q/q$. \square

Bernoulli's inequality

Theorem 2.8. Let $n \geq 2$. Assume that $x_1, x_2, \dots, x_n > -1$ and have the same sign. Then

$$\prod_{i=1}^n (1 + x_i) \geq 1 + \sum_{i=1}^n x_i.$$

Proof. The proof is by using induction. For $n = 1$, there is nothing to prove. Assume that the result is true for n . By simple calculations, we have

$$\begin{aligned} \prod_{i=1}^n (1 + x_i) &= (1 + x_1) \left(\prod_{i=2}^n (1 + x_i) \right) \\ &\geq (1 + x_1) \left(1 + \sum_{i=2}^n x_i \right) \\ &= 1 + \sum_{i=1}^n x_i + x_1 \left(\sum_{i=2}^n x_i \right) \\ &\geq 1 + \sum_{i=1}^n x_i, \end{aligned}$$

where for the last inequality, we have used the property that x_i has the same sign. Now, we complete the proof. \square

Exercise 2.9. Let $m \in \mathbb{R}$ and $x > -1$. Show that if $m \in [0, 1]$, then $(1+x)^m \leq 1+mx$ and if $m < 0$ or $m > 1$, then $(1+x)^m \geq 1+mx$.

Exercise 2.10. Show that if $b > a > 0$, and $n \in \mathbb{Z}_+$, then $a^{n+1} > b^n((n+1)a - nb)$, and $b^{n+1} > a^n((n+1)b - na)$.

Proposition 2.11. *The sequence $\{(1+1/n)^n\}$ is increasing.*

Proof. Let $b = 1 + 1/n$ and $a = 1 + 1/(n+1)$, the result follows from Exercise 2.10. \square

Exercise 2.12. Show that the sequence $\{(1+1/n)^{n+1}\}$ is decreasing.

2.3 Real numbers

Density of real numbers

One of the most remarkable properties of the real numbers is that \mathbb{R} is dense. We can see from the following proposition.

Proposition 2.13. *Let $a, b \in \mathbb{R}$. Show that there exists $r_1 \in \mathbb{Q}$ and $r_2 \in \mathbb{R} \setminus \mathbb{Q}$ such that $a < r_1 < r_2 < b$.*

Proof. Assume that $a \in \mathbb{Q}$. Let $N \gg 1$ be such that $1/N < \pi/N < b - a$, we choose $r_1 = a + 1/N$ and $r_2 = a + \pi/N$. Assume that $a \in \mathbb{R} \setminus \mathbb{Q}$, we choose $N \gg 1$ such that $N(b - a) > 10$. As a result, there must be some $n \in \mathbb{Z}_+$ such that $n \in (Na, Nb)$, and then $n/N \in (a, b)$. Taking n/N as new a and applying the previous arguments, we complete the proof. \square

Closeness of calculations for real numbers

Proposition 2.14. *There exist $a, b \in \mathbb{R} \setminus \mathbb{Q}$ such that $a^b \in \mathbb{Q}$.*

Proof. Consider $a = b = \sqrt{2}$. If $a^b \in \mathbb{Q}$, then we are done. If not, we see that $(a^b)^{\sqrt{2}} = 2$. \square

Exercise 2.15. Give the corresponding examples.

- $a, b \in \mathbb{Q}, a^b \in \mathbb{Q}$.
- $a, b \in \mathbb{Q}, a^b \in \mathbb{R} \setminus \mathbb{Q}$.
- $a, b \in \mathbb{R} \setminus \mathbb{Q}, a + b \in \mathbb{Q}$.

Chapter 3

Limits of sequences

3.1 Comparing the order

Let $a > 0$ and $b > 1$, we have

$$\ln n \ll n^a \ll b^n \ll n! \ll n^n, \quad (3.1)$$

where we call $f(n) \ll g(n)$ if

$$\lim_{n \rightarrow +\infty} \frac{f(n)}{g(n)} = 0.$$

Now, let us show (3.1). We note that

$$0 < \frac{n!}{n^n} \leq \frac{1}{n},$$

which implies that $n! \ll n^n$. Let $x_n = b^n/(n!)$. We have

$$\frac{b_{n+1}}{b_n} = \frac{b}{n+1}.$$

As a result, $b^n \ll n!$ follows from the following lemma, whose proof is left for the reader.

Lemma 3.1. *Let $\{x_n\}$ be a sequence, if*

$$\lim_{n \rightarrow +\infty} \left| \frac{x_{n+1}}{x_n} \right| < 1,$$

then $\lim_{n \rightarrow +\infty} x_n = 0$.

Denote $y_n = n^a/b^n$. By using this lemma, we obtain that $n^a \ll b^n$. Finally, we turn to the proof of $\ln n \ll n^a$. Noting that

$$\frac{\ln n}{n^a} = \frac{1}{a} \cdot \frac{\ln n^a}{n^a},$$

we only show that $\ln n/n \rightarrow 0^+$. It is a direct consequence of the following exercise.

Exercise 3.2. $\lim_{n \rightarrow +\infty} n^{1/n} = 1$.

Hint: Let $n^{1/n} = 1 + x_n$. We have

$$n = (1 + x_n)^n \geq 1 + nx_n + \frac{n(n-1)}{2}x_n^2.$$

By using Lemma 3.1, we can show $\lim_{n \rightarrow +\infty} n^2 x_n^2 = 0$ (in Homework).

3.2 Basic properties on limits

3.3 Nonexistence of limits

Proposition 3.3. $\lim_{n \rightarrow +\infty} x_n = x$ if and only if for any subsequence x_{n_k} , we have $\lim_{k \rightarrow +\infty} x_{n_k} = x$.

Proof. It follows from the definition of the limit. \square

We can use this proposition to show that some sequences diverge.

Exercise 3.4. $(-1)^n$ diverges.

Exercise 3.5. For $\{x_n\}$, we have $x_{2k} \rightarrow a$ and $x_{2k-1} \rightarrow b$, where $a, b \in [-\infty, +\infty]$. Show that x_n is convergent if and only if $a = b \in (-\infty, +\infty)$.

3.4 Monotone sequence

Example 3.6. Let $n \in \mathbb{Z}_+$. Assume that $x_n \in [0, 1]$ is the root of $\sum_{i=1}^n x^i = 1$. Show that $\lim_{n \rightarrow +\infty} x_n = \frac{1}{2}$.

Proof. It is easy to see that x_n is decreasing. On the other hand, $x_n - x_n^{n+1} = 1 - x_n$. Letting $n \rightarrow +\infty$, we see that $x_n \rightarrow \frac{1}{2}$. \square

Exercise 3.7. Assume that $0 < b < a$. We let $\{a_n\}, \{b_n\}$ be defined as follows.

$$a_{n+1} = \frac{a_n + b_n}{2}, \quad b_{n+1} = \sqrt{a_n b_n}, \quad n \in \mathbb{Z}_+,$$

where $a_1 = a, b_1 = b$. Show that $\lim_{n \rightarrow +\infty} a_n = \lim_{n \rightarrow +\infty} b_n$ and exist.

Hint: Show that a_n and b_n are monotone.

Exercise 3.8. $\{x_n\}$ is monotone. Show that $\{x_n\}$ is monotone if and only if it has a convergent subsequence.

3.5 Boundedness

Any convergent sequence is bounded.

Example 3.9. Assume that $x_n \rightarrow x > 0$. Then x_n has a lower bound but may not have a minimal point.

Proof. For n sufficiently large, we have $x_n > \frac{x}{2}$. Considering $x_n = 1 + \frac{1}{n}$, we see that it does not have a minimal point. \square

Exercise 3.10. Assume that $x_n \rightarrow +\infty$, then there must be some $n_0 \in \mathbb{Z}_+$ such that $x_{n_0} = \min_{n \in \mathbb{Z}_+} \{x_n\}$.

3.6 Some important sequence and limits

Example 3.11. Show that

$$S_n = \sum_{i=1}^n \frac{1}{i^p}$$

Is convergent if and only if $p > 1$.

Proof. Taking $p = 1$ and $p = 2$ for example. For $p = 1$, since

$$\ln(n+1) - \ln(n) < \frac{1}{n},$$

S_n diverges. For $p = 2$, we see that

$$\frac{1}{n^2} < \frac{1}{n-1} - \frac{1}{n},$$

and then S_n is convergent. □

Example 3.12. Consider

$$x_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n}.$$

Show that $x_n \rightarrow \ln 2$.

Proof. Let

$$\gamma_n = \sum_{i=1}^n \frac{1}{i} - \ln n.$$

We can show that $\gamma_n \rightarrow \gamma$ (Euler constant). Then, the result follows directly. □

3.7 Sandwich theorem

Recall that if $a_n \leq b_n \leq c_n$ and $a_n, c_n \rightarrow x$, then $b_n \rightarrow x$.

Example 3.13. Assume that $0 < a_1 \leq a_2 \leq \dots \leq a_k$. Calculate $\lim_{n \rightarrow +\infty} (a_1^n + \dots + a_k^n)^{1/n}$.

Proof. We see that

$$a_k \leq (a_1^n + \dots + a_k^n)^{1/n} \leq k^{1/n} a_k.$$

Then $\lim_{n \rightarrow +\infty} (a_1^n + \dots + a_k^n)^{1/n} = a_k$. □

Exercise 3.14. Assume that $0 < a_1 \leq a_2 \leq \dots \leq a_k$. Calculate $\lim_{n \rightarrow +\infty} (a_1^{1/n} + \dots + a_k^{1/n})^n$.

Exercise 3.15. Assume that $a_k \rightarrow a > 0$. Calculate $\lim_{n \rightarrow +\infty} (a_1^n + \dots + a_n^n)^{1/n}$.

3.8 Cauchy's proposition

It is about the limit of average for a sequence having a limit.

Proposition 3.16. If $\lim_{n \rightarrow +\infty} x_n = x$, then

$$\lim_{n \rightarrow +\infty} \frac{\sum_{i=1}^n x_i}{n} = x. \tag{3.2}$$

Proof. For any $\varepsilon > 0$, we can choose $N \in \mathbb{Z}_+$ such that $|x_n - x| < \varepsilon$ for any $n > N$. This implies that if $n > N$, there holds

$$\left| \frac{\sum_{i=1}^N x_i}{n} - x \right| \leq \frac{\sum_{i=1}^N |x_i|}{n} + \frac{n - N}{n} \varepsilon.$$

Choosing larger N without changing the notation, we can assume that if $n > N$, then

$$\left| \frac{\sum_{i=1}^N x_i}{n} - x \right| < 2\varepsilon.$$

As a result, we have (3.2). □

Exercise 3.17. Assume that $a_n \rightarrow a$ and $b_n \rightarrow b$. Show that

$$\lim_{n \rightarrow +\infty} \frac{\sum_{i=1}^{n-1} a_i b_{n-i}}{n} = ab.$$

Exercise 3.18. Assume that $x_n \rightarrow x$. Show that

$$\lim_{n \rightarrow +\infty} \frac{\sum_{k=0}^n C_n^k x_k}{2^n} = x$$

Exercise 3.19 (Stolz $\frac{0}{\infty}$). Let $\{a_n\}$ and $\{b_n\}$ be two sequences. Assume that $\{b_n\}$ is a strictly monotone and divergent sequence (namely, strictly increasing and approaching $+\infty$, or strictly decreasing and approaching $-\infty$) and the following limit exists

$$\lim_{n \rightarrow +\infty} \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = l.$$

Then

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = l.$$

Exercise 3.20 (Stolz $\frac{0}{0}$). Let $\{a_n\}$ and $\{b_n\}$ be two sequences. Assume now that $a_n, b_n \rightarrow 0$ and $\{b_n\}$ is strictly decreasing. If

$$\lim_{n \rightarrow +\infty} \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = l.$$

Then

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = l.$$

3.9 Limits with respect to the definition of e

We know that

$$e = \lim_{n \rightarrow +\infty} (1 + 1/n)^n. \tag{3.3}$$

By this, we can consider related limits.

Question 3.21. What about the limit $\lim_{n \rightarrow -\infty} (1 + 1/n)^n$.

Example 3.22. Show that

$$\lim_{n \rightarrow +\infty} \left(1 + \frac{1}{n} + \frac{1}{n^2} \right)^n = e$$

Proof. It follows from (3.3) that

$$\lim_{n \rightarrow +\infty} \left(1 + \frac{1}{n} + \frac{1}{n^2}\right)^n = \lim_{n \rightarrow +\infty} \left(1 + \frac{n+1}{n^2}\right)^{\frac{n^2}{n+1} \cdot \frac{n+1}{n}} = e.$$

□

Example 3.23. Calculate the following limits.

$$\lim_{n \rightarrow +\infty} \left(1 - \frac{1}{n}\right)^{n^2}, \quad \lim_{n \rightarrow +\infty} \left(1 - \frac{1}{n^2}\right)^n.$$

Proof. Noting that

$$\left(1 - \frac{1}{n}\right)^{n^2} = \left[\left(1 - \frac{1}{n}\right)^n\right]^n$$

and $\lim_{n \rightarrow +\infty} (1 - 1/n)^n = 1/e$, we see that the first limit is 0. On the other hand, we obtain that

$$\left(1 - \frac{1}{n^2}\right)^n = \left[\left(1 - \frac{1}{n^2}\right)^{n^2}\right]^{1/n}.$$

The term in $[\cdot]$ is bounded, and the limit is 1. □

Exercise 3.24. Calculate $\lim_{n \rightarrow +\infty} (1 + 1/n)^{n^2}$ and $\lim_{n \rightarrow +\infty} (1 + 1/n^2)^n$.

Exercise 3.25. If there exists $\{x_n\}$ such that $\lim_{n \rightarrow +\infty} x_n = 1$ and $\lim_{n \rightarrow +\infty} x_n^n = 1.001$.

Hint: Consider $x_n = 1 + \frac{a}{n}$.

Exercise 3.26. Find $\{x_n\}$ such that the following properties hold.

- (1) $\lim_{n \rightarrow +\infty} (x_n - e^n) = 0$.
- (2) $\lim_{n \rightarrow +\infty} (f(x_n) - f(e^n)) \neq 0$, where $f(x) = x \ln x$.

Hint: Consider $y_n = x_n - e^n$.

3.10 Problems associated with fixed points

Consider a sequence $\{x_n\}$ such that $x_{n+1} = f(x_n)$. If x_n is convergent with $\lim_{n \rightarrow +\infty} x_n = x$ and f is continuous, then $f(x) = x$.

Example 3.27. Let $x_{n+1} = \sqrt{2 + x_n}$ and $x_1 = 1$. Show that $\lim_{n \rightarrow +\infty} x_n = 2$.

Proof. 2 is the fixed point. We have

$$\frac{x_{n+1} - 2}{x_n - 2} = \frac{1}{2 + \sqrt{x_n + 2}}.$$

Then

$$\left| \frac{x_{n+1} - 2}{x_n - 2} \right| < \frac{1}{2}.$$

□

Exercise 3.28. Let $x_{n+1} = \frac{1}{x_n^{1/2}}$ and $x_1 = 2$. Show that $\lim_{n \rightarrow +\infty} x_n = 1$.

Exercise 3.29. Let $x_{n+1} = \frac{1}{x_n^2}$ and $x_1 = 2$. Show that x_n is divergent.

Exercise 3.30. Let $x_{n+1} = \frac{1}{x_n}$ and $x_1 = 2$. Show that x_n is divergent.

Question 3.31. Can you find something interesting in the above three exercises?

Exercise 3.32. $x_1 = \sqrt{2}$. $x_{n+1} = \sqrt{2^{x_n}}$. Consider $\lim_{n \rightarrow +\infty}$.

Hint: Draw the picture.

Chapter 4

Limits of functions

4.1 Heine criterion

Proposition 4.1. Assume that $f : [a, b] \rightarrow \mathbb{R}$ is a function. For $c \in (a, b)$, $\lim_{x \rightarrow c} f(x) = A$, if and only if for any $x_n \rightarrow c$, there holds $f(x_n) \rightarrow A$ as $n \rightarrow +\infty$.

Proof. The proof is due to the definition. \square

Exercise 4.2. Use this Heine criterion and Cauchy criterion of sequences to show the Cauchy criterion of functions.

Example 4.3. Show that the Dirichlet function does not have limits at any points in $[0, 1]$. Here Dirichlet function $D(x) : [0, 1] \rightarrow \{0, 1\}$ is defined by

$$D(x) = 1 \text{ if } x \in \mathbb{Q}, \quad D(x) = 0 \text{ if } x \notin \mathbb{Q}$$

Exercise 4.4. Show that the Dirichlet function can be represented by

$$D(x) = \lim_{m \rightarrow \infty} \left\{ \lim_{n \rightarrow \infty} [\cos(\pi m! x)]^{2n} \right\}.$$

Exercise 4.5. The limit $\lim_{x \rightarrow +\infty} (a \cos x + b \sin x)$ exists. Show that $a = b = 0$.

Exercise 4.6. Show that $\lim_{x \rightarrow 0+} \sin(\frac{1}{x})$ does not exist.

4.2 Riemannian function

Riemannian function is defined as follows.

$$R(x) = \begin{cases} \frac{1}{q}, & x = \frac{p}{q}, \text{ where } p, q \in \mathbb{Z} \text{ are relatively prime, } q > 0, \\ 0, & x \notin \mathbb{Q}. \end{cases}$$

Example 4.7. Consider the limit $\lim_{x \rightarrow x_0} R(x)$, where $x_0 \notin \mathbb{Q}$.

Proof. Assume that $x_0 \notin \mathbb{Q}$, we claim that $\lim_{x \rightarrow x_0} R(x) = 0$. For any $\varepsilon > 0$, we can choose $N \in \mathbb{Z}_+$ such that $\frac{1}{N} < \varepsilon$. Then there are only finite numbers in \mathbb{Q} such that the representation $\frac{p}{q}$ with $p, q \in \mathbb{Z}$, $(p, q) = 1$, and $q > 0$ satisfy $q < N$. Then, the result follows. \square

4.3 Some basic calculations

Example 4.8. Calculate

$$\lim_{x \rightarrow +\infty} \left(\frac{x^2 - 1}{x^2 + 1} \right)^{\frac{x-1}{x+2}}.$$

Proof. Simple calculations imply that it is 1. □

Exercise 4.9. Calculate the following limits.

(1) $\lim_{x \rightarrow +\infty} \frac{x^k}{a^x} = 0, a > 1, k > 0.$

(2) $\lim_{x \rightarrow +\infty} \frac{\ln x}{x^k} = 0, k > 0.$

(3) $\lim_{x \rightarrow +\infty} x^{1/x} = 1.$

Hint: when $x \notin \mathbb{Z}_+$, consider $[x]$.

4.4 Two important limits

We know that

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1, \quad \lim_{x \rightarrow 0} (1 + x)^{\frac{1}{x}} = 0. \quad (4.1)$$

Example 4.10. Calculate

$$\lim_{x \rightarrow \infty} \frac{\sin x}{x}, \quad \lim_{x \rightarrow +\infty} (1 + x)^{\frac{1}{x}}.$$

Exercise 4.11. By using (4.1) to calculate the following limits.

(1) $\lim_{x \rightarrow +\infty} \left(\frac{2}{\pi} \arctan x \right)^x;$

(2) $\lim_{x \rightarrow \frac{\pi}{2}-} (\sin x)^{\tan x};$

(3) $\lim_{x \rightarrow \infty} \left(\frac{x^2 - 1}{x^2 + 1} \right)^{x^2};$

(4) $\lim_{x \rightarrow \frac{\pi}{2}-} (\cos x)^{\frac{\pi}{2} - x};$

(5) $\lim_{x \rightarrow 0} \frac{\sin 2x - 2 \sin x}{x^3};$

(6) $\lim_{x \rightarrow 1} (1 - x) \tan \left(\frac{\pi}{2} x \right).$

Hint: You have to change the variables.

4.5 Order analysis

Assume that $f, g : (a - \delta, a) \cup (a + \delta)$. We have the following definitions.

$f(x) = o(g(x))$ as $x \rightarrow a$ if and only if

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 0.$$

$f(x) = O(g(x))$ as $x \rightarrow a$ if and only if there exists $M > 0$ such that

$$\left| \frac{f(x)}{g(x)} \right| \leq M.$$

Example 4.12. what is the meaning of $o(1)$ and $O(1)$?

Exercise 4.13. Consider the following limits.

(1) $\lim_{x \rightarrow 0} x \sin \frac{1}{x};$

(2) $\lim_{x \rightarrow \infty} x \sin \frac{1}{x};$

(3) $\lim_{x \rightarrow 0} x \sin x;$

(4) $\lim_{x \rightarrow \infty} x \sin x;$

(5) $\lim_{x \rightarrow 0} \frac{1}{x} \sin \frac{1}{x}$

(6) $\lim_{x \rightarrow \infty} \frac{1}{x} \sin \frac{1}{x}.$

Question 4.14. Please use the notation o , O to illustrate the above limits.

Chapter 5

Continuous function

5.1 Discontinuity

Example 5.1. Consider the following three functions and classify their continuity and discontinuity of them.

$$\begin{aligned}f(x) &= \begin{cases} e^{\frac{1}{x}}, & x \in (-\infty, 0), \\ x^2, & x \in [0, +\infty); \end{cases} \\f(x) &= \begin{cases} e^{\frac{1}{x}}, & x \in \mathbb{R} \setminus \{0\} \\ 0, & x = 0 \end{cases} \\f(x) &= \begin{cases} \frac{\ln(1+x^2)}{x}, & x \in (-1, 0) \cup (0, 1), \\ 2, & x = 0. \end{cases}\end{aligned}$$

Question 5.2. If the range of the function f on the interval $[a, b]$ is a closed interval, is f necessarily continuous on $[a, b]$?

5.2 Uniqueness

Example 5.3. Assume that $f, g \in C(\mathbb{R})$. If $f(x) = g(x)$ for any $x \in \mathbb{Q}$, then $f \equiv g$.

Proof. By using the continuity of f and g . □

Exercise 5.4. Assume that $f \in C(\mathbb{R})$ and $f(x+y) = f(x) + f(y)$. Show that $f(x) = f(1)x$.

5.3 Intermediate value theorem

Example 5.5. Assume that $0 \leq f \in C[0, 1]$ with $f(0) = f(1) = 0$. Show that for any $a \in (0, 1)$, there exists $x_0 \in [0, 1]$ such that $x_0 + a \in [0, 1]$, and

$$f(x_0) = f(x_0 + a).$$

Proof. Let $g(x) = f(x+a) - f(x)$. The result follows directly from the intermediate value theorem. □

Question 5.6. What will happen? If f does not have the property of nonnegativity.

Example 5.7. Assume that $f \in C[0, 1]$ with $f(0) = f(1)$. Show that for any $n \in \mathbb{Z}_+$, there exists ξ such that

$$f\left(\xi + \frac{1}{n}\right) = f(\xi).$$

Proof. Let $g(x) = f(x + \frac{1}{n}) - f(x)$. Consider $g(0), g(\frac{1}{n}), \dots, f(1)$. WLOG, assume that $g(0) > 0$. If there exists $1 \leq k \leq n$ such that $g(\frac{k}{n}) < 1$, then the result follows. If not, it contradicts $f(0) = f(1)$. \square

Exercise 5.8. Assume that $f \in C[0, 1]$, and $f([0, 1]) \subset [0, 1]$. Show that the figure of $y = f(x)$ has an intersection with both $y = x$ and $y = 1 - x$.

Hint: Draw the picture, and then you will find the answer.

Exercise 5.9. Let $f_n(x) = x^n + x$ with $n \in \mathbb{Z}_+$. Show that the following properties hold.

- (1) For any $n \in \mathbb{Z}_+$, the equation $f_n(x) = 1$ has a unique root in $(\frac{1}{2}, 1)$.
- (2) If $c_n \in (\frac{1}{2}, 1)$ is a root of $f_n(x) = 1$ in $(\frac{1}{2}, 1)$, then

$$\lim_{n \rightarrow \infty} c_n = 1.$$

Hint: Firstly, show that c_n is increasing. If $c_n < \alpha$ for any $n \in \mathbb{Z}_+$, there is a contradiction.

5.4 Contraction maps

Example 5.10. Let $f \in C(-\infty, +\infty)$. For any $x, y \in \mathbb{R}$, there holds $|f(x) - f(y)| \leq k|x - y|$ with $0 < k < 1$. Show the following properties.

- (1) The function $kx - f(x)$ is increasing.
- (2) There exists unique $\xi \in \mathbb{R}$, such that $f(\xi) = \xi$.

Hint: Consider the sequence $x_{n+1} = f(x_n)$. Show that x_n is a Cauchy sequence.

Exercise 5.11. Let the function f be defined on the domain $[a, b]$, with the image set $f([a, b]) \subset [a, b]$, and satisfying $|f(x) - f(y)| \leq |x - y|$ for all $x, y \in [a, b]$. For any $x_0 \in [a, b]$, define the sequence recursively by

$$x_{n+1} = \frac{1}{2}(x_n + f(x_n)).$$

Show that the sequence $\{x_n\}$ is convergent.

Hint: Use the MCT.

Exercise 5.12. Let $f \in C[0, +\infty)$. Assume that $f(0) = 0$ and $0 < f(x) < x$ when $x > 0$. For $x_{n+1} = f(x_n)$, show that $\lim_{n \rightarrow +\infty} x_n = 0$.

Hint: Draw the picture.

5.5 Boundedness, maximum, and minimum

Example 5.13. Let the function f have only first-type discontinuities on the interval $[a, b]$. Prove that f is bounded on $[a, b]$.

Proof. Prove by the definition. □

Exercise 5.14. If $f \in C[a, +\infty)$ and there exists a finite limit $\lim_{x \rightarrow +\infty} f(x)$, prove that f is bounded on $[a, +\infty)$.

Exercise 5.15. If $f \in C[a, +\infty)$ and there exists a finite limit $\lim_{x \rightarrow +\infty} f(x)$, prove that f can attain at least either a maximum or a minimum value on $[a, +\infty)$.

Hint: First consider the behavior of f on $[a, a + N]$, $N > 1$.

Exercise 5.16. If $f \in C(a, b)$ and $f(a^+) = f(b^-) = +\infty$, show that f has a minimum value on (a, b) .

Exercise 5.17. If $f \in C(a, b)$ and $f(a^+) = f(b^-)$, prove that f can attain at least either a maximum or a minimum value on (a, b) .

Exercise 5.18. If $f \in C(-\infty, +\infty)$, $\lim_{x \rightarrow \pm\infty} f(x) = +\infty$, and the minimum value of $f(x)$ is $f(a) < a$, prove that the composite function $f(f(x))$ attains its minimum value at least at two points.

Hint: Draw the picture.

5.6 Monotone functions

Exercise 5.19. Let $f \in C[a, b]$, and for any two rational numbers r_1, r_2 in $[a, b]$ with $r_1 < r_2$, it holds that $f(r_1) \leq f(r_2)$. Prove that the function f is monotonically increasing on $[a, b]$.

Exercise 5.20. Let f be a monotonically increasing function on $(-\infty, +\infty)$, and define $g(x) = f(x^+)$ at each point. Prove that g is a function that is right-continuous everywhere on $(-\infty, +\infty)$.

Exercise 5.21. Let f be a monotonically increasing function on $(\infty, +\infty)$, and for all $x, y \in \mathbb{R}$, it satisfies $f(x + y) = f(x) + f(y)$. Prove that $f(x) = f(1)x$.

Exercise 5.22. Let $f \in C[a, b]$. Prove that a necessary and sufficient condition for f to have no extreme points in (a, b) is that f is strictly monotonically increasing on $[a, b]$.

Chapter 6

Differentiation

6.1 Definitions

Example 6.1. Given that the even function $f(x)$ is differentiable at the point $x = 0$, find $f'(0)$.

Proof. Considering $f'(0-)$ and $f'(0+)$, it is easy to see that $f'(0) = 0$. \square

Exercise 6.2. Let $p(x)$ be an n -degree polynomial with n roots, and denote its distinct roots as x_1, \dots, x_k , where the multiplicity of root x_i is n_i for $i = 1, 2, \dots, k$, and $n_1 + n_2 + \dots + n_k = n$. Prove that the following holds.

$$p'(x) = p(x) \left(\sum_{i=1}^k \frac{n_i}{x - x_i} \right)$$

Example 6.3. Let the function be defined as

$$f(x) = \begin{cases} \frac{x}{1+e^{1/x}}, & x \neq 0, \\ 0, & x = 0. \end{cases}$$

Study the differentiability of $f(x)$.

Proof. Consider $\frac{f(x)-f(0)}{x}$. \square

Exercise 6.4. Let n be a natural number. Under what conditions is the function

$$f(x) = \begin{cases} x^n \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Study the differentiability of $f(x)$.

Example 6.5. Let the function $f(x)$ satisfy the functional equation $f(x+y) = f(x)f(y)$, and it is known that $f'(0) = 1$. Prove that $f(x)$ is differentiable everywhere and that $f'(x) = f(x)$ holds.

Proof. The assumption that $f(x+y) = f(x)f(y)$ implies that the differentiability can be spread. \square

Exercise 6.6. Suppose in a neighborhood U of the origin, we have $|f(x)| \leq |g(x)|$, and $g(0) = g'(0) = 0$. Find $f'(0)$.

Hint: Draw the picture.

Exercise 6.7. Assume that $f \in C[a, b]$. Show that if there is some $\alpha, M > 1$ such that for any $x, y \in [a, b]$, $|f(x) - f(y)| \leq M|x - y|^\alpha$, then $f \equiv \text{const}$.

Hint: Show that $f' \equiv 0$.

6.2 Some tricks

Example 6.8. Let $f(0) = 0$ and $f'(0)$ exist. Define the sequence

$$x_n = f\left(\frac{1}{n^2}\right) + f\left(\frac{2}{n^2}\right) + \cdots + f\left(\frac{n}{n^2}\right), \quad n \in \mathbf{N}_+$$

Try to find $\lim_{n \rightarrow \infty} x_n$.

Proof. It is easy to find that

$$f\left(\frac{k}{n^2}\right) = f'(0)\frac{k}{n^2} + o\left(\frac{k}{n^2}\right).$$

Then we can sum k up from 1 to n , and find that $x_n \rightarrow \frac{1}{2}f'(0)$. □

Exercise 6.9. Find the limits of the following sequences.

- (1) $\lim_{n \rightarrow \infty} \left(\sin \frac{1}{n^2} + \sin \frac{2}{n^2} + \cdots + \sin \frac{n}{n^2} \right).$
- (2) $\lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{n^2} \right) \left(1 + \frac{2}{n^2} \right) \cdots \left(1 + \frac{n}{n^2} \right) \right].$

Hint: Use the previous limit.

Exercise 6.10. Let $y = \frac{1+x}{\sqrt{1-x}}$, calculate $y^{(n)}(x)$, where $n \in \mathbf{N}_+$.

Hint: Take \ln .

Exercise 6.11. Let f have derivatives of all orders on \mathbb{R} . Prove that for each natural number n the following holds.

$$\frac{1}{x^{n+1}} f^{(n)}\left(\frac{1}{x}\right) = (-1)^n \left[x^{n-1} f\left(\frac{1}{x}\right) \right]^{(n)}.$$

Exercise 6.12. Using the sum of the series $1 + x + x^2 + \cdots + x^n$, find the sums of the following.

- (1) $1 + 2x + 3x^2 + \cdots + nx^{n-1}.$
- (2) $1^2 + 2^2x + 3^2x^2 + \cdots + n^2x^{n-1}.$

,

Chapter 7

Indefinite integral

7.1 Pairing method

Example 7.1. Calculate $\int \frac{dx}{1+x^4}$.

Consider $\int \frac{x^2 dx}{1+x^4}$.

Exercise 7.2. Calculate the following indefinite integrals.

(1) $\int \frac{dx}{1+x^2+x^4}$.

(2) $\int \frac{e^x dx}{e^x + e^{-x}}$.

(3) $\int \frac{b \sin x + a \cos x}{a \sin x + b \cos x} dx \quad (a \neq b)$.

(4) $\int \frac{dx}{1+x^3}$.

Hint: (1) Consider $\int \frac{x^2}{1+x^2+x^4} dx$. (2) Consider $\int \frac{e^{-x}}{e^x + e^{-x}} dx$. (3) Consider $\int \frac{\sin x}{a \sin x + b \cos x} dx$ and $\int \frac{\cos x}{a \sin x + b \cos x} dx$. Consider $\int \frac{x}{1+x^3} dx$.

Exercise 7.3. Calculate

$$\int (\cos^4 x - \sin^4 x) dx \quad \int (\cos^4 x + \sin^4 x) dx$$

and

$$\int \cos^4 x dx \quad \int \sin^4 x dx.$$

7.2 By iterations

Example 7.4. Calculate $\int \frac{dx}{(1+x^2)^n}$

Using integration by parts, we have

$$\int \frac{dx}{(1+x^2)^n} = \frac{x}{(1+x^2)^n} + \int \frac{2nx^2 dx}{(1+x^2)^{n+1}}.$$

Exercise 7.5. Calculate

$$\int \sin^m x \cos^n x dx \quad \int \tan^n x dx,$$

where $m, n \in \mathbb{Z}_+$.

7.3 Some special integrals

Exercise 7.6 (Rational functions). Calculate the following indefinite integrals.

- (1) $\int \frac{x^5 - x}{1 + x^8} dx.$
- (2) $\int \frac{dx}{x(x^n + a)} (a \neq 0).$
- (3) $\int \frac{x dx}{(x+1)(x^2+3)}.$
- (4) $\int \frac{x-1}{(x^2+2x+3)^2} dx.$
- (5) $\int \frac{1+x+x^2}{(x-2)^{10}} dx.$
- (6) $\int \frac{x^9+1}{x^3-5x^2+6x} dx.$
- (7) $\int \frac{dx}{1+x^6}.$
- (8) $\int \frac{x^2-x+3}{(x^2+x+1)(x-1)^2} dx.$

Exercise 7.7 (Triangular functions). Calculate the following indefinite integrals.

- (1) $\int \frac{dx}{1+\cos x}.$
- (2) $\int \frac{dx}{\sin x \cos^4 x}.$
- (3) $\int \frac{dx}{2+\tan^2 x}.$
- (4) $\int \tan^5 x \sec^3 x dx.$
- (5) $\int \frac{\sec x}{(1+\sec x)^2} dx.$
- (6) $\int \frac{\sin^2 x \cos x}{\sin x + \cos x} dx.$
- (7) $\int \frac{\sin x \cos x}{1+\sin^4 x} dx.$
- (8) $\int \frac{dx}{\sin^4 x \cos^2 x}.$

Example 7.8. Calculate the Poisson integral $\int \frac{1-r^2}{1-2r \cos x + r^2} dx \quad (-1 < r < 1).$

Proof. We have

$$\begin{aligned}
 \int \frac{1-r^2}{1-2r \cos x + r^2} dx &= \int \frac{1-r^2}{(1+r)^2 - 4r \cos^2 \frac{x}{2}} dx \\
 &= \int \frac{1-r^2}{(1-r)^2 \cos^2 \frac{x}{2} + (1+r)^2 \sin^2 \frac{x}{2}} dx \\
 &= \int \frac{2(1-r^2) d(\tan \frac{x}{2})}{(1-r)^2 + (1+r)^2 \tan^2 \frac{x}{2}}.
 \end{aligned}$$

□

Exercise 7.9 (Other types). Calculate the following indefinite integrals.

$$(1) \int \frac{dx}{\sqrt{\sin x \cos 7x}}$$

$$(2) \int \frac{dx}{x\sqrt{2x^2+3}}$$

$$(3) \int \sqrt{\tan^2 x + 2} dx.$$

$$(4) \int \frac{xe^x dx}{\sqrt{1+e^x}}.$$

$$(5) \int \frac{dx}{\sqrt{(x-1)^3(x-2)}}.$$

$$(6) \int \frac{\sqrt{x}-1}{\sqrt[3]{x+1}} dx.$$

Chapter 8

Definite integral

8.1 Calculate limits

Exercise 8.1. Calculate

$$\lim_{n \rightarrow +\infty} \left(\frac{1}{n+1} + \dots + \frac{1}{2n} \right).$$

Exercise 8.2. Calculate

$$\lim_{n \rightarrow +\infty} \left(\frac{1}{n^2} \sum_{k=1}^n k \ln(n+k) - \frac{n+1}{2n} \ln n \right).$$

Hint: Note that $\sum_{k=1}^n k = \frac{n(n+1)}{2}$.

Exercise 8.3. Assume that $f \in R[a, b]$. Calculate

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n (-1)^{k-1} f\left(\frac{k}{n}\right).$$

Hint: WLOG, assume that $n = 2m$.

$$\frac{1}{2m} \sum_{k=1}^{2m} (-1)^{k-1} f\left(\frac{k}{2m}\right) = \frac{1}{2m} \sum_{k=1}^{2m} f\left(\frac{k}{2m}\right) - \frac{1}{2m} \sum_{k=1}^m f\left(\frac{k}{m}\right).$$

8.2 Divide into two parts

Example 8.4. Show that

$$\lim_{n \rightarrow +\infty} \int_0^{\frac{\pi}{2}} \sin^n x dx = 0.$$

Proof. Consider $(0, \frac{\pi}{2} - \delta)$ and $(\frac{\pi}{2} - \delta, \frac{\pi}{2})$. □

Exercise 8.5. Let $f \in C[-1, 1]$. Show that

$$\lim_{h \rightarrow 0^+} \int_{-1}^1 \frac{h}{h^2 + x^2} f(x) dx = \pi f(0).$$

Hint: Consider intervals $(-1, -\delta)$, $(-\delta, \delta)$, and $(\delta, 1)$.

8.3 Approximation and Riemann-Lebesgue Lemma

Example 8.6. Show that integrals can be approximated by step functions and continuous functions.

Draw the picture.

Exercise 8.7. Assume that $f \in R[0, \pi]$. Show that

$$\lim_{n \rightarrow +\infty} \int_0^\pi f(x) \sin nx \, dx = 0.$$

8.4 Calculations by symmetry

Exercise 8.8. Using symmetry, calculate the following problems.

- (1) $I_1 = \int_0^\pi \frac{x \, dx}{1 + \cos^2 x}$
- (2) $I_2 = \int_0^1 \frac{x}{e^x + e^{1-x}} \, dx$
- (3) $I_3 = \int_{-2}^2 x \ln(1 + e^x) \, dx$
- (4) $I_4 = \int_0^{\pi/4} \ln(1 + \tan x) \, dx$
- (5) $I_5 = \int_0^{\pi/2} \frac{\sin^n x}{\sin^n x + \cos^n x} \, dx$
- (6) $I_6 = \int_0^\pi \frac{a^n \sin^2 x + b^n \cos^2 x}{a^{2n} \sin^2 x + b^{2n} \cos^2 x} \, dx$

Exercise 8.9. Let $f \in C[0, a]$, $a > 0$.

- (1) If $f(x) + f(a - x) \neq 0$ on $[0, a]$, calculate $I = \int_0^a \frac{f(x)}{f(x) + f(a - x)} \, dx$;
- (2) If $f(x)f(a - x) \equiv 1$ on $[0, a]$, calculate $I = \int_0^a \frac{dx}{1 + f(x)}$.

Exercise 8.10. Let f be a continuous function and a, b be real numbers. Prove the following equalities.

- (1) $\int_0^\pi x f(\sin x) \, dx = \frac{\pi}{2} \int_0^\pi f(\sin x) \, dx$;
- (2) $\int_1^a f\left(x^2 + \frac{a^2}{x^2}\right) \frac{dx}{x} = \int_1^a f\left(x + \frac{a^2}{x}\right) \frac{dx}{x}$;
- (3) $\int_0^{2\pi} f(a \cos x + b \sin x) \, dx = 2 \int_0^\pi f\left(\sqrt{a^2 + b^2} \cos x\right) \, dx$.

8.5 Mean value theorem

Example 8.11. Let f be a non-negative strictly increasing function that is continuous on the interval $[a, b]$. By the Mean Value Theorem for integrals, for each $p > 0$, there exists a unique $x_p \in (a, b)$ such that

$$f^p(x_p) = \frac{1}{b - a} \int_a^b f^p(t) \, dt$$

Find $\lim_{p \rightarrow \infty} x_p$.

Proof. WLOG, we assume that $a = 0$ and $b = 1$. Then

$$f(x_p) = \left(\int_0^1 f^p(t) dt \right)^{\frac{1}{p}}.$$

Letting $p \rightarrow +\infty$, the RHS goes to $f(1^-)$. Thus, $x_p \rightarrow 1$. \square

Exercise 8.12. Let $f \in C[a, b]$, and let $g(x)$ be a periodic function with period T that is nonnegative integrable on $[0, T]$. Prove that:

$$\lim_{n \rightarrow \infty} \int_a^b f(x)g(nx)dx = \frac{1}{T} \int_a^b f(x)dx \int_0^T g(x)dx.$$

Proof. Assume that $b - a = N\frac{T}{n} + c_n(a, b, T)$, where $N \in \mathbb{Z}_+$ and $0 \leq c_n(a, b, T) < \frac{T}{n}$. We have

$$\begin{aligned} \int_a^b f(x)g(nx)dx &= \sum_{i=1}^N \int_{a+\frac{(i-1)T}{n}}^{a+\frac{iT}{n}} f(x)g(nx)dx + \int_{b-c_n(a,b,T)}^b f(x)g(nx)dx \\ &= f(\xi_k) \int_{a+\frac{k(b-a)}{n}}^{a+\frac{(k+1)T}{n}} g(nx)dx + \int_{b-c_n(a,b,T)}^b f(x)g(nx)dx. \end{aligned}$$

For the last equality, we used the mean value theorem of integration. Then, we are done. \square

8.6 Some inequalities

Proposition 8.13 (Cauchy-Schwartz). *Let $f, g \in R[a, b]$. Then*

$$\left(\int_a^b f(x)g(x)dx \right)^2 \leq \int_a^b f^2(x)dx \int_a^b g^2(x)dx.$$

Hint: Consider

$$\int_a^b (f(x) - tg(x))^2 dx.$$

Example 8.14. $f \in C^1[a, b]$ and let $f(a) = 0$. Prove that

$$\int_a^b f^2(x)dx \leq \frac{(b-a)^2}{2} \int_a^b (f'(x))^2 dx.$$

Exercise 8.15. It is given that the function $f \in C[a, b]$ and $f(x) > 0$ for all x . Prove that

$$\int_a^b f(x)dx \int_a^b \frac{1}{f(x)}dx \geq (b-a)^2.$$

Exercise 8.16. Given a non-negative function $f \in R[a, b]$ with $\int_a^b f(x)dx = 1$ and k being a real number, prove that

$$\left(\int_a^b f(x) \cos kx dx \right)^2 + \left(\int_a^b f(x) \sin kx dx \right)^2 \leq 1.$$

Proposition 8.17 (Young). *Let f be continuous and differentiable on $[0, \infty)$ and strictly increasing, with $f(0) = 0$ and $a, b > 0$. Then*

$$ab \leq \int_0^a f(x)dx + \int_0^b g(y)dy.$$

where $g(y)$ is the inverse function of $f(x)$, and equality holds if and only if $b = f(a)$.

Hint: Draw the picture.

8.7 Combination of previous topics

Exercise 8.18. Calculate the following asymptotic equality

$$\int_0^1 \frac{x^{n-1}}{1+x} dx = \frac{a}{n} + \frac{b}{n^2} + o\left(\frac{1}{n^2}\right) \quad (n \rightarrow \infty)$$

for the undetermined constants a and b .

Hint: By using integration by parts.

Exercise 8.19. Prove that for every natural number n ,

$$\frac{2}{3}n\sqrt{n} < 1 + \sqrt{2} + \cdots + \sqrt{n} < \frac{4n+3}{6}\sqrt{n}.$$

Exercise 8.20. Let f be a non-increasing function on $[1, +\infty)$. Define

$$a_n = \sum_{k=1}^n f(k) - \int_1^n f(x) dx, \quad n \in \mathbb{N}_+.$$

Prove that the sequence $\{a_n\}$ converges.

Exercise 8.21. Calculate

$$\lim_{n \rightarrow \infty} \left(\frac{\sin \pi/n}{n+1} + \frac{\sin 2\pi/n}{n+\frac{1}{2}} + \cdots + \frac{\sin \pi}{n+\frac{1}{n}} \right).$$

Hint: First consider the case

$$\lim_{n \rightarrow \infty} \left(\frac{\sin \pi/n}{n} + \frac{\sin 2\pi/n}{n} + \cdots + \frac{\sin \pi}{n} \right).$$

Exercise 8.22. Assume that $f \in C^1[0, a]$, where $a > 0$. Prove that

$$|f(0)| \leq \frac{1}{a} \int_0^a |f(x)| dx + \int_0^a |f'(x)| dx.$$

Hint: WLOG, we assume that $a = 1$. We have

$$f(0) - \int_0^1 f = \int_0^1 (f - f(0)) = \int_0^1 \left(\int_0^x f'(t) dt \right) dx.$$

Exercise 8.23. Assume that $f \in C^1[0, 1]$. Prove that

$$\int_0^1 |f(x)| dx \leq \max \left\{ \int_0^1 |f'(x)| dx, \left| \int_0^1 f(x) dx \right| \right\}.$$

Hint: If $f \geq 0$ or $f \leq 0$ in $[0, 1]$, there is nothing to prove. Otherwise, we can apply the N-L formula.

Chapter 9

Review of the midterm exam

9.1 Some questions about sequences

Exercise 9.1. Let

$$x_n = (1 + q)(1 + q^2) \cdots (1 + q^n),$$

where the constant $q \in [0, 1)$. Prove that the sequence $\{x_n\}$ has a limit.

Exercise 9.2. Let

$$x_1 > 0, \quad x_{n+1} = \frac{3(1 + x_n)}{3 + x_n} \quad (n = 1, 2, \dots).$$

Find

$$\lim_{n \rightarrow +\infty} x_n.$$

9.2 Some questions about integrals

Exercise 9.3. Find the limit

$$\lim_{x \rightarrow +\infty} \frac{1}{x} \int_0^x (t - [t]) dx.$$

Hint: For x , consider $[x]$.

Exercise 9.4. Calculate

$$\int_0^\pi \sqrt{1 - \sin x} dx.$$

Hint: Note that $1 - \sin x = (\sin \frac{x}{2} - \cos \frac{x}{2})^2$.

Exercise 9.5. Calculate

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^4 x}{1 + e^{-x}} dx.$$

Hint: Consider the change of variables $y = -x$.

Exercise 9.6. Let the function $y = f(x)$ be differentiable on the interval $[a, b]$ and strictly increasing. Let $g(y)$ be the inverse function of $f(x)$. Prove the following formula:

$$\int_a^b f(x) dx = bf(b) - af(a) - \int_{f(a)}^{f(b)} g(y) dy.$$

Hint: By the change of variables and the relation of derivatives for functions and their reverse functions.

Exercise 9.7. Let $f(x)$ be continuous on $(-\infty, +\infty)$. Find

$$\frac{d}{dx} \int_0^x t^2 f(x^3 - t^3) dt.$$

Hint: By change of variables.

Exercise 9.8. 33. Let the function $f(x)$ be continuous on $[0, +\infty)$, monotonically increasing, and differentiable on $(0, +\infty)$.

1. Prove that for any $x > 0$,

$$f(x) \geq \frac{1}{x} \int_0^x f(t) dt.$$

2. Define $F(x) = \frac{1}{x} \int_0^x f(t) dt$. Prove that $F(x)$ is monotonically increasing on $(0, +\infty)$.

3. Find $\lim_{x \rightarrow 0^+} F(x)$.

Exercise 9.9. Let $f(x)$ and $g(x)$ be continuous on $[a, b]$.

1. If $f(x) \geq 0$ for $x \in [a, b]$, prove that $F(x) = \int_a^x f(t) dt$ is monotonically increasing on $[a, b]$.
Furthermore, if $F(b) = \int_a^b f(t) dt = 0$, prove that $F(x) = 0$ and $f(x) = 0$ for $x \in [a, b]$.

2. If $f(x) \geq g(x)$ and $f(x) \neq g(x)$ for $x \in [a, b]$, prove that

$$\int_a^b f(x) dx > \int_a^b g(x) dx.$$

Exercise 9.10. Calculate

$$\int \frac{dx}{x\sqrt{3x^2 - 2x - 1}}.$$

Hint: Let $x = \frac{1}{t}$.

Chapter 10

Mean value theorem of derivatives

10.1 Constructing auxiliary functions

Exercise 10.1. Let f be continuous on $[a, b]$, differentiable on (a, b) , and with $0 < a < b$. Prove that there exists $\xi \in (a, b)$ such that

$$f(b) - f(a) = \ln \frac{b}{a} \cdot \xi f'(\xi).$$

Hint: Consider $f(e^x)$.

Exercise 10.2. Let f be differentiable on $[a, b]$. Prove that there exists $\xi \in (a, b)$ such that

$$2\xi[f(b) - f(a)] = (b^2 - a^2) f'(\xi).$$

Hint: Consider $f(x^{1/2})$.

Example 10.3. Let f, g be continuous on $[a, b]$ and differentiable on (a, b) , with $g(x)$ having no zeros in the interval (a, b) . Prove that there exists $\xi \in (a, b)$ such that

$$\frac{f'(\xi)}{g'(\xi)} = \frac{f(\xi) - f(a)}{g(b) - g(\xi)}.$$

Proof. Consider the function

$$F(x) = -f(x)g(x) + g(b)f(x) + f(a)g(x).$$

It is easy to see that $F(a) = g(b)f(a) = F(b)$. Then the result follows from Rolle's theorem. \square

Exercise 10.4. Let f be continuous on $[a, +\infty)$ and differentiable on $(a, +\infty)$, with $\lim_{x \rightarrow +\infty} f(x) = f(a)$. Prove that there exists $\xi > a$ such that $f'(\xi) = 0$.

Exercise 10.5. Let $f(x)$ be differentiable on $[0, +\infty)$ and satisfy $0 \leq f(x) \leq \frac{x}{1+x^2}$. Prove that there exists $\xi > 0$ such that

$$f'(\xi) = \frac{1 - \xi^2}{(1 + \xi^2)^2}.$$

Hint: Consider $f(x) - \frac{x}{1+x^2}$.

10.2 Related calculations

Exercise 10.6. For (1) $f(x) = ax^2 + bx + c$ ($a \neq 0$) and (2) $f(x) = \frac{1}{x}$ ($x > 0$), calculate the value of θ in the formula $f(x + \Delta x) - f(x) = f'(x + \theta\Delta x)\Delta x$ and find the limit $\lim_{\Delta x \rightarrow 0} \theta$.

Exercise 10.7. Prove that when $x \geq 0$, we have

$$\sqrt{x+1} - \sqrt{x} = \frac{1}{2\sqrt{x+\theta(x)}},$$

where $\frac{1}{4} \leq \theta(x) \leq \frac{1}{2}$, and the following properties hold.

$$\lim_{x \rightarrow 0^+} \theta(x) = \frac{1}{4}, \quad \lim_{x \rightarrow +\infty} \theta(x) = \frac{1}{2}.$$

Exercise 10.8. Prove that for $|x| \leq \frac{1}{\sqrt{2}}$, the following holds.

$$2 \arcsin x \equiv \arcsin \left(2x\sqrt{1-x^2} \right).$$

10.3 Comprehensive applications

Exercise 10.9. Let f be differentiable on the interval $[a, b]$. Prove that if $f(a)$ is the maximum value of f , then $f'_+(a) \leq 0$; if $f(b)$ is the maximum value of f , then $f'_-(b) \geq 0$.

Exercise 10.10. Let the function f be twice differentiable on the interval I , and suppose that $f''(x) \equiv 0$. What kind of function is f ?

Exercise 10.11. Prove that the derivative of a differentiable function that is unbounded on a finite open interval (a, b) is also unbounded.

Exercise 10.12. Let f be differentiable on $(0, a)$ and $f(0^+) = +\infty$. Prove that $f'(x)$ does not have a lower bound on the right side of the point $x = 0$.

Exercise 10.13. Let f be continuous on $[a, b]$, differentiable on (a, b) , and satisfy $f(a) = f(b)$, but f is not a constant function. Prove that there exists $\xi \in (a, b)$ such that $f'(\xi) > 0$.

Exercise 10.14. Let f be continuous on $[0, 1]$ and twice differentiable in $(0, 1)$. It is known that the line segment connecting points $A(0, f(0))$ and $B(1, f(1))$ intersects the curve $y = f(x)$ at point $C(c, f(c))$, where $0 < c < 1$. Prove that there exists a point $\xi \in (0, 1)$ such that $f''(\xi) = 0$.

Hint: Draw the picture.

Exercise 10.15. Let f be continuous on $[a, b]$ and differentiable in (a, b) , with $f'(x)$ having no zeros. Prove that there exist $\xi, \eta \in (a, b)$ such that

$$\frac{f'(\xi)}{f'(\eta)} = \frac{e^b - e^a}{b - a} e^{-\eta}.$$

Hint: Consider the case that $\xi = \eta$.

Exercise 10.16. Let f be continuous on $[0, 1]$, differentiable in $(0, 1)$, with $f(0) = f(1) = 0$ and $f\left(\frac{1}{2}\right) = 1$. Prove that:

- (1) There exists $\eta \in (\frac{1}{2}, 1)$ such that $f(\eta) = \eta$.
- (2) For any real number λ , there exists $\xi \in (0, \eta)$ such that

$$f'(\xi) - \lambda(f(\xi) - \xi) = 1.$$

Hint: (1): IVT. (2) Consider the function $e^{-\lambda x}(f(x) - x)$.

Exercise 10.17. Let f be a differentiable function on the interval I . Prove that the necessary and sufficient condition for f' to be a constant function on I is that f is a linear function.

Proposition 10.18 (Darboux). *The derivative of a function satisfies IVT.*

Hint: Consider the function $F(x) = f(x) - \eta x$.

Chapter 11

L'Hospital's rule

Exercise 11.1. $\delta_n = e - \left(1 + \frac{1}{n}\right)^n$. Calculate the limit $\lim_{n \rightarrow +\infty} n\delta_n$.

Hint: Consider the limit

$$\lim_{x \rightarrow 0} \frac{e - (1+x)^{\frac{1}{x}}}{x}.$$

Exercise 11.2. The following limits of functions should not be evaluated using L'Hospital's Rule. Why?

1. $\lim_{x \rightarrow 1} \frac{x}{x+1}$.
2. $\lim_{x \rightarrow +\infty} \frac{x + \sin x}{x - \sin x}$.
3. $\lim_{x \rightarrow +\infty} \frac{e^x + e^{-x}}{e^x - e^{-x}}$.
4. $\lim_{x \rightarrow +\infty} \frac{x}{\sqrt{1+x^2}}$.

Exercise 11.3. Find the following limits.

1. $\lim_{x \rightarrow 0} \frac{a^{x^2} - b^{x^2}}{(a^x - b^x)^2}$.
2. $\lim_{x \rightarrow +\infty} \left[\left(x^3 - x^2 + \frac{x}{2}\right) e^{\frac{1}{x}} - \sqrt{1+x^6} \right]$.
3. $\lim_{x \rightarrow 0^+} \left(\ln \frac{1}{x}\right)^x$.
4. $\lim_{x \rightarrow +\infty} \left(\frac{\pi}{2} - \arctan x\right)^{\frac{1}{\ln x}}$.

Chapter 12

Taylor expansion and further applications of MVT

12.1 The property of best approximation

Example 12.1. Let the function f be defined in the neighborhood $(x_0 - r, x_0 + r)$ and be n -times differentiable at $x = x_0$. Let $P_n^T(x)$ be the n -th Taylor polynomial of $f(x)$ at $x = x_0$. For any other n -th degree polynomial $P_n(x)$, prove that there exists a number $\delta \in (0, r)$, depending on the choice of $P_n(x)$, such that

$$|f(x) - P_n^T(x)| \leq |f(x) - P_n(x)|$$

holds for all $x \in (x_0 - \delta, x_0 + \delta)$.

Proof. Let $P_n^T(x) = \sum_{i=0}^n a_i x^i$ and $P_n(x) = \sum_{i=0}^n b_i x^i$. WLOG, we assume that $P_n^T(x) \neq P_n(x)$. Choose $i_0 \in \{0, 1, 2, \dots, n-1\}$ such that $a_i = b_i$ when $i \leq i_0$, and $a_i \neq b_i$ when $i > i_0$. By the Peano's remainder of Taylor expansion, we have

$$\lim_{x \rightarrow 0} \frac{|f(x) - P_n^T(x)|}{|x|^{n+1}} = 0$$

and

$$\lim_{x \rightarrow 0} \frac{\left| f(x) - \sum_{i=1}^{i_0} b_i x^i \right|}{|x|^{i_0+1}} = 0.$$

By the choice of i_0 , we have

$$\lim_{x \rightarrow 0} \frac{|f(x) - P_n(x)|}{|x|^{i_0+1}} = |a_{i_0+1} - b_{i_0+1}|.$$

Then we are done. □

12.2 Asymptotic behavior

Example 12.2. Let the function $f(x)$ be differentiable on $[a, +\infty)$, and suppose that the limit $\lim_{x \rightarrow \infty} f(x)$ exists and is finite. Answer the following questions:

1. Provide a counterexample to show that $\lim_{x \rightarrow \infty} f'(x) = 0$ does not always hold.
2. If $f(x)$ is twice differentiable and $f''(x)$ is bounded on $[a, +\infty)$, prove that $\lim_{x \rightarrow \infty} f'(x) = 0$.

Proof. For the first point, we consider the function

$$f'(x) = \begin{cases} \text{linear function} & x \in [n, n + n^{-100}], \\ 0 & \text{otherwise.} \end{cases}$$

By Taylor expansion with Lagrange's remainder, we have

$$f(x+t) = f(x) + f'(x)t + \frac{t^2}{2}f''(\xi_{x,t}),$$

where $\xi_x \in [x, x+t]$. □

Example 12.3. Let f be twice differentiable on $(0, +\infty)$, and suppose that

$$M_0 = \sup\{|f(x)| \mid x \in (0, +\infty)\}, \quad \text{and} \quad M_2 = \sup\{|f''(x)| \mid x \in (0, +\infty)\}$$

are finite numbers. Prove that $M_1 = \sup\{|f'(x)| \mid x \in (0, +\infty)\}$ is also finite and satisfies the inequality

$$M_1 \leq 2\sqrt{M_0 M_2}.$$

Proof. Write

$$f(x+t) = f(x) + f'(x)t + \frac{f''(\xi)}{2}t^2,$$

where $x, t > 0$ and $\xi \in (x, x+t)$. From this, we have the estimate

$$|tf'(x)| \leq \left| f(x+t) - f(x) - \frac{t^2}{2}f''(\xi) \right| \leq 2M_0 + \frac{t^2}{2}M_2.$$

Thus, we obtain

$$|f'(x)| \leq \frac{2M_0}{t} + \frac{1}{2}tM_2.$$

This holds for every $x \in (0, +\infty)$. Taking the supremum, we get

$$M_1 \leq \frac{2M_0}{t} + \frac{t}{2}M_2.$$

Therefore, M_1 is a finite number. Since this holds for every $t > 0$, to get the best estimate, we can take $t = 2\sqrt{M_0/M_2}$ to minimize the expression on the right, yielding

$$M_1 \leq 2\sqrt{M_0 M_2}.$$

□

12.3 Existences of particular points

Example 12.4. Let f be twice differentiable on $[a, b]$. Prove that there exists $\xi \in (a, b)$ such that

$$f(a) - 2f\left(\frac{a+b}{2}\right) + f(b) = \frac{1}{4}(b-a)^2 f''(\xi).$$

Proof. Write the Taylor expansions of $f(a)$ and $f(b)$ at the point $\frac{a+b}{2}$:

$$\begin{aligned} f(a) &= f\left(\frac{a+b}{2}\right) + f'\left(\frac{a+b}{2}\right)\left(\frac{a-b}{2}\right) + \frac{1}{2}f''(\eta_1)\left(\frac{b-a}{2}\right)^2, \\ f(b) &= f\left(\frac{a+b}{2}\right) + f'\left(\frac{a+b}{2}\right)\left(\frac{b-a}{2}\right) + \frac{1}{2}f''(\eta_2)\left(\frac{b-a}{2}\right)^2, \end{aligned}$$

Then adding both equations, we have

$$f(a) - 2f\left(\frac{a+b}{2}\right) + f(b) = \frac{1}{8}(b-a)^2 [f''(\eta_1) + f''(\eta_2)].$$

By applying the Darboux theorem to f'' , there exists $\xi \in \{a, b\}$ such that

$$f''(\xi) = \frac{1}{2} [f''(\eta_1) + f''(\eta_2)].$$

□

Exercise 12.5. Calculate $\lim_{n \rightarrow +\infty} \sin(\pi en!)$.

Hint: Use Taylor's expansion to e .

Example 12.6. If the Taylor polynomial of any order of the function f at a point x_0 is zero, can we conclude that $f(x) \equiv 0$?

Let f have derivatives of all orders on $[-1, 1]$, with $f^{(n)}(0) = 0$ for all $n \in \mathbb{N}_+$, and there exists a constant $C \geq 0$ such that for all $n \in \mathbb{N}_+$ and $x \in [-1, 1]$ the inequality $|f^{(n)}(x)| \leq n!C^n$ holds. Prove that $f(x) \equiv 0$.

Hint: Consider the function $e^{-\frac{1}{x^2}}$.

Example 12.7. Let f be twice differentiable on $[a, b]$, and let $f'(a) = f'(b) = 0$. Prove that there exists $\xi \in (a, b)$ such that

$$|f''(\xi)| \geq \frac{4}{(b-a)^2} |f(b) - f(a)|.$$

Proof. WLOG, we assume that $a = 0$, $b = 1$, and $f(1) > f(0)$. If the result is not true. We see that

$$|f''(x)| < 4(f(1) - f(0)).$$

By Taylor's expansion, it follows that

$$\begin{aligned} f\left(\frac{1}{2}\right) - f(0) &= \frac{1}{2}f''(\xi_1)\left(\frac{1}{2}\right)^2, \\ f\left(\frac{1}{2}\right) - f(1) &= \frac{1}{2}f''(\xi_2)\left(\frac{1}{2}\right)^2. \end{aligned}$$

This is a contradiction. □

Exercise 12.8. Let f be twice differentiable on $[a, +\infty)$, and suppose that $f(x) \geq 0$ and $f''(x) \leq 0$. Prove that for $x \geq a$, it holds that $f'(x) \geq 0$.

Hint: Prove by contradiction.

Exercise 12.9. Let f be differentiable on (a, b) . Is it true that for every point $t \in (a, b)$, there always exist two points $x_1, x_2 \in (a, b)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(t)?$$

Let f be differentiable on (a, b) , and suppose that at some point $\xi \in (a, b)$, we have $f''(\xi) > 0$. Prove that there exist two points $x_1, x_2 \in (a, b)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(\xi)?$$

Hint: Draw the picture.

12.4 Basic calculations

Exercise 12.10. Let f be $(n+1)$ -times differentiable in $(-1, 1)$, with $f^{(n+1)}(0) \neq 0$ for $n \in \mathbb{N}_+$. For $0 < |x| < 1$, we have

$$f(x) = f(0) + f'(0)x + \cdots + \frac{f^{(n-1)}(0)}{(n-1)!}x^{n-1} + \frac{f^{(n)}(\theta x)}{n!}x^n,$$

prove that

$$\lim_{x \rightarrow 0} \theta = \frac{1}{n+1}.$$

Hint: Note that

$$\frac{f^{(n)}(0)}{n!}x^n + \frac{f^{(n+1)}(0)}{(n+1)!}x^{n+1} + o(x^{n+2}) = \frac{f^{(n)}(\theta x)}{n!}x^n.$$

Exercise 12.11. Let f be n -times differentiable in the neighborhood $O_\delta(x_0)$, and suppose that $f''(x_0) = \cdots = f^{(n-1)}(x_0) = 0$ and $f^{(n)}(x_0) \neq 0$. Prove that when $0 < |h| < \delta$, it holds that

$$f(x_0 + h) - f(x_0) = hf'(x_0 + \theta h), \quad 0 < \theta < 1,$$

and that

$$\lim_{h \rightarrow 0} \theta = \frac{1}{n^{\frac{1}{n-1}}}.$$

Exercise 12.12. Let f be twice continuously differentiable on $(-\infty, +\infty)$, and suppose that for all $x, h \in \mathbb{R}$ the following holds:

$$f(x+h) - f(x) = hf'\left(x + \frac{h}{2}\right).$$

Prove that $f(x) = ax^2 + bx + c$.

Hint: Take the derivatives wrt to h .

12.5 Domination and zero functions

Exercise 12.13. Let f be differentiable on $[0, +\infty)$, with $f(0) = 0$, and there exists a constant $c > 0$ such that

$$|f'(x)| \leq c|f(x)| \quad \forall x \in [0, +\infty).$$

Prove that $f(x) \equiv 0$.

Hint: Consider the equation $f'(x) = cf(x)$.

Exercise 12.14. Let f have derivatives of all orders on $(-1, 1)$, and for each $n \geq 0$, suppose that

$$|f^{(n)}(x)| \leq n!|x|.$$

Prove that $f(x) \equiv 0$.

Hint: Taylor expansion.

Chapter 13

Application of derivatives

13.1 Monotonicity and zero points

Example 13.1. Let

$$P_n(x) = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \cdots + \frac{x^n}{n!}, \quad n \in \mathbb{Z}_+.$$

Prove the following properties.

1. When n is even, $P_n(x) > 0$ for all $x \in \mathbb{R}$.
2. When n is odd, $P_n(x)$ has at least one real root.
3. When $x < 0$, the inequality $P_{2n}(x) > e^x > P_{2n+1}(x)$ holds.
4. When $x > 0$, the inequality $e^x > P_n(x) \geq \left(1 + \frac{x}{n}\right)^n$ holds.
5. For all $x \in \mathbb{R}$, it holds that $\lim_{n \rightarrow \infty} P_n(x) = e^x$.

Proof. For (1) and (2), you have to use the induction. For (3), we can apply Taylor expansion. As for (4), we note that For $\left(1 + \frac{1}{n}\right)^n$, we consider the binomial expansion and substitute the definition of the binomial coefficient:

$$\begin{aligned} \left(1 + \frac{x}{n}\right)^n &= 1 + \frac{n}{1} \cdot \frac{x}{n} + \frac{n(n-1)}{1 \cdot 2} \cdot \frac{x^2}{n^2} + \cdots + \frac{n(n-1) \cdots 1}{1 \cdot 2 \cdots n} \cdot \frac{x^n}{n^n} \\ &= 1 + \frac{x}{1!} \cdot \frac{n}{n} + \frac{x^2}{2!} \cdot \frac{n}{n} \cdot \frac{n-1}{n} + \cdots + \frac{x^n}{n!} \cdot \frac{n}{n} \cdot \frac{n-1}{n} \cdots \frac{1}{n}. \end{aligned}$$

If we scale the fractions $\frac{k}{n}$ on the right side to 1, we have:

$$\left(1 + \frac{x}{n}\right)^n \leq 1 + \sum_{k=1}^n \frac{x^k}{k!}.$$

For this last one, we can apply Taylor expansion. □

Exercise 13.2. Let $n \in \mathbb{Z}_+$. Prove that the equation

$$x^{n+2} - 2x^n - 1 = 0$$

has a unique positive real root.

Hint: Draw the picture.

13.2 Uniqueness of ODE

Exercise 13.3. Consider the boundary value problem for the second-order linear ordinary differential equation on the interval $[a, b]$:

$$y'' + p(x)y' + q(x)y = r(x), \quad a < x < b, \quad y(a) = A, \quad y(b) = B,$$

where p, q, r are given functions and A, B are given constants. If $q(x) < 0$ holds for all $x \in (a, b)$, then the equation has at most one solution $y(x)$.

Hint: Consider the case that $A = B = 0$.

13.3 Basic calculations

Exercise 13.4. Let $|f(x) + f'(x)| \leq 1$ and $f(x)$ be bounded on $(-\infty, +\infty)$. Prove that $|f(x)| \leq 1$.

Hint: Consider the function $g(x) = e^x f(x)$.

13.4 Proof of inequalities

Exercise 13.5. Let $f(x)$ be twice differentiable in a neighborhood of $x = 0$, and suppose that

$$\lim_{x \rightarrow 0} \left(1 + x + \frac{f(x)}{x} \right)^{\frac{1}{x}} = e^3.$$

Find $f(0)$, $f'(0)$, $f''(0)$, and $\lim_{x \rightarrow 0} \left(1 + \frac{f(x)}{x} \right)^{\frac{1}{x}}$.

Hint: Taylor expansion

Exercise 13.6. Prove that for every natural number n , the inequality

$$\frac{e}{2n+2} < e - \left(1 + \frac{1}{n} \right)^n < \frac{e}{2n+1}$$

holds.

Hint: Consider some associated functions.

Exercise 13.7. Prove that when $0 < x < \frac{\pi}{2}$, the inequality

$$\left(\frac{\sin x}{x} \right)^3 > \cos x$$

holds.

Exercise 13.8. Prove that when $0 < x < \frac{\pi}{2}$, the inequality

$$2 \sin x + \tan x \geq 3x$$

holds.

Exercise 13.9. Prove that when $0 < x < 1$, the inequality

$$\pi < \frac{\sin \pi x}{\pi(1-x)} \leq 4$$

holds.

Chapter 14

Calculus with multi-variables I

Example 14.1. Let $f(x, y) = |x - y|\varphi(x, y)$, where $\varphi(x, y)$ is defined in a neighborhood of the point $(0, 0)$. We require appropriate conditions on the function $\varphi(x, y)$ such that:

1. $f(x, y)$ is continuous at the point $(0, 0)$;
2. $f(x, y)$ has partial derivatives at the point $(0, 0)$;
3. $f(x, y)$ is differentiable at the point $(0, 0)$.

Proof. (1) Since $f(0, 0) = 0$, in the vicinity of the point $(0, 0)$,

$$|f(x, y)| \leq 2\sqrt{x^2 + y^2}|\varphi(x, y)|,$$

thus, when

$$\lim_{(x, y) \rightarrow (0, 0)} \sqrt{x^2 + y^2} \varphi(x, y) = 0,$$

$f(x, y)$ is continuous at $(0, 0)$. In particular, if $\varphi(x, y)$ is bounded in the vicinity of $(0, 0)$, then $f(x, y)$ is continuous at $(0, 0)$.

(2) The one-sided derivatives are given by

$$\begin{aligned} \left(\frac{\partial f}{\partial x} \right)_{\pm}(0, 0) &= \lim_{x \rightarrow 0^{\pm}} \frac{f(x, 0) - f(0, 0)}{x} \\ &= \lim_{x \rightarrow 0^{\pm}} \frac{\pm x \varphi(x, 0)}{x} = \pm \lim_{x \rightarrow 0^{\pm}} \varphi(x, 0). \end{aligned}$$

Thus, when $\lim_{x \rightarrow 0^+} \varphi(x, 0) = -\lim_{x \rightarrow 0^-} \varphi(x, 0)$, we have $\frac{\partial f}{\partial x}(0, 0) = \lim_{x \rightarrow 0^+} \varphi(x, 0)$. Similarly, when $\lim_{y \rightarrow 0^+} \varphi(0, y) = -\lim_{y \rightarrow 0^-} \varphi(0, y)$, we find $\frac{\partial f}{\partial y}(0, 0) = \lim_{y \rightarrow 0^+} \varphi(0, y)$. In particular, when $\lim_{(x, y) \rightarrow (0, 0)} \varphi(x, y) = 0$, we have $\frac{\partial f}{\partial x}(0, 0) = \frac{\partial f}{\partial y}(0, 0) = 0$.

(3) Since

$$\begin{aligned} &f(x, y) - \frac{\partial f}{\partial x}(0, 0)x - \frac{\partial f}{\partial y}(0, 0)y \\ &= |x - y|\varphi(x, y) - \left[\lim_{x \rightarrow 0^+} \varphi(x, 0) \right] x - \left[\lim_{y \rightarrow 0^+} \varphi(0, y) \right] y \\ &= \begin{cases} [\varphi(x, y) - \lim_{x \rightarrow 0^+} \varphi(x, 0)] x - [\varphi(x, y) + \lim_{y \rightarrow 0^+} \varphi(0, y)] y, & \text{if } x \geq y, \\ -[\varphi(x, y) + \lim_{x \rightarrow 0^+} \varphi(x, 0)] x + [\varphi(x, y) - \lim_{y \rightarrow 0^+} \varphi(0, y)] y, & \text{if } x < y, \end{cases} \end{aligned}$$

Thus, we can conclude that when

$$\lim_{(x,y) \rightarrow (0,0)} \varphi(x,y) = 0,$$

then $\frac{\partial f}{\partial x}(0,0) = \frac{\partial f}{\partial y}(0,0) = 0$ and $f(x,y)$ is differentiable at the point $(0,0)$. \square

Exercise 14.2. Why does the existence of $f_x(x_0, y_0)$ guarantee that the one-variable function $f(x, y_0)$ is continuous at the point x_0 ? Can we further assert that for y_1 sufficiently close to y_0 , the one-variable function $f(x, y_1)$ is continuous at the point x_0 ?

Exercise 14.3. Provide examples to illustrate:

1. $f(x, y)$ has partial derivatives in a neighborhood of a point but may not be continuous at that point and thus may not be differentiable;
2. $f(x, y)$ is continuous at a point but the partial derivatives may not exist at that point and thus may not be differentiable;
3. $f(x, y)$ is differentiable at a point, but the partial derivatives may not be continuous at that point.

Hint: By using the dummy variables.

Exercise 14.4. Prove: If $f_x(x, y)$ exists at the point (x_0, y_0) and $f_y(x, y)$ is continuous at the point (x_0, y_0) , then $f(x, y)$ is differentiable at the point (x_0, y_0) .

Hint: Consider the function $f(x, y_0)$. Precisely, we have

$$f(x, y) - f(x, y_0) = \int_{y_0}^y \partial_1 f(x, t) dt.$$

Exercise 14.5. Let $z = f(x, y)$ be differentiable in the open set $D = (a, b) \times (c, d)$, and the total differential dz is always zero. Does this imply that $f(x, y)$ must take a constant value in D ? Prove your conclusion.

Hint: Connectness.

Exercise 14.6. Let

$$f(x, y) = \begin{cases} \frac{\sqrt{|x-y|}}{x^2+y^2} \sin(x^2 + y^2), & x^2 + y^2 \neq 0, \\ 0, & x^2 + y^2 = 0. \end{cases}$$

Discuss:

1. Is $f(x, y)$ continuous at the point $(0,0)$?
2. Is $f(x, y)$ differentiable at the point $(0,0)$?

Exercise 14.7. Let

$$f(x, y) = \begin{cases} xy \sin \frac{1}{x^2+y^2}, & x^2 + y^2 \neq 0, \\ 0, & x^2 + y^2 = 0. \end{cases}$$

Prove:

1. Both $f_x(0,0)$ and $f_y(0,0)$ exist;

2. $f_x(0,0)$ and $f_y(0,0)$ are not continuous at the point $(0,0)$;
3. $f(x,y)$ is differentiable at the point $(0,0)$.

Exercise 14.8. Let $f(x,y,z)$ be defined in an open set D . If $f_x(x,y,z)$ and $f_y(x,y,z)$ are bounded in D , and for fixed (x,y) , $f(x,y,z)$ is a continuous function of z , prove that $f(x,y,z)$ is continuous in D .

Exercise 14.9. Find the total differential of $u = \ln(1 + x^2 + y^2)$ at the point $(x,y) = (1,2)$.

Exercise 14.10. Let $f(x,y)$ be defined on the rectangle $I = [a,b] \times [c,d]$, and f_y is continuous on I . Prove that $f(x,y)$ satisfies a uniform Lipschitz condition with respect to y , i.e., there exists $L > 0$ such that for all $(x,y_1), (x,y_2) \in I$,

$$|f(x,y_1) - f(x,y_2)| \leq L|y_1 - y_2|,$$

where L is independent of x .

Exercise 14.11. If the partial derivatives f_x and f_y of a function $f(x,y)$ exist in the region D , and for all $(x,y) \in D$, $f_x(x,y) = f_y(x,y) = 0$, prove that $f(x,y)$ is a constant function in D .

Exercise 14.12. Let $\Omega \subset \mathbb{R}^2$ be an open region, and let $u(x,y), v(x,y)$ satisfy

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}, \quad u^2 + v^2 = C,$$

where C is a constant, please show that $u(x,y)$ and $v(x,y)$ are constant functions in Ω .

Exercise 14.13. Let $f(x,y)$ be defined on $G = \{(x,y) \mid x^2 + y^2 < 1\}$. If $f(x,0)$ is continuous at the point $x = 0$ and $f_y(x,y)$ is bounded in G , prove that $f(x,y)$ is continuous at the point $(0,0)$.

Chapter 15

Calculus with multi-variables II

15.1 Homogeneous functions

A function $f(x, y)$ is called an n -th homogeneous function if it satisfies $f(tx, ty) = t^n f(x, y)$ for $t > 0$. Homogeneous functions have the following properties:

Exercise 15.1. If f has continuous partial derivatives, then a necessary and sufficient condition for f to be an n -th homogeneous function is

$$x \frac{\partial f}{\partial x}(x, y) + y \frac{\partial f}{\partial y}(x, y) = n f(x, y).$$

Exercise 15.2. If $f(x, y)$ is a twice continuously differentiable n -th homogeneous function, then

$$\left(x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} \right)^2 f(x, y) = n(n-1) f(x, y).$$

Here,

$$\left(x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} \right)^2 = x^2 \left(\frac{\partial}{\partial x} \right)^2 + 2xy \left(\frac{\partial}{\partial x} \right) \left(\frac{\partial}{\partial y} \right) + y^2 \left(\frac{\partial}{\partial y} \right)^2$$

is a shorthand notation, where x , y , $\frac{\partial}{\partial x}$, and $\frac{\partial}{\partial y}$ are treated as constants during the binomial expansion, and it does not mean applying the operator $x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y}$ to $f(x, y)$ twice in succession.

Exercise 15.3. If $f(x, y)$ is a twice continuously differentiable n -th homogeneous function, then $f_x(x, y)$ and $f_y(x, y)$ are $(n-1)$ -th homogeneous functions.

Exercise 15.4. If $f(x, y)$ is a continuous n -th homogeneous function on $\mathbb{R}^2 \setminus \{0\}$, then

$$|f(x, y)| \leq C (x^2 + y^2)^{\frac{n}{2}},$$

where C is a constant.

15.2 Nondegeneracy

Exercise 15.5. Suppose $u(x, y)$ and $v(x, y)$ have continuous first partial derivatives in the region \mathbb{R}^2 , and there exists a constant $C > 0$ such that for any two points $(x_i, y_i) \in \mathbb{R}^2$ ($i = 1, 2$), the following holds:

$$(u_1 - u_2)^2 + (v_1 - v_2)^2 \geq C \left[(x_1 - x_2)^2 + (y_1 - y_2)^2 \right],$$

where $u_i = u(x_i, y_i)$ and $v_i = v(x_i, y_i)$ for $i = 1, 2$. Then for all $(x, y) \in \mathbb{R}^2$, we have

$$\frac{\partial(u, v)}{\partial(x, y)} \neq 0.$$

15.3 Reverse problems

Exercise 15.6. Proof: A differentiable function $z = f(x, y)$ is a function of $ax + by$ (where $ab \neq 0$) if and only if

$$b \frac{\partial z}{\partial x} = a \frac{\partial z}{\partial y}.$$

15.4 The implicit function theorem

Exercise 15.7. Proof: The equation $F(x, y) = 1 - e^{-x} + y^3 e^{-y} = 0$ has a unique solution $y = y(x)$ for $x > 0$ in $\{x > 0, y \in \mathbb{R}^1\}$, and $y(x)$ is continuously differentiable.

Exercise 15.8. Let $f(x, y)$ satisfy: f_x exists on \mathbb{R}^2 , f_y exists and is continuous on \mathbb{R}^2 , and

$$|f_x| < M |f_y|, \quad f(x_0, y_0) = 0,$$

where M is a constant. Prove that $f(x, y) = 0$ uniquely determines a differentiable solution defined on \mathbb{R} as $y = y(x)$, satisfying $y(x_0) = y_0$. Is the condition $|f_x| < M |f_y|$ necessary? If we remove the condition $f(x_0, y_0) = 0$, does the conclusion still hold?