

Regression Models for Epidemiology

Ezra Morrison

Last modified: 2024-04-18: 17:38:02 (PM)

Contents

| | |
|--|------------|
| Preface | 1 |
| 1. Introduction | 2 |
| 1.1. Introduction to Epi 204 | 3 |
| 1.2. Regression models | 5 |
| 1.3. Other resources | 17 |
| | |
| I. Generalized Linear Models | 18 |
| | |
| 2. Linear (Gaussian) Models | 22 |
| 2.1. Overview | 23 |
| 2.2. Understanding Gaussian Linear Regression Models | 24 |
| 2.3. Estimating Linear Models via Maximum Likelihood | 40 |
| 2.4. Inference about Gaussian Linear Regression Models | 47 |
| 2.5. Goodness of fit | 54 |
| 2.6. Rescaling | 61 |
| 2.7. Prediction | 63 |
| 2.8. Diagnostics | 67 |
| 2.9. Model selection | 88 |
| 2.10. Categorical covariates with more than two levels | 89 |
| | |
| 3. Models for Binary Outcomes | 102 |
| 3.1. Introduction | 103 |
| 3.2. Risk Estimation and Prediction | 104 |
| 3.3. Introduction to logistic regression | 120 |

Contents

| | |
|---|------------|
| 3.4. Multiple logistic regression | 135 |
| 3.5. Fitting logistic regression models | 159 |
| 3.6. Model comparisons for logistic models | 160 |
| 3.7. Residual-based diagnostics | 165 |
| 3.8. Odds Ratios vs Probability (Risk) Ratios | 183 |
| 4. Models for Count Outcomes | 186 |
| 4.1. Introduction | 187 |
| 4.2. Inference for count regression models | 191 |
| 4.3. Prediction | 192 |
| 4.4. Diagnostics | 192 |
| 4.5. Zero-inflation | 193 |
| 4.6. Over-dispersion | 194 |
| II. Time to Event Models | 196 |
| 5. Introduction to Survival Analysis | 198 |
| 5.1. Overview | 199 |
| 5.2. Time-to-event outcome distributions | 199 |
| 5.3. Distribution functions for time-to-event variables | 200 |
| 5.4. Parametric Models for Time-to-Event Outcomes | 217 |
| 5.5. Nonparametric Survival Analysis | 220 |
| 5.6. Example: clinical trial for pediatric acute leukemia | 221 |
| 5.7. The Kaplan-Meier Product Limit Estimator | 224 |
| 5.8. Using the <code>survival</code> package in R | 230 |
| 5.9. Example: Bone Marrow Transplant Data | 241 |
| 5.10. Nelson-Aalen Estimates of Cumulative Hazard and Survival | 250 |
| 6. Proportional Hazards Models | 254 |
| 6.1. The proportional hazards model | 255 |
| 6.2. Cox Model for the <code>bmt</code> data | 266 |
| 6.3. Adjustment for Ties (optional) | 276 |

Contents

| | |
|--|------------|
| 7. Building Cox Proportional Hazards models | 279 |
| 7.1. Building Cox Proportional Hazards models | 280 |
| 7.2. Diagnostic graphs for proportional hazards assumption . . . | 283 |
| 7.3. Predictions and Residuals | 292 |
| 7.4. Goodness of Fit using the Cox-Snell Residuals | 301 |
| 7.5. Martingale Residuals | 303 |
| 7.6. Checking for Outliers and Influential Observations | 308 |
| 7.7. Stratified survival models | 319 |
| 7.8. Time-varying covariates | 331 |
| 7.9. Recurrent Events | 347 |
| 7.10. Age as the time scale | 354 |
| 8. Parametric survival models | 355 |
| 8.1. Parametric Survival Models | 356 |
| 8.2. Combining left-truncation and interval-censoring | 371 |
| References | 373 |
| Appendices | 375 |
| A. Probability | 375 |
| A.1. Random variables | 375 |
| A.2. Characteristics of probability distributions | 375 |
| A.3. Key probability distributions | 377 |
| B. Estimation | 378 |
| B.1. Probabilistic models | 378 |
| B.2. Estimands, estimates, and estimators | 379 |
| B.3. Accuracy of estimators | 382 |
| C. Introduction to Maximum Likelihood Inference | 389 |
| C.1. Introduction | 389 |
| C.2. Example: Maximum likelihood for Tropical Cyclones in Australia | 390 |

Contents

| | |
|--|------------|
| C.3. Maximum likelihood inference for univariate Gaussian models | 406 |
| D. Common Mistakes | 417 |
| D.1. Parameters versus random variables | 417 |
| E. Notation | 418 |

Preface

This web-book is derived from my lecture slides for the Spring 2023 session of Epidemiology 204: “Quantitative Epidemiology III: Statistical Models”, at UC Davis.

I have drawn these materials from many sources, including but not limited to:

- David Rocke’s materials from the 2021 edition of Epi 204¹
- *Regression methods in biostatistics: linear, logistic, survival, and repeated measures models*, 2nd edition (Vittinghoff et al. 2012)
- *An Introduction to Generalized Linear Models*, 4th edition (Dobson and Barnett 2018)

License

This book is licensed to you under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License².

The code samples in this book are licensed under Creative Commons CC0 1.0 Universal (CC0 1.0)³, i.e. public domain.

¹<https://dmrocke.ucdavis.edu/Class/EPI204-Spring-2021/EPI204-Spring-2021.html>

²<http://creativecommons.org/licenses/by-nc-nd/4.0/>

³<https://creativecommons.org/publicdomain/zero/1.0/>

1. Introduction

Configuring R

Functions from these packages will be used throughout this document:

```
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(plotly) # interactive graphics
library(dplyr) # manipulate data
library(tidyr) # Tools to help to create tidy data
library(haven) # import Stata files
library(pander) # format tables for markdown
library(knitr) # format R output for markdown
library(kableExtra) # more markdown formatting
library(parameters) # format model output tables for markdown
library(reactable) # interactive tables
library(dobson) # datasets from Dobson and Barnett 2018
library(conflicted) # check for conflicting function definitions
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
knitr::opts_chunk$set(warning = FALSE)
pander::panderOptions("table.emphasize.rownames", FALSE)
options('digits' = 4)
```

1. Introduction

```
conflicts_prefer(plotly::filter)
conflicts_prefer(ggplot2::autoplot)
```

1.1. Introduction to Epi 204

Welcome to Epidemiology 204: Quantitative Epidemiology III (Statistical Models).

In this course, we will start where Epi 203 left off: with linear regression models.

Note

Epi 203/STA 130B/STA 131B is a prerequisite for this course. If you haven't passed one of these courses, please talk to me after class.

1.1.1. What you should know

Epi 202: probability models for different data types

- binomial
- Poisson
- Gaussian
- exponential

Epi 203: inference for one or several homogenous populations

- the maximum likelihood inference framework:
 - likelihood functions
 - log-likelihood functions
 - score functions

1. Introduction

- estimating equations
- information matrices
- point estimates
- standard errors
- confidence intervals
- hypothesis tests
- p-values
- Hypothesis tests for one, two, and >2 groups:
 - t-tests/ANOVA for Gaussian models
 - chi-square tests for binomial and Poisson models
- Some linear regression

Stat 108: linear regression models

- building models for Gaussian outcomes
 - multiple predictors
 - interactions
- regression diagnostics
- fundamentals of R programming; e.g.:
 - R for Data Science (Wickham, Cetinkaya-Rundel, Golemund 2023)¹
 - Introductory Statistics with R (Dalgaard 2008)²
- RMarkdown or Quarto for formatting homework
- LaTeX for writing math in RMarkdown/Quarto

1.1.2. What we will cover in this course

- Linear (Gaussian) regression models (review and more details)

¹<https://r4ds.hadley.nz/>

²<https://link.springer.com/book/10.1007/978-0-387-79054-1>

1. Introduction

- Regression models for non-Gaussian outcomes
 - binary
 - count
 - time to event
- Statistical analysis using R

1.2. Regression models

Why do we need them?

- continuous predictors
- not enough data to analyze some subgroups individually

1.2.1. Example: Adelie penguins

```
library(ggplot2)
library(plotly)
library(dplyr)
ggpenguins <-
  palmerpenguins::penguins |>
  dplyr::filter(species == "Adelie") |>
  ggplot(
    aes(x = bill_length_mm , y = body_mass_g)) +
  geom_point() +
  xlab("Bill length (mm)") +
  ylab("Body mass (g)")
```

1. Introduction

```
ggpenguins |> print()
```



Figure 1.1.: Palmer penguins

1.2.2. Linear regression

```
ggpenguins2 =  
  ggpenguins +  
  stat_smooth(method = "lm",  
              formula = y ~ x,  
              geom = "smooth")
```

1. Introduction




Figure 1.2.: Palmer penguins with linear regression fit

1.2.3. Curved regression lines

```
ggpenguins2 = ggpenguins +  
  stat_smooth(  
    method = "lm",  
    formula = y ~ log(x),  
    geom = "smooth") +  
  xlab("Bill length (mm)") +  
  ylab("Body mass (g)")
```

```
ggpenguins2 |> print()
```

1. Introduction




1.2.4. Multiple regression

```
ggpenguins =  
  palmerpenguins::penguins |>  
  ggplot(  
    aes(x = bill_length_mm ,  
        y = body_mass_g,  
        color = species  
    )  
  ) +  
  geom_point() +  
  stat_smooth(  
    method = "lm",  
    formula = y ~ x,
```

1. Introduction

```
geom = "smooth") +  
xlab("Bill length (mm)") +  
ylab("Body mass (g)")
```

```
ggpenguins |> print()
```



1.2.5. Modeling non-Gaussian outcomes

```
library(glmx)  
data(BeetleMortality)  
beetles = BeetleMortality |>  
  mutate(  
    pct = died/n,
```

1. Introduction

```
    survived = n - died
  )

plot1 =
  beetles |>
  ggplot(aes(x = dose, y = pct)) +
  geom_point(aes(size = n)) +
  xlab("Dose (log mg/L)") +
  ylab("Mortality rate (%)") +
  scale_y_continuous(labels = scales::percent) +
  # xlab(bquote(log[10]), bquote(CS[2]))) +
  scale_size(range = c(1,2))

print(plot1)
```

1. Introduction



Figure 1.3.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.6. Why don't we use linear regression?

```
beetles_long =  
  beetles |>  
  reframe(.by = everything(),  
    outcome = c(  
      rep(1, times = died),  
      rep(0, times = survived))  
    )  
  
lm1 =
```


1. Introduction

```
beetles_long |>
lm(
  formula = outcome ~ dose,
  data = _)

range1 = range(beetles$dose) + c(-.2, .2)

f.linear = function(x) predict(lm1, newdata = data.frame(dose = x))

plot2 =
  plot1 +
  geom_function(fun = f.linear, aes(col = "Straight line")) +
  labs(colour="Model", size = "")

print(plot2)
```

1. Introduction




Figure 1.4.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.7. Zoom out

```
print(plot2 + expand_limits(x = c(1.6, 2)))
```

1. Introduction




Figure 1.5.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.8. log transformation of dose?

```
lm2 =  
  beetles_long |>  
  lm(formula = outcome ~ log(dose), data = _)  
  
f.linearlog = function(x) predict(lm2, newdata = data.frame(dose = x))  
  
plot3 = plot2 +  
  expand_limits(x = c(1.6, 2)) +  
  geom_function(fun = f.linearlog, aes(col = "Log-transform dose"))
```

1. Introduction

```
print(plot3 + expand_limits(x = c(1.6, 2)))
```




Figure 1.6.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.9. Logistic regression

```
glm1 = beetles |>  
  glm(formula = cbind(died, survived) ~ dose, family = "binomial")  
  
f = function(x) predict(glm1, newdata = data.frame(dose = x), type = "response")  
  
plot4 = plot3 + geom_function(fun = f, aes(col = "Logistic regression"))
```

1. Introduction

```
print(plot4)
```



Figure 1.7.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.10. Three parts to regression models

- What distribution does the outcome have for a specific subpopulation defined by covariates? (outcome model)
- How does the combination of covariates relate to the mean? (link function)
- How do the covariates combine? (linear predictor, interactions)

1. Introduction

1.3. Other resources

These notes represent my still-developing perspective on regression models in epidemiology. Many other statisticians and epidemiologists have published their own perspectives, and you are encouraged to explore your many options and find ones that resonate with you. I have attempted to cite my sources throughout these notes. Here are some additional resources:

- Dunn, Smyth, et al. (2018) is a recent textbook on GLMs. It doesn't cover time-to-event models, and it doesn't use the modern `tidyverse` packages (`ggplot2`, `dplyr`, etc.), but otherwise it seems great.

Part I.

Generalized Linear Models

This section is primarily adapted starting from the textbook “An Introduction to Generalized Linear Models” (4th edition, 2018) by Annette J. Dobson and Adrian G. Barnett:

<https://doi.org/10.1201/9781315182780>

The type of predictive model one uses depends on several issues; one is the type of response.

- Measured values such as quantity of a protein, age, weight usually can be handled in an ordinary linear regression model, possibly after a log transformation.
- Patient survival, which may be censored, calls for a different method (survival analysis, Cox regression).
- If the response is binary, then can we use logistic regression models
- If the response is a count, we can use Poisson regression
- If the count has a higher variance than is consistent with the Poisson, we can use a negative binomial or over-dispersed Poisson
- Other forms of response can generate other types of generalized linear models

We need a linear predictor of the same form as in linear regression x . In theory, such a linear predictor can generate any type of number as a prediction, positive, negative, or zero

We choose a suitable distribution for the type of data we are predicting (normal for any number, gamma for positive numbers, binomial for binary responses, Poisson for counts)

We create a link function which maps the mean of the distribution onto the set of all possible linear prediction results, which is the whole real line $(-\infty, \infty)$. The inverse of the link function takes the linear predictor to the actual prediction.

- Ordinary linear regression has identity link (no transformation by the link function) and uses the normal distribution
- If one is predicting an inherently positive quantity, one may want to use the log link since μ is always positive.
- An alternative to using a generalized linear model with a log link, is to transform the data using the log. This is a device that works well with measurement data and may be usable in other cases, but it cannot be used for 0/1 data or for count data that may be 0.

Table 1.1.: R `glm()` Families

| Family | Links |
|------------------|---|
| gaussian | identity , log, inverse |
| binomial | logit , probit, cauchit, log, cloglog |
| gamma | inverse , identity, log |
| inverse.gaussian | 1/μ^2 , inverse, identity, log |
| Poisson | log , identity, sqrt |
| quasi | identity , logit, probit, cloglog, inverse, log, 1/ μ^2 and sqrt |
| quasibinomial | logit , probit, identity, cloglog, inverse, log, 1/ μ^2 and sqrt |
| quasipoisson | log , identity, logit, probit, cloglog, inverse, 1/ μ^2 and sqrt |

Table 1.2.: R `glm()` Link Functions; $\eta = X\beta = g(\mu)$

| Name | Domain | Range | Link Function | Inverse Link Function |
|----------|---------------------|---------------------|-----------------------|-----------------------|
| identity | $(-\infty, \infty)$ | $(-\infty, \infty)$ | $\eta = \mu$ | $\mu = \eta$ |
| log | $(0, \infty)$ | $(-\infty, \infty)$ | $\eta = \log \{\mu\}$ | $\mu = \exp \{\eta\}$ |
| inverse | $(0, \infty)$ | $(0, \infty)$ | $\eta = 1/\mu$ | $\mu = 1/\eta$ |

| Name | Domain | Range | Link Function | Inverse Link Function |
|---------|---------------|---------------------|-------------------------------------|---|
| logit | $(0, 1)$ | $(-\infty, \infty)$ | $\eta = \log \{\mu / (1 - \mu)\}$ | $\mu = \exp \{\eta\} / (1 + \exp \{\eta\})$ |
| probit | $(0, 1)$ | $(-\infty, \infty)$ | $\eta = \Phi^{-1}(\mu)$ | $\mu = \Phi(\eta)$ |
| cloglog | $(0, 1)$ | $(-\infty, \infty)$ | $\eta = \log \{-\log \{1 - \mu\}\}$ | $\mu = \exp \{-\exp \{\eta\}\}$ |
| 1/mu^2 | $(0, \infty)$ | $(0, \infty)$ | $\eta = 1/\mu^2$ | $\mu = 1/\sqrt{\eta}$ |
| sqrt | $(0, \infty)$ | $(0, \infty)$ | $\eta = \sqrt{\mu}$ | $\mu = \eta^2$ |

2. Linear (Gaussian) Models

Note

This content is adapted from:

- Dobson and Barnett (2018), Chapters 2-6
- Dunn, Smyth, et al. (2018), Chapters 2-3
- Vittinghoff et al. (2012), Chapter 4

There are numerous textbooks specifically for linear regression, including:

- Kutner et al. (2005)
- Chatterjee and Hadi (2015)

Functions from these packages will be used throughout this document:

```
library(knitr) # compiling .qmd files
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics:
# - `ggeasy::easy_labs()` applies same logic as `ggplot2::labs()` but uses as default
library(dplyr) # manipulate data
library(haven) # import Stata files
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
```

2. Linear (Gaussian) Models

```
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
library(magrittr) # `>%` and other additional piping tools
```

2.1. Overview

2.1.1. Why this course includes linear regression

- This course is about *generalized linear models* (for non-Gaussian outcomes)
- UC Davis STA 108 (“Applied Statistical Methods: Regression Analysis”) is a prerequisite for this course, so everyone here should have some understanding of linear regression already.
- We will review linear regression to:
 - make sure everyone is caught up
 - to provide an epidemiological perspective on model interpretation.

2.1.2. Chapter overview

- Section 2.2: how to interpret linear regression models
- Section 2.3: how to estimate linear regression models
- Section 2.4: how to quantify uncertainty about our estimates
- Section 2.8: how to tell if your model is insufficiently complex

2.2. Understanding Gaussian Linear Regression Models

2.2.1. Motivating example: birthweights and gestational age

Suppose we want to learn about the distributions of birthweights (*outcome* Y) for (human) babies born at different gestational ages (*covariate* A) and with different chromosomal sexes (*covariate* S) (Dobson and Barnett (2018) Example 2.2.2).

2.2.1.1. Data as table

```
library(dobson)
data("birthweight", package = "dobson")
birthweight |> knitr::kable()
```

Table 2.1.: `birthweight` data (Dobson and Barnett (2018) Example 2.2.2)

| boys gestational age | boys weight | girls gestational age | girls weight |
|----------------------|-------------|-----------------------|--------------|
| 40 | 2968 | 40 | 3317 |
| 38 | 2795 | 36 | 2729 |
| 40 | 3163 | 40 | 2935 |
| 35 | 2925 | 38 | 2754 |
| 36 | 2625 | 42 | 3210 |
| 37 | 2847 | 39 | 2817 |
| 41 | 3292 | 40 | 3126 |
| 40 | 3473 | 37 | 2539 |
| 37 | 2628 | 36 | 2412 |
| 38 | 3176 | 38 | 2991 |
| 40 | 3421 | 39 | 2875 |
| 38 | 2975 | 40 | 3231 |

2. Linear (Gaussian) Models

Table 2.1.: `birthweight` data (Dobson and Barnett (2018) Example 2.2.2)

| boys gestational age | boys weight | girls gestational age | girls weight |
|----------------------|-------------|-----------------------|--------------|
|----------------------|-------------|-----------------------|--------------|

2.2.1.2. Reshape data for graphing

```
bw =  
  birthweight |>  
  pivot_longer(  
    cols = everything(),  
    names_to = c("sex", ".value"),  
    names_sep = "s "  
  ) |>  
  rename(age = `gestational age`) |>  
  mutate(  
    sex = sex |>  
    case_match(  
      "boy" ~ "male",  
      "girl" ~ "female") |>  
    factor(levels = c("female", "male"))  
  )  
bw
```

Table 2.2.: `birthweight` data reshaped

| sex | age | weight |
|--------|-----|--------|
| male | 40 | 2968 |
| female | 40 | 3317 |
| male | 38 | 2795 |
| female | 36 | 2729 |
| male | 40 | 3163 |

2. Linear (Gaussian) Models

Table 2.2.: `birthweight` data reshaped

| sex | age | weight |
|--------|-----|--------|
| female | 40 | 2935 |
| male | 35 | 2925 |
| female | 38 | 2754 |
| male | 36 | 2625 |
| female | 42 | 3210 |
| male | 37 | 2847 |
| female | 39 | 2817 |
| male | 41 | 3292 |
| female | 40 | 3126 |
| male | 40 | 3473 |
| female | 37 | 2539 |
| male | 37 | 2628 |
| female | 36 | 2412 |
| male | 38 | 3176 |
| female | 38 | 2991 |
| male | 40 | 3421 |
| female | 39 | 2875 |
| male | 38 | 2975 |
| female | 40 | 3231 |

2.2.1.3. Data as graph

```
plot1 = bw |>
  ggplot(aes(
    x = age,
    y = weight,
    linetype = sex,
    shape = sex,
    col = sex)) +
```

2. Linear (Gaussian) Models

```
theme_bw() +  
  xlab("Gestational age (weeks)") +  
  ylab("Birthweight (grams)") +  
  theme(legend.position = "bottom") +  
  # expand_limits(y = 0, x = 0) +  
  geom_point(alpha = .7)  
print(plot1 + facet_wrap(~ sex))
```




Figure 2.1.: `birthweight` data (Dobson and Barnett (2018) Example 2.2.2)

2. Linear (Gaussian) Models

2.2.1.4. Data notation

Let's define some notation to represent this data.

- Y : birthweight (measured in grams)
- A : chromosomal sex:
 - $A = 0$ if female (XX)
 - $A = 1$ if male (XY)
- S : gestational age at birth (measured in weeks).

Note

Female is the **reference level** for the categorical variable A (chromosomal sex). The choice of a reference level is arbitrary and does not limit what we can do with the resulting model; it only makes it more computationally convenient to make inferences about comparisons involving that reference group.

2.2.2. Parallel lines regression

We don't have enough data to model the distribution of birth weight separately for each combination of gestational age and sex, so let's instead consider a (relatively) simple model for how that distribution varies with gestational age and sex:

$$p(Y = y|A = a, S = s) \sim_{\text{iid}} N(\mu(a, s), \sigma^2)$$

$$\begin{aligned}\mu(a, s) &\stackrel{\text{def}}{=} \mathbb{E}[Y|A = a, S = s] \\ &= \beta_0 + \beta_A a + \beta_S s\end{aligned}\tag{2.1}$$

2. Linear (Gaussian) Models

Table 2.3 shows the parameter estimates from R. Figure 2.2 shows the estimated model, superimposed on the data.

```
bw_lm1 = lm(  
  formula = weight ~ sex + age,  
  data = bw)  
  
bw_lm1 |>  
  parameters() |>  
  print_md(  
    include_reference = TRUE,  
    # show_sigma = TRUE,  
    select = "{estimate}")
```

Table 2.3.: Estimate of Equation 2.1 for birthweight data

| Parameter | Estimate |
|--------------|----------|
| (Intercept) | -1773.32 |
| sex (female) | 0.00 |
| sex (male) | 163.04 |
| age | 120.89 |

```
bw =  
  bw |>  
  mutate(`E[Y|X=x]` = fitted(bw_lm1)) |>  
  arrange(sex, age)  
  
plot2 =  
  plot1 %+% bw +  
  geom_line(aes(y = `E[Y|X=x]`))  
  
print(plot2)
```

2. Linear (Gaussian) Models




Figure 2.2.: Parallel-slopes model of birthweight

2.2.2.1. Model assumptions and predictions

To learn what this model is assuming, let's plug in a few values.

Exercise 2.1. According to this model, what's the mean birthweight for a female born at 36 weeks?

```
coef(bw_lm1)
```

| (Intercept) | sexmale | age |
|-------------|-----------|-----------|
| -1773.32184 | 163.03930 | 120.89433 |

2. Linear (Gaussian) Models

Solution.

```
pred_female = coef(bw_lm1)["(Intercept)"] + coef(bw_lm1)["age"]*36  
  
# print(pred_female)  
### built-in prediction:  
# predict(bw_lm1, newdata = tibble(sex = "female", age = 36))
```

$$\begin{aligned} E[Y|A = 0, S = 36] &= \beta_0 + \beta_S \cdot 0 + \beta_A \cdot 36 \\ &= 2578.873934 \end{aligned}$$

Exercise 2.2. What's the mean birthweight for a male born at 36 weeks?

```
coef(bw_lm1)
```

| (Intercept) | sexmale | age |
|-------------|-----------|-----------|
| -1773.32184 | 163.03930 | 120.89433 |

Solution.

```
pred_male =  
  coef(bw_lm1)["(Intercept)"] +  
  coef(bw_lm1)["sexmale"] +  
  coef(bw_lm1)["age"]*36
```

2. Linear (Gaussian) Models

$$\begin{aligned} E[Y|A = 1, S = 36] &= \beta_0 + \beta_S \cdot 1 + \beta_A \cdot 36 \\ &= 2741.91323693 \end{aligned}$$

Exercise 2.3. What's the difference in mean birthweights between males born at 36 weeks and females born at 36 weeks?

```
coef(bw_lm1)
```

| (Intercept) | sexmale | age |
|-------------|-----------|-----------|
| -1773.32184 | 163.03930 | 120.89433 |

Solution.

$$\begin{aligned} E[Y|A = 1, S = 36] - E[Y|A = 0, S = 36] \\ &= 2741.91323693 - 2578.873934 \\ &= 163.03930293 \end{aligned}$$

Shortcut:

$$\begin{aligned} E[Y|A = 1, S = 36] - E[Y|A = 0, S = 36] \\ &= (\beta_0 + \beta_S \cdot 1 + \beta_A \cdot 36) - (\beta_0 + \beta_S \cdot 0 + \beta_A \cdot 36) \\ &= \beta_S \\ &= 163.03930293 \end{aligned}$$

Note that age doesn't show up in this difference: in other words, according to this model, the difference between females and males with the same gestational age is the same for every age.

2. Linear (Gaussian) Models

That’s an assumption of the model; it’s built-in to the parametric structure, even before we plug in the estimated values of those parameters.

That’s why the lines are parallel.

2.2.3. Interactions

What if we don’t like that parallel lines assumption?

Then we need to allow an “interaction” between age A and sex S :

$$E[Y|A = a, S = s] = \beta_0 + \beta_A a + \beta_S s + \beta_{AS}(a \cdot s) \quad (2.2)$$

Now, the slope of mean birthweight $E[Y|A, S]$ with respect to gestational age A depends on the value of sex S .

```
bw_lm2 = lm(weight ~ sex + age + sex:age, data = bw)
bw_lm2 |>
  parameters() |>
  print_md(
    include_reference = TRUE,
    # show_sigma = TRUE,
    select = "{estimate}")
```

Table 2.4.: Birthweight model with interaction term

| Parameter | Estimate |
|-------------------------|----------|
| (Intercept) | -2141.67 |
| sex (female) | 0.00 |
| sex (male) | 872.99 |
| age | 130.40 |
| sex (male) \times age | -18.42 |

2. Linear (Gaussian) Models

```
bw =  
  bw |>  
  mutate(  
    predlm2 = predict(bw_lm2)  
  ) |>  
  arrange(sex, age)  
  
plot1_interact =  
  plot1 %>% bw +  
  geom_line(aes(y = predlm2))  
  
print(plot1_interact)
```




Figure 2.3.: Birthweight model with interaction term

2. Linear (Gaussian) Models

Now we can see that the lines aren't parallel.

Here's another way we could rewrite this model (by collecting terms involving S):

$$E[Y|A, S] = \beta_0 + \beta_A A + (\beta_S + \beta_{AS} A)S$$

i Note

If you want to understand a coefficient in a model with interactions, collect terms for the corresponding variable, and you will see what other variables are interacting with the variable you are interested in.

In this case, the coefficient S is interacting with A . So the slope of Y with respect to X_2 depends on the value of X_2 .

According to this model, there is no such thing as “*the* slope of birthweight with respect to age”. There are two slopes, one for each sex.¹ We can only talk about “the slope of birthweight with respect to age among males” and “the slope of birthweight with respect to age among females”.

Then: that coefficient is the difference in means per unit change in its corresponding coefficient, when the other collected variables are set to 0.

To learn what this model is assuming, let's plug in a few values.

¹using the definite article “the” would mean there is only one slope.

2. Linear (Gaussian) Models

Exercise 2.4. According to this model, what's the mean birthweight for a female born at 36 weeks?

```
pred_female = coef(bw_lm2)["(Intercept)"] + coef(bw_lm2)["age"]*36
```

$$E[Y|A = 0, X_2 = 36] = \beta_0 + \beta_S \cdot 0 + \beta_A \cdot 36 + \beta_{AS} \cdot (0 \cdot 36) = 2552.73333333$$

What's the mean birthweight for a male born at 36 weeks?

```
pred_male =  
  coef(bw_lm2)["(Intercept)"] +  
  coef(bw_lm2)["sexmale"] +  
  coef(bw_lm2)["age"]*36 +  
  coef(bw_lm2)["sexmale:age"] * 36
```

$$\begin{aligned} E[Y|A = 0, X_2 = 36] &= \beta_0 + \beta_S \cdot 1 + \beta_A \cdot 36 + \beta_{AS} \cdot 1 \cdot 36 \\ &= 2762.70689655 \end{aligned}$$

Exercise 2.5. What's the difference in mean birthweights between males born at 36 weeks and females born at 36 weeks?

Solution.

$$\begin{aligned} &E[Y|S = 1, A = 36] - E[Y|S = 0, A = 36] \\ &= (\beta_0 + \beta_S \cdot 1 + \beta_A \cdot 36 + \beta_{AS} \cdot 1 \cdot 36) \\ &\quad - (\beta_0 + \beta_S \cdot 0 + \beta_A \cdot 36 + \beta_{AS} \cdot 0 \cdot 36) \\ &= \beta_S + \beta_{AS} \cdot 36 \\ &= 209.97356322 \end{aligned}$$

2. Linear (Gaussian) Models

Note that age now does show up in the difference: in other words, according to this model, the difference in mean birthweights between females and males with the same gestational age can vary by gestational age.

That's how the lines in the graph ended up non-parallel.

2.2.4. Stratified regression

We could re-write the interaction model as a stratified model, with a slope and intercept for each sex:

Let:

$$\mathbb{E}[Y|A = a, S = s] = \beta_M s + \beta_{AM}(a \cdot s) + \beta_F(1 - s) + \beta_{AF}(a \cdot (1 - s)) \quad (2.3)$$

Compare this stratified model with our interaction model, Equation 2.2:

$$\mathbb{E}[Y|A = a, S = s] = \beta_0 + \beta_A a + \beta_S s + \beta_{AS}(a \cdot s)$$

In the stratified model, the intercept term β_0 has been relabeled as β_F .

```
bw_lm2 = lm(weight ~ sex + age + sex:age, data = bw)
bw_lm2 |>
  parameters() |>
  print_md(
    include_reference = TRUE,
    # show_sigma = TRUE,
    select = "{estimate}")
```

2. Linear (Gaussian) Models

Table 2.5.: Birthweight model with interaction term

| Parameter | Estimate |
|-------------------------|----------|
| (Intercept) | -2141.67 |
| sex (female) | 0.00 |
| sex (male) | 872.99 |
| age | 130.40 |
| sex (male) \times age | -18.42 |

```
bw_lm_strat =  
  bw |>  
  lm(  
    formula = weight ~ sex + sex:age - 1,  
    data = _)  
  
bw_lm_strat |>  
  parameters() |>  
  print_md(  
    # show_sigma = TRUE,  
    select = "{estimate}")
```

Table 2.6.: Birthweight model - stratified betas

| Parameter | Estimate |
|---------------------------|----------|
| sex (female) | -2141.67 |
| sex (male) | -1268.67 |
| sex (female) \times age | 130.40 |
| sex (male) \times age | 111.98 |

2. Linear (Gaussian) Models

2.2.5. Curved-line regression

If we transform some of our covariates (X s) and plot the resulting model on the original covariate scale, we end up with curved regression lines:

```
bw_lm3 = lm(weight ~ sex:log(age) - 1, data = bw)
library(palmerpenguins)

ggpenguins <-
  palmerpenguins::penguins |>
  dplyr::filter(species == "Adelie") |>
  ggplot(
    aes(x = bill_length_mm , y = body_mass_g)) +
  geom_point() +
  xlab("Bill length (mm)") +
  ylab("Body mass (g)")

ggpenguins2 = ggpenguins +
  stat_smooth(
    method = "lm",
    formula = y ~ log(x),
    geom = "smooth") +
  xlab("Bill length (mm)") +
  ylab("Body mass (g)")

ggpenguins2 |> print()
```

2. Linear (Gaussian) Models




Figure 2.4.: palmerpenguins model with bill_length entering on log scale

2.3. Estimating Linear Models via Maximum Likelihood

2.3.1. Review of one-sample inference

Previously, we learned how to fit outcome-only models of the form $p(X = x|\theta)$ to iid data $\mathbf{x} = (x_1, \dots, x_n)$ using maximum likelihood estimation:

2. Linear (Gaussian) Models

$$\begin{aligned}\mathcal{L}(\mathbf{x}|\theta) &\stackrel{\text{def}}{=} p(X_1 = x_1, \dots, X_n = x_n|\theta) \\ &= \prod_{i=1}^n p(X = x_i|\theta)\end{aligned}\tag{2.4}$$

$$\begin{aligned}\ell(x|\theta) &\stackrel{\text{def}}{=} \log \{\mathcal{L}(x|\theta)\} \\ &= \sum_{i=1}^n \log \{p(X = x_i|\theta)\}\end{aligned}\tag{2.5}$$

$$\hat{\theta}_{\text{ML}} \stackrel{\text{def}}{=} \arg \max_{\theta} \ell(x|\theta)\tag{2.6}$$

Recall: the second equality in Equation 2.4 relies on the assumption that the X_i s are mutually independent.

We learned how to quantify our uncertainty about these maximum likelihood estimates; with sufficient sample size, $\hat{\theta}_{\text{ML}}$ has the approximate distribution:

$$\hat{\theta}_{\text{ML}} \sim N(\theta, \mathcal{J}(\theta)^{-1})$$

Recall:

- $\mathcal{J}(\theta) \stackrel{\text{def}}{=} \mathbb{E} [I(\tilde{X}; \theta)]$
- $I(\tilde{X}; \theta) \stackrel{\text{def}}{=} -\ell''(\tilde{X}; \theta)$

We can estimate $\mathcal{J}(\theta)$ using either $\mathcal{J}(\hat{\theta}_{\text{ML}})$ or $I(\tilde{x}; \hat{\theta}_{\text{ML}})$.

So we can estimate the standard error of $\hat{\theta}_k$ as:

$$\widehat{\text{SE}}(\hat{\theta}_k) = \sqrt{\left(\left(\hat{\mathcal{J}}(\hat{\theta}_{\text{ML}})\right)_{kk}\right)^{-1}}$$

2. Linear (Gaussian) Models

Then an asymptotic approximation of a 95% confidence interval for θ_k is

$$\hat{\theta}_{\text{ML}} \pm z_{0.975} \times \widehat{\text{SE}}(\hat{\theta}_k)$$

where z_β the β quantile of the standard Gaussian distribution.

2.3.2. MLEs for Linear Regression

Let's use maximum likelihood again:

$$\mathcal{L}(\mathbf{y}|\mathbf{x}, \beta, \sigma^2) = \prod_{i=1}^n (2\pi\sigma^2)^{-1/2} \exp\left\{-\frac{1}{2\sigma^2}(y_i - x'_i\beta)^2\right\}$$

$$\ell(\mathbf{y}|\mathbf{x}, \beta, \sigma^2) \propto -\frac{n}{2}\log\{\sigma^2\} - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - x'_i\beta)^2$$

$$\ell'(\mathbf{y}|\mathbf{x}, \beta, \sigma^2) \propto -\frac{n}{2}\log\{\sigma^2\} - \frac{1}{2\sigma^2} \frac{d}{d\beta} \left(\sum_{i=1}^n (y_i - x'_i\beta)^2 \right)$$

A few tools from linear algebra will make this analysis go easier (see Fieller (2016), Section 7.2 for details).

$$f_\beta(\mathbf{x}) = (f_\beta(x_1), f_\beta(x_2), \dots, f_\beta(x_n))^\top$$

Let \mathbf{x} and β be vectors of length p , or in other words, matrices of length $p \times 1$:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix} \quad \beta = \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{bmatrix}$$

2. Linear (Gaussian) Models

Then

$$x' \equiv x^\top \equiv [x_1, x_2, \dots, x_p]$$

and

$$x'\beta = [x_1, x_2, \dots, x_p] \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{bmatrix} = x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p$$

If $f(\beta)$ is a function that takes β as input and outputs a scalar, such as $f(\beta) = x'\beta$, then:

$$\frac{d}{d\beta}f(\beta) = \begin{bmatrix} \frac{d}{d\beta_1}f(\beta) \\ \frac{d}{d\beta_2}f(\beta) \\ \vdots \\ \frac{d}{d\beta_p}f(\beta) \end{bmatrix}$$

In particular, if $f(\beta) = x'\beta$, then:

$$\frac{d}{d\beta}x'\beta = \begin{bmatrix} \frac{d}{d\beta_1}(x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p) \\ \frac{d}{d\beta_2}(x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p) \\ \vdots \\ \frac{d}{d\beta_p}(x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p) \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix} = \mathbf{x}$$

In general:

$$\frac{d}{d\beta}x'\beta = x$$

2. Linear (Gaussian) Models

This looks a lot like non-vector calculus, except that you have to transpose the coefficient.

Similarly,

$$\frac{d}{d\beta} \beta' \beta = 2\beta$$

This is like taking the derivative of x^2 .

And finally, if S is a $p \times p$ matrix, then:

$$\frac{d}{d\beta} \beta' S \beta = 2S\beta$$

Again, this is like taking the derivative of cx^2 with respect to x in non-vector calculus.

Thus:

$$\sum_{i=1}^n (y_i - f_{\beta}(x_i))^2 = (\mathbf{y} - X\beta)'(\mathbf{y} - X\beta)$$

$$(\mathbf{y} - X\beta)' = (\mathbf{y}' - (X\beta)') = (\mathbf{y}' - \beta' X')$$

So

$$\begin{aligned} (\mathbf{y} - X\beta)'(\mathbf{y} - X\beta) &= (\mathbf{y}' - \beta' X')(\mathbf{y} - X\beta) \\ &= \mathbf{y}'\mathbf{y} - \beta' X'\mathbf{y} - \mathbf{y}' X\beta + \beta' X' X\beta \\ &= \mathbf{y}'\mathbf{y} - 2\mathbf{y}' X\beta + \beta' X' X\beta \end{aligned}$$

So

2. Linear (Gaussian) Models

$$\begin{aligned}
 \frac{d}{d\beta} \left(\sum_{i=1}^n (y_i - x_i' \beta)^2 \right) &= \frac{d}{d\beta} (\mathbf{y} - X\beta)' (\mathbf{y} - X\beta) \\
 &= \frac{d}{d\beta} (y' y - 2y' X\beta + \beta' X' X\beta) \\
 &= (-2X'y + 2X' X\beta)
 \end{aligned}$$

So if $\ell(\beta, \sigma^2) = 0$, then

$$\begin{aligned}
 0 &= (-2X'y + 2X' X\beta) \\
 2X'y &= 2X' X\beta \\
 X'y &= X' X\beta \\
 (X' X)^{-1} X'y &= \beta
 \end{aligned}$$

The second derivative matrix $\ell''_{\beta, \beta'}(\beta, \sigma^2; \mathbf{X}, \mathbf{y})$ is negative definite at $\beta = (X' X)^{-1} X'y$, so $\hat{\beta}_{ML} = (X' X)^{-1} X'y$ is the MLE for β .

Similarly (not shown):

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} (Y - X\hat{\beta})' (Y - X\hat{\beta})$$

And

$$\begin{aligned}
 J_{\beta} &= E[-\ell''_{\beta, \beta'}(Y|X, \beta, \sigma^2)] \\
 &= \frac{1}{\sigma^2} X' X
 \end{aligned}$$

So:

$$Var(\hat{\beta}) \approx (J_{\beta})^{-1} = \sigma^2 (X' X)^{-1}$$

2. Linear (Gaussian) Models

and

$$\hat{\beta} \sim N(\beta, \mathcal{J}_{\beta}^{-1})$$

These are all results you have hopefully seen before.

In the Gaussian linear regression case, we also have exact results:

$$\frac{\hat{\beta}_j}{\hat{\mathcal{J}}(\beta)_j} \sim t_{n-p}$$

In our model 2 above, $\hat{\mathcal{J}}(\beta)$ is:

```
bw_lm2 |> vcov()
```

| | (Intercept) | sexmale | age | sexmale:age |
|-------------|--------------|--------------|--------------|--------------|
| (Intercept) | 1353968.361 | -1353968.361 | -34870.96582 | 34870.96582 |
| sexmale | -1353968.361 | 2596387.127 | 34870.96582 | -67210.97445 |
| age | -34870.966 | 34870.966 | 899.89589 | -899.89589 |
| sexmale:age | 34870.966 | -67210.974 | -899.89589 | 1743.54829 |

If we take the square roots of the diagonals, we get the standard errors listed in the model output:

```
bw_lm2 |> vcov() |> diag() |> sqrt()
```

| (Intercept) | sexmale | age | sexmale:age |
|-------------|-------------|-----------|-------------|
| 1163.601461 | 1611.330856 | 29.998265 | 41.755817 |

2. Linear (Gaussian) Models

```
bw_lm2 |> parameters() |> print_md()
```

Table 2.8.: Estimated model for `birthweight` data with interaction term

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|------------------|-------------|---------|---------------------|-------|--------|
| (Intercept) | -2141.67 | 1163.60 | (-4568.90, 285.56) | -1.84 | 0.081 |
| sex (male) | 872.99 | 1611.33 | (-2488.18, 4234.17) | 0.54 | 0.594 |
| age | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

So we can do confidence intervals, hypothesis tests, and p-values exactly as in the one-variable case we looked at previously.

2.4. Inference about Gaussian Linear Regression Models

2.4.1. Motivating example: birthweight data

Research question: is there really an interaction between sex and age?

$$H_0 : \beta_{AS} = 0$$

$$H_A : \beta_{AS} \neq 0$$

$$P(|\hat{\beta}_{AS}| > |-18.41724138| \mid H_0) = ?$$

2. Linear (Gaussian) Models

2.4.2. Wald tests and CIs

R can give you Wald tests for single coefficients and corresponding CIs:

```
bw_lm2 |>
  parameters() |>
  print_md(
    include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---------------------|-------------|---------|------------------------|-------|-----------|
| (Intercept) | -2141.67 | 1163.60 | (-4568.90, 285.56) | -1.84 | 0.081 |
| sex (female) | 0.00 | | | | |
| sex (male) | 872.99 | 1611.33 | (-2488.18, 4234.17) | 0.54 | 0.594 |
| age | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

To understand what's happening, let's replicate these results by hand for the interaction term.

2.4.3. P-values

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
    include_reference = TRUE)
```

2. Linear (Gaussian) Models

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|------------------|-------------|-------|------------------|-------|-------|
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

```
beta_hat = coef(summary(bw_lm2))["sexmale:age", "Estimate"]
se_hat = coef(summary(bw_lm2))["sexmale:age", "Std. Error"]
dfresid = bw_lm2$df.residual
t_stat = abs(beta_hat)/se_hat
pval_t =
  pt(-t_stat, df = dfresid, lower.tail = TRUE) +
  pt(t_stat, df = dfresid, lower.tail = FALSE)
```

$$\begin{aligned}
& P(|\hat{\beta}_{AS}| > |-18.41724138| | H_0) \\
&= P\left(\left|\frac{\hat{\beta}_{AS}}{\hat{SE}(\hat{\beta}_{AS})}\right| > \left|\frac{-18.41724138}{41.75581745}\right| \middle| H_0\right) \\
&= P(|T_{20}| > 0.44107007 | H_0) \\
&= 0.66389342
\end{aligned}$$

This matches the result in the table above.

2.4.4. Confidence intervals

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
    include_reference = TRUE)
```

2. Linear (Gaussian) Models

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---------------------|-------------|-------|------------------|-------|-------|
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

```
q_t = qt(
  p = 0.975,
  df = dfresid,
  lower.tail = TRUE)

q_t = qt(
  p = 0.025,
  df = dfresid,
  lower.tail = TRUE)

confint_radius_t =
  se_hat * q_t

confint_t = beta_hat + c(-1,1) * confint_radius_t

print(confint_t)
```

```
[1] 68.683868 -105.518350
```

This also matches.

2.4.5. Gaussian approximations

Here are the asymptotic (Gaussian approximation) equivalents:

2. Linear (Gaussian) Models

2.4.6. P-values

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
    include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---------------------|-------------|-------|------------------|-------|-------|
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

```
pval_z = pnorm(abs(t_stat), lower = FALSE) * 2
print(pval_z)
```

```
[1] 0.65916227
```

2.4.7. Confidence intervals

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
    include_reference = TRUE)
```


2. Linear (Gaussian) Models

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---------------------|-------------|-------|------------------|-------|-------|
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

```
confint_radius_z = se_hat * qnorm(0.975, lower = TRUE)
confint_z =
  beta_hat + c(-1,1) * confint_radius_z
print(confint_z)
```

```
[1] -100.257140    63.422657
```

2.4.8. Likelihood ratio statistics

```
logLik(bw_lm2)
```

```
'log Lik.' -156.57929 (df=5)
```

```
logLik(bw_lm1)
```

```
'log Lik.' -156.69545 (df=4)
```

```
llR = (logLik(bw_lm2) - logLik(bw_lm1)) |> as.numeric()
delta_df = (bw_lm1$df.residual - df.residual(bw_lm2))
```

```
x_max = 1
```

2. Linear (Gaussian) Models

```
d_LLR = function(x, df = delta_df) dchisq(x, df = df)

chisq_plot =
  ggplot() +
  geom_function(fun = d_LLR) +
  stat_function(fun = d_LLR, xlim = c(1LR, x_max), geom = "area", fill = "gray") +
  geom_segment(aes(x = 1LR, xend = 1LR, y = 0, yend = d_LLR(1LR)), col = "red") +
  xlim(0.0001, x_max) +
  ylim(0, 4) +
  ylab("p(X=x)") +
  xlab("log(likelihood ratio) statistic [x]") +
  theme_classic()
chisq_plot |> print()
```




Figure 2.5.: Chi-square distribution

2. Linear (Gaussian) Models

Now we can get the p-value:

```
pchisq(
  q = 2*LLR,
  df = delta_df,
  lower = FALSE) |>
print()
```

```
[1] 0.62980648
```

In practice you don't have to do this by hand; there are functions to do it for you:

```
# built in
library(lmtest)
lrtest(bw_lm2, bw_lm1)
```

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|------------|----|------------|------------|
| 5 | -156.57929 | NA | NA | NA |
| 4 | -156.69545 | -1 | 0.23232327 | 0.62980648 |

2.5. Goodness of fit

2.5.1. AIC and BIC

When we use likelihood ratio tests, we are comparing how well different models fit the data.

2. Linear (Gaussian) Models

Likelihood ratio tests require “nested” models: one must be a special case of the other.

If we have non-nested models, we can instead use the Akaike Information Criterion (AIC) or Bayesian Information Criterion (BIC):

- $\text{AIC} = -2 * \ell(\hat{\theta}) + 2 * p$
- $\text{BIC} = -2 * \ell(\hat{\theta}) + p * \log(n)$

where ℓ is the log-likelihood of the data evaluated using the parameter estimates $\hat{\theta}$, p is the number of estimated parameters in the model (including $\hat{\sigma}^2$), and n is the number of observations.

You can calculate these criteria using the `logLik()` function, or use the built-in R functions:

2.5.1.1. AIC in R

```
-2 * logLik(bw_lm2) |> as.numeric() +  
  2*(length(coef(bw_lm2))+1) # sigma counts as a parameter here
```

```
[1] 323.15858
```

```
AIC(bw_lm2)
```

```
[1] 323.15858
```

2.5.1.2. BIC in R

2. Linear (Gaussian) Models

```
-2 * logLik(bw_lm2) |> as.numeric() +  
  (length(coef(bw_lm2))+1) * log(nobs(bw_lm2))
```

```
[1] 329.04884
```

```
BIC(bw_lm2)
```

```
[1] 329.04884
```

Large values of AIC and BIC are worse than small values. There are no hypothesis tests or p-values associated with these criteria.

2.5.2. (Residual) Deviance

Let q be the number of distinct covariate combinations in a data set.

```
bw.X.unique =  
  bw |>  
  count(sex, age)  
  
n_unique.bw = nrow(bw.X.unique)
```

For example, in the `birthweight` data, there are $q = 12$ unique patterns:

```
bw.X.unique |>  
  pander(  
    row.names = rownames(bw.X.unique))
```

2. Linear (Gaussian) Models

| | sex | age | n |
|-----------|--------|-----|---|
| 1 | female | 36 | 2 |
| 2 | female | 37 | 1 |
| 3 | female | 38 | 2 |
| 4 | female | 39 | 2 |
| 5 | female | 40 | 4 |
| 6 | female | 42 | 1 |
| 7 | male | 35 | 1 |
| 8 | male | 36 | 1 |
| 9 | male | 37 | 2 |
| 10 | male | 38 | 3 |
| 11 | male | 40 | 4 |
| 12 | male | 41 | 1 |

i Note

If a given covariate pattern has more than one observation in a dataset, those observations are called **replicates**.

Then the most complicated model we could fit would have one parameter (a mean) for each of these combinations, plus a variance parameter:

```
lm_max =  
  bw |>  
  mutate(age = factor(age)) |>  
  lm(  
    formula = weight ~ sex:age - 1,  
    data = _)  
  
lm_max |>  
  parameters() |>  
  print_md()
```

2. Linear (Gaussian) Models

| Parameter | Coefficient | SE | 95% CI | t(12) | p |
|-------------------------|-------------|--------|-----------------------|-------|-----------|
| sex (male) × age35 | 2925.00 | 187.92 | (2515.55, 3334.45) | 15.56 | < .001 |
| sex (female) × age36 | 2570.50 | 132.88 | (2280.98, 2860.02) | 19.34 | < .001 |
| sex (male) × age36 | 2625.00 | 187.92 | (2215.55, 3034.45) | 13.97 | < .001 |
| sex (female) × age37 | 2539.00 | 187.92 | (2129.55, 2948.45) | 13.51 | < .001 |
| sex (male) × age37 | 2737.50 | 132.88 | (2447.98, 3027.02) | 20.60 | < .001 |
| sex (female) × age38 | 2872.50 | 132.88 | (2582.98, 3162.02) | 21.62 | < .001 |
| sex (male) × age38 | 2982.00 | 108.50 | (2745.60, 3218.40) | 27.48 | < .001 |
| sex (female) × age39 | 2846.00 | 132.88 | (2556.48, 3135.52) | 21.42 | < .001 |
| sex (female) × age40 | 3152.25 | 93.96 | (2947.52, 3356.98) | 33.55 | < .001 |
| sex (male) × age40 | 3256.25 | 93.96 | (3051.52, 3460.98) | 34.66 | < .001 |
| sex (male) × age41 | 3292.00 | 187.92 | (2882.55, 3701.45) | 17.52 | < .001 |
| sex (female) × age42 | 3210.00 | 187.92 | (2800.55, 3619.45) | 17.08 | < .001 |

We call this model the **full**, **maximal**, or **saturated** model for this dataset.

We can calculate the log-likelihood of this model as usual:

2. Linear (Gaussian) Models

```
logLik(lm_max)
```

```
'log Lik.' -151.4016 (df=13)
```

We can compare this model to our other models using chi-square tests, as usual:

```
lrtest(lm_max, bw_lm2)
```

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|------------|----|-----------|------------|
| 13 | -151.40160 | NA | NA | NA |
| 5 | -156.57929 | -8 | 10.355374 | 0.24096372 |

The likelihood ratio statistic for this test is

$$\lambda = 2 * (\ell_{\text{full}} - \ell) = 10.35537421$$

where:

- ℓ_{max} is the log-likelihood of the full model: -151.40160056
- ℓ is the log-likelihood of our comparison model (two slopes, two intercepts): -156.57928767

This statistic is called the **deviance** or **residual deviance** for our two-slopes and two-intercepts model; it tells us how much the likelihood of that model deviates from the likelihood of the maximal model.

The corresponding p-value tells us whether there we have enough evidence to detect that our two-slopes, two-intercepts model is a worse fit for the data than the maximal model; in other words, it tells us if there's evidence that we missed any important patterns. (Remember, a nonsignificant p-value could mean that we didn't miss anything and a more complicated model is unnecessary, or it could mean we just don't have enough data to tell the difference between these models.)

2. Linear (Gaussian) Models

2.5.3. Null Deviance

Similarly, the *least* complicated model we could fit would have only one mean parameter, an intercept:

$$E[Y|X = x] = \beta_0$$

We can fit this model in R like so:

```
lm0 = lm(weight ~ 1, data = bw)

lm0 |> parameters() |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(23) | p |
|-------------|-------------|-------|--------------------|-------|--------|
| (Intercept) | 2967.67 | 57.58 | (2848.56, 3086.77) | 51.54 | < .001 |

This model also has a likelihood:

```
logLik(lm0)
```

```
'log Lik.' -168.95497 (df=2)
```

And we can compare it to more complicated models using a likelihood ratio test:

```
lrtest(bw_lm2, lm0)
```

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|------------|----|-------|------------|
| 5 | -156.57929 | NA | NA | NA |

2. Linear (Gaussian) Models

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|------------|----|-----------|------------|
| 2 | -168.95497 | -3 | 24.751358 | 1.74e-05 |

The likelihood ratio statistic for the test comparing the null model to the maximal model is

$$\lambda = 2 * (\ell_{\text{full}} - \ell_0) = 35.10673188$$

where:

- ℓ_0 is the log-likelihood of the null model: -168.9549665
- ℓ_{full} is the log-likelihood of the maximal model: -151.40160056

In R, this test is:

```
lrtest(lm_max, lm0)
```

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|------------|-----|-----------|------------|
| 13 | -151.40160 | NA | NA | NA |
| 2 | -168.95497 | -11 | 35.106732 | 0.00023795 |

This log-likelihood ratio statistic is called the **null deviance**. It tells us whether we have enough data to detect a difference between the null and full models.

2.6. Rescaling

2.6.1. Rescale age

2. Linear (Gaussian) Models

```
bw =  
  bw |>  
  mutate(  
    `age - mean` = age - mean(age),  
    `age - 36wks` = age - 36  
  )  
  
lm1c = lm(weight ~ sex + `age - 36wks`, data = bw)  
  
lm2c = lm(weight ~ sex + `age - 36wks` + sex:`age - 36wks`, data = bw)  
  
parameters(lm2c, ci_method = "wald") |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|-----------------------------|-------------|--------|-----------------------|-------|-----------|
| (Intercept) | 2552.73 | 97.59 | (2349.16, 2756.30) | 26.16 | < .001 |
| sex (male) | 209.97 | 129.75 | (-60.68, 480.63) | 1.62 | 0.121 |
| age - 36wks | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| sex (male) × age - 36wks | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

Compare with what we got without rescaling:

```
parameters(bw_lm2, ci_method = "wald") |> print_md()
```

2. Linear (Gaussian) Models

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|------------------|-------------|---------|---------------------|-------|--------|
| (Intercept) | -2141.67 | 1163.60 | (-4568.90, 285.56) | -1.84 | 0.081 |
| sex (male) | 872.99 | 1611.33 | (-2488.18, 4234.17) | 0.54 | 0.594 |
| age | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| sex (male) × age | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

2.7. Prediction

2.7.1. Prediction for linear models

Definition 2.1 (Predicted value). In a regression model $p(y|x)$, the **predicted value** of y given x is the estimated mean of Y given X :

$$\hat{y} \stackrel{\text{def}}{=} \hat{E}[Y|X = x]$$

For linear models, the predicted value can be straightforwardly calculated by multiplying each predictor value x_j by its corresponding coefficient β_j and adding up the results:

$$\begin{aligned} \hat{Y} &= \hat{E}[Y|X = x] \\ &= x' \hat{\beta} \\ &= \hat{\beta}_0 \cdot 1 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_p x_p \end{aligned}$$

2. Linear (Gaussian) Models

2.7.2. Example: prediction for the birthweight data

```
X = c(1,1,40)
sum(X * coef(bw_lm1))
```

```
[1] 3225.4905
```

R has built-in functions for prediction:

```
x = tibble(age = 40, sex = "male")
bw_lm1 |> predict(newdata = x)
```

```
1
3225.4905
```

If you don't provide `newdata`, R will use the covariate values from the original dataset:

```
predict(bw_lm1)
```

| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 3225.4905 | 3062.4512 | 2983.7019 | 2578.8739 | 3225.4905 | 3062.4512 | 2621.0189 | 2820.6626 |
| 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| 2741.9132 | 3304.2399 | 2862.8076 | 2941.5569 | 3346.3849 | 3062.4512 | 3225.4905 | 2699.7683 |
| 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 |
| 2862.8076 | 2578.8739 | 2983.7019 | 2820.6626 | 3225.4905 | 2941.5569 | 2983.7019 | 3062.4512 |

These special predictions are called the *fitted values* of the dataset:

2. Linear (Gaussian) Models

Definition 2.2. For a given dataset (\tilde{Y}, \tilde{X}) and corresponding fitted model $p_{\hat{\beta}}(y|x)$, the **fitted value** of y_i is the predicted value of y when $X = x_i$ using the estimate parameters $\hat{\beta}$.

R has an extra function to get these values:

```
fitted(bw_lm1)
```

| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 3225.4905 | 3062.4512 | 2983.7019 | 2578.8739 | 3225.4905 | 3062.4512 | 2621.0189 | 2820.6626 |
| 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| 2741.9132 | 3304.2399 | 2862.8076 | 2941.5569 | 3346.3849 | 3062.4512 | 3225.4905 | 2699.7683 |
| 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 |
| 2862.8076 | 2578.8739 | 2983.7019 | 2820.6626 | 3225.4905 | 2941.5569 | 2983.7019 | 3062.4512 |

2.7.3. Quantifying uncertainty in predictions

```
bw_lm1 |>  
  predict(  
    newdata = x,  
    se.fit = TRUE)
```

```
$fit
```

```
      1  
3225.4905
```

```
$se.fit
```

```
[1] 61.459866
```

```
$df
```

2. Linear (Gaussian) Models

```
[1] 21
```

```
$residual.scale
```

```
[1] 177.11588
```

This is a `list()`; you can extract the elements with `$` or `magrittr::use_series()`:

```
bw_lm1 |>
  predict(
    newdata = x,
    se.fit = TRUE) |>
  use_series(se.fit)
```

```
[1] 61.459866
```

You can get **confidence intervals** for $\mathbb{E}[Y|X = x]$:

```
bw_lm1 |> predict(
  newdata = x,
  interval = "confidence")
```

| fit | lwr | upr |
|-----------|-----------|-----------|
| 3225.4905 | 3097.6778 | 3353.3033 |

You can also get **prediction intervals** for the value of an individual outcome Y :

```
bw_lm1 |>
  predict(newdata = x, interval = "predict")
```

2. Linear (Gaussian) Models

| fit | lwr | upr |
|-----------|-----------|-----------|
| 3225.4905 | 2835.6123 | 3615.3688 |

The warning from the last command is: “predictions on current data refer to *future* responses” (since you already know what happened to the current data, and thus don’t need to predict it).

See `?predict.lm` for more.

2.8. Diagnostics

2.8.1. Residuals

Definition 2.3 (Residual noise). The **residual noise** in a probability model $p(Y)$ is the difference between an observed value y and its distributional mean:

$$\epsilon(y) \stackrel{\text{def}}{=} y - \mathbb{E}[Y] \quad (2.7)$$

This is the same notation that we used for **errors**. Each observation y can be viewed as an estimate of $\mathbb{E}[Y]$ (albeit an imprecise one, individually, since $n = 1$). Likewise, $\mathbb{E}[Y]$ can be viewed as an estimate of Y , before y is observed.

We can rearrange Equation 2.7 to view y as the sum of its mean plus the residual noise:

$$y = \mathbb{E}[Y|X] + \epsilon(y)$$

2. Linear (Gaussian) Models

Theorem 2.1 (Residuals in Gaussian models). *If Y has a Gaussian distribution, then $\epsilon(Y)$ also has a Gaussian distribution, and vice versa.*

Proof. Left to the reader. □

Definition 2.4 (Residual errors of a fitted model value). The **residual of a fitted value** \hat{y} (shorthand: “residual”) is its **error**:

$$\begin{aligned} e(\hat{y}) &\stackrel{\text{def}}{=} \epsilon(\hat{y}) \\ &= y - \hat{y} \end{aligned}$$

$e(\hat{y})$ can be seen as the maximum likelihood estimate of the residual noise:

$$\begin{aligned} e(\hat{y}) &= y - \hat{y} \\ &= \hat{\epsilon}_{ML} \end{aligned}$$

Example 2.1 (residuals in `birthweight` data). R provides a function for residuals:

```
resid(bw_lm2)
```

| | | | | | |
|------------|-------------|-------------|------------|------------|-------------|
| 1 | 2 | 3 | 4 | 5 | 6 |
| 176.266667 | -140.733333 | -144.133333 | -59.533333 | 177.466667 | -126.933333 |
| 7 | 8 | 9 | 10 | 11 | 12 |
| -68.933333 | 242.666667 | -139.333333 | 51.666667 | 156.666667 | -125.133333 |
| 13 | 14 | 15 | 16 | 17 | 18 |

2. Linear (Gaussian) Models

| | | | | | |
|------------|-------------|------------|-------------|-------------|------------|
| 274.275862 | -137.706897 | -27.689655 | -246.689655 | -191.672414 | 189.327586 |
| 19 | 20 | 21 | 22 | 23 | 24 |
| -11.672414 | -242.637931 | -47.637931 | 262.362069 | 210.362069 | -30.620690 |

Exercise 2.6. Check R's output by computing the residuals directly.

Solution.

```
bw$weight - fitted(bw_lm2)
```

| | | | | | |
|------------|-------------|-------------|-------------|-------------|-------------|
| 1 | 2 | 3 | 4 | 5 | 6 |
| 176.266667 | -140.733333 | -144.133333 | -59.533333 | 177.466667 | -126.933333 |
| 7 | 8 | 9 | 10 | 11 | 12 |
| -68.933333 | 242.666667 | -139.333333 | 51.666667 | 156.666667 | -125.133333 |
| 13 | 14 | 15 | 16 | 17 | 18 |
| 274.275862 | -137.706897 | -27.689655 | -246.689655 | -191.672414 | 189.327586 |
| 19 | 20 | 21 | 22 | 23 | 24 |
| -11.672414 | -242.637931 | -47.637931 | 262.362069 | 210.362069 | -30.620690 |

This matches R's output!

2.8.1.1. General characteristics of residuals

For unbiased estimators $\hat{\theta}$:

$$\mathbb{E}[e(y)] = 0$$

2. Linear (Gaussian) Models

Proof.

$$\begin{aligned}\mathbb{E}[e(y)] &= \mathbb{E}[y - \hat{y}] \\ &= \mathbb{E}[y] - \mathbb{E}[\hat{y}] \\ &= \mathbb{E}[y] - \mathbb{E}[\hat{\mathbf{E}}[y]] \\ &= \mathbb{E}[y] - \mathbb{E}[\hat{\mathbf{E}}[y]] \\ &= \mathbb{E}[y] - \mathbb{E}[Ey] \\ &= \mathbb{E}[y] - \mathbb{E}[y] \\ &= 0\end{aligned}$$

□

$$\widehat{\text{SD}}(e(y)) \approx \hat{\sigma}$$

Proof.

$$\begin{aligned}\text{Var}(e(y)) &= \text{Var}(y - \hat{y}) \\ &= \text{Var}(y) + \text{Var}(\hat{y}) - 2\text{Cov}(y, \hat{y}) \\ &\approx \text{Var}(y) + 0 - 2 \cdot 0 \\ &= \text{Var}(y)\end{aligned}$$

□

2.8.1.2. Characteristics of residuals in Gaussian models

With enough data and a correct model, the residuals will be approximately Gaussian distributed, with variance σ^2 , which we can estimate using $\hat{\sigma}^2$: that is:

$$e_i \sim_{\text{iid}} N(0, \hat{\sigma}^2)$$

2. Linear (Gaussian) Models

Definition 2.5 (Standardized residuals).

$$r_i = \frac{e_i}{\widehat{SD}(e_i)}$$

Hence, with enough data and a correct model, the standardized residuals will be approximately standard Gaussian; that is,

$$r_i \sim_{\text{iid}} N(0, 1)$$

2.8.2. Marginal distributions of residuals

To look for problems with our model, we can check whether the residuals e_i and standardized residuals r_i look like they have the distributions that they are supposed to have, according to the model.

2.8.2.1. Standardized residuals in R

```
rstandard(bw_lm2)
```

| | | | | | |
|--------------|--------------|--------------|--------------|--------------|--------------|
| 1 | 2 | 3 | 4 | 5 | 6 |
| 1.159816604 | -0.926010912 | -0.874791659 | -0.347225534 | 1.035066486 | -0.734731462 |
| 7 | 8 | 9 | 10 | 11 | 12 |
| -0.399008578 | 1.437516422 | -0.825387176 | 0.306064623 | 0.928066921 | -0.876159249 |
| 13 | 14 | 15 | 16 | 17 | 18 |
| 1.914278022 | -0.865592113 | -0.164299301 | -1.463757406 | -1.110159894 | 1.096578735 |
| 19 | 20 | 21 | 22 | 23 | 24 |
| -0.067606211 | -1.461586459 | -0.286958245 | 1.580399427 | 1.267165237 | -0.198054338 |

2. Linear (Gaussian) Models

```
resid(bw_lm2)/sigma(bw_lm2)
```


| | 1 | 2 | 3 | 4 | 5 | 6 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|
| | 0.975933108 | -0.779196215 | -0.798020946 | -0.329617347 | 0.982577131 | -0.702789954 |
| | 7 | 8 | 9 | 10 | 11 | 12 |
| | -0.381662191 | 1.343569030 | -0.771444855 | 0.286062087 | 0.867414071 | -0.692823920 |
| | 13 | 14 | 15 | 16 | 17 | 18 |
| | 1.518579205 | -0.762439786 | -0.153308914 | -1.365843052 | -1.061229886 | 1.048247313 |
| | 19 | 20 | 21 | 22 | 23 | 24 |
| | -0.064626485 | -1.343409930 | -0.263756245 | 1.452616279 | 1.164708630 | -0.169537130 |

These are not quite the same, because R is doing something more complicated and precise to get the standard errors. Let's not worry about those details for now; the difference is pretty small in this case:

```
rstandard_compare_plot =  
  tibble(  
    x = resid(bw_lm2)/sigma(bw_lm2),  
    y = rstandard(bw_lm2)) |>  
  ggplot(aes(x = x, y = y)) +  
  geom_point() +  
  theme_bw() +  
  coord_equal() +  
  xlab("resid(bw_lm2)/sigma(bw_lm2)") +  
  ylab("rstandard(bw_lm2)") +  
  geom_abline(  
    aes(  
      intercept = 0,  
      slope = 1,  
      col = "x=y")) +  
  labs(colour="") +  
  scale_colour_manual(values="red")
```

2. Linear (Gaussian) Models

```
print(rstandard_compare_plot)
```



Let's add these residuals to the `tibble` of our dataset:

```
bw =  
  bw |>  
  mutate(  
    fitted_lm2 = fitted(bw_lm2),  
  
    resid_lm2 = resid(bw_lm2),  
    # resid_lm2 = weight - fitted_lm2,  
  
    std_resid_lm2 = rstandard(bw_lm2),  
    # std_resid_lm2 = resid_lm2 / sigma(bw_lm2)  
  )
```

2. Linear (Gaussian) Models

```
bw |>
  select(
    sex,
    age,
    weight,
    fitted_lm2,
    resid_lm2,
    std_resid_lm2
  )
```

| sex | age | weight | fitted_lm2 | resid_lm2 | std_resid_lm2 |
|--------|-----|--------|------------|-------------|---------------|
| female | 36 | 2729 | 2552.7333 | 176.266667 | 1.15981660 |
| female | 36 | 2412 | 2552.7333 | -140.733333 | -0.92601091 |
| female | 37 | 2539 | 2683.1333 | -144.133333 | -0.87479166 |
| female | 38 | 2754 | 2813.5333 | -59.533333 | -0.34722553 |
| female | 38 | 2991 | 2813.5333 | 177.466667 | 1.03506649 |
| female | 39 | 2817 | 2943.9333 | -126.933333 | -0.73473146 |
| female | 39 | 2875 | 2943.9333 | -68.933333 | -0.39900858 |
| female | 40 | 3317 | 3074.3333 | 242.666667 | 1.43751642 |
| female | 40 | 2935 | 3074.3333 | -139.333333 | -0.82538718 |
| female | 40 | 3126 | 3074.3333 | 51.666667 | 0.30606462 |
| female | 40 | 3231 | 3074.3333 | 156.666667 | 0.92806692 |
| female | 42 | 3210 | 3335.1333 | -125.133333 | -0.87615925 |
| male | 35 | 2925 | 2650.7241 | 274.275862 | 1.91427802 |
| male | 36 | 2625 | 2762.7069 | -137.706897 | -0.86559211 |
| male | 37 | 2847 | 2874.6897 | -27.689655 | -0.16429930 |
| male | 37 | 2628 | 2874.6897 | -246.689655 | -1.46375741 |
| male | 38 | 2795 | 2986.6724 | -191.672414 | -1.11015989 |
| male | 38 | 3176 | 2986.6724 | 189.327586 | 1.09657874 |
| male | 38 | 2975 | 2986.6724 | -11.672414 | -0.06760621 |
| male | 40 | 2968 | 3210.6379 | -242.637931 | -1.46158646 |
| male | 40 | 3163 | 3210.6379 | -47.637931 | -0.28695825 |

2. Linear (Gaussian) Models

| sex | age | weight | fitted_lm2 | resid_lm2 | std_resid_lm2 |
|------|-----|--------|------------|------------|---------------|
| male | 40 | 3473 | 3210.6379 | 262.362069 | 1.58039943 |
| male | 40 | 3421 | 3210.6379 | 210.362069 | 1.26716524 |
| male | 41 | 3292 | 3322.6207 | -30.620690 | -0.19805434 |


Now let's build histograms:

2.8.2.2. Marginal distribution of (nonstandardized) residuals

```
resid_marginal_hist =  
  bw |>  
  ggplot(aes(x = resid_lm2)) +  
  geom_histogram()
```

```
print(resid_marginal_hist)
```


2. Linear (Gaussian) Models




Hard to tell with this small amount of data, but I'm a bit concerned that the histogram doesn't show a bell-curve shape.

Marginal distribution of standardized residuals

```
std_resid_marginal_hist =  
  bw |>  
  ggplot(aes(x = std_resid_lm2)) +  
  geom_histogram()
```

```
print(std_resid_marginal_hist)
```

2. Linear (Gaussian) Models



This looks similar, although the scale of the x-axis got narrower, because we divided by $\hat{\sigma}$ (roughly speaking).

Still hard to tell if the distribution is Gaussian.

2.8.3. QQ plot of standardized residuals

Another way to assess normality is the QQ plot of the standardized residuals versus normal quantiles:


```
library(ggfortify)
# needed to make ggplot2::autoplot() work for `lm` objects

qqplot_lm2_auto =
  bw_lm2 |>
  autoplot(
```

2. Linear (Gaussian) Models

```
which = 2, # options are 1:6; can do multiple at once
ncol = 1) +
theme_classic()
```

```
print(qqplot_lm2_auto)
```



If the Gaussian model were correct, these points should follow the dotted line.

i Note

Fig 2.4 panel (c) in Dobson and Barnett (2018) is a little different; they didn't specify how they produced it, but other statistical analysis systems do things differently from R.

2. Linear (Gaussian) Models

2.8.3.1. QQ plot - how it's built

Let's construct it by hand:

```
bw = bw |>
  mutate(
    p = (rank(std_resid_lm2) - 1/2)/n(), # "Blom's method"
    expected_quantiles_lm2 = qnorm(p)
  )

qqplot_lm2 =
  bw |>
  ggplot(
    aes(
      x = expected_quantiles_lm2,
      y = std_resid_lm2,
      col = sex,
      shape = sex)
  ) +
  geom_point() +
  theme_classic() +
  theme(legend.position='none') + # removing the plot legend
  ggtitle("Normal Q-Q") +
  xlab("Theoretical Quantiles") +
  ylab("Standardized residuals")
```

We find the expected line like so:

```
ps <- c(.25, .75) # reference probabilities
a <- quantile(rstandard(bw_lm2), ps) # empirical quantiles
b <- qnorm(ps) # theoretical quantiles


qq_slope = diff(a)/diff(b)
```

2. Linear (Gaussian) Models

```
qq_intcpt = a[1] - b[1] * qq_slope

qqplot_lm2 =
  qqplot_lm2 +
  geom_abline(slope = qq_slope, intercept = qq_intcpt)

print(qqplot_lm2)
```



2.8.4. Conditional distributions of residuals

If our Gaussian linear regression model is correct, the residuals e_i and standardized residuals r_i should have:

- an approximately Gaussian distribution, with:
- a mean of 0

2. Linear (Gaussian) Models

- a constant variance

This should be true **regardless** of the value of X .

But if we didn't correctly guess the functional form of linear component of the mean,


$$E[Y|X = x] = \beta_0 + \beta_1 X_1 + \dots + \beta_p X_p$$

Then the the residuals might have nonzero mean or nonconstant variance for some values of x .

2.8.4.1. Residuals versus fitted values

To look for these issues, we can plot the residuals e_i against the fitted values \hat{y}_i :

```
autoplot(bw_lm2, which = 1, ncol = 1) |> print()
```



2. Linear (Gaussian) Models

If the model is correct, the blue line should stay flat and close to 0, and the cloud of dots should have the same vertical spread regardless of the fitted value.

If not, we probably need to change the functional form of linear component of the mean,

$$E[Y|X = x] = \beta_0 + \beta_1 X_1 + \dots + \beta_p X_p$$

2.8.4.2. Scale-location plot

We can also plot the square roots of the absolute values of the standardized residuals against the fitted values:

```
autoplot(bw_lm2, which = 3, ncol = 1) |> print()
```




Figure 2.6.: Scale-location plot of `birthweight` data


2. Linear (Gaussian) Models

Here, the blue line doesn't need to be near 0, but it should be flat. If not, the residual variance σ^2 might not be constant, and we might need to transform our outcome Y (or use a model that allows non-constant variance).

2.8.4.3. Residuals versus leverage

We can also plot our standardized residuals against “leverage”, which roughly speaking is a measure of how unusual each x_i value is. Very unusual x_i values can have extreme effects on the model fit, so we might want to remove those observations as outliers, particularly if they have large residuals.

```
autoplot(bw_lm2, which = 5, ncol = 1) |> print()
```



The blue line should be relatively flat and close to 0 here.

2.8.5. Diagnostics constructed by hand


```
bw =  
  bw |>  
  mutate(  
    predlm2 = predict(bw_lm2),  
    residlm2 = weight - predlm2,  
    std_resid = residlm2 / sigma(bw_lm2),  
    # std_resid_builtin = rstandard(bw_lm2), # uses leverage  
    sqrt_abs_std_resid = std_resid |> abs() |> sqrt()  
  )
```

Residuals vs fitted

```
resid_vs_fit = bw |>  
  ggplot(  
    aes(x = predlm2, y = residlm2, col = sex, shape = sex)  
  ) +  
  geom_point() +  
  theme_classic() +  
  geom_hline(yintercept = 0)
```

2. Linear (Gaussian) Models


```
print(resid_vs_fit)
```



Standardized residuals vs fitted

```
bw |>
  ggplot(
    aes(x = predlm2, y = std_resid, col = sex, shape = sex)
  ) +
  geom_point() +
  theme_classic() +
  geom_hline(yintercept = 0)
```


2. Linear (Gaussian) Models



Standardized residuals vs gestational age

```
bw |>
  ggplot(
    aes(x = age, y = std_resid, col = sex, shape = sex)
  ) +
  geom_point() +
  theme_classic() +
  geom_hline(yintercept = 0)
```

2. Linear (Gaussian) Models




`sqrt(abs(rstandard()))` vs fitted

Compare with `autoplot(bw_lm2, 3)`

```
bw |>
  ggplot(
    aes(x = predlm2, y = sqrt_abs_std_resid, col = sex, shape = sex)
  ) +
  geom_point() +
  theme_classic() +
  geom_hline(yintercept = 0)
```

2. Linear (Gaussian) Models



2.9. Model selection

If we have a lot of covariates in our dataset, we might want to choose a small subset to use in our model.

There are a few possible metrics to consider for choosing a “best” model.

2.9.1. Mean squared error

We might want to minimize the **mean squared error**, $E[(y - \hat{y})^2]$, for new observations that weren't in our data set when we fit the model.

Unfortunately,

$$\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

2. Linear (Gaussian) Models

gives a biased estimate of $E[(y - \hat{y})^2]$ for new data. If we want an unbiased estimate, we will have to be clever.

2.9.2. Cross-validation

```
data("carbohydrate", package = "dobson")
library(cvTools)
full.model <- lm(carbohydrate ~ ., data = carbohydrate)
temp =
  cvFit(full.model, data = carbohydrate, K = 5, R = 10,
y = carbohydrate$carbohydrate)
```

2.10. Categorical covariates with more than two levels

2.10.1. Example: birthweight

In the birthweight example, the variable `sex` had only two observed values:

```
unique(bw$sex)
```

```
[1] female male
Levels: female male
```

If there are more than two observed values, we can't just use a single variable with 0s and 1s.

2. Linear (Gaussian) Models

2.10.2.

For example, here's the (in)famous `iris` data:

```
iris |> tibble()
```

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|---------|
| 5.1 | 3.5 | 1.4 | 0.2 | setosa |
| 4.9 | 3.0 | 1.4 | 0.2 | setosa |
| 4.7 | 3.2 | 1.3 | 0.2 | setosa |
| 4.6 | 3.1 | 1.5 | 0.2 | setosa |
| 5.0 | 3.6 | 1.4 | 0.2 | setosa |
| 5.4 | 3.9 | 1.7 | 0.4 | setosa |
| 4.6 | 3.4 | 1.4 | 0.3 | setosa |
| 5.0 | 3.4 | 1.5 | 0.2 | setosa |
| 4.4 | 2.9 | 1.4 | 0.2 | setosa |
| 4.9 | 3.1 | 1.5 | 0.1 | setosa |
| 5.4 | 3.7 | 1.5 | 0.2 | setosa |
| 4.8 | 3.4 | 1.6 | 0.2 | setosa |
| 4.8 | 3.0 | 1.4 | 0.1 | setosa |
| 4.3 | 3.0 | 1.1 | 0.1 | setosa |
| 5.8 | 4.0 | 1.2 | 0.2 | setosa |
| 5.7 | 4.4 | 1.5 | 0.4 | setosa |
| 5.4 | 3.9 | 1.3 | 0.4 | setosa |
| 5.1 | 3.5 | 1.4 | 0.3 | setosa |
| 5.7 | 3.8 | 1.7 | 0.3 | setosa |
| 5.1 | 3.8 | 1.5 | 0.3 | setosa |
| 5.4 | 3.4 | 1.7 | 0.2 | setosa |
| 5.1 | 3.7 | 1.5 | 0.4 | setosa |
| 4.6 | 3.6 | 1.0 | 0.2 | setosa |
| 5.1 | 3.3 | 1.7 | 0.5 | setosa |
| 4.8 | 3.4 | 1.9 | 0.2 | setosa |

2. Linear (Gaussian) Models

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|------------|
| 5.0 | 3.0 | 1.6 | 0.2 | setosa |
| 5.0 | 3.4 | 1.6 | 0.4 | setosa |
| 5.2 | 3.5 | 1.5 | 0.2 | setosa |
| 5.2 | 3.4 | 1.4 | 0.2 | setosa |
| 4.7 | 3.2 | 1.6 | 0.2 | setosa |
| 4.8 | 3.1 | 1.6 | 0.2 | setosa |
| 5.4 | 3.4 | 1.5 | 0.4 | setosa |
| 5.2 | 4.1 | 1.5 | 0.1 | setosa |
| 5.5 | 4.2 | 1.4 | 0.2 | setosa |
| 4.9 | 3.1 | 1.5 | 0.2 | setosa |
| 5.0 | 3.2 | 1.2 | 0.2 | setosa |
| 5.5 | 3.5 | 1.3 | 0.2 | setosa |
| 4.9 | 3.6 | 1.4 | 0.1 | setosa |
| 4.4 | 3.0 | 1.3 | 0.2 | setosa |
| 5.1 | 3.4 | 1.5 | 0.2 | setosa |
| 5.0 | 3.5 | 1.3 | 0.3 | setosa |
| 4.5 | 2.3 | 1.3 | 0.3 | setosa |
| 4.4 | 3.2 | 1.3 | 0.2 | setosa |
| 5.0 | 3.5 | 1.6 | 0.6 | setosa |
| 5.1 | 3.8 | 1.9 | 0.4 | setosa |
| 4.8 | 3.0 | 1.4 | 0.3 | setosa |
| 5.1 | 3.8 | 1.6 | 0.2 | setosa |
| 4.6 | 3.2 | 1.4 | 0.2 | setosa |
| 5.3 | 3.7 | 1.5 | 0.2 | setosa |
| 5.0 | 3.3 | 1.4 | 0.2 | setosa |
| 7.0 | 3.2 | 4.7 | 1.4 | versicolor |
| 6.4 | 3.2 | 4.5 | 1.5 | versicolor |
| 6.9 | 3.1 | 4.9 | 1.5 | versicolor |
| 5.5 | 2.3 | 4.0 | 1.3 | versicolor |
| 6.5 | 2.8 | 4.6 | 1.5 | versicolor |
| 5.7 | 2.8 | 4.5 | 1.3 | versicolor |
| 6.3 | 3.3 | 4.7 | 1.6 | versicolor |

2. Linear (Gaussian) Models

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|------------|
| 4.9 | 2.4 | 3.3 | 1.0 | versicolor |
| 6.6 | 2.9 | 4.6 | 1.3 | versicolor |
| 5.2 | 2.7 | 3.9 | 1.4 | versicolor |
| 5.0 | 2.0 | 3.5 | 1.0 | versicolor |
| 5.9 | 3.0 | 4.2 | 1.5 | versicolor |
| 6.0 | 2.2 | 4.0 | 1.0 | versicolor |
| 6.1 | 2.9 | 4.7 | 1.4 | versicolor |
| 5.6 | 2.9 | 3.6 | 1.3 | versicolor |
| 6.7 | 3.1 | 4.4 | 1.4 | versicolor |
| 5.6 | 3.0 | 4.5 | 1.5 | versicolor |
| 5.8 | 2.7 | 4.1 | 1.0 | versicolor |
| 6.2 | 2.2 | 4.5 | 1.5 | versicolor |
| 5.6 | 2.5 | 3.9 | 1.1 | versicolor |
| 5.9 | 3.2 | 4.8 | 1.8 | versicolor |
| 6.1 | 2.8 | 4.0 | 1.3 | versicolor |
| 6.3 | 2.5 | 4.9 | 1.5 | versicolor |
| 6.1 | 2.8 | 4.7 | 1.2 | versicolor |
| 6.4 | 2.9 | 4.3 | 1.3 | versicolor |
| 6.6 | 3.0 | 4.4 | 1.4 | versicolor |
| 6.8 | 2.8 | 4.8 | 1.4 | versicolor |
| 6.7 | 3.0 | 5.0 | 1.7 | versicolor |
| 6.0 | 2.9 | 4.5 | 1.5 | versicolor |
| 5.7 | 2.6 | 3.5 | 1.0 | versicolor |
| 5.5 | 2.4 | 3.8 | 1.1 | versicolor |
| 5.5 | 2.4 | 3.7 | 1.0 | versicolor |
| 5.8 | 2.7 | 3.9 | 1.2 | versicolor |
| 6.0 | 2.7 | 5.1 | 1.6 | versicolor |
| 5.4 | 3.0 | 4.5 | 1.5 | versicolor |
| 6.0 | 3.4 | 4.5 | 1.6 | versicolor |
| 6.7 | 3.1 | 4.7 | 1.5 | versicolor |
| 6.3 | 2.3 | 4.4 | 1.3 | versicolor |
| 5.6 | 3.0 | 4.1 | 1.3 | versicolor |

2. Linear (Gaussian) Models

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|------------|
| 5.5 | 2.5 | 4.0 | 1.3 | versicolor |
| 5.5 | 2.6 | 4.4 | 1.2 | versicolor |
| 6.1 | 3.0 | 4.6 | 1.4 | versicolor |
| 5.8 | 2.6 | 4.0 | 1.2 | versicolor |
| 5.0 | 2.3 | 3.3 | 1.0 | versicolor |
| 5.6 | 2.7 | 4.2 | 1.3 | versicolor |
| 5.7 | 3.0 | 4.2 | 1.2 | versicolor |
| 5.7 | 2.9 | 4.2 | 1.3 | versicolor |
| 6.2 | 2.9 | 4.3 | 1.3 | versicolor |
| 5.1 | 2.5 | 3.0 | 1.1 | versicolor |
| 5.7 | 2.8 | 4.1 | 1.3 | versicolor |
| 6.3 | 3.3 | 6.0 | 2.5 | virginica |
| 5.8 | 2.7 | 5.1 | 1.9 | virginica |
| 7.1 | 3.0 | 5.9 | 2.1 | virginica |
| 6.3 | 2.9 | 5.6 | 1.8 | virginica |
| 6.5 | 3.0 | 5.8 | 2.2 | virginica |
| 7.6 | 3.0 | 6.6 | 2.1 | virginica |
| 4.9 | 2.5 | 4.5 | 1.7 | virginica |
| 7.3 | 2.9 | 6.3 | 1.8 | virginica |
| 6.7 | 2.5 | 5.8 | 1.8 | virginica |
| 7.2 | 3.6 | 6.1 | 2.5 | virginica |
| 6.5 | 3.2 | 5.1 | 2.0 | virginica |
| 6.4 | 2.7 | 5.3 | 1.9 | virginica |
| 6.8 | 3.0 | 5.5 | 2.1 | virginica |
| 5.7 | 2.5 | 5.0 | 2.0 | virginica |
| 5.8 | 2.8 | 5.1 | 2.4 | virginica |
| 6.4 | 3.2 | 5.3 | 2.3 | virginica |
| 6.5 | 3.0 | 5.5 | 1.8 | virginica |
| 7.7 | 3.8 | 6.7 | 2.2 | virginica |
| 7.7 | 2.6 | 6.9 | 2.3 | virginica |
| 6.0 | 2.2 | 5.0 | 1.5 | virginica |
| 6.9 | 3.2 | 5.7 | 2.3 | virginica |

2. Linear (Gaussian) Models

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|-----------|
| 5.6 | 2.8 | 4.9 | 2.0 | virginica |
| 7.7 | 2.8 | 6.7 | 2.0 | virginica |
| 6.3 | 2.7 | 4.9 | 1.8 | virginica |
| 6.7 | 3.3 | 5.7 | 2.1 | virginica |
| 7.2 | 3.2 | 6.0 | 1.8 | virginica |
| 6.2 | 2.8 | 4.8 | 1.8 | virginica |
| 6.1 | 3.0 | 4.9 | 1.8 | virginica |
| 6.4 | 2.8 | 5.6 | 2.1 | virginica |
| 7.2 | 3.0 | 5.8 | 1.6 | virginica |
| 7.4 | 2.8 | 6.1 | 1.9 | virginica |
| 7.9 | 3.8 | 6.4 | 2.0 | virginica |
| 6.4 | 2.8 | 5.6 | 2.2 | virginica |
| 6.3 | 2.8 | 5.1 | 1.5 | virginica |
| 6.1 | 2.6 | 5.6 | 1.4 | virginica |
| 7.7 | 3.0 | 6.1 | 2.3 | virginica |
| 6.3 | 3.4 | 5.6 | 2.4 | virginica |
| 6.4 | 3.1 | 5.5 | 1.8 | virginica |
| 6.0 | 3.0 | 4.8 | 1.8 | virginica |
| 6.9 | 3.1 | 5.4 | 2.1 | virginica |
| 6.7 | 3.1 | 5.6 | 2.4 | virginica |
| 6.9 | 3.1 | 5.1 | 2.3 | virginica |
| 5.8 | 2.7 | 5.1 | 1.9 | virginica |
| 6.8 | 3.2 | 5.9 | 2.3 | virginica |
| 6.7 | 3.3 | 5.7 | 2.5 | virginica |
| 6.7 | 3.0 | 5.2 | 2.3 | virginica |
| 6.3 | 2.5 | 5.0 | 1.9 | virginica |
| 6.5 | 3.0 | 5.2 | 2.0 | virginica |
| 6.2 | 3.4 | 5.4 | 2.3 | virginica |
| 5.9 | 3.0 | 5.1 | 1.8 | virginica |

2. Linear (Gaussian) Models

```
summary(iris)
```

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|---------------|
| Min. | Min. | Min. | Min. | setosa :50 |
| :4.3000 | :2.0000 | :1.000 | :0.1000 | |
| 1st | 1st | 1st | 1st | versicolor:50 |
| Qu.:5.1000 | Qu.:2.8000 | Qu.:1.600 | Qu.:0.3000 | |
| Median | Median | Median | Median | virginica |
| :5.8000 | :3.0000 | :4.350 | :1.3000 | :50 |
| Mean | Mean | Mean | Mean | NA |
| :5.8433 | :3.0573 | :3.758 | :1.1993 | |
| 3rd | 3rd | 3rd | 3rd | NA |
| Qu.:6.4000 | Qu.:3.3000 | Qu.:5.100 | Qu.:1.8000 | |
| Max. | Max. | Max. | Max. | NA |
| :7.9000 | :4.4000 | :6.900 | :2.5000 | |

2.10.3.

There are three species:

```
iris$Species |> unique()
```

```
[1] setosa    versicolor virginica  
Levels: setosa versicolor virginica
```

2.10.4.

If we want to model `Sepal.Length` by species, we could create a variable X that represents “setosa” as $X = 1$, “virginica” as $X = 2$, and “versicolor” as $X = 3$.

2. Linear (Gaussian) Models

```
data(iris) # this step is not always necessary, but ensures you're starting
# from the original version of a dataset stored in a loaded package

iris =
  iris |>
  tibble() |>
  mutate(
    X = case_when(
      Species == "setosa" ~ 1,
      Species == "virginica" ~ 2,
      Species == "versicolor" ~ 3
    )
  )

iris |>
  distinct(Species, X) |>
  print()
```

```
# A tibble: 3 x 2
  Species      X
  <fct>    <dbl>
1 setosa      1
2 versicolor  3
3 virginica   2
```

Then we could fit a model like:

```
iris_lm1 = lm(Sepal.Length ~ X, data = iris)
iris_lm1 |> parameters() |> print_md()
```

2. Linear (Gaussian) Models


| Parameter | Coefficient | SE | 95% CI | t(148) | p |
|-------------|-------------|------|--------------|--------|--------|
| (Intercept) | 4.91 | 0.16 | (4.60, 5.23) | 30.83 | < .001 |
| X | 0.47 | 0.07 | (0.32, 0.61) | 6.30 | < .001 |

2.10.5. Let's see how that model looks:

```
iris_plot1 = iris |>
  ggplot(
    aes(
      x = X,
      y = Sepal.Length)
  ) +
  geom_point(alpha = .1) +
  geom_abline(
    intercept = coef(iris_lm1)[1],
    slope = coef(iris_lm1)[2] +
  theme_bw(base_size = 18)
```

```
print(iris_plot1)
```

2. Linear (Gaussian) Models



We have forced the model to use a straight line for the three estimated means. Maybe not a good idea?

2.10.6. Let's see what R does with categorical variables by default:

```
iris_lm2 = lm(Sepal.Length ~ Species, data = iris)
iris_lm2 |> parameters() |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(147) | p |
|-------------------------|-------------|------|--------------|--------|--------|
| (Intercept) | 5.01 | 0.07 | (4.86, 5.15) | 68.76 | < .001 |
| Species (versicolor) | 0.93 | 0.10 | (0.73, 1.13) | 9.03 | < .001 |

2. Linear (Gaussian) Models

| Parameter | Coefficient | SE | 95% CI | t(147) | p |
|---------------------|-------------|------|--------------|--------|--------|
| Species (virginica) | 1.58 | 0.10 | (1.38, 1.79) | 15.37 | < .001 |

2.10.7. Re-parametrize with no intercept

If you don't want the default and offset option, you can use “-1” like we've seen previously:

```
iris_lm2b = lm(Sepal.Length ~ Species - 1, data = iris)
iris_lm2b |> parameters() |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(147) | p |
|----------------------|-------------|------|--------------|--------|--------|
| Species (setosa) | 5.01 | 0.07 | (4.86, 5.15) | 68.76 | < .001 |
| Species (versicolor) | 5.94 | 0.07 | (5.79, 6.08) | 81.54 | < .001 |
| Species (virginica) | 6.59 | 0.07 | (6.44, 6.73) | 90.49 | < .001 |


2.10.8. Let's see what these new models look like:

```
iris_plot2 =
  iris |>
  mutate(
    predlm2 = predict(iris_lm2) |>
    arrange(X) |>
    ggplot(aes(x = X, y = Sepal.Length)) +
    geom_point(alpha = .1) +
    geom_line(aes(y = predlm2), col = "red") +
    geom_abline(
```


2. Linear (Gaussian) Models

```
intercept = coef(iris_lm1)[1],  
slope = coef(iris_lm1)[2] +  
theme_bw(base_size = 18)
```

```
print(iris_plot2)
```



2.10.9. Let's see how R did that:

```
formula(iris_lm2)
```

Sepal.Length ~ Species

2. Linear (Gaussian) Models

```
model.matrix(iris_lm2) |> as_tibble() |> unique()
```

| (Intercept) | Speciesversicolor | Speciesvirginica |
|-------------|-------------------|------------------|
| 1 | 0 | 0 |
| 1 | 1 | 0 |
| 1 | 0 | 1 |

This is called a “corner point parametrization”.

```
formula(iris_lm2b)
```

Sepal.Length ~ Species - 1

```
model.matrix(iris_lm2b) |> as_tibble() |> unique()
```

| Speciessetosa | Speciesversicolor | Speciesvirginica |
|---------------|-------------------|------------------|
| 1 | 0 | 0 |
| 0 | 1 | 0 |
| 0 | 0 | 1 |

This can be called a “group point parametrization”.

There are more options; see Dobson & Barnett §6.4.1.

3. Models for Binary Outcomes

Logistic regression and variations

Acknowledgements

This content is adapted from:

- Dobson and Barnett (2018), Chapter 7
- Vittinghoff et al. (2012), Chapter 5
- <https://dmrocke.ucdavis.edu/Class/EPI204-Spring-2021/EPI204-Spring-2021.html>

Configuring R

Functions from these packages will be used throughout this document:

```
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(dplyr) # manipulate data
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
```

3. Models for Binary Outcomes

```
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)
```

3.1. Introduction

3.1.1. What is logistic regression?

Logistic regression is a framework for modeling **binary** outcomes, conditional on one or more *predictors* (a.k.a. *covariates*).

Exercise 3.1 (Examples of binary outcomes). What are some examples of binary outcomes in the health sciences?

Solution. Examples of binary outcomes include:

- exposure (exposed vs unexposed)
- disease (diseased vs healthy)
- recovery (recovered vs unrecovered)
- relapse (relapse vs remission)
- return to hospital (returned vs not)
- vital status (dead vs alive)

Logistic regression uses the **Bernoulli** distribution to model the outcome variable, conditional on one or more covariates.

3. Models for Binary Outcomes

Exercise 3.2. Write down a mathematical definition of the Bernoulli distribution.

Solution. See Definition A.8.

3.1.2. Logistic regression versus linear regression

Logistic regression differs from linear regression, which uses the Gaussian (“normal”) distribution to model the outcome variable, conditional on the covariates.

Exercise 3.3. Recall: what kinds of outcomes is linear regression used for?

Solution. Linear regression is typically used for numerical outcomes that aren’t event counts or waiting times for an event. Examples of outcomes that are often analyzed using linear regression include include weight, height, and income.

3.2. Risk Estimation and Prediction

In Epi 203, you have already seen methods for modeling binary outcomes using one covariate that is also binary (such as exposure/non-exposure). In this section, we review one-covariate analyses, with a special focus on risk ratios and odds ratios, which are important concepts for interpreting logistic regression.

Example 3.1 (Oral Contraceptive Use and Heart Attack).

- Research question: how does oral contraceptive (OC) use affect the risk of myocardial infarction (MI; a.k.a. heart attack)?

3. Models for Binary Outcomes

This was an issue when oral contraceptives were first developed, because the original formulations used higher concentrations of hormones. Modern OCs don't have this issue.

Table 3.1 contains simulated data for an imaginary follow-up (a.k.a. *prospective*) study in which two groups are identified, one using OCs and another not using OCs, and both groups are tracked for three years to determine how many in each groups have MIs.

```
library(dplyr)
oc_mi =
  tribble(
    ~OC, ~MI, ~Total,
    "OC use", 13, 5000,
    "No OC use", 7, 10000
  ) |>
  mutate(`No MI` = Total - MI) |>
  relocate(`No MI`, .after = MI)

totals =
  oc_mi |>
  summarize(across(c(MI, `No MI`, Total), sum)) |>
  mutate(OC = "Total")

oc_mi = bind_rows(oc_mi, totals)

pander(oc_mi)
```

Table 3.1.: Simulated data from study of oral contraceptive use and heart attack risk

| OC | MI | No MI | Total |
|--------|----|-------|-------|
| OC use | 13 | 4987 | 5000 |

3. Models for Binary Outcomes

Table 3.1.: Simulated data from study of oral contraceptive use and heart attack risk

| | OC | MI | No MI | Total |
|-----------|----|----|-------|-------|
| No OC use | | 7 | 9993 | 10000 |
| Total | | 20 | 14980 | 15000 |

Exercise 3.4. Review: estimate the probabilities of MI for OC users and non-OC users in Example 3.1.

Solution.

$$\hat{p}(MI|OC) = \frac{13}{5000} = 0.0026$$

$$\hat{p}(MI|\neg OC) = \frac{7}{10000} = 7 \times 10^{-4}$$

i Two meanings of “controls”

Depending on context, “controls” can mean either individuals who don’t experience an *exposure* of interest, or individuals who don’t experience an *outcome* of interest.

Definition 3.1 (cases and controls in retrospective studies). In *retrospective studies*, participants who experience the outcome of interest are called **cases**, while participants who don’t experience that outcome are called **controls**.

Definition 3.2 (treatment groups and control groups in prospective studies). In *prospective studies*, the group of participants who experience the treatment or exposure of interest is called the **treatment group**, while the participants who receive the baseline or comparison treatment (for example, clinical trial participants who receive a placebo or a standard-of-care treatment rather than an experimental

treatment) are called **controls**.

3.2.1. Comparing probabilities

3.2.1.1. Risk differences

The simplest comparison of two probabilities, π_1 , and π_2 , is the difference of their values:

Definition 3.3 (Risk difference). The **risk difference** of two probabilities, π_1 , and π_2 , is the difference of their values:

$$\delta(\pi_1, \pi_2) \stackrel{\text{def}}{=} \pi_1 - \pi_2$$

Example 3.2. In Example 3.1, the estimated risk difference in MI risk between OC users and OC non-users is:

$$\begin{aligned} \hat{\delta}(\pi(OC), \pi(\neg OC)) &= \delta(\hat{\pi}(OC), \hat{\pi}(\neg OC)) \\ &= \hat{\pi}(OC) - \hat{\pi}(\neg OC) \\ &= 0.0026 - 7 \times 10^{-4} \\ &= 0.0019 \end{aligned}$$

3.2.1.2. Risk ratios

Definition 3.4 (Relative risk ratios). The **relative risk** of probability π_1 compared to another probability π_2 , also called the **risk ratio**, **relative risk ratio**, **probability ratio**, or **rate ratio**, is the ratio of those probabilities:

$$\rho(\pi_1, \pi_2) = \frac{\pi_1}{\pi_2}$$

3. Models for Binary Outcomes

Example 3.3. Above, we estimated that:

$$\hat{p}(MI|OC) = 0.0026$$

$$\hat{p}(MI|\neg OC) = 7 \times 10^{-4}$$

So we might estimate that the *relative risk* of MI for OC versus non-OC is:

$$\text{rr} = (13/5000)/(7/10000)$$

$$\begin{aligned}\hat{\rho}(OC, \neg OC) &= \frac{\hat{p}(MI|OC)}{\hat{p}(MI|\neg OC)} \\ &= \frac{0.0026}{7 \times 10^{-4}} \\ &= 3.71428571\end{aligned}$$

We might summarize this result by saying that “the estimated probability of MI among OC users was 3.71428571 as high as the estimated probability among OC non-users.

Sometimes, we divide the risk difference by the comparison probability, or equivalently, subtract 1 from the risk ratio; the result is called the **relative risk difference**:

$$\begin{aligned}\xi(\pi_1, \pi_2) &= \frac{\pi_1 - \pi_2}{\pi_2} \\ &= \frac{\pi_1}{\pi_2} - 1\end{aligned}$$

Risk differences, risk ratios, and relative risk differences are defined by two probabilities, plus a choice of which probability is the **baseline** or

3. Models for Binary Outcomes

reference probability (i.e., which probability is the subtrahend of the risk difference or the denominator of the risk ratio). To switch which one is the reference probability, invert the ratio and multiply the difference by -1.

Example 3.4. Above, we estimated that the risk ratio of OC versus non-OC is:

$$\rho(OC, \neg OC) = 3.71428571$$

In comparison, the risk ratio for non-OC versus OC is:

$$\begin{aligned}\rho(\neg OC, OC) &= \frac{\hat{p}(MI|\neg OC)}{\hat{p}(MI|OC)} \\ &= \frac{7 \times 10^{-4}}{0.0026} \\ &= 0.26923077 \\ &= \frac{1}{\rho(OC, \neg OC)}\end{aligned}$$

3.2.2. Odds and probabilities

In logistic regression, we will make use of a transformation (rescaling) of probability, called *odds*.

Definition 3.5 (Odds). The **odds** of an outcome, denoted ω (“omega”), is the probability that the outcome occurs, divided by the probability that it doesn’t occur.

That is, if the probability of an outcome is π , then the corresponding odds of that outcome is

3. Models for Binary Outcomes


$$\omega(\pi) \stackrel{\text{def}}{=} \frac{\pi}{1 - \pi}$$

This function, which transforms probabilities into odds, is called the **odds function** (see `?@fig-odds-probs`).

```
odds = function(pi) pi/(1-pi)
library(ggplot2)
odds_plot = ggplot() +
  geom_function(fun = odds, aes(col = "odds function")) +
  xlim(0, .5) +
  xlab("Probability") +
  ylab("Odds") +
  geom_abline(aes(intercept = 0, slope = 1, col = "y=x")) +
  theme_bw() +
  labs(colour = "") +
  theme(legend.position = "bottom")
```

```
print(odds_plot)
```

3. Models for Binary Outcomes



Example 3.5 (Calculating odds). In Exercise 3.4, we estimated that the probability of MI, given OC use, is $\pi(OC) \stackrel{\text{def}}{=} \Pr(MI|OC) = 0.0026$. If this estimate is correct, then the odds of MI, given OC use, is:

$$\begin{aligned}
 \omega(OC) &\stackrel{\text{def}}{=} \frac{\Pr(MI|OC)}{\Pr(\neg MI|OC)} \\
 &= \frac{\Pr(MI|OC)}{1 - \Pr(MI|OC)} \\
 &= \frac{\pi(OC)}{1 - \pi(OC)} \\
 &= \frac{0.0026}{1 - 0.0026} \\
 &= 0.00260678
 \end{aligned}$$

3. Models for Binary Outcomes

Exercise 3.5 (Calculating odds). Estimate the odds of MI, for non-OC users.

Solution.

$$\omega(\neg OC) = 7.00490343 \times 10^{-4}$$

3.2.2.1. A shortcut for calculating odds

The usual estimate for a probability of an event is “# events/# observations”. We often denote # events as x and # observations as n . So:

$$\hat{\pi} = \frac{x}{n}$$

Thus, the usual estimate for the probability of a nonevent is:

$$\begin{aligned} 1 - \hat{\pi} &= 1 - \frac{x}{n} \\ &= \frac{n}{n} - \frac{x}{n} \\ &= \frac{n - x}{n} \end{aligned}$$

Thus, the estimated odds is:

$$\begin{aligned} \frac{\hat{\pi}}{1 - \hat{\pi}} &= \frac{\left(\frac{x}{n}\right)}{\left(\frac{n-x}{n}\right)} \\ &= \frac{x}{n - x} \end{aligned}$$

That is, odds can be calculated directly as “# events” divided by “# nonevents” (without needing to calculate $\hat{\pi}$ and $1 - \hat{\pi}$ first).

Example 3.6 (calculating odds using the shortcut). In Example 3.5, we calculated

$$\omega(OC) = 0.00260678$$

3. Models for Binary Outcomes

Let's recalculate this result using our shortcut:

$$\begin{aligned}\omega(OC) &= \frac{13}{5000 - 13} \\ &= 0.00260678\end{aligned}$$

Same answer!

3.2.2.2. Odds of rare events

For rare events (small π), odds and probabilities are nearly equal, because $1 - \pi \approx 1$ (see [?@fig-odds-probs](#)).

For example, in Example [3.5](#), the probability and odds differ by $6.77762182 \times 10^{-6}$.

Exercise 3.6. What odds value corresponds to the probability $\pi = 0.2$, and what is the numerical difference between these two values?

Solution.

$$\omega = \frac{\pi}{1 - \pi} = \frac{.2}{.8} = .25$$

3.2.3. The inverse odds function

Definition 3.6 (inverse odds function). The **inverse odds function**,

$$\pi(\omega) \stackrel{\text{def}}{=} \frac{\omega}{1 + \omega}$$

converts odds into their corresponding probabilities (Figure [3.1](#)).

3. Models for Binary Outcomes

The inverse-odds function takes an odds as input and produces a probability as output. Its domain of inputs is $[0, \infty)$ and its range of outputs is $[0, 1]$.

```
odds_inv = function(omega) (1 + omega^-1)^-1
ggplot() +
  geom_function(fun = odds_inv, aes(col = "inverse-odds")) +
  xlab("Odds") +
  ylab("Probability") +
  xlim(0,5) +
  ylim(0,1) +
  geom_abline(aes(intercept = 0, slope = 1, col = "x=y"))
```




Figure 3.1.: The inverse odds function, $\pi(\omega)$

3. Models for Binary Outcomes

! Important

An equivalent expression for the inverse odds function is

$$\pi(\omega) = (1 - \omega^{-1})^{-1} \quad (3.1)$$

Exercise 3.7. Prove that Equation 3.1 is equivalent to Definition 3.6.

Exercise 3.8. What probability corresponds to an odds of $\omega = 1$, and what is the numerical difference between these two values?

Solution.

$$\begin{aligned} \pi(1) &= \frac{1}{1+1} = \frac{1}{2} = .5 \\ 1 - \pi(1) &= 1 - .5 = .5 \end{aligned}$$

3.2.4. Odds ratios

Now that we have defined odds, we can introduce another way of comparing event probabilities: odds ratios.

Definition 3.7 (Odds ratio). The **odds ratio** for two odds ω_1, ω_2 is their ratio:

$$\theta(\omega_1, \omega_2) = \frac{\omega_1}{\omega_2}$$

Example 3.7 (Calculating odds ratios). In Example 3.1, the odds ratio for OC users versus OC-non-users is:

3. Models for Binary Outcomes

$$\begin{aligned}\theta(\omega(OC), \omega(\neg OC)) &= \frac{\omega(OC)}{\omega(\neg OC)} \\ &= \frac{0.0026}{7 \times 10^{-4}} \\ &= 3.71428571\end{aligned}$$

When the outcome is rare (i.e., its probability is small) for both groups being compared in an odds ratio, the odds of the outcome will be similar to the probability of the outcome, and thus the risk ratio will be similar to the odds ratio.

For example, in Example 3.1, the outcome is rare for both OC and non-OC participants, so the odds for both groups are similar to the corresponding probabilities, and the odds ratio is similar the risk ratio.

3.2.4.1. A shortcut for calculating odds ratio estimates

The general form of a two-by-two table is shown in Table 3.2.

| | Event | Non-Event | Total |
|-------------|-------|-----------|---------|
| Exposed | a | b | a+b |
| Non-exposed | c | d | c+d |
| Total | a+c | b+d | a+b+c+d |

From this table, we have:

- $\hat{\pi}(Event|Exposed) = a/(a+b)$
- $\hat{\pi}(\neg Event|Exposed) = b/(a+b)$
- $\hat{\omega}(Event|Exposed) = \frac{(\frac{a}{a+b})}{(\frac{b}{a+b})} = \frac{a}{b}$

3. Models for Binary Outcomes

- $\hat{\omega}(Event|\neg Exposed) = \frac{c}{d}$ (see Exercise 3.9)
- $\theta(Exposed, \neg Exposed) = \frac{\frac{a}{b}}{\frac{c}{d}} = \frac{ad}{bc}$

Exercise 3.9. Given Table 3.2, show that $\hat{\omega}(Event|\neg Exposed) = \frac{c}{d}$.

3.2.4.2. Properties of odds ratios

Odds ratios have a special property: we can swap a covariate with the outcome, and the odds ratio remains the same.

Example 3.8. In Example 3.1, we have:

3. Models for Binary Outcomes

$$\begin{aligned}
\theta(MI; OC) &\stackrel{\text{def}}{=} \frac{\omega(MI|OC)}{\omega(MI|\neg OC)} \\
&\stackrel{\text{def}}{=} \frac{\left(\frac{\Pr(MI|OC)}{\Pr(\neg MI|OC)} \right)}{\left(\frac{\Pr(MI|\neg OC)}{\Pr(\neg MI|\neg OC)} \right)} \\
&= \frac{\left(\frac{\Pr(MI, OC)}{\Pr(\neg MI, OC)} \right)}{\left(\frac{\Pr(MI, \neg OC)}{\Pr(\neg MI, \neg OC)} \right)} \\
&= \left(\frac{\Pr(MI, OC)}{\Pr(\neg MI, OC)} \right) \left(\frac{\Pr(\neg MI, \neg OC)}{\Pr(MI, \neg OC)} \right) \\
&= \left(\frac{\Pr(MI, OC)}{\Pr(MI, \neg OC)} \right) \left(\frac{\Pr(\neg MI, \neg OC)}{\Pr(\neg MI, OC)} \right) \\
&= \left(\frac{\Pr(OC, MI)}{\Pr(\neg OC, MI)} \right) \left(\frac{\Pr(\neg OC, \neg MI)}{\Pr(OC, \neg MI)} \right) \\
&= \left(\frac{\Pr(OC|MI)}{\Pr(\neg OC|MI)} \right) \left(\frac{\Pr(\neg OC|\neg MI)}{\Pr(OC|\neg MI)} \right) \\
&= \frac{\left(\frac{\Pr(OC|MI)}{\Pr(\neg OC|MI)} \right)}{\left(\frac{\Pr(OC|\neg MI)}{\Pr(\neg OC|\neg MI)} \right)} \\
&\stackrel{\text{def}}{=} \frac{\omega(OC|MI)}{\omega(OC|\neg MI)} \\
&\stackrel{\text{def}}{=} \theta(OC; MI)
\end{aligned}$$

Exercise 3.10. For Table 3.2, show that $\hat{\theta}(Exposed, Unexposed) = \hat{\theta}(Event, \neg Event)$.

3.2.5. Effect of study design

Table 3.1 simulates a follow-up study in which two populations were followed and the number of MI's was observed. The risks are $P(MI|OC)$ and $P(MI|\neg OC)$ and we can estimate these risks from the data.

3. Models for Binary Outcomes

But suppose we had a case-control study in which we had 100 women with MI and selected a comparison group of 100 women without MI (matched as groups on age, etc.). Then MI is not random, and we cannot compute $P(MI|OC)$ and we cannot compute the risk ratio. However, the odds ratio however can be computed.

The disease odds ratio is the odds for the disease in the exposed group divided by the odds for the disease in the unexposed group, and we cannot validly compute and use these separate parts.

But we can validly compute and use the exposure odds ratio, which is the odds for exposure in the disease group divided by the odds for exposure in the non-diseased group (because exposure can be treated as random):

$$\hat{\theta}(OC|MI) = \frac{\hat{\omega}(OC|MI)}{\hat{\omega}(OC|\neg MI)}$$

And these two odds ratios, $\hat{\theta}(MI|OC)$ and $\hat{\theta}(OC|MI)$ are mathematically equivalent, as we saw in Section 3.2.4.2.

Exercise 3.11. Calculate the odds ratio of MI with respect to OC use, assuming that Table 3.1 comes from a case-control study. Confirm that the result is the same as in Example 3.7.

Solution.

- $\omega(OC|MI) = P(OC|MI)/(1-P(OC|MI)) = \frac{13}{7} = 1.85714286$
- $\omega(OC|\neg MI) = P(OC|\neg MI)/(1-P(OC|\neg MI)) = \frac{4987}{9993} = 0.49904933$
- $\theta(OC, MI) = \frac{\omega(OC|MI)}{\omega(OC|\neg MI)} = \frac{13/7}{4987/9993} = 3.72136125$

This is the same estimate we calculated in Example 3.7.

3. Models for Binary Outcomes

3.2.5.1. Cross-Sectional Studies

- If a cross-sectional study is a probability sample of a population (which it rarely is) then we can estimate risks.
- If it is a sample, but not an unbiased probability sample, then we need to treat it in the same way as a case-control study.
- We can validly estimate odds ratios in either case.
- But we can usually not validly estimate risks and risk ratios.

3.3. Introduction to logistic regression

- In Example 3.1, we estimated the risk and the odds of MI for two discrete cases, as to whether or not the individual used oral contraceptives.
- If the predictor is quantitative (dose) or there is more than one predictor, the task becomes more difficult.
- In this case, we will use logistic regression, which is a generalization of the linear regression models you have been using that can account for a binary response instead of a continuous one.

3. Models for Binary Outcomes

3.3.1. Binary outcomes models - one group, no covariates

$$\begin{aligned}p(Y = 1) &= \pi \\p(Y = 0) &= 1 - \pi \\p(Y = y) &= \pi^y(1 - \pi)^{1-y} \\ \mathbf{y} &= (y_1, \dots, y_n) \\ \mathcal{L}(\pi; \mathbf{y}) &= \pi^{\sum y_i} (1 - \pi)^{n - \sum y_i} \\ \ell(\pi, \mathbf{y}) &= \left(\sum y_i \right) \log \{ \pi \} + \left(n - \sum y_i \right) \log \{ 1 - \pi \} \\ &= \left(\sum y_i \right) (\log \{ \pi \} - \log \{ 1 - \pi \}) + n \cdot \log \{ 1 - \pi \} \\ &= \left(\sum y_i \right) \log \left\{ \frac{\pi}{1 - \pi} \right\} + n \cdot \log \{ 1 - \pi \}\end{aligned}$$

3.3.2. Binary outcomes - general

$$\begin{aligned}p(Y_i = 1) &= \pi_i \\p(Y_i = 0) &= 1 - \pi_i \\p(Y_i = y) &= (\pi_i)^y (1 - \pi_i)^{1-y} \\ \mathbf{y} &= (y_1, \dots, y_n) \\ \mathcal{L}(\pi; \mathbf{y}) &= \prod_{i=1}^n (\pi_i)^{y_i} (1 - \pi_i)^{1-y_i} \\ \ell(\pi, \mathbf{y}) &= \sum_{i=1}^n y_i \log \{ \pi_i \} + (1 - y_i) \log \{ 1 - \pi_i \}\end{aligned}$$

3.3.3. Modeling π_i as a function of X_i

If there are only a few distinct X_i values, we can model each one separately.

Otherwise, we need regression.

3. Models for Binary Outcomes

$$\begin{aligned}\pi(x) &\equiv \mathbb{E}(Y = 1|X = x) \\ &= f(x^\top \beta)\end{aligned}$$

Typically, we use $f(\eta) = \text{expit}(\eta)$.

Definition 3.8 (expit function). The **expit** function (Figure 3.2), also known as the **inverse-logit** or **logistic** function, is:

$$\begin{aligned}\text{expit}(\eta) &\stackrel{\text{def}}{=} \frac{\exp\{\eta\}}{1 + \exp\{\eta\}} \\ &= (1 + \exp\{-\eta\})^{-1}\end{aligned}$$

```
expit = function(eta) exp(eta)/(1+exp(eta))
library(ggplot2)
expit_plot =
  ggplot() +
  geom_function(fun = expit) +
  xlim(-5, 5) +
  ylim(0,1) +
  ylab(expression(expit(eta))) +
  xlab(expression(eta)) +
  theme_bw()
print(expit_plot)
```

3. Models for Binary Outcomes




Figure 3.2.: The expit function

Definition 3.9 (logit function). The inverse of the `expit` function is the `logit` function:

$$g(p) = f^{-1}(p) = \text{logit}(p) = \log \left\{ \frac{p}{1-p} \right\}$$

```
logit = function(p) log(odds(p))

logit_plot =
  ggplot() +
  geom_function(fun = logit) +
  xlim(.01, .99) +
  ylab("logit(p)") +
  xlab("p") +
```


3. Models for Binary Outcomes

```
theme_bw()
print(logit_plot)
```




Figure 3.3.: the logit function

3.3.4. Diagram of expit and logit

$$\begin{array}{c}
 \text{logit}(\pi) \\
 \left[\pi \stackrel{\text{def}}{=} \Pr(Y = 1) \right] \xrightleftharpoons[\frac{\omega}{1+\omega}]{\frac{\pi}{1-\pi}} \left[\omega \stackrel{\text{def}}{=} \text{odds}(Y = 1) \right] \xrightleftharpoons[\exp\{\eta\}]{\log\{\omega\}} \left[\eta \stackrel{\text{def}}{=} \log\text{-odds}(Y = 1) \right] \\
 \underbrace{\hspace{15em}}_{\text{expit}(\eta)}
 \end{array}$$

3.3.5. Meet the beetles

```
library(glmx)

data(BeetleMortality, package = "glmx")
beetles = BeetleMortality |>
  mutate(
    pct = died/n,
    survived = n - died
  )

plot1 =
  beetles |>
  ggplot(aes(x = dose, y = pct)) +
  geom_point(aes(size = n)) +
  xlab("Dose (log mg/L)") +
  ylab("Mortality rate (%)") +
  scale_y_continuous(labels = scales::percent) +
  scale_size(range = c(1,2)) +
  theme_bw(base_size = 18)

print(plot1)
```

3. Models for Binary Outcomes



Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

3.3.6. Why don't we use linear regression?

```
beetles_glm_grouped = beetles |>
  glm(formula = cbind(died, survived) ~ dose, family = "binomial")

lm1 =
  beetles |>
  reframe(
    .by = everything(),
    outcome = c(
      rep(1, times = died),
      rep(0, times = survived))
  )
```

3. Models for Binary Outcomes

```
) |>
lm(
  formula = outcome ~ dose,
  data = _)


lm2 =
beetles |>
reframe(
  .by = everything(),
  outcome = c(
    rep(1, times = died),
    rep(0, times = survived))
) |>
lm(
  formula = outcome ~ log(dose),
  data = _)

range1 = range(beetles$dose) + c(-.2, .2)
f = function(x) predict(beetles_glm_grouped, newdata = data.frame(dose = x), type =
f.linear = function(x) predict(lm1, newdata = data.frame(dose = x))
f.linearlog = function(x) predict(lm2, newdata = data.frame(dose = x))

plot2 =
  plot1 +
  geom_function(
    fun = f.linear,
    aes(col = "Straight line")) +
  labs(colour="Model", size = "")

plot2 |> print()
```


3. Models for Binary Outcomes



3.3.7. Zoom out

```
(plot2 + expand_limits(x = c(1.6, 2))) |> print()
```


3. Models for Binary Outcomes



3.3.8. log transformation of dose?

```
plot3 = plot2 +  
  expand_limits(x = c(1.6, 2)) +  
  geom_function(fun = f.linearlog, aes(col = "Log-transform dose"))  
(plot3 + expand_limits(x = c(1.6, 2))) |> print()
```


3. Models for Binary Outcomes



3.3.9. Logistic regression

```
plot4 = plot3 + geom_function(fun = f, aes(col = "Logistic regression"))
plot4 |> print()
```

3. Models for Binary Outcomes



3.3.10. Three parts to regression models

- What distribution does the outcome have for a specific subpopulation defined by covariates? (outcome model)
- How does the combination of covariates relate to the mean? (link function)
- How do the covariates combine? (linear predictor, interactions)

3.3.11. Logistic regression in R

```
beetles_glm_grouped =  
  beetles |>
```


3. Models for Binary Outcomes

```
glm(  
  formula = cbind(died, survived) ~ dose,  
  family = "binomial")  
  
beetles_glm_grouped |>  
  parameters() |>  
  print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-------------|----------|------|------------------|--------|--------|
| (Intercept) | -60.72 | 5.18 | (-71.44, -51.08) | -11.72 | < .001 |
| dose | 34.27 | 2.91 | (28.85, 40.30) | 11.77 | < .001 |

Fitted values:

```
fitted.values(beetles_glm_grouped)
```

| | | | | | | | |
|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 0.0586 | 0.1640 | 0.3621 | 0.6053 | 0.7952 | 0.9032 | 0.9552 | 0.9790 |

```
predict(beetles_glm_grouped, type = "response")
```

| | | | | | | | |
|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 0.0586 | 0.1640 | 0.3621 | 0.6053 | 0.7952 | 0.9032 | 0.9552 | 0.9790 |

```
predict(beetles_glm_grouped, type = "link")
```

| | | | | | | | |
|---------|---------|---------|--------|--------|--------|--------|--------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| -2.7766 | -1.6286 | -0.5662 | 0.4277 | 1.3564 | 2.2337 | 3.0596 | 3.8444 |

3. Models for Binary Outcomes

```
fit_y = beetles$n * fitted.values(beetles_glm_grouped)
```

3.3.12. Individual observations

```
beetles_long =  
  beetles |>  
  reframe(  
    .by = everything(),  
    outcome = c(  
      rep(1, times = died),  
      rep(0, times = survived))  
  )  
beetles_long |> tibble() |> print()
```

```
# A tibble: 481 x 6  
  dose died    n pct survived outcome  
  <dbl> <int> <int> <dbl>    <int>    <dbl>  
1  1.69     6   59 0.102      53      1  
2  1.69     6   59 0.102      53      1  
3  1.69     6   59 0.102      53      1  
4  1.69     6   59 0.102      53      1  
5  1.69     6   59 0.102      53      1  
6  1.69     6   59 0.102      53      1  
7  1.69     6   59 0.102      53      0  
8  1.69     6   59 0.102      53      0  
9  1.69     6   59 0.102      53      0  
10 1.69     6   59 0.102      53      0  
# i 471 more rows
```

Here's the model with individual data

3. Models for Binary Outcomes

```
beetles_glm_ungrouped =  
  beetles_long |>  
  glm(  
    formula = outcome ~ dose,  
    family = "binomial")  
  
beetles_glm_ungrouped |> parameters() |> print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-------------|----------|------|------------------|--------|--------|
| (Intercept) | -60.72 | 5.18 | (-71.44, -51.08) | -11.72 | < .001 |
| dose | 34.27 | 2.91 | (28.85, 40.30) | 11.77 | < .001 |

Here's the previous version again:

```
beetles_glm_grouped |> parameters() |> print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-------------|----------|------|------------------|--------|--------|
| (Intercept) | -60.72 | 5.18 | (-71.44, -51.08) | -11.72 | < .001 |
| dose | 34.27 | 2.91 | (28.85, 40.30) | 11.77 | < .001 |

They seem the same! But not quite:

```
logLik(beetles_glm_grouped)
```

```
'log Lik.' -18.72 (df=2)
```

```
logLik(beetles_glm_ungrouped)
```

3. Models for Binary Outcomes

'log Lik.' -186.2 (df=2)

The difference is due to the binomial coefficient $\binom{n}{x}$ which isn't included in the individual-observations (Bernoulli) version of the model.

3.4. Multiple logistic regression

3.4.1. Coronary heart disease (WCGS) study data

Let's use the data from the following study to explore multiple logistic regression:

3.4.1.1. Summary of study

From Vittinghoff et al. (2012):

“The **Western Collaborative Group Study (WCGS)** was a large epidemiological study designed to investigate the association between the “type A” behavior pattern and coronary heart disease (CHD) (Rosenman et al. 1964).”

From Wikipedia, “Type A and Type B personality theory”:

“The hypothesis describes Type A individuals as outgoing, ambitious, rigidly organized, highly status-conscious, impatient, anxious, proactive, and concerned with time management....

The hypothesis describes Type B individuals as a contrast to those of Type A. Type B personalities, by definition, are noted to live at lower stress levels. They typically work steadily and may enjoy achievement, although they have a greater tendency to disregard physical or mental stress when they do not achieve.”

3. Models for Binary Outcomes

3.4.1.2. Study design

from ?faraway::wcss:

3154 healthy young men aged 39-59 from the San Francisco area were assessed for their personality type. All were free from coronary heart disease at the start of the research. Eight and a half years later change in CHD status was recorded.

Details (from faraway::wcss)

The WCSS began in 1960 with 3,524 male volunteers who were employed by 11 California companies. Subjects were 39 to 59 years old and free of heart disease as determined by electrocardiogram. After the initial screening, the study population dropped to 3,154 and the number of companies to 10 because of various exclusions. The cohort comprised both blue- and white-collar employees.

At baseline the following information was collected:

- socio-demographic including:
 - age
 - education
 - marital status
 - income
 - occupation
- physical and physiological including:
 - height
 - weight
 - blood pressure
 - electrocardiogram
 - corneal arcus;
- biochemical including:
 - cholesterol and lipoprotein fractions;
 - medical and family history and use of medications;
 - behavioral data including

3. Models for Binary Outcomes

- Type A interview,
- smoking,
- exercise
- alcohol use.

Later surveys added data on:

- anthropometry
- triglycerides
- Jenkins Activity Survey
- caffeine use

Average follow-up continued for 8.5 years with repeat examinations.

Reference: Coronary Heart Disease in the Western Collaborative Group Study Final Follow-up Experience of 8 1/2 Years Ray H. Rosenman, MD; Richard J. Brand, PhD; C. David Jenkins, PhD; Meyer Friedman, MD; Reuben Straus, MD; Moses Wurm, MD JAMA. 1975;233(8):872-877. doi:10.1001/jama.1975.03260080034016.

3.4.2. Load the data

Here, I load the data:

```
### load the data directly from a UCSF website:
# library(haven)
# url = paste0(
#   # I'm breaking up the url into two chunks for readability
#   "https://regression.ucsf.edu/sites/g/files/",
#   "tkssra6706/f/wysiwyg/home/data/wcgs.dta")
# wcgs = haven::read_dta(url)

# I presaved the data in my project's `data` folder
```

3. Models for Binary Outcomes

```
library(here) # provides the `here()` function
library(fs) # provides the `path()` function
here::here() |>
  fs::path('data/wcgs.rda') |>
  load()
```

3.4.3. Now let's do some data cleaning

```
library(arsenal) # provides `set_labels()`

wcgs = wcgs |>
  mutate(
    age = age |>
      arsenal::set_labels("Age (years)"),

    arcus =
      arcus |>
      as.logical() |>
      arsenal::set_labels("Arcus Senilis"),

    time169 =
      time169 |>
      as.numeric() |>
      arsenal::set_labels("Observation (follow up) time (days)",

    dibpat =
      dibpat |>
      as_factor() |>
      relevel(ref = "Type A") |>
      arsenal::set_labels("Behavioral Pattern"),
```

3. Models for Binary Outcomes

```
typchd69 = typchd69 |>
  labelled(
    label = "Type of CHD Event",
    labels =
      c(
        "None" = 0,
        "infdeath" = 1,
        "silent" = 2,
        "angina" = 3)),

# turn stata-style labelled variables in to R-style factors:
across(
  where(is.labelled),
  haven::as_factor)
)
```

3.4.4. What's in the data

Here's a table of the data:

```
wcgs |>
  select(-c(id, uni, t1)) |>
  tableby(chd69 ~ ., data = _) |>
  summary(
    pfootnote = TRUE,
    title =
      "Baseline characteristics by CHD status at end of follow-up")
```


3. Models for Binary Outcomes

Table 3.6.: Baseline characteristics by CHD status at end of follow-up

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|------------------------------------|--------------------|--------------------|--------------------|-------------------------|
| Age (years) | | | | < 0.001 ¹ |
| Mean (SD) | 46.082 (5.457) | 48.490 (5.801) | 46.279 (5.524) | |
| Range | 39.000 - 59.000 | 39.000 - 59.000 | 39.000 - 59.000 | |
| Arcus Senilis | | | | < 0.001 ² |
| N-Miss | 0 | 2 | 2 | |
| FALSE | 2058 (71.0%) | 153 (60.0%) | 2211 (70.1%) | |
| TRUE | 839 (29.0%) | 102 (40.0%) | 941 (29.9%) | |
| Behavioral Pattern | | | | < 0.001 ² |
| A1 | 234 (8.1%) | 30 (11.7%) | 264 (8.4%) | |
| A2 | 1177 (40.6%) | 148 (57.6%) | 1325 (42.0%) | |
| B3 | 1155 (39.9%) | 61 (23.7%) | 1216 (38.6%) | |
| B4 | 331 (11.4%) | 18 (7.0%) | 349 (11.1%) | |
| Body Mass Index (kg/m2) | | | | < 0.001 ¹ |
| Mean (SD) | 24.471 (2.561) | 25.055 (2.579) | 24.518 (2.567) | |
| Range | 11.191 - 37.653 | 19.225 - 38.947 | 11.191 - 38.947 | |

3. Models for Binary Outcomes

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|--------------------------------------|----------------------|----------------------|----------------------|-------------------------|
| Total Cholesterol | | | | < 0.001 ¹ |
| N-Miss | 12 | 0 | 12 | |
| Mean (SD) | 224.261 (42.217) | 250.070 (49.396) | 226.372 (43.420) | |
| Range | 103.000 - 400.000 | 155.000 - 645.000 | 103.000 - 645.000 | |
| Diastolic Blood Pressure | | | | < 0.001 ¹ |
| Mean (SD) | 81.723 (9.621) | 85.315 (10.311) | 82.016 (9.727) | |
| Range | 58.000 - 150.000 | 64.000 - 122.000 | 58.000 - 150.000 | |
| Behavioral Pattern | | | | < 0.001 ² |
| Type A | 1411 (48.7%) | 178 (69.3%) | 1589 (50.4%) | |
| Type B | 1486 (51.3%) | 79 (30.7%) | 1565 (49.6%) | |
| Height (inches) | | | | 0.290 ¹ |
| Mean (SD) | 69.764 (2.539) | 69.938 (2.410) | 69.778 (2.529) | |
| Range | 60.000 - 78.000 | 63.000 - 77.000 | 60.000 - 78.000 | |
| Ln of Systolic Blood Pressure | | | | < 0.001 ¹ |
| Mean (SD) | 4.846 (0.110) | 4.900 (0.125) | 4.850 (0.112) | |
| Range | 4.585 - 5.438 | 4.605 - 5.298 | 4.585 - 5.438 | |

3. Models for Binary Outcomes

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|--|-----------------------|-----------------------|-----------------------|-------------------------|
| Ln of Weight | | | | < 0.001 ¹ |
| Mean (SD) | 5.126 (0.123) | 5.155 (0.118) | 5.128 (0.123) | |
| Range | 4.357 - 5.670 | 4.868 - 5.768 | 4.357 - 5.768 | |
| Cigarettes per day | | | | < 0.001 ¹ |
| Mean (SD) | 11.151 (14.329) | 16.665 (15.657) | 11.601 (14.518) | |
| Range | 0.000 - 99.000 | 0.000 - 60.000 | 0.000 - 99.000 | |
| Systolic Blood Pressure | | | | < 0.001 ¹ |
| Mean (SD) | 128.034 (14.746) | 135.385 (17.473) | 128.633 (15.118) | |
| Range | 98.000 - 230.000 | 100.000 - 200.000 | 98.000 - 230.000 | |
| Current smoking | | | | < 0.001 ² |
| No | 1554 (53.6%) | 98 (38.1%) | 1652 (52.4%) | |
| Yes | 1343 (46.4%) | 159 (61.9%) | 1502 (47.6%) | |
| Observation (follow up) time (days) | | | | < 0.001 ¹ |
| Mean (SD) | 2775.158 (562.205) | 1654.700 (859.297) | 2683.859 (666.524) | |
| Range | 238.000 - 3430.000 | 18.000 - 3229.000 | 18.000 - 3430.000 | |

3. Models for Binary Outcomes

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|----------------------------|---------------------|----------------------|---------------------|-------------------------|
| Type of CHD Event | | | | |
| None | 0 (0.0%) | 0 (0.0%) | 0 (0.0%) | |
| infdeath | 2897 (100.0%) | 0 (0.0%) | 2897 (91.9%) | |
| silent | 0 (0.0%) | 135 (52.5%) | 135 (4.3%) | |
| angina | 0 (0.0%) | 71 (27.6%) | 71 (2.3%) | |
| 4 | 0 (0.0%) | 51 (19.8%) | 51 (1.6%) | |
| Weight (lbs) | | | | < 0.001 ¹ |
| Mean (SD) | 169.554 (21.010) | 174.463 (21.573) | 169.954 (21.096) | |
| Range | 78.000 - 290.000 | 130.000 - 320.000 | 78.000 - 320.000 | |
| Weight Category | | | | < 0.001 ² |
| < 140 | 217 (7.5%) | 15 (5.8%) | 232 (7.4%) | |
| 140-170 | 1440 (49.7%) | 98 (38.1%) | 1538 (48.8%) | |
| 170-200 | 1049 (36.2%) | 122 (47.5%) | 1171 (37.1%) | |
| > 200 | 191 (6.6%) | 22 (8.6%) | 213 (6.8%) | |
| RECODE of age (Age) | | | | < 0.001 ² |
| 35-40 | 512 (17.7%) | 31 (12.1%) | 543 (17.2%) | |
| 41-45 | 1036 (35.8%) | 55 (21.4%) | 1091 (34.6%) | |
| 46-50 | 680 (23.5%) | 70 (27.2%) | 750 (23.8%) | |

3. Models for Binary Outcomes

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|-------|----------------|----------------|-------------------|------------|
| 51-55 | 463 (16.0%) | 65 (25.3%) | 528 (16.7%) | |
| 56-60 | 206 (7.1%) | 36 (14.0%) | 242 (7.7%) | |

1. Linear Model ANOVA
2. Pearson's Chi-squared test

3.4.5. Data by age and personality type

For now, we will look at the interaction between age and personality type (dibpat). To make it easier to visualize the data, we summarize the event rates for each combination of age:

```
chd_grouped_data =
  wgs |>
  summarize(
    .by = c(age, dibpat),
    n = n(),
    `p(chd)` = mean(chd69 == "Yes") |>
      labelled(label = "CHD Event by 1969"),
    `odds(chd)` = `p(chd)`/(1-`p(chd)`),
    `logit(chd)` = log(`odds(chd)`)
  )

chd_grouped_data
```

| age | dibpat | n | p(chd) | odds(chd) | logit(chd) |
|-----|--------|----|----------|-----------|------------|
| 50 | Type A | 76 | 0.105263 | 0.1176 | -2.140 |
| 51 | Type A | 67 | 0.164179 | 0.1964 | -1.627 |

3. Models for Binary Outcomes

| age | dibpat | n | p(chd) | odds(chd) | logit(chd) |
|-----|--------|-----|----------|-----------|------------|
| 59 | Type A | 30 | 0.233333 | 0.3043 | -1.190 |
| 44 | Type A | 113 | 0.079646 | 0.0865 | -2.447 |
| 47 | Type A | 72 | 0.097222 | 0.1077 | -2.228 |
| 40 | Type A | 133 | 0.067669 | 0.0726 | -2.623 |
| 41 | Type A | 108 | 0.064815 | 0.0693 | -2.669 |
| 43 | Type A | 97 | 0.072165 | 0.0778 | -2.554 |
| 54 | Type A | 53 | 0.132075 | 0.1522 | -1.883 |
| 48 | Type A | 80 | 0.150000 | 0.1765 | -1.735 |
| 39 | Type A | 128 | 0.085938 | 0.0940 | -2.364 |
| 49 | Type A | 67 | 0.238806 | 0.3137 | -1.159 |
| 55 | Type A | 55 | 0.163636 | 0.1957 | -1.631 |
| 56 | Type A | 49 | 0.244898 | 0.3243 | -1.126 |
| 42 | Type A | 101 | 0.039604 | 0.0412 | -3.188 |
| 45 | Type A | 77 | 0.090909 | 0.1000 | -2.303 |
| 46 | Type A | 91 | 0.065934 | 0.0706 | -2.651 |
| 57 | Type A | 31 | 0.129032 | 0.1481 | -1.909 |
| 53 | Type A | 62 | 0.112903 | 0.1273 | -2.061 |
| 52 | Type A | 65 | 0.200000 | 0.2500 | -1.386 |
| 58 | Type A | 34 | 0.147059 | 0.1724 | -1.758 |
| 45 | Type B | 109 | 0.045872 | 0.0481 | -3.035 |
| 41 | Type B | 125 | 0.040000 | 0.0417 | -3.178 |
| 47 | Type B | 75 | 0.013333 | 0.0135 | -4.304 |
| 39 | Type B | 138 | 0.057971 | 0.0615 | -2.788 |
| 49 | Type B | 67 | 0.074627 | 0.0806 | -2.518 |
| 51 | Type B | 56 | 0.107143 | 0.1200 | -2.120 |
| 42 | Type B | 121 | 0.008264 | 0.0083 | -4.787 |
| 50 | Type B | 59 | 0.050847 | 0.0536 | -2.927 |
| 44 | Type B | 122 | 0.032787 | 0.0339 | -3.384 |
| 56 | Type B | 27 | 0.000000 | 0.0000 | -Inf |
| 40 | Type B | 144 | 0.020833 | 0.0213 | -3.850 |
| 58 | Type B | 22 | 0.045455 | 0.0476 | -3.045 |
| 48 | Type B | 84 | 0.083333 | 0.0909 | -2.398 |

3. Models for Binary Outcomes

| age | dibpat | n | p(chd) | odds(chd) | logit(chd) |
|-----|--------|-----|----------|-----------|------------|
| 43 | Type B | 118 | 0.050847 | 0.0536 | -2.927 |
| 53 | Type B | 43 | 0.116279 | 0.1316 | -2.028 |
| 54 | Type B | 54 | 0.074074 | 0.0800 | -2.526 |
| 46 | Type B | 79 | 0.063291 | 0.0676 | -2.695 |
| 52 | Type B | 48 | 0.041667 | 0.0435 | -3.135 |
| 55 | Type B | 25 | 0.040000 | 0.0417 | -3.178 |
| 57 | Type B | 32 | 0.093750 | 0.1034 | -2.269 |
| 59 | Type B | 17 | 0.235294 | 0.3077 | -1.179 |

3.4.6. Graphical exploration

3.4.6.1. Probability scale

```
library(ggplot2)
library(ggeasy)
library(scales)
chd_plot_probs =
  chd_grouped_data |>
  ggplot(
    aes(
      x = age,
      y = `p(chd)`,
      col = dibpat)
  ) +
  geom_point(aes(size = n), alpha = .7) +
  scale_size(range = c(1,4)) +
  geom_line() +
  theme_bw() +
  ylab("P(CHD Event by 1969)") +
  scale_y_continuous(
```

3. Models for Binary Outcomes

```
labels = scales::label_percent(),
sec.axis = sec_axis(
  ~ odds(.),
  name = "odds(CHD Event by 1969)") +
ggeasy::easy_labs() +
theme(legend.position = "bottom")
```

```
print(chd_plot_probs)
```




Figure 3.4.: CHD rates by age group, probability scale

3.4.6.2. Odds scale

3. Models for Binary Outcomes

```
trans_odds = trans_new(  
  name = "odds",  
  transform = odds,  
  inverse = odds_inv)  
  
chd_plot_probs +  
  scale_y_continuous(  
    trans = trans_odds,  
    name = paste(chd_plot_probs$labels$y, "(odds spacing)"),  
    sec.axis = sec_axis(  
      ~ odds(.),  
      name = "odds(CHD Event by 1969)")  
)  
  
chd_plot_odds =  
  chd_grouped_data |>  
  ggplot(  
    aes(  
      x = age,  
      y = `odds(chd)`,  
      col = dibpat)  
  ) +  
  geom_point(aes(size = n), alpha = .7) +  
  scale_size(range = c(1,4)) +  
  geom_line() +  
  theme_bw() +  
  ylab("odds(CHD Event by 1969)") +  
  ggeasy::easy_labs()
```

3. Models for Binary Outcomes





Figure 3.5.: CHD rates by age group, odds scale

```
print(chd_plot_odds)
```

3. Models for Binary Outcomes




3.4.6.3. Log-odds (logit) scale

```
chd_plot_logit =  
  chd_grouped_data |>  
  ggplot(  
    aes(  
      x = age,  
      y = `logit(chd)`,  
      col = dibpat)  
    ) +  
  geom_point(aes(size = n), alpha = .7) +  
  scale_size(range = c(1,4)) +  
  geom_line() +  
  theme_bw() +
```

3. Models for Binary Outcomes

```
ylab("log{odds(CHD Event by 1969)}") +  
ggeasy::easy_labs()
```

```
print(chd_plot_logit)
```



3.4.7. Logistic regression models for CHD data

Here, we fit stratified models for CHD by personality type.

```
chd_glm_strat = glm(  
  "formula" = chd69 == "Yes" ~ dibpat + dibpat:age - 1,  
  "data" = wgs,  
  "family" = binomial(link = "logit")  
)
```

3. Models for Binary Outcomes

```
chd_glm_strat |> parameters() |> print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-----------------------|----------|------|----------------|-------|--------|
| dibpat (Type A) | -5.50 | 0.67 | (-6.83, -4.19) | -8.18 | < .001 |
| dibpat (Type B) | -5.80 | 0.98 | (-7.73, -3.90) | -5.95 | < .001 |
| dibpat (Type A) × age | 0.07 | 0.01 | (0.05, 0.10) | 5.24 | < .001 |
| dibpat (Type B) × age | 0.06 | 0.02 | (0.02, 0.10) | 3.01 | 0.003 |

We can get the corresponding odds ratios (e^β s) by passing `exponentiate = TRUE` to `parameters()`:

```
chd_glm_strat |>
  parameters(exponentiate = TRUE) |>
  print_md()
```

| Parameter | Odds Ratio | SE | 95% CI | z | p |
|-----------------------|------------|----------|------------------|-------|--------|
| dibpat (Type A) | 4.09e-03 | 2.75e-03 | (1.08e-03, 0.02) | -8.18 | < .001 |
| dibpat (Type B) | 3.02e-03 | 2.94e-03 | (4.40e-04, 0.02) | -5.95 | < .001 |
| dibpat (Type A) × age | 1.07 | 0.01 | (1.05, 1.10) | 5.24 | < .001 |
| dibpat (Type B) × age | 1.06 | 0.02 | (1.02, 1.11) | 3.01 | 0.003 |

3.4.8. Models superimposed on data

We can graph our fitted models on each scale (probability, odds, log-odds).

3.4.8.1. probability scale


```
curve_type_A = function(x)
{
  chd_glm_strat |> predict(
    type = "response",
    newdata = tibble(age = x, dibpat = "Type A"))
}

curve_type_B = function(x)
{
  chd_glm_strat |> predict(
    type = "response",
    newdata = tibble(age = x, dibpat = "Type B"))
}

chd_plot_probs_2 =
  chd_plot_probs +
  geom_function(
    fun = curve_type_A,
    aes(col = "Type A")
  ) +
  geom_function(
    fun = curve_type_B,
    aes(col = "Type B")
  )
```

3. Models for Binary Outcomes

```
print(chd_plot_probs_2)
```




3.4.8.2. odds scale

```
curve_type_A = function(x)
{
  chd_glm_strat |> predict(
    type = "link",
    newdata = tibble(age = x, dibpat = "Type A")) |> exp()
}
curve_type_B = function(x)
{
  chd_glm_strat |> predict(
```

3. Models for Binary Outcomes

```
    type = "link",  
    newdata = tibble(age = x, dibpat = "Type B")) |> exp()  
}  
  
chd_plot_odds_2 =  
  chd_plot_odds +  
  geom_function(  
    fun = curve_type_A,  
    aes(col = "Type A")  
  ) +  
  geom_function(  
    fun = curve_type_B,  
    aes(col = "Type B")  
  )  
)
```

```
print(chd_plot_odds_2)
```



3. Models for Binary Outcomes


3.4.8.3. log-odds (logit) scale

```
curve_type_A = function(x)
{
  chd_glm_strat |> predict(
    type = "link",
    newdata = tibble(age = x, dibpat = "Type A"))
}
curve_type_B = function(x)
{
  chd_glm_strat |> predict(
    type = "link",
    newdata = tibble(age = x, dibpat = "Type B"))
}

chd_plot_logit_2 =
  chd_plot_logit +
  geom_function(
    fun = curve_type_A,
    aes(col = "Type A")
  ) +
  geom_function(
    fun = curve_type_B,
    aes(col = "Type B")
  )

print(chd_plot_logit_2)
```

3. Models for Binary Outcomes



3.4.9. reference-group and contrast parametrization

We can also use the corner-point parametrization (with reference groups and contrasts):

```
chd_glm_contrasts =  
  wgs |>  
  glm(  
    "data" = _,  
    "formula" = chd69 == "Yes" ~ dibpat*age,  
    "family" = binomial(link = "logit")  
  )  
  
chd_glm_contrasts |>
```

3. Models for Binary Outcomes

```
parameters() |>
print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-----------------------|----------|------|----------------|-------|--------|
| (Intercept) | -5.50 | 0.67 | (-6.83, -4.19) | -8.18 | < .001 |
| dibpat (Type B) | -0.30 | 1.18 | (-2.63, 2.02) | -0.26 | 0.797 |
| age | 0.07 | 0.01 | (0.05, 0.10) | 5.24 | < .001 |
| dibpat (Type B) × age | -0.01 | 0.02 | (-0.06, 0.04) | -0.42 | 0.674 |

Compare with what we had before:

```
chd_glm_strat |>
parameters() |>
print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-----------------------|----------|------|----------------|-------|--------|
| dibpat (Type A) | -5.50 | 0.67 | (-6.83, -4.19) | -8.18 | < .001 |
| dibpat (Type B) | -5.80 | 0.98 | (-7.73, -3.90) | -5.95 | < .001 |
| dibpat (Type A) × age | 0.07 | 0.01 | (0.05, 0.10) | 5.24 | < .001 |
| dibpat (Type B) × age | 0.06 | 0.02 | (0.02, 0.10) | 3.01 | 0.003 |

Exercise 3.12. If I give you model 1, how would you get the coefficients of model 2?

3.5. Fitting logistic regression models

3.5.1.

In general, the estimating equation $\ell'(\beta; \mathbf{x}) = 0$ cannot be solved analytically.

Instead, we have to use a variant of the Newton-Raphson method, which was discussed briefly in Epi 203. We won't go over it in this class; if you need to learn it, see Dobson and Barnett (2018), Chapter 4.

For now, all you need to know is that we make an iterative series of guesses, and each guess helps us make the next guess better (higher log-likelihood).

You can see some information about this process like so:

```
options(digits = 8)
temp =
  wgs |>
  glm(
    control = glm.control(trace = TRUE),
    "data" = _,
    "formula" = chd69 == "Yes" ~ dibpat*age,
    "family" = binomial(link = "logit")
  )
```

```
Deviance = 1775.7899 Iterations - 1
Deviance = 1708.5396 Iterations - 2
Deviance = 1704.0434 Iterations - 3
Deviance = 1703.9833 Iterations - 4
Deviance = 1703.9832 Iterations - 5
Deviance = 1703.9832 Iterations - 6
```

3. Models for Binary Outcomes

After each iteration of the fitting procedure, the deviance $(2(\ell_{\text{full}} - \ell(\hat{\beta})))$ is printed. You can see that the algorithm took six iterations to converge to a solution where the likelihood wasn't changing much anymore.

3.6. Model comparisons for logistic models

3.6.1. Deviance test

We can compare the maximized log-likelihood of our model, $\ell(\hat{\beta}; \mathbf{x})$, versus the log-likelihood of the full model (aka saturated model aka maximal model), ℓ_{full} , which has one parameter per covariate pattern. With enough data, $2(\ell_{\text{full}} - \ell(\hat{\beta}; \mathbf{x})) \sim \chi^2(N - p)$, where N is the number of distinct covariate patterns and p is the number of β parameters in our model. A significant p-value for this **deviance** statistic indicates that there's some detectable pattern in the data that our model isn't flexible enough to catch.

Caution

The deviance statistic needs to have a large amount of data **for each covariate pattern** for the χ^2 approximation to hold. A guideline from Dobson is that if there are q distinct covariate patterns x_1, \dots, x_q , with n_1, \dots, n_q observations per pattern, then the expected frequencies $n_k \cdot \pi(x_k)$ should be at least 1 for every pattern $k \in 1 : q$.

If you have covariates measured on a continuous scale, you may not be able to use the deviance tests to assess goodness of fit.

3.6.2. Hosmer-Lemeshow test

If our covariate patterns produce groups that are too small, a reasonable solution is to make bigger groups by merging some of the covariate-pattern

3. Models for Binary Outcomes

groups together.

Hosmer and Lemeshow (1980) proposed that we group the patterns by their predicted probabilities according to the model of interest. For example, you could group all of the observations with predicted probabilities of 10% or less together, then group the observations with 11%-20% probability together, and so on; $g = 10$ categories in all.

Then we can construct a statistic

$$X^2 = \sum_{c=1}^g \frac{(o_c - e_c)^2}{e_c}$$

where o_c is the number of events *observed* in group c , and e_c is the number of events expected in group c (based on the sum of the fitted values $\hat{\pi}_i$ for observations in group c).

If each group has enough observations in it, you can compare X^2 to a χ^2 distribution; by simulation, the degrees of freedom has been found to be approximately $g - 2$.

For our CHD model, this procedure would be:

```
wcgs =  
  wcgs |>  
  mutate(  
    pred_probs_glm1 = chd_glm_strat |> fitted(),  
    pred_prob_cats1 =  
      pred_probs_glm1 |>  
      cut(breaks = seq(0, 1, by = .1),  
          include.lowest = TRUE))  
  
HL_table =  
  wcgs |>  
  summarize(  
    .by = pred_prob_cats1,
```

3. Models for Binary Outcomes

```
n = n(),
o = sum(chd69 == "Yes"),
e = sum(pred_probs_glm1)
)

HL_table |> pander()
```

| pred_prob_cats1 | n | o | e |
|-----------------|------|-----|-------|
| (0.1,0.2] | 785 | 116 | 108 |
| (0.2,0.3] | 64 | 12 | 13.77 |
| [0,0.1] | 2305 | 129 | 135.2 |

```
X2 = HL_table |>
  summarize(
    `X^2` = sum((o-e)^2/e)
  ) |>
  pull(`X^2`)
print(X2)
```

```
[1] 1.1102871
```

```
pval1 = pchisq(X2, lower = FALSE, df = nrow(HL_table) - 2)
```

Our statistic is $X^2 = 1.11028711$; $p(\chi^2(1) > 1.11028711) = 0.29201955$, which is our p-value for detecting a lack of goodness of fit.

Unfortunately that grouping plan left us with just three categories with any observations, so instead of grouping by 10% increments of predicted probability, typically analysts use deciles of the predicted probabilities:

3. Models for Binary Outcomes

```
wcgs =  
  wcgs |>  
  mutate(  
    pred_probs_glm1 = chd_glm_strat |> fitted(),  
    pred_prob_cats1 =  
      pred_probs_glm1 |>  
      cut(breaks = quantile(pred_probs_glm1, seq(0, 1, by = .1)),  
          include.lowest = TRUE))  
  
HL_table =  
  wcgs |>  
  summarize(  
    .by = pred_prob_cats1,  
    n = n(),  
    o = sum(chd69 == "Yes"),  
    e = sum(pred_probs_glm1)  
  )  
  
HL_table |> pander()
```

| pred_prob_cats1 | n | o | e |
|-----------------|-----|----|-------|
| (0.114,0.147] | 275 | 48 | 36.81 |
| (0.147,0.222] | 314 | 51 | 57.19 |
| (0.0774,0.0942] | 371 | 27 | 32.56 |
| (0.0942,0.114] | 282 | 30 | 29.89 |
| (0.0633,0.069] | 237 | 17 | 15.97 |
| (0.069,0.0774] | 306 | 20 | 22.95 |
| (0.0487,0.0633] | 413 | 27 | 24.1 |
| (0.0409,0.0487] | 310 | 14 | 14.15 |
| [0.0322,0.0363] | 407 | 16 | 13.91 |
| (0.0363,0.0409] | 239 | 7 | 9.48 |

3. Models for Binary Outcomes

```
X2 = HL_table |>
  summarize(
    `X^2` = sum((o-e)^2/e)
  ) |>
  pull(`X^2`)

print(X2)
```

```
[1] 6.7811383
```

```
pval1 = pchisq(X2, lower = FALSE, df = nrow(HL_table) - 2)
```


Now we have more evenly split categories. The p-value is 0.56041994, still not significant.

Graphically, we have compared:

```
HL_plot =
  HL_table |>
  ggplot(aes(x = pred_prob_cats1)) +
  geom_line(aes(y = e, x = pred_prob_cats1, group = "Expected", col = "Expected")) +
  geom_point(aes(y = e, size = n, col = "Expected")) +
  geom_point(aes(y = o, size = n, col = "Observed")) +
  geom_line(aes(y = o, col = "Observed", group = "Observed")) +
  scale_size(range = c(1,4)) +
  theme_bw() +
  ylab("number of CHD events") +
  theme(axis.text.x = element_text(angle = 45))

print(HL_plot)
```

3. Models for Binary Outcomes



3.6.3. Comparing models

- $AIC = -2 * \ell(\hat{\theta}) + 2 * p$ [lower is better]
- $BIC = -2 * \ell(\hat{\theta}) + p * \log(n)$ [lower is better]
- likelihood ratio [higher is better]

3.7. Residual-based diagnostics

3.7.1. Logistic regression residuals only work for grouped data

Residuals only work if there is more than one observation for most covariate patterns.

3. Models for Binary Outcomes

Here we will create the grouped-data version of our CHD model from the WCGS study:

```
wcgs_grouped =
  wcgs |>
  summarize(
    .by = c(dibpat, age),
    n = n(),
    chd = sum(chd69 == "Yes"),
    `!chd` = sum(chd69 == "No")
  )

chd_glm_strat_grouped = glm(
  "formula" = cbind(chd, `!chd`) ~ dibpat + dibpat:age - 1,
  "data" = wcgs_grouped,
  "family" = binomial(link = "logit")
)

chd_glm_strat_grouped |> parameters() |> print_md()
```

| Parameter | Log-Odds | SE | 95% CI | z | p |
|-----------------------|----------|------|----------------|-------|--------|
| dibpat (Type A) | -5.50 | 0.67 | (-6.83, -4.19) | -8.18 | < .001 |
| dibpat (Type B) | -5.80 | 0.98 | (-7.73, -3.90) | -5.95 | < .001 |
| dibpat (Type A) × age | 0.07 | 0.01 | (0.05, 0.10) | 5.24 | < .001 |
| dibpat (Type B) × age | 0.06 | 0.02 | (0.02, 0.10) | 3.01 | 0.003 |

3.7.2. (Response) residuals

$$e_k \stackrel{\text{def}}{=} \bar{y}_k - \hat{\pi}(x_k)$$

3. Models for Binary Outcomes

(k indexes the covariate patterns)

We can graph these residuals e_k against the fitted values $\hat{\pi}(x_k)$:

```
wcgs_grouped =  
  wcgs_grouped |>  
  mutate(  
    fitted = chd_glm_strat_grouped |> fitted(),  
    fitted_logit = fitted |> logit(),  
    response_resids =  
      chd_glm_strat_grouped |> resid(type = "response")  
  )  
  
wcgs_response_resid_plot =  
  wcgs_grouped |>  
  ggplot(  
    mapping = aes(  
      x = fitted,  
      y = response_resids  
    )  
  ) +  
  geom_point(  
    aes(col = dibpat)  
  ) +  
  geom_hline(yintercept = 0) +  
  geom_smooth(  
    se = TRUE,  
    method.args = list(  
      span=2/3,  
      degree=1,  
      family="symmetric",  
      iterations=3),  
    method = stats::loess) ①
```


① Don't worry about these options for now; I chose them to match

3. Models for Binary Outcomes

`autoplot()` as closely as I can. `plot.glm` and `autoplot` use `stats::lowess` instead of `stats::loess`; `stats::lowess` is older, hard to use with `geom_smooth`, and hard to match exactly with `stats::loess`; see <https://support.bioconductor.org/p/2323/>.]

```
wcgs_response_resid_plot |> print()
```

3. Models for Binary Outcomes



We can see a slight fan-shape here: observations on the right have larger variance (as expected since $\text{var}(\bar{y}) = \pi(1 - \pi)/n$ is maximized when $\pi = 0.5$).

3. Models for Binary Outcomes

3.7.3. Pearson residuals

The fan-shape in the response residuals plot isn't necessarily a concern here, since we haven't made an assumption of constant residual variance, as we did for linear regression.

However, we might want to divide by the standard error in order to make the graph easier to interpret. Here's one way to do that:

The Pearson (chi-squared) residual for covariate pattern k is:

$$X_k = \frac{\bar{y}_k - \hat{\pi}_k}{\sqrt{\hat{\pi}_k(1 - \hat{\pi}_k)/n_k}}$$

where


$$\begin{aligned}\hat{\pi}_k &\stackrel{\text{def}}{=} \hat{\pi}(x_k) \\ &\stackrel{\text{def}}{=} \hat{P}(Y = 1|X = x_k) \\ &\stackrel{\text{def}}{=} \text{expit}(x'_k \hat{\beta}) \\ &\stackrel{\text{def}}{=} \text{expit}(\hat{\beta}_0 + \sum_{j=1}^p \hat{\beta}_j x_{kj})\end{aligned}$$

Let's take a look at the Pearson residuals for our CHD model from the WCGS data (graphed against the fitted values on the logit scale):

```
library(ggfortify)
```

```
autoplot(chd_glm_strat_grouped, which = 1, ncol = 1) |> print()
```

3. Models for Binary Outcomes



The fan-shape is gone, and these residuals don't show any obvious signs of model fit issues.


3. Models for Binary Outcomes

3.7.3.1. Pearson residuals plot for beetles data

If we create the same plot for the `beetles` model, we see some strong evidence of a lack of fit:

```
autoplot(beetles_glm_grouped, which = 1, ncol = 1) |> print()
```

3. Models for Binary Outcomes




3.7.3.2. Pearson residuals with individual (ungrouped) data

What happens if we try to compute residuals without grouping the data by covariate pattern?

3. Models for Binary Outcomes

```
library(ggfortify)
```

```
autoplot(chd_glm_strat, which = 1, ncol = 1) |> print()
```



3. Models for Binary Outcomes

Meaningless.

3.7.3.3. Residuals plot by hand (*optional section*)

If you want to check your understanding of what these residual plots are, try building them yourself:


```
wcgs_grouped =  
  wcgs_grouped |>  
  mutate(  
    fitted = chd_glm_strat_grouped |> fitted(),  
    fitted_logit = fitted |> logit(),  
    resids = chd_glm_strat_grouped |> resid(type = "pearson")  
  )  
  
wcgs_resid_plot1 =  
  wcgs_grouped |>  
  ggplot(  
    mapping = aes(  
      x = fitted_logit,  
      y = resids  
    )  
  ) +  
  geom_point(  
    aes(col = dibpat)  
  ) +  
  geom_hline(yintercept = 0) +  
  geom_smooth(se = FALSE,  
             method.args = list(  
               span=2/3,  
               degree=1,  
             )  
  )
```

3. Models for Binary Outcomes

```
family="symmetric",
iterations=3,
surface="direct"
# span = 2/3,
# iterations = 3
),
method = stats::loess)
# plot.glm and autoplot use stats::lowess, which is hard to use with
# geom_smooth and hard to match exactly;
# see https://support.bioconductor.org/p/2323/

wcgs_resid_plot1 |> print()
```

3. Models for Binary Outcomes



3. Models for Binary Outcomes

3.7.4. Pearson chi-squared goodness of fit test

The Pearson chi-squared goodness of fit statistic is:

$$X^2 = \sum_{k=1}^m X_k^2$$

Under the null hypothesis that the model in question is correct (i.e., sufficiently complex), $X^2 \sim \chi^2(N - p)$.

```
X = chd_glm_strat_grouped |>
  resid(type = "pearson")

chisq_stat = sum(X^2)

pval = pchisq(
  chisq_stat,
  lower = FALSE,
  df = length(X) - length(coef(chd_glm_strat_grouped)))
```

For our CHD model, the p-value for this test is 0.26523556; no significant evidence of a lack of fit at the 0.05 level.

3.7.4.1. Standardized Pearson residuals

Especially for small data sets, we might want to adjust our residuals for leverage (since outliers in X add extra variance to the residuals):

$$r_{P_k} = \frac{X_k}{\sqrt{1 - h_k}}$$

where h_k is the leverage of X_k . The functions `autoplot()` and `plot.lm()` use these for some of their graphs.

3. Models for Binary Outcomes

3.7.5. Deviance residuals

For large sample sizes, the Pearson and deviance residuals will be approximately the same. For small sample sizes, the deviance residuals from covariate patterns with small sample sizes can be unreliable (high variance).

$$d_k = \text{sign}(y_k - n_k \hat{\pi}_k) \left\{ \sqrt{2[\ell_{\text{full}}(x_k) - \ell(\hat{\beta}; x_k)]} \right\}$$

3.7.5.1. Standardized deviance residuals


$$r_{D_k} = \frac{d_k}{\sqrt{1 - h_k}}$$

3.7.6. Diagnostic plots

Let's take a look at the full set of `autoplot()` diagnostics now for our CHD model:

```
chd_glm_strat_grouped |> autoplot(which = 1:6) |> print()
```


3. Models for Binary Outcomes



Things look pretty good here. The QQ plot is still usable; with large samples; the residuals should be approximately Gaussian.


3. Models for Binary Outcomes

3.7.6.1. Beetles

Let's look at the beetles model diagnostic plots for comparison:

```
beetles_glm_grouped |> autoplot(which = 1:6) |> print()
```

3. Models for Binary Outcomes



Hard to tell much from so little data, but there might be some issues here.

3.8. Odds Ratios vs Probability (Risk) Ratios

3.8.0.1. Case 1: rare events

For rare events, odds ratios and probability (a.k.a. risk, a.k.a. prevalence) ratios will be close:

$$\pi_1 = .01 \quad \pi_2 = .02$$

```
pi1 = .01  
pi2 = .02  
pi2/pi1
```

```
[1] 2
```

```
odds(pi2)/odds(pi1)
```

```
[1] 2.0204082
```

3.8.0.2. Case 2: frequent events

$$\pi_1 = .4 \quad \pi_2 = .5$$

For more frequently-occurring outcomes, this won't be the case:

```
pi1 = .4  
pi2 = .5  
pi2/pi1
```

```
[1] 1.25
```

3. Models for Binary Outcomes

```
odds(pi2)/odds(pi1)
```

```
[1] 1.5
```

If you want risk ratios, you can sometimes get them by changing the link function:

```
data(anthers, package = "dobson")
anthers.sum<-aggregate(
  anthers[c("n","y")],
  by=anthers[c("storage")],FUN=sum)

anthers_glm_log = glm(
  formula = cbind(y,n-y)~storage,
  data=anthers.sum,
  family=binomial(link="log"))

anthers_glm_log |> parameters() |> print_md()
```

| Parameter | Log-Risk | SE | 95% CI | z | p |
|-------------|----------|------|----------------|-------|--------|
| (Intercept) | -0.80 | 0.12 | (-1.04, -0.58) | -6.81 | < .001 |
| storage | 0.17 | 0.07 | (0.02, 0.31) | 2.31 | 0.021 |

Now $\exp\{\beta\}$ gives us risk ratios instead of odds ratios:

```
anthers_glm_log |> parameters(exponentiate = TRUE) |> print_md()
```

3. Models for Binary Outcomes

| Parameter | Risk Ratio | SE | 95% CI | z | p |
|-------------|------------|------|--------------|-------|--------|
| (Intercept) | 0.45 | 0.05 | (0.35, 0.56) | -6.81 | < .001 |
| storage | 1.18 | 0.09 | (1.03, 1.36) | 2.31 | 0.021 |

Let's compare this model with a logistic model:

```
anthers_glm_logit = glm(  
  formula = cbind(y, n - y) ~ storage,  
  data = anthers.sum,  
  family = binomial(link = "logit"))  
  
anthers_glm_logit |> parameters(exponentiate = TRUE) |> print_md()
```

| Parameter | Odds Ratio | SE | 95% CI | z | p |
|-------------|------------|------|--------------|-------|-------|
| (Intercept) | 0.76 | 0.20 | (0.45, 1.27) | -1.05 | 0.296 |
| storage | 1.49 | 0.26 | (1.06, 2.10) | 2.29 | 0.022 |

[to add: fitted plots on each outcome scale]

When I try to use `link = "log"` in practice, I often get errors about not finding good starting values for the estimation procedure. This is likely because the model is producing fitted probabilities greater than 1.

When this happens, you can try to fit Poisson regression models instead (we will see those soon!). But then the outcome distribution isn't quite right, and you won't get warnings about fitted probabilities greater than 1. In my opinion, the Poisson model for binary outcomes is confusing and not very appealing.

4. Models for Count Outcomes

Poisson regression and variations

Acknowledgements

This content is adapted from:

- Dobson and Barnett (2018), Chapter 9
- Vittinghoff et al. (2012), Chapter 8

Configuring R

Functions from these packages will be used throughout this document:

```
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(dplyr) # manipulate data
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
```

4. Models for Count Outcomes

```
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
pander::panderOptions("table.emphasize.rownames", FALSE)
options('digits' = 4)
```

4.1. Introduction

4.1.1. Examples of count outcomes

- Cyclones per season
- Seconds of tooth-brushing per session (if rounded)
- Infections per person-year
- Visits to ER per person-month
- Car accidents per 1000 miles driven

Note

In many count outcomes, there is some sense of “exposure magnitude” or “duration of observation”: person-year, time at risk, session, miles driven, etc.

4. Models for Count Outcomes

4.1.2. Poisson distribution

$$P(Y = y) = \frac{\mu^y e^{-\mu}}{y!}$$

4.1.2.1. Properties

- $\mathbb{E}[Y] = \mu$
- $\text{Var}[Y] = \mu$

4.1.3. Accounting for exposure

If the exposures/observation durations, denoted $T = t$, are not all equal, we model

$$\mu = \lambda t$$

λ is interpreted as the “expected event rate per unit of exposure”; that is,

$$\lambda = \frac{\mathbb{E}[Y|T = t]}{t}$$

! Important

The exposure magnitude, T , is *similar* to a covariate in linear or logistic regression. However, there is an important difference: in count regression, **there is no intercept corresponding to $\mathbb{E}[Y|T = 0]$** . In other words, this model assumes that if there is no exposure, there can't be any events.

4. Models for Count Outcomes

4.1.4. Adding covariates

With covariates, λ becomes a function of the covariates $\tilde{X} = (X_1, \dots, X_n)$, with a $\log \{\}$ link function (and thus an $\exp \{\}$ inverse-link). That is:

$$\begin{aligned}\mathbb{E}[Y|\tilde{X} = \tilde{x}, T = t] &= \mu(\tilde{x}, t) \\ \mu(\tilde{x}, t) &= \lambda(\tilde{x}) \cdot t \\ \lambda(\tilde{x}) &= \exp \{\eta(\tilde{x})\} \\ \eta(\tilde{x}) &= \tilde{x}'\tilde{\beta} = \beta_0 + \beta_1 x_1 + \dots + \beta_p x_p\end{aligned}$$

Therefore,

$$\begin{aligned}\log \left\{ \left\{ \mathbb{E}[Y|\tilde{X} = \tilde{x}, T = t] \right\} \right\} &= \log \left\{ \left\{ \mu(\tilde{x}) \right\} \right\} \\ &= \log \left\{ \left\{ \lambda(\tilde{x}) \cdot t \right\} \right\} \\ &= \log \left\{ \lambda(\tilde{x}) \right\} + \log \left\{ t \right\} \\ &= \log \left\{ \exp \left\{ \eta(\tilde{x}) \right\} \right\} + \log \left\{ t \right\} \\ &= \eta(\tilde{x}) + \log \left\{ t \right\} \\ &= \tilde{x}'\tilde{\beta} + \log \left\{ t \right\} \\ &= (\beta_0 + \beta_1 x_1 + \dots + \beta_p x_p) + \log \left\{ t \right\}\end{aligned}$$

In contrast with the X s, T enters this expression with a $\log \{\}$ transformation and without a corresponding β coefficient.

i Note

Terms that enter the linear component of a model without a coefficient, such as $\log \{t\}$ here, are called **offsets**.

4.1.5. Rate ratios

Differences on the log-rate scale become ratios on the rate scale.

💡 Tip

$$\exp\{a - b\} = \frac{\exp\{a\}}{\exp\{b\}}$$

Therefore, according to this model, **differences of δ in covariate x_j correspond to rate ratios of $\exp\{\beta_j \cdot \delta\}$.**

That is, letting \tilde{X}_{-j} denote vector \tilde{X} with element j removed:

$$\begin{aligned} & \left\{ \begin{array}{l} \log\{\mathbb{E}[Y|X_j = a, \tilde{X}_{-j} = \tilde{x}_{-j}, T = t]\} \\ -\log\{\mathbb{E}[Y|X_j = b, \tilde{X}_{-j} = \tilde{x}_{-j}, T = t]\} \end{array} \right\} \\ &= \left\{ \begin{array}{l} \log\{t\} + \beta_0 + \beta_1 x_1 + \dots + \beta_j(a) + \dots + \beta_p x_p \\ -\log\{t\} + \beta_0 + \beta_1 x_1 + \dots + \beta_j(b) + \dots + \beta_p x_p \end{array} \right\} \\ &= \beta_j(a - b) \end{aligned}$$

And accordingly,

$$\frac{\mathbb{E}[Y|X_j = a, \tilde{X}_{-j} = \tilde{x}_{-j}, T = t]}{\mathbb{E}[Y|X_j = b, \tilde{X}_{-j} = \tilde{x}_{-j}, T = t]} = \exp\{\beta_j(a - b)\}$$

4.2. Inference for count regression models

4.2.1. Confidence intervals for regression coefficients and rate ratios

As usual:

$$\beta \in \left[\hat{\beta} \pm z_{1-\frac{\alpha}{2}} \cdot \widehat{\text{se}}(\hat{\beta}) \right]$$

Rate ratios: exponentiate CI endpoints

$$\exp\{\beta\} \in \left[\exp\left\{ \hat{\beta} \pm z_{1-\frac{\alpha}{2}} \cdot \widehat{\text{se}}(\hat{\beta}) \right\} \right]$$

4.2.2. Hypothesis tests for regression coefficients

$$t = \frac{\hat{\beta} - \beta_0}{\widehat{\text{se}}(\hat{\beta})}$$

Compare t or $|t|$ to the tails of the standard Gaussian distribution, according to the null hypothesis.

4.2.3. Comparing nested models

log(likelihood ratio) tests, as usual.

4.3. Prediction

$$\begin{aligned}
 \hat{y} &\stackrel{\text{def}}{=} \hat{\mathbb{E}}[Y | \tilde{X} = \tilde{x}, T = t] \\
 &= \hat{\mu}(\tilde{x}, t) \\
 &= \hat{\lambda}(\tilde{x}) \cdot t \\
 &= \exp \{ \hat{\eta}(\tilde{x}) \} \cdot t \\
 &= \exp \{ \tilde{x}' \hat{\beta} \} \cdot t
 \end{aligned}$$

4.4. Diagnostics

4.4.1. Residuals

4.4.1.1. Observation residuals

$$e \stackrel{\text{def}}{=} y - \hat{y}$$

4.4.1.2. Pearson residuals

$$r = \frac{e}{\widehat{\text{se}}(e)} \approx \frac{e}{\sqrt{\hat{y}}}$$

4.4.1.3. Standardized Pearson residuals

$$r_p = \frac{r}{\sqrt{1 - h}}$$

where h is the “leverage” (which we will continue to leave undefined).

4.4.1.4. Deviance residuals

$$d_k = \text{sign}(y - \hat{y}) \left\{ \sqrt{2[\ell_{\text{full}}(y) - \ell(\hat{\beta}; y)]} \right\}$$

i Note

$$\text{sign}(x) \stackrel{\text{def}}{=} \frac{x}{|x|}$$

In other words:

- $\text{sign}(x) = -1$ if $x < 0$
- $\text{sign}(x) = 0$ if $x = 0$
- $\text{sign}(x) = 1$ if $x > 0$

4.5. Zero-inflation

4.5.1. Models for zero-inflated counts

We assume a latent (unobserved) binary variable, Z , which we model using logistic regression:

$$P(Z = 1|X = x) = \pi(x) = \text{expit}(\gamma_0 + \gamma_1 x_1 + \dots)$$

According to this model, if $Z = 1$, then Y will always be zero, regardless of X and T :

$$P(Y = 0|Z = 1, X = x, T = t) = 1$$

Otherwise (if $Z = 0$), Y will have a Poisson distribution, conditional on X and T , as above.

4. Models for Count Outcomes

Even though we never observe Z , we can estimate the parameters γ_0 - γ_p , via maximum likelihood:

$$P(Y = y|X = x, T = t) = P(Y = y, Z = 1|...) + P(Y = y, Z = 0|...)$$

(by the Law of Total Probability)

where

$$P(Y = y, Z = z|...) = P(Y = y|Z = z, ...)P(Z = z|...)$$

4.5.1.1. Exercise

Expand $P(Y = 0|X = x, T = t)$, $P(Y = 1|X = x, T = t)$ and $P(Y = y|X = x, T = t)$ into expressions involving $P(Z = 1|X = x, T = t)$ and $P(Y = y|Z = 0, X = x, T = t)$.

4.5.1.2. Exercise

Derive the expected value and variance of Y , conditional on X and T , as functions of $P(Z = 1|X = x, T = t)$ and $\mathbb{E}[Y|Z = 0, X = x, T = t]$.

4.6. Over-dispersion

4.6.1. Negative binomial models

The Poisson distribution model **forces** the variance to equal the mean. In practice, many count distributions will have a variance substantially larger than the mean (or occasionally smaller).

4. Models for Count Outcomes

When we encounter this, we can try to reduce the residual variance by adding more covariates. However, there are also alternatives to the Poisson model.

Most notably, the negative binomial model:

$$P(Y = y) = \frac{\mu^y}{y!} \cdot \frac{\Gamma(\rho + y)}{\Gamma(\rho) \cdot (\rho + \mu)^y} \cdot \left(1 + \frac{\mu}{\rho}\right)^{-\rho}$$

where ρ is an overdispersion parameter and $\Gamma(x) = (x - 1)!$ for integers x .

You don't need to memorize or understand this expression, but as $\rho \rightarrow \infty$, the second term converges to 1 and the third term converges to $\exp\{-\mu\}$, which brings us back to the Poisson distribution.

For this distribution, $\mathbb{E}[Y] = \mu$ and $\text{Var}(Y) = \mu + \frac{\mu^2}{\rho} > \mu$.

We can still model μ as a function of X and T as before, and we can combine this model with zero-inflation by using it in place of the Poisson distribution for $P(Y = y|Z = 0, X = x, T = t)$.

4.6.2. Quasipoisson

An alternative to Negative binomial is the “quasipoisson” distribution. I've never used it, but it seems to be a method-of-moments type approach rather than maximum likelihood. It models the variance as $\text{Var}(Y) = \mu\theta$, and estimates θ accordingly.

See `?quasipoisson` in R for more.

Part II.

Time to Event Models

In many health sciences applications, binary outcomes are *incompletely observed*. For example, if we are studying whether cancer patients experience a relapse after a initial remission, we may may not be able to follow patients to the end of their lives; instead, we may only know whether each patient has relapsed before the end of the study. If a patient has not relapsed by that point, we might not know if they will relapse at some other date or if they will stay cancer-free for the rest of their lives.¹ Their recurrence status at end-of-life is *missing data*. If some study participants withdraw from a study before the end date in the study design, there will be even more missing data. All of this missing data will make logistic regression difficult for this type of data.

However, these outcome observations are not *entirely* missing. We know that those patients stayed relapse free *at least* until the time point when we last saw them. If we also know the *time-to-event* for the participants who did experience events while under study, we can analyze *time-to-event-or-study-exit*, combined with the indicator of which of these two cases occurred, using *survival analysis*. The survival analysis framework is the subject of the rest of these course notes.

¹Binary outcomes are typically defined *for a specific time-point*. It is important to clearly define whether we are interested in outcome status at end of study, at end of life, or at some other time.

5. Introduction to Survival Analysis

Configuring R

Functions from these packages will be used throughout this document:

```
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(scales) # scales formatting
library(dplyr) # manipulate data
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by default
options('digits' = 4)
```

5.1. Overview

5.1.1. Time-to-event outcomes

Survival analysis is a framework for modeling *time-to-event* outcomes. It is used in:

- clinical trials, where the event is often death or recurrence of disease.
- engineering reliability analysis, where the event is failure of a device or system.
- insurance, particularly life insurance, where the event is death.

i Note

The term *Survival analysis* is a bit misleading. Survival outcomes can sometimes be analyzed using binomial models (logistic regression). *Time-to-event models* might be a better name.

5.2. Time-to-event outcome distributions

5.2.1. Distributions of Time-to-Event Data

- The distribution of event times is asymmetric and can be long-tailed, and starts at 0 (that is, $P(T < 0) = 0$).
- The base distribution is not normal, but exponential.
- There are usually **censored** observations, which are ones in which the failure time is not observed.
- Often, these are **right-censored**, meaning that we know that the event occurred after some known time t , but we don't know the actual event time, as when a patient is still alive at the end of the study.

5. Introduction to Survival Analysis

- Observations can also be **left-censored**, meaning we know the event has already happened at time t , or **interval-censored**, meaning that we only know that the event happened between times t_1 and t_2 .
- Analysis is difficult if censoring is associated with treatment.

5.2.2. Right Censoring

- Patients are in a clinical trial for cancer, some on a new treatment and some on standard of care.
- Some patients in each group have died by the end of the study. We know the survival time (measured for example from time of diagnosis—each person on their own clock).
- Patients still alive at the end of the study are right censored.
- Patients who are lost to follow-up or withdraw from the study may be right-censored.

5.2.3. Left and Interval Censoring

- An individual tests positive for HIV.
- If the event is infection with HIV, then we only know that it has occurred before the testing time t , so this is left censored.
- If an individual has a negative HIV test at time t_1 and a positive HIV test at time t_2 , then the infection event is interval censored.

5.3. Distribution functions for time-to-event variables

5.3.1. The Probability Density Function (PDF)

For a time-to-event variable T with a continuous distribution, the **probability density function** is defined as usual:

5. Introduction to Survival Analysis

$$f(t) \stackrel{\text{def}}{=} p(t) \stackrel{\text{def}}{=} p(T = t)$$


Typically, this density is assumed to be 0 for all $t < 0$; that is, $f(t) = 0, \forall t < 0$. In other words, the range of T is typically $[0, \infty)$.

Example 5.1 (exponential distribution). Recall from Epi 202: the pdf of the exponential distribution family of models is:

$$p(T = t) = \mathbb{1}_{t \geq 0} \cdot \lambda e^{-\lambda t}$$

where $\lambda > 0$.

Here are some examples of exponential pdfs:



5.3.2. The Cumulative Distribution Function (CDF)

The **cumulative distribution function** is defined as:


$$\begin{aligned} F(t) &\stackrel{\text{def}}{=} \Pr(T \leq t) \\ &= \int_{u=0}^t f(u) du \end{aligned}$$

Example 5.2 (exponential distribution). Recall from Epi 202: the cdf of the exponential distribution family of models is:

$$P(T \leq t) = \mathbb{1}_{t \geq 0} \cdot (1 - e^{-\lambda t})$$

where $\lambda > 0$.

Here are some examples of exponential cdfs:



5.3.3. The Survival Function

For survival data, a more important quantity is the **survival function**:

$$\begin{aligned} S(t) &\stackrel{\text{def}}{=} \Pr(T > t) \\ &= \int_{u=t}^{\infty} p(u) du \\ &= 1 - F(t) \end{aligned}$$

The survival function $S(t)$ is the probability that the event time is later than t . If the event in a clinical trial is death, then $S(t)$ is the expected fraction of the original population at time 0 who have survived up to time t and are still alive at time t ; that is, if X_t represents survival status at time t , with $X_t = 1$ denoting alive at time t and $X_t = 0$ denoting deceased at time t , then:

$$S(t) = \mathbb{E}[X_t]$$


Example 5.3 (exponential distribution). Since $S(t) = 1 - F(t)$, the survival function of the exponential distribution family of models is:

$$P(T > t) = \begin{cases} e^{-\lambda t}, & t \geq 0 \\ 1, & t \leq 0 \end{cases}$$

where $\lambda > 0$.

Here are some examples of exponential pdfs:

5. Introduction to Survival Analysis



5.3.4. The Hazard Function

Another important quantity is the **hazard function**:

Definition 5.1 (Hazard function). The hazard function for a random variable T at value t is the conditional density of T at t , given $T \geq t$; that is:

$$h(t) \stackrel{\text{def}}{=} p(T = t | T \geq t)$$

If T represents the time at which an event occurs, then $h(t)$ is the probability that the event occurs at time t , given that it has not occurred prior to time t .

5. Introduction to Survival Analysis

The hazard function has an important relationship to the density and survival functions, which we can use to derive the hazard function for a given probability distribution.

Theorem 5.1.

$$h(t) = \frac{f(t)}{S(t)}$$

Proof.

Lemma 5.1 (Joint probability of a variable with itself).

$$p(T = t, T \geq t) = p(T = t)$$

Proof. Recall from Epi 202: if A and B are statistical events and $A \subseteq B$, then $p(A, B) = p(A)$. In particular, $\{T = t\} \subseteq \{T \geq t\}$, so $p(T = t, T \geq t) = p(T = t)$. \square

Hence:

$$\begin{aligned} h(t) &= p(T = t | T \geq t) \\ &= \frac{p(T = t, T \geq t)}{p(T \geq t)} \\ &= \frac{p(T = t)}{p(T \geq t)} \\ &= \frac{f(t)}{S(t)} \end{aligned}$$

\square

5. Introduction to Survival Analysis

Example 5.4 (exponential distribution). The hazard function of the exponential distribution family of models is:

$$\begin{aligned} P(T = t | T \geq t) &= \frac{f(t)}{S(t)} \\ &= \frac{\mathbb{1}_{t \geq 0} \cdot \lambda e^{-\lambda t}}{e^{-\lambda t}} \\ &= \mathbb{1}_{t \geq 0} \cdot \lambda \end{aligned}$$

Figure 5.1 shows some examples of exponential hazard functions:




Figure 5.1.: Examples of hazard functions for exponential distributions

We can also view the hazard function as the derivative of the negative of the logarithm of the survival function:

Theorem 5.2.

$$h(t) = \frac{d}{dt} \{-\log \{S(t)\}\}$$

Proof.

$$\begin{aligned} h(t) &= \frac{f(t)}{S(t)} \\ &= \frac{-S'(t)}{S(t)} \\ &= -\frac{S'(t)}{S(t)} \\ &= -\frac{d}{dt} \log \{S(t)\} \\ &= \frac{d}{dt} \{-\log \{S(t)\}\} \end{aligned}$$

□

5.3.5. The Cumulative Hazard Function

Since $h(t) = \frac{d}{dt} \{-\log \{S(t)\}\}$ (see Theorem 5.2), we also have:

Corollary 5.1.

$$S(t) = \exp \left\{ - \int_{u=0}^t h(u) du \right\} \quad (5.1)$$

The integral in Equation 5.1 is important enough to have its own name: **cumulative hazard**.

Definition 5.2 (cumulative hazard). The **cumulative hazard function** $H(t)$ is defined as:

$$H(t) \stackrel{\text{def}}{=} \int_{u=0}^t h(u) du$$


5. Introduction to Survival Analysis

As we will see below, $H(t)$ is tractable to estimate, and we can then derive an estimate of the hazard function using an approximate derivative of the estimated cumulative hazard.

Example 5.5. The cumulative hazard function of the exponential distribution family of models is:

$$H(t) = \mathbb{1}_{t \geq 0} \cdot \lambda t$$

Here are some examples of exponential cumulative hazard functions:



5.3.6. Some Key Mathematical Relationships among Survival Concepts

Diagram:

5. Introduction to Survival Analysis

$$h(t) \xrightarrow{\int_{u=0}^t h(u)du} H(t) \xrightarrow{\exp\{-H(t)\}} S(t) \xrightarrow{1-S(t)} F(t)$$

$$h(t) \xleftarrow{\frac{d}{dt} H(t)} H(t) \xleftarrow{-\log\{S(t)\}} S(t) \xleftarrow{1-F(t)} F(t)$$

Identities:

$$\begin{aligned} S(t) &= 1 - F(t) \\ &= \exp\{-H(t)\} \\ S'(t) &= -f(t) \\ H(t) &= -\log\{S(t)\} \\ H'(t) &= h(t) \\ h(t) &= \frac{f(t)}{S(t)} \\ &= -\frac{d}{dt}\log\{S(t)\} \\ f(t) &= h(t) \cdot S(t) \end{aligned}$$

Some proofs (others left as exercises):

$$\begin{aligned} S'(t) &= \frac{d}{dt}(1 - F(t)) \\ &= -F'(t) \\ &= -f(t) \end{aligned}$$

$$\begin{aligned} \frac{d}{dt}\log\{S(t)\} &= \frac{S'(t)}{S(t)} \\ &= -\frac{f(t)}{S(t)} \\ &= -h(t) \end{aligned}$$

5. Introduction to Survival Analysis

$$\begin{aligned} H(t) &\stackrel{\text{def}}{=} \int_{u=0}^t h(u) du \\ &= \int_0^t -\frac{d}{du} \log \{S(u)\} du \\ &= [-\log \{S(u)\}]_{u=0}^{u=t} \\ &= [\log \{S(u)\}]_{u=t}^{u=0} \\ &= \log \{S(0)\} - \log \{S(t)\} \\ &= \log \{1\} - \log \{S(t)\} \\ &= 0 - \log \{S(t)\} \\ &= -\log \{S(t)\} \end{aligned}$$

Equivalently:

$$S(t) = \exp \{-H(t)\}$$

5.3.6.1. Example: Time to death the US in 2004

Daily hazard rates for US Females in 2004

The first day is the most dangerous:



Figure 5.2.: Daily Hazard Rates in 2004 for US Females

Daily hazard rates for US Males and Females in 2004

5. Introduction to Survival Analysis

Exercise: hypothesize why these curves differ where they do?




Figure 5.3.: Daily Hazard Rates in 2004 for US Males and Females 1-40

Survival curve for US females

Exercise: compare and contrast this curve with the corresponding hazard curve.

5. Introduction to Survival Analysis




Figure 5.4.: Survival Curve in 2004 for US Females

5.3.6.2. Likelihood with censoring *

i Note

This subsection was not presented in class in 2023; it is not necessary to understand for the qualifying exam.

If an event time T is observed exactly as $T = t$, then the likelihood of that observation is just its probability density function:

5. Introduction to Survival Analysis

$$\begin{aligned}
\mathcal{L}(t) &= p(T = t) \\
&\stackrel{\text{def}}{=} f_T(t) \\
&= h_T(t)S_T(t) \\
\ell(t) &\stackrel{\text{def}}{=} \log \{\mathcal{L}(t)\} \\
&= \log \{h_T(t)S_T(t)\} \\
&= \log \{h_T(t)\} + \log \{S_T(t)\} \\
&= \log \{h_T(t)\} - H_T(t)
\end{aligned}$$

If instead the event time T is censored and only known to be after time y , then the likelihood of that censored observation is instead the survival function evaluated at the censoring time:

$$\begin{aligned}
\mathcal{L}(y) &= p_T(T > y) \\
&\stackrel{\text{def}}{=} S_T(y) \\
\ell(y) &\stackrel{\text{def}}{=} \log \{\mathcal{L}(y)\} \\
&= \log \{S(y)\} \\
&= -H(y)
\end{aligned}$$

What's written above is incomplete. We also observed whether or not the observation was censored. Let C denote the time when censoring would occur (if the event did not occur first); let $f_C(y)$ and $S_C(y)$ be the corresponding density and survival functions for the censoring event.

Let Y denote the time when observation ended (either by censoring or by the event of interest occurring), and let D be an indicator variable for the event occurring at Y (so $D = 0$ represents a censored observation and $D = 1$ represents an uncensored observation). In other words, let $Y \stackrel{\text{def}}{=} \min(T, C)$ and $D \stackrel{\text{def}}{=} \mathbb{1}\{T \leq C\}$.

Then the complete likelihood of the observed data (Y, D) is:

5. Introduction to Survival Analysis

$$\begin{aligned}\mathcal{L}(y, d) &= p(Y = y, D = d) \\ &= [p(T = y, C > y)]^d \cdot [p(T > y, C = y)]^{1-d}\end{aligned}$$

Typically, survival analyses assume that C and T are mutually independent; this assumption is called “non-informative” censoring.

Then the joint likelihood $p(Y, D)$ factors into the product $p(Y), p(D)$, and the likelihood reduces to:

$$\begin{aligned}\mathcal{L}(y, d) &= [p(T = y, C > y)]^d \cdot [p(T > y, C = y)]^{1-d} \\ &= [p(T = y)p(C > y)]^d \cdot [p(T > y)p(C = y)]^{1-d} \\ &= [f_T(y)S_C(y)]^d \cdot [S(y)f_C(y)]^{1-d} \\ &= [f_T(y)^d S_C(y)^d] \cdot [S_T(y)^{1-d} f_C(y)^{1-d}] \\ &= (f_T(y)^d \cdot S_T(y)^{1-d}) \cdot (f_C(y)^{1-d} \cdot S_C(y)^d)\end{aligned}$$

The corresponding log-likelihood is:

$$\begin{aligned}\ell(y, d) &= \log \{ \mathcal{L}(y, d) \} \\ &= \log \{ (f_T(y)^d \cdot S_T(y)^{1-d}) \cdot (f_C(y)^{1-d} \cdot S_C(y)^d) \} \\ &= \log \{ f_T(y)^d \cdot S_T(y)^{1-d} \} + \log \{ f_C(y)^{1-d} \cdot S_C(y)^d \}\end{aligned}$$

Let

- θ_T represent the parameters of $p_T(t)$,
- θ_C represent the parameters of $p_C(c)$,
- $\theta = (\theta_T, \theta_C)$ be the combined vector of all parameters.

Then corresponding score function is:

5. Introduction to Survival Analysis

$$\begin{aligned}\ell'(y, d) &= \frac{d}{d\theta} [\log \{f_T(y)^d \cdot S_T(y)^{1-d}\} + \log \{f_C(y)^{1-d} \cdot S_C(y)^d\}] \\ &= \left(\frac{d}{d\theta} \log \{f_T(y)^d \cdot S_T(y)^{1-d}\} \right) + \left(\frac{d}{d\theta} \log \{f_C(y)^{1-d} \cdot S_C(y)^d\} \right)\end{aligned}$$

As long as θ_C and θ_T don't share any parameters, then if censoring is non-informative, the partial derivative with respect to θ_T is:

$$\begin{aligned}\ell'_{\theta_T}(y, d) &\stackrel{\text{def}}{=} \frac{d}{d\theta_T} \ell(y, d) \\ &= \left(\frac{d}{d\theta_T} \log \{f_T(y)^d \cdot S_T(y)^{1-d}\} \right) + \left(\frac{d}{d\theta_T} \log \{f_C(y)^{1-d} \cdot S_C(y)^d\} \right) \\ &= \left(\frac{d}{d\theta_T} \log \{f_T(y)^d \cdot S_T(y)^{1-d}\} \right) + 0 \\ &= \frac{d}{d\theta_T} \log \{f_T(y)^d \cdot S_T(y)^{1-d}\}\end{aligned}$$

Thus, the MLE for θ_T won't depend on θ_C , and we can ignore the distribution of C when estimating the parameters of $f_T(t) = p(T = t)$.

Then:

$$\begin{aligned}\mathcal{L}(y, d) &= f_T(y)^d \cdot S_T(y)^{1-d} \\ &= (h_T(y)^d S_T(y)^d) \cdot S_T(y)^{1-d} \\ &= h_T(y)^d \cdot S_T(y)^d \cdot S_T(y)^{1-d} \\ &= h_T(y)^d \cdot S_T(y) \\ &= S_T(y) \cdot h_T(y)^d\end{aligned}$$

That is, if the event occurred at time y (i.e., if $d = 1$), then the likelihood of $(Y, D) = (y, d)$ is equal to the hazard function at y times the survival

5. Introduction to Survival Analysis

function at y . Otherwise, the likelihood is equal to just the survival function at y .

The corresponding log-likelihood is:

$$\begin{aligned}\ell(y, d) &= \log \{ \mathcal{L}(y, d) \} \\ &= \log \{ S_T(y) \cdot h_T(y)^d \} \\ &= \log \{ S_T(y) \} + \log \{ h_T(y)^d \} \\ &= \log \{ S_T(y) \} + d \cdot \log \{ h_T(y) \} \\ &= -H_T(y) + d \cdot \log \{ h_T(y) \}\end{aligned}$$

In other words, the log-likelihood contribution from a single observation $(Y, D) = (y, d)$ is equal to the negative cumulative hazard at y , plus the log of the hazard at y if the event occurred at time y .

i Note

End of extra section.

5.4. Parametric Models for Time-to-Event Outcomes

5.4.1. Exponential Distribution

- The exponential distribution is the base distribution for survival analysis.
- The distribution has a constant hazard λ
- The mean survival time is λ^{-1}

5.4.1.1. Mathematical details of exponential distribution

$$\begin{aligned}
 f(t) &= \lambda e^{-\lambda t} \\
 E(t) &= \lambda^{-1} \\
 Var(t) &= \lambda^{-2} \\
 F(t) &= 1 - e^{-\lambda x} \\
 S(t) &= e^{-\lambda x} \\
 \ln(S(t)) &= -\lambda x \\
 h(t) &= -\frac{f(t)}{S(t)} = -\frac{\lambda e^{-\lambda t}}{e^{-\lambda t}} = \lambda
 \end{aligned}$$

5.4.1.2. Estimation of λ

- Suppose we have m exponential survival times of t_1, t_2, \dots, t_m and k right-censored values at u_1, u_2, \dots, u_k .
- A survival time of $t_i = 10$ means that subject i died at time 10. A right-censored time $u_i = 10$ means that at time 10, subject i was still alive and that we have no further follow-up.
- For the moment we will assume that the survival distribution is exponential and that all the subjects have the same parameter λ .

We have m exponential survival times of t_1, t_2, \dots, t_m and k right-censored values at u_1, u_2, \dots, u_k . The log-likelihood of an observed survival time t_i is

$$\log \{ \lambda e^{-\lambda t_i} \} = \log \{ \lambda \} - \lambda t_i$$

and the likelihood of a censored value is the probability of that outcome (survival greater than u_j) so the log-likelihood is

$$\log \{ \lambda e^{-\lambda u_j} \} = -\lambda u_j.$$

Let $T = \sum t_i$ and $U = \sum u_j$. Then:

5. Introduction to Survival Analysis

$$\begin{aligned}
 \ell(\lambda) &= \sum_{i=1}^m (\ln \lambda - \lambda t_i) + \sum_{j=1}^k (-\lambda u_j) \\
 &= m \ln \lambda - (T + U)\lambda \\
 \ell'(\lambda) &= m\lambda^{-1} - (T + U) \\
 \hat{\lambda} &= \frac{m}{T + U} \\
 \ell'' &= -m/\lambda^2 \\
 &< 0 \\
 \hat{E}[T] &= \hat{\lambda}^{-1} \\
 &= \frac{T + U}{m}
 \end{aligned}$$

5.4.1.3. Fisher Information and Standard Error

$$\begin{aligned}
 E[-\ell''] &= m/\lambda^2 \\
 \text{Var}(\hat{\lambda}) &\approx (E[-\ell''])^{-1} \\
 &= \lambda^2/m \\
 \text{SE}(\hat{\lambda}) &= \sqrt{\text{Var}(\hat{\lambda})} \\
 &\approx \lambda/\sqrt{m}
 \end{aligned}$$

$\hat{\lambda}$ depends on the censoring times of the censored observations, but $\text{Var}(\hat{\lambda})$ only depends on the number of uncensored observations, m , and not on the number of censored observations (k).

5.4.1.4. Other Parametric Survival Distributions

- Any density on $[0, \infty)$ can be a survival distribution, but the most useful ones are all skew right.

5. Introduction to Survival Analysis

- The commonest generalization of the exponential is the Weibull.
- Other common choices are the gamma, log-normal, log-logistic, Gompertz, inverse Gaussian, and Pareto.
- Most of what we do going forward is non-parametric or semi-parametric, but sometimes these parametric distributions provide a useful approach.

5.4.2. Weibull Distribution

$$p(t) = \alpha \lambda x^{\alpha-1} e^{-\lambda x^\alpha}$$

$$h(t) = \alpha \lambda x^{\alpha-1}$$

$$S(t) = e^{-\lambda x^\alpha}$$

$$E(T) = \Gamma(1 + 1/\alpha) \cdot \lambda^{-1/\alpha}$$

When $\alpha = 1$ this is the exponential. When $\alpha > 1$ the hazard is increasing and when $\alpha < 1$ the hazard is decreasing. This provides more flexibility than the exponential.

We will see more of this distribution later.

5.5. Nonparametric Survival Analysis

5.5.1. Basic ideas

- Mostly, we work without a parametric model.
- The first task is to estimate a survival function from data listing survival times, and censoring times for censored data.
- For example one patient may have relapsed at 10 months. Another might have been followed for 32 months without a relapse having occurred (censored).

5. Introduction to Survival Analysis

- The minimum information we need for each patient is a time and a censoring variable which is 1 if the event occurred at the indicated time and 0 if this is a censoring time.

5.6. Example: clinical trial for pediatric acute leukemia

5.6.1. Overview of study

This is from a clinical trial in 1963 for 6-MP treatment vs. placebo for Acute Leukemia in 42 children.

- Pairs of children:
 - matched by remission status at the time of treatment (`remstat`: 1 = partial, 2 = complete)
 - randomized to 6-MP (exit times in `t2`) or placebo (exit times in `t1`)
- Followed until relapse or end of study.
- All of the placebo group relapsed, but some of the 6-MP group were censored (which means they were still in remission); indicated by `relapse` variable (0 = censored, 1 = relapse).
- 6-MP = 6-Mercaptopurine (Purinethol) is an anti-cancer (“antineoplastic” or “cytotoxic”) chemotherapy drug used currently for Acute lymphoblastic leukemia (ALL). It is classified as an antimetabolite.

5.6.2. Study design

Clinical trial in 1963 for 6-MP treatment vs. placebo for Acute Leukemia in 42 children. Pairs of children matched by remission status at the time of treatment (1 = partial or 2 = complete) and randomized to 6-MP or placebo. Followed until relapse or end of study. All of the placebo group relapsed, but some of the 6-MP group were censored.

```
library(KMsurv)
data(drug6mp)
drug6mp |> tibble() |> print()
```

```
# A tibble: 21 x 5
   pair remstat   t1   t2 relapse
   <int>   <int> <int> <int>   <int>
1     1     1     1     1     10     1
2     2     2     2    22     7     1
3     3     2     3    32     0
4     4     2    12    23     1
5     5     2     8    22     1
6     6     1    17     6     1
7     7     2     2    16     1
8     8     2    11    34     0
9     9     2     8    32     0
10    10     2    12    25     0
# i 11 more rows
```

5.6.3. Data documentation for drug6mp

```
library(printr) # inserts help-file output into markdown output
library(KMsurv)
?drug6mp
```

5. Introduction to Survival Analysis

data from Section 1.2

Description:

The 'drug6mp' data frame has 21 rows and 5 columns.

Format:

This data frame contains the following columns:

pair pair number

remstat Remission status at randomization (1=partial, 2=complete)

t1 Time to relapse for placebo patients, months

t2 Time to relapse for 6-MP patients, months

relapse Relapse indicator (0=censored, 1=relapse) for 6-MP patients

5.6.4. Descriptive Statistics

- The average time in each group is not useful. Some of the 6-MP patients have not relapsed at the time recorded, while all of the placebo patients have relapsed.
- The median time is not really useful either because so many of the 6-MP patients have not relapsed (12/21).
- Both are biased down in the 6-MP group. Remember that lower times are worse since they indicate sooner recurrence.
- We can compute the average hazard rate, which is the estimate of the exponential parameter: number of relapses divided by the sum of the times.

5. Introduction to Survival Analysis

- For the placebo, that is just the reciprocal of the mean time = $1/8.667 = 0.115$.
- For the 6-MP group this is $9/359 = 0.025$
- The estimated average hazard in the placebo group is 4.6 times as large (if the hazard is constant over time).

5.7. The Kaplan-Meier Product Limit Estimator

- The estimated survival function for the placebo patients is easy to compute. For any time t in months, $S(t)$ is the fraction of patients with times greater than t .
- For the 6-MP patients, we cannot ignore the censored data because we know that the time to relapse is greater than the censoring time.
- For any time t in months, we know that 6-MP patients with times greater than t have not relapsed, and those with relapse time less than t have relapsed, but we don't know if patients with censored time less than t have relapsed or not.
- The procedure we usually use is the Kaplan-Meier product-limit estimator of the survival function.
- The Kaplan-Meier estimator is a step function (like the empirical cdf), which changes value only at the event times, not at the censoring times.
- At each event time t , we compute the at-risk group size Y , which is all those observations whose event time or censoring time is at least t .
- If d of the observations have an event time (not a censoring time) of t , then the group of survivors immediately following time t is reduced by the fraction

$$\frac{Y-d}{Y} = 1 - \frac{d}{Y}$$

5. Introduction to Survival Analysis

If the event times are t_i with events per time of d_i ($1 \leq i \leq k$), then

$$\hat{S}(t) = \prod_{t_i < t} [1 - d_i/Y_i]$$

where Y_i is the set of observations whose time (event or censored) is $\geq t_i$, the group at risk at time t_i .

If there are no censored data, and there are n data points, then just after (say) the third event time

$$\begin{aligned}\hat{S}(t) &= \prod_{t_i < t} [1 - d_i/Y_i] \\ &= \left[\frac{n - d_1}{n}\right] \left[\frac{n - d_1 - d_2}{n - d_1}\right] \left[\frac{n - d_1 - d_2 - d_3}{n - d_1 - d_2}\right] \\ &= \frac{n - d_1 - d_2 - d_3}{n} \\ &= 1 - \frac{d_1 + d_2 + d_3}{n} \\ &= 1 - \hat{F}(t)\end{aligned}$$

where $\hat{F}(t)$ is the usual empirical CDF estimate.

5.7.1. Kaplan-Meier curve for drug6mp data

Here is the Kaplan-Meier estimated survival curve for the patients who received 6-MP in the `drug6mp` dataset (we will see code to produce figures like this one shortly):

5. Introduction to Survival Analysis




Figure 5.5.: Kaplan-Meier Survival Curve for 6-MP Patients

5.7.2. Kaplan-Meier calculations

Let's compute these estimates and build the chart by hand:

```
library(KMsurv)
library(dplyr)
data(drug6mp)

drug6mp.v2 =
  drug6mp |>
  as_tibble() |>
  mutate(
    remstat = remstat |>
```

5. Introduction to Survival Analysis

```
      case_match(
        1 ~ "partial",
        2 ~ "complete"
      ),
      # renaming to "outcome" while relabeling is just a style choice:
      outcome = relapse |>
        case_match(
          0 ~ "censored",
          1 ~ "relapsed"
        )
    )

km.6mp =
  drug6mp.v2 |>
  summarize(
    .by = t2,
    Relapses = sum(outcome == "relapsed"),
    Censored = sum(outcome == "censored")) |>
  # here we add a start time row, so the graph starts at time 0:
  bind_rows(
    tibble(
      t2 = 0,
      Relapses = 0,
      Censored = 0)
  ) |>
  # sort in time order:
  arrange(t2) |>
  mutate(
    Exiting = Relapses + Censored,
    `Study Size` = sum(Exiting),
    Exited = cumsum(Exiting) |> dplyr::lag(default = 0),
    `At Risk` = `Study Size` - Exited,
    Hazard = Relapses / `At Risk`,
```


5. Introduction to Survival Analysis

```

`KM Factor` = 1 - Hazard,
`Cumulative Hazard` = cumsum(`Hazard`),
`KM Survival Curve` = cumprod(`KM Factor`)
)

library(pander)
pander(km.6mp)

```

| t2 | Relapse | Censored | Editing | Study Size | At Exit | Risk | Hazard | KM Factor | Cumulative Hazard | KM Survival Curve |
|----|---------|----------|---------|------------|---------|------|--------|-----------|-------------------|-------------------|
| 0 | 0 | 0 | 0 | 21 | 0 | 21 | 0 | 1 | 0 | 1 |
| 6 | 3 | 1 | 4 | 21 | 0 | 21 | 0.1429 | 0.8571 | 0.1429 | 0.8571 |
| 7 | 1 | 0 | 1 | 21 | 4 | 17 | 0.0588 | 0.9412 | 0.2017 | 0.8067 |
| 9 | 0 | 1 | 1 | 21 | 5 | 16 | 0 | 1 | 0.2017 | 0.8067 |
| 10 | 1 | 1 | 2 | 21 | 6 | 15 | 0.0667 | 0.9333 | 0.2683 | 0.7529 |
| 11 | 0 | 1 | 1 | 21 | 8 | 13 | 0 | 1 | 0.2683 | 0.7529 |
| 13 | 1 | 0 | 1 | 21 | 9 | 12 | 0.0833 | 0.9167 | 0.3517 | 0.6902 |
| 16 | 1 | 0 | 1 | 21 | 10 | 11 | 0.0909 | 0.9091 | 0.4426 | 0.6275 |
| 17 | 0 | 1 | 1 | 21 | 11 | 10 | 0 | 1 | 0.4426 | 0.6275 |
| 19 | 0 | 1 | 1 | 21 | 12 | 9 | 0 | 1 | 0.4426 | 0.6275 |
| 20 | 0 | 1 | 1 | 21 | 13 | 8 | 0 | 1 | 0.4426 | 0.6275 |
| 22 | 1 | 0 | 1 | 21 | 14 | 7 | 0.1429 | 0.8571 | 0.5854 | 0.5378 |
| 23 | 1 | 0 | 1 | 21 | 15 | 6 | 0.1667 | 0.8333 | 0.7521 | 0.4482 |
| 25 | 0 | 1 | 1 | 21 | 16 | 5 | 0 | 1 | 0.7521 | 0.4482 |
| 32 | 0 | 2 | 2 | 21 | 17 | 4 | 0 | 1 | 0.7521 | 0.4482 |
| 34 | 0 | 1 | 1 | 21 | 19 | 2 | 0 | 1 | 0.7521 | 0.4482 |
| 35 | 0 | 1 | 1 | 21 | 20 | 1 | 0 | 1 | 0.7521 | 0.4482 |

5. Introduction to Survival Analysis

5.7.2.1. Summary

For the 6-MP patients at time 6 months, there are 21 patients at risk. At $t = 6$ there are 3 relapses and 1 censored observations.


The Kaplan-Meier factor is $(21 - 3)/21 = 0.857$. The number at risk for the next time ($t = 7$) is $21 - 3 - 1 = 17$.

At time 7 months, there are 17 patients at risk. At $t = 7$ there is 1 relapse and 0 censored observations. The Kaplan-Meier factor is $(17 - 1)/17 = 0.941$. The Kaplan Meier estimate is $0.857 \times 0.941 = 0.807$. The number at risk for the next time ($t = 9$) is $17 - 1 = 16$.

Now, let's graph this estimated survival curve using `ggplot()`:

```
library(ggplot2)
conflicts_prefer(dplyr::filter)
km.6mp |>
  ggplot(aes(x = t2, y = `KM Survival Curve`)) +
  geom_step() +
  geom_point(data = km.6mp |> filter(Censored > 0), shape = 3) +
  expand_limits(y = c(0,1), x = 0) +
  xlab('Time since diagnosis (months)') +
  ylab("KM Survival Curve") +
  scale_y_continuous(labels = scales::percent)
```

5. Introduction to Survival Analysis



5.8. Using the survival package in R

We don't have to do these calculations by hand every time; the `survival` package and several others have functions available to automate many of these tasks (full list: <https://cran.r-project.org/web/views/Survival.html>).

5.8.1. The `Surv` function

To use the `survival` package, the first step is telling R how to combine the exit time and exit reason (censoring versus event) columns. The `Surv()` function accomplishes this task.

5.8.1.1. Example: `Surv()` with `drug6mp` data

```

1 library(survival)
2 drug6mp.v3 =
3   drug6mp.v2 |>
4   mutate(
5     surv2 = Surv(
6       time = t2,
7       event = (outcome == "relapsed")))
8
9 print(drug6mp.v3)

```

```

# A tibble: 21 x 7
   pair remstat    t1    t2 relapse outcome  surv2
   <int> <chr>   <int> <int>   <int> <chr>   <Surv>
1     1 1 partial     1    10     1 relapsed    10
2     2 2 complete   22     7     1 relapsed     7
3     3 3 complete    3    32     0 censored   32+
4     4 4 complete   12    23     1 relapsed    23
5     5 5 complete    8    22     1 relapsed    22
6     6 6 partial   17     6     1 relapsed     6
7     7 7 complete    2    16     1 relapsed    16
8     8 8 complete   11    34     0 censored   34+
9     9 9 complete    8    32     0 censored   32+
10    10 10 complete   12    25     0 censored   25+
# i 11 more rows

```

The output of `Surv()` is a vector of objects with class `Surv`. When we print this vector:

- observations where the event was observed are printed as the event time (for example, `surv2 = 10` on line 1)

- observations where the event was right-censored are printed as the censoring time with a plus sign (+; for example, `surv2 = 32+` on line 3).

5.8.2. The `survfit` function

Once we have constructed our `Surv` variable, we can calculate the Kaplan-Meier estimate of the survival curve using the `survfit()` function.

i Note

The documentation for `?survfit` isn't too helpful; the `survfit.formula` documentation is better.

5.8.2.1. Example: `survfit()` with `drug6mp` data

Here we use `survfit()` to create a `survfit` object, which contains the Kaplan-Meier estimate:

```
drug6mp.km_model = survfit(  
  formula = surv2 ~ 1,  
  data = drug6mp.v3)
```

`print.survfit()` just gives some summary statistics:

```
print(drug6mp.km_model)
```

```
Call: survfit(formula = surv2 ~ 1, data = drug6mp.v3)
```

```
      n events median 0.95LCL 0.95UCL  
[1,] 21      9     23     16     NA
```

5. Introduction to Survival Analysis

`summary.survfit()` shows us the underlying Kaplan-Meier table:

```
summary(drug6mp.km_model)
```

```
Call: survfit(formula = surv2 ~ 1, data = drug6mp.v3)
```

| time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
|------|--------|---------|----------|---------|--------------|--------------|
| 6 | 21 | 3 | 0.857 | 0.0764 | 0.720 | 1.000 |
| 7 | 17 | 1 | 0.807 | 0.0869 | 0.653 | 0.996 |
| 10 | 15 | 1 | 0.753 | 0.0963 | 0.586 | 0.968 |
| 13 | 12 | 1 | 0.690 | 0.1068 | 0.510 | 0.935 |
| 16 | 11 | 1 | 0.627 | 0.1141 | 0.439 | 0.896 |
| 22 | 7 | 1 | 0.538 | 0.1282 | 0.337 | 0.858 |
| 23 | 6 | 1 | 0.448 | 0.1346 | 0.249 | 0.807 |

`summary.survfit()` shows us the underlying Kaplan-Meier table:

```
summary(drug6mp.km_model)
```

```
Call: survfit(formula = surv2 ~ 1, data = drug6mp.v3)
```

| time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
|------|--------|---------|----------|---------|--------------|--------------|
| 6 | 21 | 3 | 0.857 | 0.0764 | 0.720 | 1.000 |
| 7 | 17 | 1 | 0.807 | 0.0869 | 0.653 | 0.996 |
| 10 | 15 | 1 | 0.753 | 0.0963 | 0.586 | 0.968 |
| 13 | 12 | 1 | 0.690 | 0.1068 | 0.510 | 0.935 |
| 16 | 11 | 1 | 0.627 | 0.1141 | 0.439 | 0.896 |
| 22 | 7 | 1 | 0.538 | 0.1282 | 0.337 | 0.858 |
| 23 | 6 | 1 | 0.448 | 0.1346 | 0.249 | 0.807 |

We can specify which time points we want using the `times` argument:

5. Introduction to Survival Analysis

```
summary(  
  drug6mp.km_model,  
  times = c(0, drug6mp.v3$t2))
```

Call: `survfit(formula = surv2 ~ 1, data = drug6mp.v3)`

| time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
|------|--------|---------|----------|---------|--------------|--------------|
| 0 | 21 | 0 | 1.000 | 0.0000 | 1.000 | 1.000 |
| 6 | 21 | 3 | 0.857 | 0.0764 | 0.720 | 1.000 |
| 6 | 21 | 0 | 0.857 | 0.0764 | 0.720 | 1.000 |
| 6 | 21 | 0 | 0.857 | 0.0764 | 0.720 | 1.000 |
| 6 | 21 | 0 | 0.857 | 0.0764 | 0.720 | 1.000 |
| 7 | 17 | 1 | 0.807 | 0.0869 | 0.653 | 0.996 |
| 9 | 16 | 0 | 0.807 | 0.0869 | 0.653 | 0.996 |
| 10 | 15 | 1 | 0.753 | 0.0963 | 0.586 | 0.968 |
| 10 | 15 | 0 | 0.753 | 0.0963 | 0.586 | 0.968 |
| 11 | 13 | 0 | 0.753 | 0.0963 | 0.586 | 0.968 |
| 13 | 12 | 1 | 0.690 | 0.1068 | 0.510 | 0.935 |
| 16 | 11 | 1 | 0.627 | 0.1141 | 0.439 | 0.896 |
| 17 | 10 | 0 | 0.627 | 0.1141 | 0.439 | 0.896 |
| 19 | 9 | 0 | 0.627 | 0.1141 | 0.439 | 0.896 |
| 20 | 8 | 0 | 0.627 | 0.1141 | 0.439 | 0.896 |
| 22 | 7 | 1 | 0.538 | 0.1282 | 0.337 | 0.858 |
| 23 | 6 | 1 | 0.448 | 0.1346 | 0.249 | 0.807 |
| 25 | 5 | 0 | 0.448 | 0.1346 | 0.249 | 0.807 |
| 32 | 4 | 0 | 0.448 | 0.1346 | 0.249 | 0.807 |
| 32 | 4 | 0 | 0.448 | 0.1346 | 0.249 | 0.807 |
| 34 | 2 | 0 | 0.448 | 0.1346 | 0.249 | 0.807 |
| 35 | 1 | 0 | 0.448 | 0.1346 | 0.249 | 0.807 |

```
?summary.survfit
```

Summary of a Survival Curve

5. Introduction to Survival Analysis

Description:

Returns a list containing the survival curve, confidence limits for the curve, and other information.

Usage:

```
## S3 method for class 'survfit'
summary(object, times, censored=FALSE, scale=1,
        extend=FALSE, rmean=getOption('survfit.rmean'), ...)
```

Arguments:

object: the result of a call to the 'survfit' function.

times: vector of times; the returned matrix will contain 1 row for each time. The vector will be sorted into increasing order; missing values are not allowed. If 'censored=T', the default 'times' vector contains all the unique times in 'fit', otherwise the default 'times' vector uses only the event (death) times.

censored: logical value: should the censoring times be included in the output? This is ignored if the 'times' argument is present.

scale: numeric value to rescale the survival time, e.g., if the input data to 'survfit' were in days, 'scale = 365.25' would scale the output to years.

extend: logical value: if TRUE, prints information for all specified 'times', even if there are no subjects left at the end of the specified 'times'. This is only used if the 'times' argument is present.

5. Introduction to Survival Analysis

rmean: Show restricted mean: see 'print.survfit' for details

...: for future methods

5.8.3. Plotting estimated survival functions

We can plot `survfit` objects with `plot()`, `autoplot()`, or `ggsurvplot()`:

```
library(ggfortify)
autoplot(drug6mp.km_model)
```




Figure 5.6.: Kaplan-Meier Survival Curve for 6-MP Patients

5. Introduction to Survival Analysis

```
# not shown:  
# plot(drug6mp.km_model)  
  
# library(survminer)  
# ggsurvplot(drug6mp.km_model)
```

5.8.3.1. quantiles of survival curve

We can extract quantiles with `quantile()`:

```
1 drug6mp.km_model |>  
2   quantile(p = c(.25, .5)) |>  
3   as_tibble() |>  
4   mutate(p = c(.25, .5)) |>  
5   relocate(p, .before = everything())
```

| p | quantile | lower | upper |
|------|----------|-------|-------|
| 0.25 | 13 | 6 | NA |
| 0.50 | 23 | 16 | NA |

5.8.4. Two-sample tests

5.8.4.1. The survdiff function

```
?survdiff
```

Test Survival Curve Differences

5. Introduction to Survival Analysis

Description:

Tests if there is a difference between two or more survival curves using the G-rho family of tests, or for a single curve against a known alternative.

Usage:

```
survdiff(formula, data, subset, na.action, rho=0, timefix=TRUE)
```

5.8.4.2. Example: `survdiff()` with drug6mp data

Now we are going to compare the placebo and 6-MP data. We need to reshape the data to make it usable with the standard `survival` workflow:

```
library(survival)

drug6mp.v4 =
  drug6mp.v3 |>
  select(pair, remstat, t1, t2, outcome) |>
  # here we are going to change the data from a wide format to long:
  pivot_longer(
    cols = c(t1, t2),
    names_to = "treatment",
    values_to = "exit_time") |>
  mutate(
    treatment = treatment |>
      case_match(
        "t1" ~ "placebo",
        "t2" ~ "6-MP"
      ),
    outcome = if_else(
      treatment == "placebo",
```

5. Introduction to Survival Analysis

```
    "relapsed",  
    outcome  
  ),  
  surv = Surv(  
    time = exit_time,  
    event = (outcome == "relapsed"))  
)
```


Using this long data format, we can fit a Kaplan-Meier curve for each treatment group simultaneously:

```
drug6mp.km_model2 =  
  survfit(  
    formula = surv ~ treatment,  
    data = drug6mp.v4)
```

We can plot the curves in the same graph:

```
drug6mp.km_model2 |> autoplot()
```

5. Introduction to Survival Analysis



We can also perform something like a t-test, where the null hypothesis is that the curves are the same:

```
survdif(  
  formula = surv ~ treatment,  
  data = drug6mp.v4)
```

Call:

```
survdif(formula = surv ~ treatment, data = drug6mp.v4)
```

| | N | Observed | Expected | (O-E) ² /E | (O-E) ² /V |
|-------------------|----|----------|----------|-----------------------|-----------------------|
| treatment=6-MP | 21 | 9 | 19.3 | 5.46 | 16.8 |
| treatment=placebo | 21 | 21 | 10.7 | 9.77 | 16.8 |

Chisq= 16.8 on 1 degrees of freedom, p= 4e-05

5. Introduction to Survival Analysis

By default, `survdif()` ignores any pairing, but we can use `strata()` to perform something similar to a paired t-test:

```
survdif(  
  formula = surv ~ treatment + strata(pair),  
  data = drug6mp.v4)
```

Call:

```
survdif(formula = surv ~ treatment + strata(pair), data = drug6mp.v4)
```

| | N | Observed | Expected | (O-E) ² /E | (O-E) ² /V |
|-------------------|----|----------|----------|-----------------------|-----------------------|
| treatment=6-MP | 21 | 9 | 16.5 | 3.41 | 10.7 |
| treatment=placebo | 21 | 21 | 13.5 | 4.17 | 10.7 |

Chisq= 10.7 on 1 degrees of freedom, p= 0.001

Interestingly, accounting for pairing reduces the significance of the difference.

5.9. Example: Bone Marrow Transplant Data

(Copelan et al., 1991)

* Treatment

- allogeneic (from a donor) bone marrow transplant therapy

* Inclusion criteria

- acute myeloid leukemia (AML)
- acute lymphoblastic leukemia (ALL).

5. Introduction to Survival Analysis

* Possible intermediate events

- **graft vs. host disease (GVHD)**: an immunological rejection response to the transplant
- **platelet recovery**: a return of platelet count to normal levels.

One or the other, both in either order, or neither may occur.

End point events

- relapse of the disease
- death

Any or all of these events may be censored.

5.9.1. `KMsurv::bmt` data in R

```
library(printr) # inserts help-file output into markdown output
library(KMsurv)
?bmt
```

data from Section 1.3

Description:

The 'bmt' data frame has 137 rows and 22 columns.

Format:

This data frame contains the following columns:

group Disease Group 1-ALL, 2-AML Low Risk, 3-AML High Risk

5. Introduction to Survival Analysis

t1 Time To Death Or On Study Time

t2 Disease Free Survival Time (Time To Relapse, Death Or End Of Study)

d1 Death Indicator 1-Dead 0-Alive

d2 Relapse Indicator 1-Relapsed, 0-Disease Free

d3 Disease Free Survival Indicator 1-Dead Or Relapsed, 0-Alive
Disease Free)

ta Time To Acute Graft-Versus-Host Disease

da Acute GVHD Indicator 1-Developed Acute GVHD 0-Never Developed
Acute GVHD)

tc Time To Chronic Graft-Versus-Host Disease

dc Chronic GVHD Indicator 1-Developed Chronic GVHD 0-Never
Developed Chronic GVHD

tp Time To Chronic Graft-Versus-Host Disease

dp Platelet Recovery Indicator 1-Platelets Returned To Normal,
0-Platelets Never Returned to Normal

z1 Patient Age In Years

z2 Donor Age In Years

z3 Patient Sex: 1-Male, 0-Female

5. Introduction to Survival Analysis

z4 Donor Sex: 1-Male, 0-Female

z5 Patient CMV Status: 1-CMV Positive, 0-CMV Negative

z6 Donor CMV Status: 1-CMV Positive, 0-CMV Negative

z7 Waiting Time to Transplant In Days

z8 FAB: 1-FAB Grade 4 Or 5 and AML, 0-Otherwise

z9 Hospital: 1-The Ohio State University, 2-Alfred , 3-St.
Vincent, 4-Hahnemann

z10 MTX Used as a Graft-Versus-Host- Prophylactic: 1-Yes 0-No

Source:

Klein and Moeschberger (1997) *_Survival Analysis Techniques for
Censored and truncated data_*, Springer.

Examples:

```
data(bmt)
```

5.9.2. Analysis plan

- We concentrate for now on disease-free survival (t2 and d3) for the three risk groups, ALL, AML Low Risk, and AML High Risk.
- We will construct the Kaplan-Meier survival curves, compare them, and test for differences.
- We will construct the cumulative hazard curves and compare them.
- We will estimate the hazard functions, interpret, and compare them.

5.9.3. Survival Function Estimate and Variance

$$\hat{S}(t) = \prod_{t_i < t} \left[1 - \frac{d_i}{Y_i} \right]$$

where Y_i is the group at risk at time t_i .

The estimated variance of $\hat{S}(t)$ is (Greenwood's formula)

$$\hat{\text{Var}}(\hat{S}(t)) = \hat{S}(t)^2 \sum_{t_i < t} \frac{d_i}{Y_i(Y_i - d_i)}$$

which we can use for confidence intervals for a survival function or a difference of survival functions.

Kaplan-Meier survival curves

```
library(KMsurv)
library(survival)
data(bmt)

bmt =
  bmt |>
  as_tibble() |>
  mutate(
    group =
      group |>
      factor(
        labels = c("ALL", "Low Risk AML", "High Risk AML"),
        surv = Surv(t2, d3))

km_model1 = survfit(
  formula = surv ~ group,
  data = bmt)
```

5. Introduction to Survival Analysis

```
library(ggfortify)
autoplot(
  km_model1,
  conf.int = TRUE,
  ylab = "Pr(disease-free survival)",
  xlab = "Time since transplant (days)" +
  theme_bw() +
  theme(legend.position="bottom")
)
```




Figure 5.7.: Disease-Free Survival by Disease Group

5.9.3.1. Understanding Greenwood's formula (optional)

To see where Greenwood's formula comes from, let $x_i = Y_i - d_i$. We approximate the solution treating each time as independent, with Y_i fixed

5. Introduction to Survival Analysis

and ignore randomness in times of failure and we treat x_i as independent binomials $\text{Bin}(Y_i, p_i)$. Letting $S(t)$ be the “true” survival function

$$\hat{S}(t) = \prod_{t_i < t} x_i / Y_i$$

$$S(t) = \prod_{t_i < t} p_i$$

$$\begin{aligned} \frac{\hat{S}(t)}{S(t)} &= \prod_{t_i < t} \frac{x_i}{p_i Y_i} = \prod_{t_i < t} \frac{\hat{p}_i}{p_i} \\ &= \prod_{t_i < t} \left(1 + \frac{\hat{p}_i - p_i}{p_i} \right) \\ &\approx 1 + \sum_{t_i < t} \frac{\hat{p}_i - p_i}{p_i} \end{aligned}$$

$$\begin{aligned} \text{Var} \left(\frac{\hat{S}(t)}{S(t)} \right) &\approx \text{Var} \left(1 + \sum_{t_i < t} \frac{\hat{p}_i - p_i}{p_i} \right) \\ &= \sum_{t_i < t} \frac{1}{p_i^2} \frac{p_i(1-p_i)}{Y_i} \\ &= \sum_{t_i < t} \frac{(1-p_i)}{p_i Y_i} \approx \sum_{t_i < t} \frac{(1-x_i/Y_i)}{x_i} \\ &= \sum_{t_i < t} \frac{Y_i - x_i}{x_i Y_i} = \sum_{t_i < t} \frac{d_i}{Y_i(Y_i - d_i)} \\ \text{Var}(\hat{S}(t)) &\approx \hat{S}(t)^2 \sum_{t_i < t} \frac{d_i}{Y_i(Y_i - d_i)} \end{aligned}$$

5.9.4. Test for differences among the disease groups

Here we compute a chi-square test for association between disease group (group) and disease-free survival:

```
survdif(surv ~ group, data = bmt)
```

Call:

```
survdif(formula = surv ~ group, data = bmt)
```

| | N | Observed | Expected | (O-E)^2/E | (O-E)^2/V |
|---------------------|----|----------|----------|-----------|-----------|
| group=ALL | 38 | 24 | 21.9 | 0.211 | 0.289 |
| group=Low Risk AML | 54 | 25 | 40.0 | 5.604 | 11.012 |
| group=High Risk AML | 45 | 34 | 21.2 | 7.756 | 10.529 |

Chisq= 13.8 on 2 degrees of freedom, p= 0.001

5.9.5. Cumulative Hazard

$$\begin{aligned}
 h(t) &\stackrel{\text{def}}{=} P(T = t | T \geq t) \\
 &= \frac{p(T = t)}{P(T \geq t)} \\
 &= -\frac{d}{dt} \log \{S(t)\}
 \end{aligned}$$

The **cumulative hazard** (or **integrated hazard**) function is

$$H(t) \stackrel{\text{def}}{=} \int_0^t h(t) dt$$

Since $h(t) = -\frac{d}{dt} \log \{S(t)\}$ as shown above, we have:

5. Introduction to Survival Analysis

$$H(t) = -\log \{S\}(t)$$

So we can estimate $H(t)$ as:

$$\begin{aligned}\hat{H}(t) &= -\log \{ \hat{S}(t) \} \\ &= -\log \left\{ \prod_{t_i < t} \left[1 - \frac{d_i}{Y_i} \right] \right\} \\ &= -\sum_{t_i < t} \log \left\{ 1 - \frac{d_i}{Y_i} \right\}\end{aligned}$$

This is the **Kaplan-Meier (product-limit) estimate of cumulative hazard**.

5.9.5.1. Example: Cumulative Hazard Curves for Bone-Marrow Transplant (bmt) data

```
autoplot(  
  fun = "cumhaz",  
  km_model1,  
  conf.int = FALSE,  
  ylab = "Cumulative hazard (disease-free survival)",  
  xlab = "Time since transplant (days)" +  
  theme_bw() +  
  theme(legend.position="bottom")
```




Figure 5.8.: Disease-Free Cumulative Hazard by Disease Group

5.10. Nelson-Aalen Estimates of Cumulative Hazard and Survival

The point hazard at time t_i can be estimated by d_i/Y_i , which leads to the Nelson-Aalen estimator of the cumulative hazard:

$$\hat{H}_{NA}(t) \stackrel{\text{def}}{=} \sum_{t_i < t} \frac{d_i}{Y_i}$$

5. Introduction to Survival Analysis

The variance of this estimator is approximately:

$$\begin{aligned}\hat{\text{Var}}\left(\hat{H}_{NA}(t)\right) &= \sum_{t_i < t} \frac{(d_i/Y_i)(1 - d_i/Y_i)}{Y_i} \\ &\approx \sum_{t_i < t} \frac{d_i}{Y_i^2}\end{aligned}$$

Since $S(t) = \exp\{-H(t)\}$, the Nelson-Aalen cumulative hazard estimate can be converted into an alternate estimate of the survival function:

$$\begin{aligned}\hat{S}_{NA}(t) &= \exp\left\{-\hat{H}_{NA}(t)\right\} \\ &= \exp\left\{-\sum_{t_i < t} \frac{d_i}{Y_i}\right\} \\ &= \prod_{t_i < t} \exp\left\{-\frac{d_i}{Y_i}\right\}\end{aligned}$$

Compare these with the corresponding Kaplan-Meier estimates:

$$\begin{aligned}\hat{H}_{KM}(t) &= -\sum_{t_i < t} \log\left\{1 - \frac{d_i}{Y_i}\right\} \\ \hat{S}_{KM}(t) &= \prod_{t_i < t} \left[1 - \frac{d_i}{Y_i}\right]\end{aligned}$$

The product limit estimate and the Nelson-Aalen estimate often do not differ by much. The latter is considered more accurate in small samples and also directly estimates the cumulative hazard. The "fleming-harrington" method for `survfit()` reduces to Nelson-Aalen when the data are unweighted. We can also estimate the cumulative hazard as the negative log of the KM survival function estimate.

5.10.1. Application to bmt dataset

```
na_fit = survfit(  
  formula = surv ~ group,  
  type = "fleming-harrington",  
  data = bmt)  
  
km_fit = survfit(  
  formula = surv ~ group,  
  type = "kaplan-meier",  
  data = bmt)  
  
km_and_na =  
  bind_rows(  
    .id = "model",  
    "Kaplan-Meier" = km_fit |> fortify(surv.connect = TRUE),  
    "Nelson-Aalen" = na_fit |> fortify(surv.connect = TRUE)  
  ) |>  
  as_tibble()
```

```
km_and_na |>  
  ggplot(aes(x = time, y = surv, col = model)) +  
  geom_step() +  
  facet_grid(. ~ strata) +  
  theme_bw() +  
  ylab("S(t) = P(T>=t)") +  
  xlab("Survival time (t, days)") +  
  theme(legend.position = "bottom")
```

5. Introduction to Survival Analysis




Figure 5.9.: Kaplan-Meier and Nelson-Aalen Survival Function Estimates, stratified by disease group

The Kaplan-Meier and Nelson-Aalen survival estimates are very similar for this dataset.

6. Proportional Hazards Models

Configuring R

Functions from these packages will be used throughout this document:

```
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(dplyr) # manipulate data
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
conflicts_prefer(dplyr::filter)
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
pander::panderOptions("table.emphasize.rownames", FALSE)
options('digits' = 4)
```

6.1. The proportional hazards model

6.1.1. Background on the Proportional Hazards Model

The exponential distribution has constant hazard:

$$\begin{aligned}f(t) &= \lambda e^{-\lambda t} \\S(t) &= e^{-\lambda t} \\h(t) &= \lambda\end{aligned}$$

Let's make two generalizations. First, we let the hazard depend on some covariates x_1, x_2, \dots, x_p ; we will indicate this dependence by extending our notation for hazard:

$$h(t|x) \stackrel{\text{def}}{=} p(T = t | T \geq t, X = x)$$

Second, we let the base hazard depend on t , but not on the covariates (for now). We can do this using either parametric or semi-parametric approaches.

6.1.2. Cox's Proportional Hazards Model

The generalization is that the hazard function is

$$\begin{aligned}h(t|x) &= h_0(t)\theta(x) \\ \theta(x) &= \exp\{\eta(x)\} \\ \eta(x) &= x'\beta \\ &\stackrel{\text{def}}{=} \beta_1 x_1 + \dots + \beta_p x_p\end{aligned}$$

6. Proportional Hazards Models

The relationship between $h(t|x)$ and $\eta(x)$ has a log link (that is, $\log \{h(t|x)\} = \log \{h_0(t)\} + \eta(x)$), as in a generalized linear model.

This model is **semi-parametric**, because the linear predictor depends on estimated parameters but the base hazard function is unspecified. There is no constant term in $\eta(x)$, because it is absorbed in the base hazard.

Alternatively, we could define $\beta_0(t) = \log \{h_0(t)\}$, and then $\eta(x, t) = \beta_0(t) + \beta_1 x_1 + \dots + \beta_p x_p$.

For two different individuals with covariate patterns x_1 and x_2 , the ratio of the hazard functions (a.k.a. **hazard ratio**, a.k.a. **relative hazard**) is:

$$\begin{aligned} \frac{h(t|x_1)}{h(t|x_2)} &= \frac{h_0(t)\theta(x_1)}{h_0(t)\theta(x_2)} \\ &= \frac{\theta(x_1)}{\theta(x_2)} \end{aligned}$$

Under the proportional hazards model, this ratio (a.k.a. proportion) does not depend on t . This property is a structural limitation of the model; it is called the **proportional hazards assumption**.

Definition 6.1 (proportional hazards). A conditional probability distribution $p(T|X)$ has **proportional hazards** if the hazard ratio $h(t|x_1)/h(t|x_2)$ does not depend on t . Mathematically, it can be written as:

$$\frac{h(t|x_1)}{h(t|x_2)} = \theta(x_1, x_2)$$

As we saw above, Cox's proportional hazards model has this property, with $\theta(x_1, x_2) = \frac{\theta(x_1)}{\theta(x_2)}$.

6. Proportional Hazards Models

i Note

We are using two similar notations, $\theta(x_1, x_2)$ and $\theta(x)$. We can link these notations if we define $\theta(x) \stackrel{\text{def}}{=} \theta(x, 0)$ and $\theta(0) = 1$.

It also has additional notable properties:

$$\begin{aligned} \frac{h(t|x_1)}{h(t|x_2)} &= \frac{\theta(x_1)}{\theta(x_2)} \\ &= \frac{\exp\{\eta(x_1)\}}{\exp\{\eta(x_2)\}} \\ &= \exp\{\eta(x_1) - \eta(x_2)\} \\ &= \exp\{x_1'\beta - x_2'\beta\} \\ &= \exp\{(x_1 - x_2)'\beta\} \end{aligned}$$

Hence on the log scale, we have:

$$\begin{aligned} \log \left\{ \frac{h(t|x_1)}{h(t|x_2)} \right\} &= \eta(x_1) - \eta(x_2) \\ &= x_1'\beta - x_2'\beta \\ &= (x_1 - x_2)'\beta \end{aligned}$$

If only one covariate x_j is changing, then:

$$\begin{aligned} \log \left\{ \frac{h(t|x_1)}{h(t|x_2)} \right\} &= (x_{1j} - x_{2j}) \cdot \beta_j \\ &\propto (x_{1j} - x_{2j}) \end{aligned}$$

That is, under Cox's model $h(t|x) = h_0(t)\exp\{x'\beta\}$, the log of the hazard ratio is proportional to the difference in x_j , with the proportionality coefficient equal to β_j .

6. Proportional Hazards Models

Further,

$$\log \{h(t|x)\} = \log \{h_0(t)\} + x'\beta$$

That is, the covariate effects are additive on the log-hazard scale.

See also:

https://en.wikipedia.org/wiki/Proportional_hazards_model#Why_it_is_called_%22proportional%22

6.1.3. Additional properties of the proportional hazards model

If $h(t|x) = h_0(t)\theta(x)$, then:

6.1.3.1. Cumulative hazards are also proportional to $H_0(t)$

$$\begin{aligned} H(t|x) &\stackrel{\text{def}}{=} \int_{u=0}^t h(u)du \\ &= \int_{u=0}^t h_0(u)\theta(x)du \\ &= \theta(x) \int_{u=0}^t h_0(u)du \\ &= \theta(x)H_0(t) \end{aligned}$$

where $H_0(t) \stackrel{\text{def}}{=} H(t|0) = \int_{u=0}^t h_0(u)du$.

6. Proportional Hazards Models

6.1.3.2. Survival functions are exponential multiples of $S_0(t)$

$$\begin{aligned} S(t|x) &= \exp \{-H(t|x)\} \\ &= \exp \{-\theta(x) \cdot H_0(t)\} \\ &= (\exp \{-H_0(t)\})^{\theta(x)} \\ &= (S_0(t))^{\theta(x)} \end{aligned}$$

where $S_0(t) \stackrel{\text{def}}{=} P(T \geq t | X = 0)$ is the survival function for an individual whose covariates are all equal to their default values.

6.1.4. Testing the proportional hazards assumption

The Nelson-Aalen estimate of the cumulative hazard is usually used for estimates of the hazard and often the cumulative hazard.

If the hazards of the three groups are proportional, that means that the ratio of the hazards is constant over t . We can test this using the ratios of the estimated cumulative hazards, which also would be proportional, as shown above.

```
library(KMsurv)
library(survival)
data(bmt)

bmt =
  bmt |>
  as_tibble() |>
  mutate(
    group =
      group |>
      factor(
        labels = c("ALL", "Low Risk AML", "High Risk AML")))
```


6. Proportional Hazards Models

```
nafit = survfit(  
  formula = Surv(t2,d3) ~ group,  
  type = "fleming-harrington",  
  data = bmt)  
  
bmt_curves = tibble(timevec = 1:1000)  
sf1 <- with(nafit[1], stepfun(time,c(1,surv)))  
sf2 <- with(nafit[2], stepfun(time,c(1,surv)))  
sf3 <- with(nafit[3], stepfun(time,c(1,surv)))  
  
bmt_curves =  
  bmt_curves |>  
  mutate(  
    cumhaz1 = -log(sf1(timevec)),  
    cumhaz2 = -log(sf2(timevec)),  
    cumhaz3 = -log(sf3(timevec)))
```

```
library(ggplot2)  
bmt_rel_hazard_plot =  
  bmt_curves |>  
  ggplot(  
    aes(  
      x = timevec,  
      y = cumhaz1/cumhaz2)  
    ) +  
  geom_line(aes(col = "ALL/Low Risk AML")) +  
  ylab("Hazard Ratio") +  
  xlab("Time") +  
  ylim(0,6) +  
  geom_line(aes(y = cumhaz3/cumhaz1, col = "High Risk AML/ALL")) +  
  geom_line(aes(y = cumhaz3/cumhaz2, col = "High Risk AML/Low Risk AML")) +  
  theme_bw() +
```

6. Proportional Hazards Models

```
labs(colour = "Comparison") +  
theme(legend.position="bottom")  
  
print(bmt_rel_hazard_plot)
```




Figure 6.1.: Hazard Ratios by Disease Group

We can zoom in on 30-300 days to take a closer look:

```
bmt_rel_hazard_plot + xlim(c(30,300))
```

6. Proportional Hazards Models




Figure 6.2.: Hazard Ratios by Disease Group (30-300 Days)

6.1.5. Smoothed hazard functions

The Nelson-Aalen estimate of the cumulative hazard is usually used for estimates of the hazard. Since the hazard is the derivative of the cumulative hazard, we need a smooth estimate of the cumulative hazard, which is provided by smoothing the step-function cumulative hazard.

The R package `muhaz` handles this for us. What we are looking for is whether the hazard function is more or less the same shape, increasing, decreasing, constant, etc. Are the hazards “proportional”?

```
plot(  
  survfit(Surv(t2,d3)~group,data=bmt),  
  col=1:3,
```

6. Proportional Hazards Models

```
lwd=2,  
fun="cumhaz",  
mark.time = TRUE)  
legend("bottomright",c("ALL","Low Risk AML","High Risk AML"),col=1:3,lwd=2)
```




Figure 6.3.: Disease-Free Cumulative Hazard by Disease Group

```
library(muhaz)  
  
muhas(bmt$t2,bmt$d3,bmt$group=="High Risk AML") |> plot(lwd=2,col=3)  
muhas(bmt$t2,bmt$d3,bmt$group=="ALL") |> lines(lwd=2,col=1)  
muhas(bmt$t2,bmt$d3,bmt$group=="Low Risk AML") |> lines(lwd=2,col=2)  
legend("topright",c("ALL","Low Risk AML","High Risk AML"),col=1:3,lwd=2)
```

6. Proportional Hazards Models




Figure 6.4.: Smoothed Hazard Rate Estimates by Disease Group

Group 3 was plotted first because it has the highest hazard.

We will see that except for an initial blip in the high risk AML group, the hazards look roughly proportional. They are all strongly decreasing.

6.1.6. Fitting the Proportional Hazards Model

How do we fit a proportional hazards regression model? We need to estimate the coefficients of the covariates, and we need to estimate the base hazard $h_0(t)$. For the covariates, supposing for simplicity that there are no tied event times, let the event times for the whole data set be t_1, t_2, \dots, t_D . Let the risk set at time t_i be $R(t_i)$ and

6. Proportional Hazards Models

$$\begin{aligned}\eta(x) &= \beta_1 x_1 + \cdots + \beta_p x_p \\ \theta(x) &= e^{\eta(x)} \\ h(t|X=x) &= h_0(t)e^{\eta(x)} = \theta(x)h_0(t)\end{aligned}$$

Conditional on a single failure at time t , the probability that the event is due to subject $f \in R(t)$ is approximately

$$\begin{aligned}\Pr(f \text{ fails} | 1 \text{ failure at } t) &= \frac{h_0(t)e^{\eta(x_f)}}{\sum_{k \in R(t)} h_0(t)e^{\eta(x_k)}} \\ &= \frac{\theta(x_f)}{\sum_{k \in R(t)} \theta(x_k)}\end{aligned}$$

The logic behind this has several steps. We first fix (ex post) the failure times and note that in this discrete context, the probability p_j that a subject j in the risk set fails at time t is just the hazard of that subject at that time.

If all of the p_j are small, the chance that exactly one subject fails is

$$\sum_{k \in R(t)} p_k \left[\prod_{m \in R(t), m \neq k} (1 - p_m) \right] \approx \sum_{k \in R(t)} p_k$$

If subject i is the one who experiences the event of interest at time t_i , then the **partial likelihood** is

$$\mathcal{L}^*(\beta|T) = \prod_i \frac{\theta(x_i)}{\sum_{k \in R(t_i)} \theta(x_k)}$$

and we can numerically maximize this with respect to the coefficients β that specify $\eta(x) = x'\beta$. When there are tied event times adjustments

6. Proportional Hazards Models

need to be made, but the likelihood is still similar. Note that we don't need to know the base hazard to solve for the coefficients.

Once we have coefficient estimates $\hat{\beta} = (\hat{\beta}_1, \dots, \hat{\beta}_p)$, this also defines $\hat{\eta}(x)$ and $\hat{\theta}(x)$ and then the estimated base cumulative hazard function is

$$\hat{H}(t) = \sum_{t_i < t} \frac{d_i}{\sum_{k \in R(t_i)} \theta(x_k)}$$

which reduces to the Nelson-Aalen estimate when there are no covariates. There are numerous other estimates that have been proposed as well.

6.2. Cox Model for the bmt data

6.2.1. Fit the model

```
bmt.cox <- coxph(Surv(t2, d3) ~ group, data = bmt)
summary(bmt.cox)
```

Call:

```
coxph(formula = Surv(t2, d3) ~ group, data = bmt)
```

```
n= 137, number of events= 83
```

| | | | coef | exp(coef) | se(coef) | z | Pr(> z) | |
|---|----------|--|--------|-----------|----------|-------|----------|---|
| groupLow | Risk AML | | -0.574 | 0.563 | 0.287 | -2.00 | 0.046 | * |
| groupHigh | Risk AML | | 0.383 | 1.467 | 0.267 | 1.43 | 0.152 | |
| --- | | | | | | | | |
| Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1 | | | | | | | | |

```
exp(coef) exp(-coef) lower .95 upper .95
```

6. Proportional Hazards Models

| | | | | |
|--------------------|-------|-------|-------|-------|
| groupLow Risk AML | 0.563 | 1.776 | 0.321 | 0.989 |
| groupHigh Risk AML | 1.467 | 0.682 | 0.869 | 2.478 |

Concordance= 0.625 (se = 0.03)
Likelihood ratio test= 13.4 on 2 df, p=0.001
Wald test = 13 on 2 df, p=0.001
Score (logrank) test = 13.8 on 2 df, p=0.001

The table provides hypothesis tests comparing groups 2 and 3 to group 1. Group 3 has the highest hazard, so the most significant comparison is not directly shown.

The coefficient 0.3834 is on the log-hazard-ratio scale, as in log-risk-ratio. The next column gives the hazard ratio 1.4673, and a hypothesis (Wald) test.

The (not shown) group 3 vs. group 2 log hazard ratio is $0.3834 + 0.5742 = 0.9576$. The hazard ratio is then $\exp(0.9576)$ or 2.605.

Inference on all coefficients and combinations can be constructed using `coef(bmt.cox)` and `vcov(bmt.cox)` as with logistic and poisson regression.

Concordance is agreement of first failure between pairs of subjects and higher predicted risk between those subjects, omitting non-informative pairs.

The Rsquare value is Cox and Snell's pseudo R-squared and is not very useful.

`summary()` prints three tests for whether the model with the group covariate is better than the one without

- **Likelihood ratio test** (chi-squared)
- **Wald test** (also chi-squared), obtained by adding the squares of the z-scores
- **Score** = log-rank test, as with comparison of survival functions.

6. Proportional Hazards Models

The likelihood ratio test is probably best in smaller samples, followed by the Wald test.

6.2.2. Survival Curves from the Cox Model

We can take a look at the resulting group-specific curves:

```
#| fig-cap: "Survival Functions for Three Groups by KM and Cox Model"


km_fit = survfit(Surv(t2, d3) ~ group, data = as.data.frame(bmt))

cox_fit = survfit(
  bmt.cox,
  newdata =
    data.frame(
      group = unique(bmt$group),
      row.names = unique(bmt$group)))

library(survminer)

list(KM = km_fit, Cox = cox_fit) |>
  survminer::ggsurvplot(
    # facet.by = "group",
    legend = "bottom",
    legend.title = "",
    combine = TRUE,
    fun = 'pct',
    size = .5,
    ggtheme = theme_bw(),
    conf.int = FALSE,
    censor = FALSE) |>
  suppressWarnings() # ggsurvplot() throws some warnings that aren't too worrying
```

6. Proportional Hazards Models



When we use `survfit()` with a Cox model, we have to specify the covariate levels we are interested in; the argument `newdata` should include a `data.frame` with the same named columns as the predictors in the Cox model and one or more levels of each.

Otherwise (that is, if the `newdata` argument is missing), a curve is produced for a single “pseudo” subject with covariate values equal to the means component of the fit.

The resulting curve(s) almost never make sense, but the default remains due to an unwarranted attachment to the option shown by some users and by other packages.

Two particularly egregious examples are factor variables and interactions. Suppose one were studying interspecies transmission of a virus, and the data set has a factor variable with levels (“pig”, “chicken”) and about equal numbers of observations for each. The “mean” covariate level will be 0.5 – is this a flying pig?

6. Proportional Hazards Models

6.2.3. Examining survfit

```
survfit(Surv(t2, d3)~group,data=bmt)
```

```
Call: survfit(formula = Surv(t2, d3) ~ group, data = bmt)
```

| | n | events | median | 0.95LCL | 0.95UCL |
|---------------------|----|--------|--------|---------|---------|
| group=ALL | 38 | 24 | 418 | 194 | NA |
| group=Low Risk AML | 54 | 25 | 2204 | 704 | NA |
| group=High Risk AML | 45 | 34 | 183 | 115 | 456 |

```
survfit(Surv(t2, d3)~group,data=bmt) |> summary()
```

```
Call: survfit(formula = Surv(t2, d3) ~ group, data = bmt)
```

| group=ALL | | | | | | |
|-----------|--------|---------|----------|---------|--------------|--------------|
| time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
| 1 | 38 | 1 | 0.974 | 0.0260 | 0.924 | 1.000 |
| 55 | 37 | 1 | 0.947 | 0.0362 | 0.879 | 1.000 |
| 74 | 36 | 1 | 0.921 | 0.0437 | 0.839 | 1.000 |
| 86 | 35 | 1 | 0.895 | 0.0498 | 0.802 | 0.998 |
| 104 | 34 | 1 | 0.868 | 0.0548 | 0.767 | 0.983 |
| 107 | 33 | 1 | 0.842 | 0.0592 | 0.734 | 0.966 |
| 109 | 32 | 1 | 0.816 | 0.0629 | 0.701 | 0.949 |
| 110 | 31 | 1 | 0.789 | 0.0661 | 0.670 | 0.930 |
| 122 | 30 | 2 | 0.737 | 0.0714 | 0.609 | 0.891 |
| 129 | 28 | 1 | 0.711 | 0.0736 | 0.580 | 0.870 |
| 172 | 27 | 1 | 0.684 | 0.0754 | 0.551 | 0.849 |
| 192 | 26 | 1 | 0.658 | 0.0770 | 0.523 | 0.827 |
| 194 | 25 | 1 | 0.632 | 0.0783 | 0.495 | 0.805 |
| 230 | 23 | 1 | 0.604 | 0.0795 | 0.467 | 0.782 |

6. Proportional Hazards Models

| | | | | | | |
|-----|----|---|-------|--------|-------|-------|
| 276 | 22 | 1 | 0.577 | 0.0805 | 0.439 | 0.758 |
| 332 | 21 | 1 | 0.549 | 0.0812 | 0.411 | 0.734 |
| 383 | 20 | 1 | 0.522 | 0.0817 | 0.384 | 0.709 |
| 418 | 19 | 1 | 0.494 | 0.0819 | 0.357 | 0.684 |
| 466 | 18 | 1 | 0.467 | 0.0818 | 0.331 | 0.658 |
| 487 | 17 | 1 | 0.439 | 0.0815 | 0.305 | 0.632 |
| 526 | 16 | 1 | 0.412 | 0.0809 | 0.280 | 0.605 |
| 609 | 14 | 1 | 0.382 | 0.0803 | 0.254 | 0.577 |
| 662 | 13 | 1 | 0.353 | 0.0793 | 0.227 | 0.548 |

group=Low Risk AML

| time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
|------|--------|---------|----------|---------|--------------|--------------|
| 10 | 54 | 1 | 0.981 | 0.0183 | 0.946 | 1.000 |
| 35 | 53 | 1 | 0.963 | 0.0257 | 0.914 | 1.000 |
| 48 | 52 | 1 | 0.944 | 0.0312 | 0.885 | 1.000 |
| 53 | 51 | 1 | 0.926 | 0.0356 | 0.859 | 0.998 |
| 79 | 50 | 1 | 0.907 | 0.0394 | 0.833 | 0.988 |
| 80 | 49 | 1 | 0.889 | 0.0428 | 0.809 | 0.977 |
| 105 | 48 | 1 | 0.870 | 0.0457 | 0.785 | 0.965 |
| 211 | 47 | 1 | 0.852 | 0.0483 | 0.762 | 0.952 |
| 219 | 46 | 1 | 0.833 | 0.0507 | 0.740 | 0.939 |
| 248 | 45 | 1 | 0.815 | 0.0529 | 0.718 | 0.925 |
| 272 | 44 | 1 | 0.796 | 0.0548 | 0.696 | 0.911 |
| 288 | 43 | 1 | 0.778 | 0.0566 | 0.674 | 0.897 |
| 381 | 42 | 1 | 0.759 | 0.0582 | 0.653 | 0.882 |
| 390 | 41 | 1 | 0.741 | 0.0596 | 0.633 | 0.867 |
| 414 | 40 | 1 | 0.722 | 0.0610 | 0.612 | 0.852 |
| 421 | 39 | 1 | 0.704 | 0.0621 | 0.592 | 0.837 |
| 481 | 38 | 1 | 0.685 | 0.0632 | 0.572 | 0.821 |
| 486 | 37 | 1 | 0.667 | 0.0642 | 0.552 | 0.805 |
| 606 | 36 | 1 | 0.648 | 0.0650 | 0.533 | 0.789 |
| 641 | 35 | 1 | 0.630 | 0.0657 | 0.513 | 0.773 |
| 704 | 34 | 1 | 0.611 | 0.0663 | 0.494 | 0.756 |
| 748 | 33 | 1 | 0.593 | 0.0669 | 0.475 | 0.739 |

6. Proportional Hazards Models

| | | | | | | |
|------|----|---|-------|--------|-------|-------|
| 1063 | 26 | 1 | 0.570 | 0.0681 | 0.451 | 0.720 |
| 1074 | 25 | 1 | 0.547 | 0.0691 | 0.427 | 0.701 |
| 2204 | 6 | 1 | 0.456 | 0.1012 | 0.295 | 0.704 |

group=High Risk AML

| time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
|------|--------|---------|----------|---------|--------------|--------------|
| 2 | 45 | 1 | 0.978 | 0.0220 | 0.936 | 1.000 |
| 16 | 44 | 1 | 0.956 | 0.0307 | 0.897 | 1.000 |
| 32 | 43 | 1 | 0.933 | 0.0372 | 0.863 | 1.000 |
| 47 | 42 | 2 | 0.889 | 0.0468 | 0.802 | 0.986 |
| 48 | 40 | 1 | 0.867 | 0.0507 | 0.773 | 0.972 |
| 63 | 39 | 1 | 0.844 | 0.0540 | 0.745 | 0.957 |
| 64 | 38 | 1 | 0.822 | 0.0570 | 0.718 | 0.942 |
| 74 | 37 | 1 | 0.800 | 0.0596 | 0.691 | 0.926 |
| 76 | 36 | 1 | 0.778 | 0.0620 | 0.665 | 0.909 |
| 80 | 35 | 1 | 0.756 | 0.0641 | 0.640 | 0.892 |
| 84 | 34 | 1 | 0.733 | 0.0659 | 0.615 | 0.875 |
| 93 | 33 | 1 | 0.711 | 0.0676 | 0.590 | 0.857 |
| 100 | 32 | 1 | 0.689 | 0.0690 | 0.566 | 0.838 |
| 105 | 31 | 1 | 0.667 | 0.0703 | 0.542 | 0.820 |
| 113 | 30 | 1 | 0.644 | 0.0714 | 0.519 | 0.801 |
| 115 | 29 | 1 | 0.622 | 0.0723 | 0.496 | 0.781 |
| 120 | 28 | 1 | 0.600 | 0.0730 | 0.473 | 0.762 |
| 157 | 27 | 1 | 0.578 | 0.0736 | 0.450 | 0.742 |
| 162 | 26 | 1 | 0.556 | 0.0741 | 0.428 | 0.721 |
| 164 | 25 | 1 | 0.533 | 0.0744 | 0.406 | 0.701 |
| 168 | 24 | 1 | 0.511 | 0.0745 | 0.384 | 0.680 |
| 183 | 23 | 1 | 0.489 | 0.0745 | 0.363 | 0.659 |
| 242 | 22 | 1 | 0.467 | 0.0744 | 0.341 | 0.638 |
| 268 | 21 | 1 | 0.444 | 0.0741 | 0.321 | 0.616 |
| 273 | 20 | 1 | 0.422 | 0.0736 | 0.300 | 0.594 |
| 318 | 19 | 1 | 0.400 | 0.0730 | 0.280 | 0.572 |
| 363 | 18 | 1 | 0.378 | 0.0723 | 0.260 | 0.550 |
| 390 | 17 | 1 | 0.356 | 0.0714 | 0.240 | 0.527 |

6. Proportional Hazards Models

| | | | | | | |
|-----|----|---|-------|--------|-------|-------|
| 422 | 16 | 1 | 0.333 | 0.0703 | 0.221 | 0.504 |
| 456 | 15 | 1 | 0.311 | 0.0690 | 0.201 | 0.481 |
| 467 | 14 | 1 | 0.289 | 0.0676 | 0.183 | 0.457 |
| 625 | 13 | 1 | 0.267 | 0.0659 | 0.164 | 0.433 |
| 677 | 12 | 1 | 0.244 | 0.0641 | 0.146 | 0.409 |

```
survfit(bmt.cox)
```

```
Call: survfit(formula = bmt.cox)
```

| | n | events | median | 0.95LCL | 0.95UCL |
|------|-----|--------|--------|---------|---------|
| [1,] | 137 | 83 | 422 | 268 | NA |

```
survfit(bmt.cox, newdata = tibble(group = unique(bmt$group)))
```

```
Call: survfit(formula = bmt.cox, newdata = tibble(group = unique(bmt$group)))
```

| | n | events | median | 0.95LCL | 0.95UCL |
|---|-----|--------|--------|---------|---------|
| 1 | 137 | 83 | 422 | 268 | NA |
| 2 | 137 | 83 | NA | 625 | NA |
| 3 | 137 | 83 | 268 | 162 | 467 |

```
bmt.cox |>
  survfit(newdata = tibble(group = unique(bmt$group))) |>
  summary()
```

```
Call: survfit(formula = bmt.cox, newdata = tibble(group = unique(bmt$group)))
```

| time | n.risk | n.event | survival1 | survival2 | survival3 |
|------|--------|---------|-----------|-----------|-----------|
| 1 | 137 | 1 | 0.993 | 0.996 | 0.989 |
| 2 | 136 | 1 | 0.985 | 0.992 | 0.978 |

6. Proportional Hazards Models

| | | | | | |
|-----|-----|---|-------|-------|-------|
| 10 | 135 | 1 | 0.978 | 0.987 | 0.968 |
| 16 | 134 | 1 | 0.970 | 0.983 | 0.957 |
| 32 | 133 | 1 | 0.963 | 0.979 | 0.946 |
| 35 | 132 | 1 | 0.955 | 0.975 | 0.935 |
| 47 | 131 | 2 | 0.941 | 0.966 | 0.914 |
| 48 | 129 | 2 | 0.926 | 0.957 | 0.893 |
| 53 | 127 | 1 | 0.918 | 0.953 | 0.882 |
| 55 | 126 | 1 | 0.911 | 0.949 | 0.872 |
| 63 | 125 | 1 | 0.903 | 0.944 | 0.861 |
| 64 | 124 | 1 | 0.896 | 0.940 | 0.851 |
| 74 | 123 | 2 | 0.881 | 0.931 | 0.830 |
| 76 | 121 | 1 | 0.873 | 0.926 | 0.819 |
| 79 | 120 | 1 | 0.865 | 0.922 | 0.809 |
| 80 | 119 | 2 | 0.850 | 0.913 | 0.788 |
| 84 | 117 | 1 | 0.843 | 0.908 | 0.778 |
| 86 | 116 | 1 | 0.835 | 0.903 | 0.768 |
| 93 | 115 | 1 | 0.827 | 0.899 | 0.757 |
| 100 | 114 | 1 | 0.820 | 0.894 | 0.747 |
| 104 | 113 | 1 | 0.812 | 0.889 | 0.737 |
| 105 | 112 | 2 | 0.797 | 0.880 | 0.717 |
| 107 | 110 | 1 | 0.789 | 0.875 | 0.707 |
| 109 | 109 | 1 | 0.782 | 0.870 | 0.697 |
| 110 | 108 | 1 | 0.774 | 0.866 | 0.687 |
| 113 | 107 | 1 | 0.766 | 0.861 | 0.677 |
| 115 | 106 | 1 | 0.759 | 0.856 | 0.667 |
| 120 | 105 | 1 | 0.751 | 0.851 | 0.657 |
| 122 | 104 | 2 | 0.735 | 0.841 | 0.637 |
| 129 | 102 | 1 | 0.727 | 0.836 | 0.627 |
| 157 | 101 | 1 | 0.720 | 0.831 | 0.617 |
| 162 | 100 | 1 | 0.712 | 0.826 | 0.607 |
| 164 | 99 | 1 | 0.704 | 0.821 | 0.598 |
| 168 | 98 | 1 | 0.696 | 0.815 | 0.588 |
| 172 | 97 | 1 | 0.688 | 0.810 | 0.578 |
| 183 | 96 | 1 | 0.680 | 0.805 | 0.568 |

6. Proportional Hazards Models

| | | | | | |
|-----|----|---|-------|-------|-------|
| 192 | 95 | 1 | 0.672 | 0.800 | 0.558 |
| 194 | 94 | 1 | 0.664 | 0.794 | 0.549 |
| 211 | 93 | 1 | 0.656 | 0.789 | 0.539 |
| 219 | 92 | 1 | 0.648 | 0.783 | 0.530 |
| 230 | 90 | 1 | 0.640 | 0.778 | 0.520 |
| 242 | 89 | 1 | 0.632 | 0.773 | 0.511 |
| 248 | 88 | 1 | 0.624 | 0.767 | 0.501 |
| 268 | 87 | 1 | 0.616 | 0.761 | 0.492 |
| 272 | 86 | 1 | 0.608 | 0.756 | 0.482 |
| 273 | 85 | 1 | 0.600 | 0.750 | 0.473 |
| 276 | 84 | 1 | 0.592 | 0.745 | 0.464 |
| 288 | 83 | 1 | 0.584 | 0.739 | 0.454 |
| 318 | 82 | 1 | 0.576 | 0.733 | 0.445 |
| 332 | 81 | 1 | 0.568 | 0.727 | 0.436 |
| 363 | 80 | 1 | 0.560 | 0.722 | 0.427 |
| 381 | 79 | 1 | 0.552 | 0.716 | 0.418 |
| 383 | 78 | 1 | 0.544 | 0.710 | 0.409 |
| 390 | 77 | 2 | 0.528 | 0.698 | 0.392 |
| 414 | 75 | 1 | 0.520 | 0.692 | 0.383 |
| 418 | 74 | 1 | 0.512 | 0.686 | 0.374 |
| 421 | 73 | 1 | 0.504 | 0.680 | 0.366 |
| 422 | 72 | 1 | 0.496 | 0.674 | 0.357 |
| 456 | 71 | 1 | 0.488 | 0.667 | 0.349 |
| 466 | 70 | 1 | 0.480 | 0.661 | 0.340 |
| 467 | 69 | 1 | 0.472 | 0.655 | 0.332 |
| 481 | 68 | 1 | 0.464 | 0.649 | 0.324 |
| 486 | 67 | 1 | 0.455 | 0.642 | 0.315 |
| 487 | 66 | 1 | 0.447 | 0.636 | 0.307 |
| 526 | 65 | 1 | 0.439 | 0.629 | 0.299 |
| 606 | 63 | 1 | 0.431 | 0.623 | 0.291 |
| 609 | 62 | 1 | 0.423 | 0.616 | 0.283 |
| 625 | 61 | 1 | 0.415 | 0.609 | 0.275 |
| 641 | 60 | 1 | 0.407 | 0.603 | 0.267 |
| 662 | 59 | 1 | 0.399 | 0.596 | 0.260 |

6. Proportional Hazards Models

| | | | | | |
|------|----|---|-------|-------|-------|
| 677 | 58 | 1 | 0.391 | 0.589 | 0.252 |
| 704 | 57 | 1 | 0.383 | 0.582 | 0.244 |
| 748 | 56 | 1 | 0.374 | 0.575 | 0.237 |
| 1063 | 47 | 1 | 0.365 | 0.567 | 0.228 |
| 1074 | 46 | 1 | 0.356 | 0.559 | 0.220 |
| 2204 | 9 | 1 | 0.313 | 0.520 | 0.182 |

6.3. Adjustment for Ties (optional)

6.3.1.

At each time t_i at which more than one of the subjects has an event, let d_i be the number of events at that time, D_i the set of subjects with events at that time, and let s_i be a covariate vector for an artificial subject obtained by adding up the covariate values for the subjects with an event at time t_i . Let

$$\bar{\eta}_i = \beta_1 s_{i1} + \cdots + \beta_p s_{ip}$$

and $\bar{\theta}_i = \exp \{\bar{\eta}_i\}$.

Let s_i be a covariate vector for an artificial subject obtained by adding up the covariate values for the subjects with an event at time t_i . Note that

$$\begin{aligned} \bar{\eta}_i &= \sum_{j \in D_i} \beta_1 x_{j1} + \cdots + \beta_p x_{jp} \\ &= \beta_1 s_{i1} + \cdots + \beta_p s_{ip} \\ \bar{\theta}_i &= \exp \{\bar{\eta}_i\} \\ &= \prod_{j \in D_i} \theta_j \end{aligned}$$

6. Proportional Hazards Models

6.3.1.1. Breslow's method for ties

Breslow's method estimates the partial likelihood as

$$\begin{aligned} L(\beta|T) &= \prod_i \frac{\bar{\theta}_i}{[\sum_{k \in R(t_i)} \theta_k]^{d_i}} \\ &= \prod_i \prod_{j \in D_i} \frac{\theta_j}{\sum_{k \in R(t_i)} \theta_k} \end{aligned}$$

This method is equivalent to treating each event as distinct and using the non-ties formula. It works best when the number of ties is small. It is the default in many statistical packages, including PROC PHREG in SAS.

6.3.1.2. Efron's method for ties

The other common method is Efron's, which is the default in R.

$$L(\beta|T) = \prod_i \frac{\bar{\theta}_i}{\prod_{j=1}^{d_i} [\sum_{k \in R(t_i)} \theta_k - \frac{j-1}{d_i} \sum_{k \in D_i} \theta_k]}$$

This is closer to the exact discrete partial likelihood when there are many ties.

The third option in R (and an option also in SAS as **discrete**) is the “exact” method, which is the same one used for matched logistic regression.

6.3.1.3. Example: Breslow's method

Suppose as an example we have a time t where there are 20 individuals at risk and three failures. Let the three individuals have risk parameters

6. Proportional Hazards Models

$\theta_1, \theta_2, \theta_3$ and let the sum of the risk parameters of the remaining 17 individuals be θ_R . Then the factor in the partial likelihood at time t using Breslow's method is

$$\left(\frac{\theta_1}{\theta_R + \theta_1 + \theta_2 + \theta_3} \right) \left(\frac{\theta_2}{\theta_R + \theta_1 + \theta_2 + \theta_3} \right) \left(\frac{\theta_3}{\theta_R + \theta_1 + \theta_2 + \theta_3} \right)$$

If on the other hand, they had died in the order 1, 2, 3, then the contribution to the partial likelihood would be:

$$\left(\frac{\theta_1}{\theta_R + \theta_1 + \theta_2 + \theta_3} \right) \left(\frac{\theta_2}{\theta_R + \theta_2 + \theta_3} \right) \left(\frac{\theta_3}{\theta_R + \theta_3} \right)$$

as the risk set got smaller with each failure. The exact method roughly averages the results for the six possible orderings of the failures.

6.3.1.4. Example: Efron's method

But we don't know the order they failed in, so instead of reducing the denominator by one risk coefficient each time, we reduce it by the same fraction. This is Efron's method.

$$\left(\frac{\theta_1}{\theta_R + \theta_1 + \theta_2 + \theta_3} \right) \left(\frac{\theta_2}{\theta_R + 2(\theta_1 + \theta_2 + \theta_3)/3} \right) \left(\frac{\theta_3}{\theta_R + (\theta_1 + \theta_2 + \theta_3)/3} \right)$$

7. Building Cox Proportional Hazards models

Configuring R

Functions from these packages will be used throughout this document:

```
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggfortify) # help with graphics
library(ggeasy) # help with graphics
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(dplyr) # manipulate data
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(fs) # filesystem path manipulations
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
conflicts_prefer(dplyr::filter)
conflicts_prefer(ggplot2::autoplot)
```

Here are some R settings I use in this document:

7. Building Cox Proportional Hazards models

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
pander::panderOptions("table.emphasize.rownames", FALSE)
options('digits' = 4)
legend_text_size = 9
```

7.1. Building Cox Proportional Hazards models

7.1.1. hodg Lymphoma Data Set from KMsurv

7.1.1.1. Participants

43 bone marrow transplant patients at Ohio State University (Avalos 1993)

7.1.1.2. Variables

- **dtype**: Disease type (Hodgkin's or non-Hodgkins lymphoma)
- **gtype**: Bone marrow graft type:
- **allogeneic**: from HLA-matched sibling
- **autologous**: from self (prior to chemo)
- **time**: time to study exit
- **delta**: study exit reason (death/relapse vs censored)
- **wtime**: waiting time to transplant (in months)
- **score**: Karnofsky score:
- 80–100: Able to carry on normal activity and to work; no special care needed.
- 50–70: Unable to work; able to live at home and care for most personal needs; varying amount of assistance needed.
- 10–60: Unable to care for self; requires equivalent of institutional or hospital care; disease may be progressing rapidly.

7. Building Cox Proportional Hazards models

7.1.1.3. Data

```
data(hodg, package = "KMsurv")
hodg2 = hodg |>
  as_tibble() |>
  mutate(
    # We add factor labels to the categorical variables:
    gtype = gtype |>
      case_match(
        1 ~ "Allogenic",
        2 ~ "Autologous"),
    dtype = dtype |>
      case_match(
        1 ~ "Non-Hodgkins",
        2 ~ "Hodgkins") |>
      factor() |>
      relevel(ref = "Non-Hodgkins"),
    delta = delta |>
      case_match(
        1 ~ "dead",
        0 ~ "alive"),
    surv = Surv(
      time = time,
      event = delta == "dead")
  )
hodg2 |> print()
```

```
# A tibble: 43 x 7
  gtype      dtype      time delta score wtime  surv
  <chr>    <fct>    <int> <chr>  <int> <int> <Surv>
1 Allogenic Non-Hodgkins    28 dead    90    24    28
2 Allogenic Non-Hodgkins    32 dead    30     7    32
```

7. Building Cox Proportional Hazards models

```

3 Allogenic Non-Hodgkins    49 dead    40      8    49
4 Allogenic Non-Hodgkins    84 dead    60     10   84
5 Allogenic Non-Hodgkins   357 dead    70     42  357
6 Allogenic Non-Hodgkins   933 alive    90      9  933+
7 Allogenic Non-Hodgkins  1078 alive   100     16 1078+
8 Allogenic Non-Hodgkins  1183 alive    90     16 1183+
9 Allogenic Non-Hodgkins  1560 alive    80     20 1560+
10 Allogenic Non-Hodgkins  2114 alive    80     27 2114+
# i 33 more rows

```

7.1.2. Proportional hazards model

```

hodg.cox1 = coxph(
  formula = surv ~ gtype * dtype + score + wtime,
  data = hodg2)

summary(hodg.cox1)

```

Call:

```
coxph(formula = surv ~ gtype * dtype + score + wtime, data = hodg2)
```

n= 43, number of events= 26

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|-------------------------------|---------|-----------|----------|-------|-------------|
| gtypeAutologous | 0.6394 | 1.8953 | 0.5937 | 1.08 | 0.2815 |
| dtypeHodgkins | 2.7603 | 15.8050 | 0.9474 | 2.91 | 0.0036 ** |
| score | -0.0495 | 0.9517 | 0.0124 | -3.98 | 6.8e-05 *** |
| wttime | -0.0166 | 0.9836 | 0.0102 | -1.62 | 0.1046 |
| gtypeAutologous:dtypeHodgkins | -2.3709 | 0.0934 | 1.0355 | -2.29 | 0.0220 * |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

7. Building Cox Proportional Hazards models

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|-------------------------------|-----------|------------|-----------|-----------|
| gtypeAutologous | 1.8953 | 0.5276 | 0.5920 | 6.068 |
| dtypeHodgkins | 15.8050 | 0.0633 | 2.4682 | 101.207 |
| score | 0.9517 | 1.0507 | 0.9288 | 0.975 |
| wtime | 0.9836 | 1.0167 | 0.9641 | 1.003 |
| gtypeAutologous:dtypeHodgkins | 0.0934 | 10.7074 | 0.0123 | 0.711 |

Concordance= 0.776 (se = 0.059)

Likelihood ratio test= 32.1 on 5 df, p=6e-06

Wald test = 27.2 on 5 df, p=5e-05

Score (logrank) test = 37.7 on 5 df, p=4e-07

7.2. Diagnostic graphs for proportional hazards assumption

7.2.1. Analysis plan

- **survival function** for the four combinations of disease type and graft type.
- **observed (nonparametric) vs. expected (semiparametric) survival functions.**
- **complementary log-log survival** for the four groups.

7.2.2. Kaplan-Meier survival functions

```
km_model = survfit(  
  formula = surv ~ dtype + gtype,  
  data = hodg2)
```


7. Building Cox Proportional Hazards models

```
km_model |>
  autoplot(conf.int = FALSE) +
  theme_bw() +
  theme(
    legend.position="bottom",
    legend.title = element_blank(),
    legend.text = element_text(size = legend_text_size)
  ) +
  guides(col=guide_legend(ncol=2)) +
  ylab('Survival probability, S(t)') +
  xlab("Time since transplant (days)")
```




Figure 7.1.: Kaplan-Meier Survival Curves for HOD/NHL and Allo/Auto Grafts

7. Building Cox Proportional Hazards models

7.2.3. Observed and expected survival curves

```
# we need to create a tibble of covariate patterns;
# we will set score and wtime to mean values for disease and graft types:
means = hodg2 |>
  summarize(
    .by = c(dtype, gtype),
    score = mean(score),
    wtime = mean(wtime)) |>
  arrange(dtype, gtype) |>
  mutate(strata = paste(dtype, gtype, sep = ",")) |>
  as.data.frame()

# survfit.coxph() will use the rownames of its `newdata`
# argument to label its output:
rownames(means) = means$strata

cox_model =
  hodg.cox1 |>
  survfit(
    data = hodg2, # ggsurvplot() will need this
    newdata = means)

# I couldn't find a good function to reformat `cox_model` for ggplot,
# so I made my own:
stack_surv_ph = function(cox_model)
{
  cox_model$surv |>
    as_tibble() |>
    mutate(time = cox_model$time) |>
    pivot_longer(
      cols = -time,
      names_to = "strata",
```

7. Building Cox Proportional Hazards models

```
    values_to = "surv") |>
  mutate(
    cumhaz = -log(surv),
    model = "Cox PH")
}

km_and_cph =
  km_model |>
  fortify(surv.connect = TRUE) |>
  mutate(
    strata = trimws(strata),
    model = "Kaplan-Meier",
    cumhaz = -log(surv)) |>
  bind_rows(stack_surv_ph(cox_model))
```

```
km_and_cph |>
  ggplot(aes(x = time, y = surv, col = model)) +
  geom_step() +
  facet_wrap(~strata) +
  theme_bw() +
  ylab("S(t) = P(T>=t)") +
  xlab("Survival time (t, days)") +
  theme(legend.position = "bottom")
```

7. Building Cox Proportional Hazards models




Figure 7.2.: Observed and expected survival curves for `bmt` data

7.2.4. Cumulative hazard (log-scale) curves

Also known as “complementary log-log (clog-log) survival curves”.

```
na_model = survfit(  
  formula = surv ~ dtype + gtype,  
  data = hodg2,  
  type = "fleming")  
  
na_model |>  
  survminer::ggsurvplot(  
    legend = "bottom",  
    legend.title = "",
```

7. Building Cox Proportional Hazards models

```
ylab = "log(Cumulative Hazard)",
xlab = "Time since transplant (days, log-scale)",
fun = 'cloglog',
size = .5,
ggtheme = theme_bw(),
conf.int = FALSE,
censor = TRUE) |>
magrittr::extract2("plot") +
guides(
  col =
    guide_legend(
      ncol = 2,
      label.theme =
        element_text(
          size = legend_text_size)))
```

7. Building Cox Proportional Hazards models




Figure 7.3.: Complementary log-log survival curves - Nelson-Aalen estimates

Let's compare these empirical (i.e., non-parametric) curves with the fitted curves from our `coxph()` model:

```
cox_model |>
  survminer::ggsurvplot(
    facet_by = "",
    legend = "bottom",
    legend.title = "",
    ylab = "log(Cumulative Hazard)",
    xlab = "Time since transplant (days, log-scale)",
    fun = 'cloglog',
    size = .5,
    ggtheme = theme_bw(),
```

7. Building Cox Proportional Hazards models

```
  censor = FALSE, # doesn't make sense for cox model
  conf.int = FALSE) |>
magrittr::extract2("plot") +
guides(
  col =
    guide_legend(
      ncol = 2,
      label.theme =
        element_text(
          size = legend_text_size)))
```



Figure 7.4.: Complementary log-log survival curves - PH estimates

Now let's overlay these cumulative hazard curves:

7. Building Cox Proportional Hazards models

```
na_and_cph =  
  na_model |>  
  fortify(fun = "cumhaz") |>  
  # `fortify.survfit()` doesn't name cumhaz correctly:  
  rename(cumhaz = surv) |>  
  mutate(  
    surv = exp(-cumhaz),  
    strata = trimws(strata)) |>  
  mutate(model = "Nelson-Aalen") |>  
  bind_rows(stack_surv_ph(cox_model))  
  
na_and_cph |>  
  ggplot(  
    aes(  
      x = time,  
      y = cumhaz,  
      col = model)) +  
  geom_step() +  
  facet_wrap(~strata) +  
  theme_bw() +  
  scale_y_continuous(  
    trans = "log10",  
    name = "Cumulative hazard H(t) (log-scale)") +  
  scale_x_continuous(  
    trans = "log10",  
    name = "Survival time (t, days, log-scale)") +  
  theme(legend.position = "bottom")
```


7. Building Cox Proportional Hazards models



Figure 7.5.: Observed and expected cumulative hazard curves for `bmt` data (cloglog format)

7.3. Predictions and Residuals

7.3.1. Review: Predictions in Linear Regression

- In linear regression, we have a linear predictor for each data point i

$$\begin{aligned}\eta_i &= \beta_0 + \beta_1 x_{1i} + \dots + \beta_p x_{pi} \\ \hat{y}_i &= \hat{\eta}_i = \hat{\beta}_0 + \hat{\beta}_1 x_{1i} + \dots + \hat{\beta}_p x_{pi} \\ y_i &\sim N(\eta_i, \sigma^2)\end{aligned}$$

7. Building Cox Proportional Hazards models

- \hat{y}_i estimates the conditional mean of y_i given the covariate values \tilde{x}_i . This together with the prediction error says that we are predicting the distribution of values of y .

7.3.2. Review: Residuals in Linear Regression

- The usual residual is $r_i = y_i - \hat{y}_i$, the difference between the actual value of y and a prediction of its mean.
- The residuals are also the quantities the sum of whose squares is being minimized by the least squares/MLE estimation.

7.3.3. Predictions and Residuals in survival models

- In survival analysis, the equivalent of y_i is the event time t_i , which is unknown for the censored observations.
- The expected event time can be tricky to calculate:

$$\hat{E}[T|X = x] = \int_{t=0}^{\infty} \hat{S}(t) dt$$

7.3.4. Wide prediction intervals

The nature of time-to-event data results in very wide prediction intervals:

- Suppose a cancer patient is predicted to have a mean lifetime of 5 years after diagnosis and suppose the distribution is exponential.
- If we want a 95% interval for survival, the lower end is at the 0.025 percentage point of the exponential which is `qexp(.025, rate = 1/5) = 0.12658904` years, or 1/40 of the mean lifetime.

7. Building Cox Proportional Hazards models

- The upper end is at the 0.975 point which is `qexp(.975, rate = 1/5) = 18.44439727` years, or 3.7 times the mean lifetime.
- Saying that the survival time is somewhere between 6 weeks and 18 years does not seem very useful, but it may be the best we can do.
- For survival analysis, something is like a residual if it is small when the model is accurate or if the accumulation of them is in some way minimized by the estimation algorithm, but there is no exact equivalence to linear regression residuals.
- And if there is, they are mostly quite large!

7.3.5. Types of Residuals in Time-to-Event Models

- It is often hard to make a decision from graph appearances, though the process can reveal much.
- Some diagnostic tests are based on residuals as with other regression methods:
- **Schoenfeld residuals** (via `cox.zph`) for proportionality.
- **Cox-Snell residuals** for goodness of fit.
- **martingale residuals** for non-linearity.
- **dfbeta** for influence.

7.3.6. Schoenfeld residuals

- There is a Schoenfeld residual for each subject i with an event (not censored) and for each predictor x_k .
- At the event time t for that subject, there is a risk set R , and each subject j in the risk set has a risk coefficient θ_j and also a value x_{jk} of the predictor.
- The Schoenfeld residual is the difference between x_{ik} and the risk-weighted average of all the x_{jk} over the risk set.

7. Building Cox Proportional Hazards models

$$r_{ik}^S = x_{ik} - \frac{\sum_{k \in R} x_{jk} \theta_k}{\sum_{k \in R} \theta_k}$$

This residual measures how typical the individual subject is with respect to the covariate at the time of the event. Since subjects should fail more or less uniformly according to risk, the Schoenfeld residuals should be approximately level over time, not increasing or decreasing.

We can test this with the correlation with time on some scale, which could be the time itself, the log time, or the rank in the set of failure times.

The default is to use the KM curve as a transform, which is similar to the rank but deals better with censoring.

The `cox.zph()` function implements a score test proposed in Grambsch and Therneau (1994).

```
hodg.zph = cox.zph(hodg.cox1)
print(hodg.zph)
```

| | chisq | df | p |
|-------------|---------|----|-------|
| gtype | 0.5400 | 1 | 0.462 |
| dtype | 1.8012 | 1 | 0.180 |
| score | 3.8805 | 1 | 0.049 |
| wtime | 0.0173 | 1 | 0.895 |
| gtype:dtype | 4.0474 | 1 | 0.044 |
| GLOBAL | 13.7573 | 5 | 0.017 |


7.3.6.1. gtype

```
ggcoxzph(hodg.zph, var = "gtype")
```

7. Building Cox Proportional Hazards models

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.4624




7.3.6.2. dtype

```
ggcoxzph(hodg.zph, var = "dtype")
```

7. Building Cox Proportional Hazards models

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.1796




7.3.6.3. score

```
ggcoxzph(hodg.zph, var = "score")
```

7. Building Cox Proportional Hazards models

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.0489




7.3.6.4. wtime

```
ggcoxzph(hodg.zph, var = "wtime")
```

7. Building Cox Proportional Hazards models

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.8954



7.3.6.5. gtype:dtype

```
ggcoxzph(hodg.zph, var = "gtype:dtype")
```


7. Building Cox Proportional Hazards models

Global Schoenfeld Test p: 0.01723



7.3.6.6. Conclusions

- From the correlation test, the Karnofsky score and the interaction with graft type disease type induce modest but statistically significant non-proportionality.
- The sample size here is relatively small (26 events in 43 subjects). If the sample size is large, very small amounts of non-proportionality can induce a significant result.
- As time goes on, autologous grafts are over-represented at their own event times, but those from HOD patients become less represented.
- Both the statistical tests and the plots are useful.

7.4. Goodness of Fit using the Cox-Snell Residuals

(references: Klein & Moeschberger textbook, §11.2, and Dobson & Barnett textbook, §10.6)

Suppose that an individual has a survival time T which has survival function $S(t)$, meaning that $\Pr(T > t) = S(t)$. Then $S(T)$ has a uniform distribution on $(0, 1)$.

$$\begin{aligned}\Pr(S(T_i) \leq u) &= \Pr(T_i > S_i^{-1}(u)) \\ &= S_i(S_i^{-1}(u)) \\ &= u\end{aligned}$$

Also, if U has a uniform distribution on $(0, 1)$, then what is the distribution of $-\ln(U)$?

$$\begin{aligned}\Pr(-\ln(U) < x) &= \Pr(U > \exp\{-x\}) \\ &= 1 - e^{-x}\end{aligned}$$

which is the CDF of an exponential distribution with parameter $\lambda = 1$.

So,

$$r_i^{CS} \stackrel{\text{def}}{=} -\ln[\hat{S}(t_i|x_i)] = \hat{H}(t_i|\tilde{x}_i)$$

should have an exponential distribution with constant hazard $\lambda = 1$ if the estimate \hat{S}_i is accurate, which means that these values should look like a censored sample from this exponential distribution. These values are called **generalized residuals** or **Cox-Snell residuals**.

7. Building Cox Proportional Hazards models

```
hodg2 = hodg2 |>
  mutate(cs = predict(hodg.cox1, type = "expected"))

surv.csr = survfit(
  data = hodg2,
  formula = Surv(time = cs, event = delta == "dead") ~ 1,
  type = "fleming-harrington")

autoplot(surv.csr, fun = "cumhaz") +
  geom_abline(aes(intercept = 0, slope = 1), col = "red") +
  theme_bw()
```




Figure 7.6.: Cumulative Hazard of Cox-Snell Residuals

The line with slope 1 and intercept 0 fits the curve relatively well, so we

7. Building Cox Proportional Hazards models

don't see lack of fit using this procedure.

7.5. Martingale Residuals

The **martingale residuals** are a slight modification of the Cox-Snell residuals. If the censoring indicator is δ_i , then

$$r_i^M = \delta_i - r_i^{CS}$$

These residuals can be interpreted as an estimate of the excess number of events seen in the data but not predicted by the model. We will use these to examine the functional forms of continuous covariates.

7.5.1. Using Martingale Residuals

Martingale residuals can be used to examine the functional form of a numeric variable.

- We fit the model without that variable and compute the martingale residuals.
- We then plot these martingale residuals against the values of the variable.
- We can see curvature, or a possible suggestion that the variable can be discretized.

Let's use this to examine the `score` and `wtime` variables in the `wtime` data set.

Karnofsky score

7. Building Cox Proportional Hazards models

```
hodg2 = hodg2 |>
  mutate(
    mres =
      hodg.cox1 |>
      update(. ~ . - score) |>
      residuals(type="martingale"))

hodg2 |>
  ggplot(aes(x = score, y = mres)) +
  geom_point() +
  geom_smooth(method = "loess", aes(col = "loess")) +
  geom_smooth(method = 'lm', aes(col = "lm")) +
  theme_classic() +
  xlab("Karnofsky Score") +
  ylab("Martingale Residuals") +
  guides(col=guide_legend(title = ""))
```

7. Building Cox Proportional Hazards models




Figure 7.7.: Martingale Residuals vs. Karnofsky Score

The line is almost straight. It could be some modest transformation of the Karnofsky score would help, but it might not make much difference.

Waiting time

```
hodg2$mres =  
  hodg.cox1 |>  
  update(. ~ . - wtime) |>  
  residuals(type="martingale")  
  
hodg2 |>  
  ggplot(aes(x = wtime, y = mres)) +
```

7. Building Cox Proportional Hazards models

```
geom_point() +  
geom_smooth(method = "loess", aes(col = "loess")) +  
geom_smooth(method = 'lm', aes(col = "lm")) +  
theme_classic() +  
xlab("Waiting Time") +  
ylab("Martingale Residuals") +  
guides(col=guide_legend(title = ""))
```




Figure 7.8.: Martingale Residuals vs. Waiting Time

The line could suggest a step function. To see where the drop is, we can look at the largest waiting times and the associated martingale residual.

The martingale residuals are all negative for `wtime > 83` and positive for the next smallest value. A reasonable cut-point is 80 days.

7. Building Cox Proportional Hazards models

Updating the model

Let's reformulate the model with dichotomized `wtime`.

```
hodg2 =  
  hodg2 |>  
  mutate(  
    wt2 = cut(  
      wtime, c(0, 80, 200),  
      labels=c("short", "long"))  
  )  
  
hodg.cox2 =  
  coxph(  
    formula =  
      Surv(time, event = delta == "dead") ~  
      gtype*dtype + score + wt2,  
    data = hodg2)
```

```
hodg.cox1 |> drop1(test="Chisq")
```

Table 7.1.: Model summary table with waiting time on continuous scale

| | Df | AIC | LRT | Pr(>Chi) |
|-------------|----|-------|--------|----------|
| | NA | 152.4 | NA | NA |
| score | 1 | 167.6 | 17.236 | 0.0000 |
| wtime | 1 | 153.6 | 3.279 | 0.0702 |
| gtype:dtype | 1 | 155.8 | 5.436 | 0.0197 |

```
hodg.cox2 |> drop1(test="Chisq")
```


7. Building Cox Proportional Hazards models

Table 7.2.: Model summary table with dichotomized waiting time

| | Df | AIC | LRT | Pr(>Chi) |
|-------------|----|-------|--------|----------|
| | NA | 149.0 | NA | NA |
| score | 1 | 168.6 | 21.604 | 0.0000 |
| wt2 | 1 | 153.6 | 6.608 | 0.0102 |
| gtype:dtype | 1 | 152.0 | 4.970 | 0.0258 |

The new model has better (lower) AIC.

7.6. Checking for Outliers and Influential Observations

We will check for outliers using the deviance residuals. The martingale residuals show excess events or the opposite, but highly skewed, with the maximum possible value being 1, but the smallest value can be very large negative. Martingale residuals can detect unexpectedly long-lived patients, but patients who die unexpectedly early show up only in the deviance residual. Influence will be examined using dfbeta in a similar way to linear regression, logistic regression, or Poisson regression.

7.6.1. Deviance Residuals

$$r_i^D = \text{sign}(r_i^M) \sqrt{-2 [r_i^M + \delta_i \ln(\delta_i - r_i^M)]}$$
$$r_i^D = \text{sign}(r_i^M) \sqrt{-2 [r_i^M + \delta_i \ln(r_i^{CS})]}$$

Roughly centered on 0 with approximate standard deviation 1.

7. Building Cox Proportional Hazards models

7.6.2.

```
hodg.mart = residuals(hodg.cox2,type="martingale")
hodg.dev = residuals(hodg.cox2,type="deviance")
hodg.dfb = residuals(hodg.cox2,type="dfbeta")
hodg.preds = predict(hodg.cox2) #linear predictor
```

```
plot(hodg.preds,
     hodg.mart,
     xlab="Linear Predictor",
     ylab="Martingale Residual")
```




Figure 7.9.: Martingale Residuals vs. Linear Predictor

The smallest three martingale residuals in order are observations 1, 29,

7. Building Cox Proportional Hazards models

and 18.

```
plot(hodg.preds,hodg.dev,xlab="Linear Predictor",ylab="Deviance Residual")
```



Figure 7.10.: Deviance Residuals vs. Linear Predictor

The two largest deviance residuals are observations 1 and 29. Worth examining.

7.6.3. dfbeta

- dfbeta is the approximate change in the coefficient vector if that observation were dropped
- dfbetas is the approximate change in the coefficients, scaled by the standard error for the coefficients.

7. Building Cox Proportional Hazards models

7.6.3.1. Graft type

```
plot(hodg.dfb[,1],xlab="Observation Order",ylab="dfbeta for Graft Type")
```




Figure 7.11.: dfbeta Values by Observation Order for Graft Type

The smallest dfbeta for graft type is observation 1.

7.6.3.2. Disease type

```
plot(hodg.dfb[,2],  
      xlab="Observation Order",  
      ylab="dfbeta for Disease Type")
```

7. Building Cox Proportional Hazards models




Figure 7.12.: dfbeta Values by Observation Order for Disease Type

The smallest two dfbeta values for disease type are observations 1 and 16.

7.6.3.3. Karnofsky score

```
plot(hodg.dfb[,3],  
     xlab="Observation Order",  
     ylab="dfbeta for Karnofsky Score")
```

7. Building Cox Proportional Hazards models




Figure 7.13.: dfbeta Values by Observation Order for Karnofsky Score

The two highest dfbeta values for score are observations 1 and 18. The next three are observations 17, 29, and 19. The smallest value is observation 2.

7.6.3.4. Waiting time (dichotomized)

```
plot(  
  hodg.dfb[,4],  
  xlab="Observation Order",  
  ylab="dfbeta for `Waiting Time < 80`")
```

7. Building Cox Proportional Hazards models




Figure 7.14.: dfbeta Values by Observation Order for Waiting Time (dichotomized)

The two large values of dfbeta for dichotomized waiting time are observations 15 and 16. This may have to do with the discretization of waiting time.

7.6.3.5. Interaction: graft type and disease type

```
plot(hodg.dfb[,5],  
     xlab="Observation Order",  
     ylab="dfbeta for dtype:grtype")
```

7. Building Cox Proportional Hazards models




Figure 7.15.: dfbeta Values by Observation Order for dtype:gtype

The two largest values are observations 1 and 16. The smallest value is observation 35.

Table 7.3.: Observations to Examine by Residuals and Influence

| Diagnostic | Observations to Examine |
|----------------------------|-------------------------|
| Martingale Residuals | 1, 29, 18 |
| Deviance Residuals | 1, 29 |
| Graft Type Influence | 1 |
| Disease Type Influence | 1, 16 |
| Karnofsky Score Influence | 1, 18 (17, 29, 19) |
| Waiting Time Influence | 15, 16 |
| Graft by Disease Influence | 1, 16, 35 |

7. Building Cox Proportional Hazards models

The most important observations to examine seem to be 1, 15, 16, 18, and 29.

7.6.4.

```
with(hodg,summary(time[delta==1]))
```

| Min. | 1st Qu. | Median | Mean | 3rd Qu. | Max. |
|------|---------|--------|-------|---------|------|
| 2 | 41.25 | 62.5 | 97.62 | 83.25 | 524 |

```
with(hodg,summary(wtime))
```

| Min. | 1st Qu. | Median | Mean | 3rd Qu. | Max. |
|------|---------|--------|------|---------|------|
| 5 | 16 | 24 | 37.7 | 55.5 | 171 |

```
with(hodg,summary(score))
```

| Min. | 1st Qu. | Median | Mean | 3rd Qu. | Max. |
|------|---------|--------|-------|---------|------|
| 20 | 60 | 80 | 76.28 | 90 | 100 |

```
hodg.cox2
```

Call:

```
coxph(formula = Surv(time, event = delta == "dead") ~ gtype *  
      dtype + score + wt2, data = hodg2)
```

7. Building Cox Proportional Hazards models

| | coef | exp(coef) | se(coef) | z | p |
|-------------------------------|-------|-----------|----------|----|-------|
| gtypeAutologous | 0.67 | 1.94 | 0.59 | 1 | 0.263 |
| dtypeHodgkins | 2.33 | 10.25 | 0.73 | 3 | 0.002 |
| score | -0.06 | 0.95 | 0.01 | -4 | 8e-06 |
| wt2long | -2.06 | 0.13 | 1.05 | -2 | 0.050 |
| gtypeAutologous:dtypeHodgkins | -2.07 | 0.13 | 0.93 | -2 | 0.026 |

Likelihood ratio test=35 on 5 df, p=1e-06
n= 43, number of events= 26

```

hodg2[c(1,15,16,18,29),] |>
  select(gtype, dtype, time, delta, score, wtime) |>
  mutate(
    comment =
      c(
        "early death, good score, low risk",
        "high risk grp, long wait, poor score",
        "high risk grp, short wait, poor score",
        "early death, good score, med risk grp",
        "early death, good score, med risk grp"
      )
  )

```

| gtype | dtype | time | delta | score | wtime | comment |
|------------|--------------|------|-------|-------|-------|---------------------------------------|
| Allogenic | Non-Hodgkins | 28 | dead | 90 | 24 | early death, good score, low risk |
| Allogenic | Hodgkins | 77 | dead | 60 | 102 | high risk grp, long wait, poor score |
| Allogenic | Hodgkins | 79 | dead | 70 | 71 | high risk grp, short wait, poor score |
| Autologous | Non-Hodgkins | 53 | dead | 90 | 17 | early death, good score, med risk grp |

7. Building Cox Proportional Hazards models

| gtype | dtype | time | delta | score | wtime | comment |
|------------|----------|------|-------|-------|-------|--|
| Autologous | Hodgkins | 30 | dead | 90 | 73 | early death, good score, med risk grp |

7.6.5. Action Items

- Unusual points may need checking, particularly if the data are not completely cleaned. In this case, observations 15 and 16 may show some trouble with the dichotomization of waiting time, but it still may be useful.
- The two largest residuals seem to be due to unexpectedly early deaths, but unfortunately this can occur.
- If hazards don't look proportional, then we may need to use strata, between which the base hazards are permitted to be different. For this problem, the natural strata are the two diseases, because they could need to be managed differently anyway.
- A main point that we want to be sure of is the relative risk difference by disease type and graft type.

```
hodg.cox2 |>
  predict(
    reference = "zero",
    newdata = means |>
      mutate(
        wt2 = "short",
        score = 0),
    type = "lp") |>
  data.frame('linear predictor' = _) |>
  pander()
```

7. Building Cox Proportional Hazards models

Table 7.8.: Linear Risk Predictors for Lymphoma

| | linear.predictor |
|------------------------|------------------|
| Non-Hodgkins,Allogenic | 0 |
| Non-Hodgkins,Allogenic | 0.6651 |
| Hodgkins,Autologous | |
| Hodgkins,Allogenic | 2.327 |
| Hodgkins,Autologous | 0.9256 |

For Non-Hodgkin's, the allogenic graft is better. For Hodgkin's, the autologous graft is much better.

7.7. Stratified survival models

7.7.1. Revisiting the leukemia dataset (anderson)

We will analyze remission survival times on 42 leukemia patients, half on new treatment, half on standard treatment.

This is the same data as the `drug6mp` data from `KMsurv`, but with two other variables and without the pairing. This version comes from the Kleinbaum and Klein survival textbook (e.g., p281):

```
anderson =  
  paste0(  
    "http://web1.sph.emory.edu/dkleinb/allDatasets/",  
    "surv2datasets/anderson.dta") |>  
  haven::read_dta() |>  
  mutate(  
    status = status |>  
    case_match(  
      "standard" ~ 1,  
      "new" ~ 2  
    )  
  )
```

7. Building Cox Proportional Hazards models

```
    1 ~ "relapse",
    0 ~ "censored"
  ),

  sex = sex |>
    case_match(
      0 ~ "female",
      1 ~ "male"
    ) |>
    factor() |>
    relevel(ref = "female"),

  rx = rx |>
    case_match(
      0 ~ "new",
      1 ~ "standard"
    ) |>
    factor() |> relevel(ref = "standard"),

  surv = Surv(
    time = survt,
    event = (status == "relapse"))
)

print(anderson)
```

7.7.2. Cox semi-parametric proportional hazards model

```
anderson.cox1 = coxph(
  formula = surv ~ rx + sex + logwbc,
  data = anderson)
```

7. Building Cox Proportional Hazards models

```
summary(anderson.cox1)
```

Call:

```
coxph(formula = surv ~ rx + sex + logwbc, data = anderson)
```

n= 42, number of events= 30

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|---------|--------|-----------|----------|-------|-------------|
| rxnew | -1.504 | 0.222 | 0.462 | -3.26 | 0.0011 ** |
| sexmale | 0.315 | 1.370 | 0.455 | 0.69 | 0.4887 |
| logwbc | 1.682 | 5.376 | 0.337 | 5.00 | 5.8e-07 *** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|---------|-----------|------------|-----------|-----------|
| rxnew | 0.222 | 4.498 | 0.090 | 0.549 |
| sexmale | 1.370 | 0.730 | 0.562 | 3.338 |
| logwbc | 5.376 | 0.186 | 2.779 | 10.398 |

Concordance= 0.851 (se = 0.041)

Likelihood ratio test= 47.2 on 3 df, p=3e-10

Wald test = 33.5 on 3 df, p=2e-07

Score (logrank) test = 48 on 3 df, p=2e-10

7.7.2.1. Test the proportional hazards assumption

```
cox.zph(anderson.cox1)
```


| | chisq | df | p |
|----|-------|----|------|
| rx | 0.036 | 1 | 0.85 |

7. Building Cox Proportional Hazards models

```
sex      5.420  1 0.02
logwbc   0.142  1 0.71
GLOBAL   5.879  3 0.12
```

7.7.2.2. Graph the K-M survival curves

```
anderson_km_model = survfit(  
  formula = surv ~ sex,  
  data = anderson)  
  
anderson_km_model |>  
  autoplot(conf.int = FALSE) +  
  theme_bw() +  
  theme(legend.position="bottom")
```



7. Building Cox Proportional Hazards models

The survival curves cross, which indicates a problem in the proportionality assumption by sex.

7.7.3. Graph the Nelson-Aalen cumulative hazard

We can also look at the log-hazard (“cloglog survival”) plots:

```
anderson_na_model = survfit(  
  formula = surv ~ sex,  
  data = anderson,  
  type = "fleming")  
  
anderson_na_model |>  
  autoplot(  
    fun = "cumhaz",  
    conf.int = FALSE) +  
  theme_classic() +  
  theme(legend.position="bottom") +  
  ylab("log(Cumulative Hazard)") +  
  scale_y_continuous(  
    trans = "log10",  
    name = "Cumulative hazard (H(t), log scale)") +  
  scale_x_continuous(  
    breaks = c(1,2,5,10,20,50),  
    trans = "log"  
  )  
)
```


7. Building Cox Proportional Hazards models




Figure 7.16.: Cumulative hazard (cloglog scale) for `anderson` data

This can be fixed by using strata or possibly by other model alterations.

7.7.4. The Stratified Cox Model

- In a stratified Cox model, each stratum, defined by one or more factors, has its own base survival function $h_0(t)$.
- But the coefficients for each variable not used in the strata definitions are assumed to be the same across strata.
- To check if this assumption is reasonable one can include interactions with strata and see if they are significant (this may generate a warning and NA lines but these can be ignored).
- Since the `sex` variable shows possible non-proportionality, we try stratifying on `sex`.

7. Building Cox Proportional Hazards models

```
anderson.coxph.strat =  
coxph(  
  formula =  
    surv ~ rx + logwbc + strata(sex),  
  data = anderson)  
  
summary(anderson.coxph.strat)
```

Call:

```
coxph(formula = surv ~ rx + logwbc + strata(sex), data = anderson)
```

n= 42, number of events= 30

| | coef | exp(coef) | se(coef) | z | Pr(> z) | |
|--------|--------|-----------|----------|-------|----------|-----|
| rxnew | -0.998 | 0.369 | 0.474 | -2.11 | 0.035 | * |
| logwbc | 1.454 | 4.279 | 0.344 | 4.22 | 2.4e-05 | *** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|--------|-----------|------------|-----------|-----------|
| rxnew | 0.369 | 2.713 | 0.146 | 0.932 |
| logwbc | 4.279 | 0.234 | 2.180 | 8.398 |

Concordance= 0.812 (se = 0.059)

Likelihood ratio test= 32.1 on 2 df, p=1e-07

Wald test = 22.8 on 2 df, p=1e-05

Score (logrank) test = 30.8 on 2 df, p=2e-07

Let's compare this to a model fit only on the subset of males:

```
anderson.coxph.male =  
coxph(  
  formula =
```

7. Building Cox Proportional Hazards models

```
formula = surv ~ rx + logwbc,  
subset = sex == "male",  
data = anderson)  
  
summary(anderson.coxph.male)
```

Call:

```
coxph(formula = surv ~ rx + logwbc, data = anderson, subset = sex ==  
      "male")
```

n= 20, number of events= 14

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|--------|--------|-----------|----------|-------|-----------|
| rxnew | -1.978 | 0.138 | 0.739 | -2.68 | 0.0075 ** |
| logwbc | 1.743 | 5.713 | 0.536 | 3.25 | 0.0011 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|--------|-----------|------------|-----------|-----------|
| rxnew | 0.138 | 7.227 | 0.0325 | 0.589 |
| logwbc | 5.713 | 0.175 | 1.9991 | 16.328 |

Concordance= 0.905 (se = 0.043)

Likelihood ratio test= 29.2 on 2 df, p=5e-07

Wald test = 15.3 on 2 df, p=5e-04

Score (logrank) test = 26.4 on 2 df, p=2e-06

```
anderson.coxph.female =  
coxph(  
  formula =  
    surv ~ rx + logwbc,  
  subset = sex == "female",
```

7. Building Cox Proportional Hazards models

```
data = anderson)

summary(anderson.coxph.female)
```

Call:

```
coxph(formula = surv ~ rx + logwbc, data = anderson, subset = sex ==
      "female")
```

n= 22, number of events= 16

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|--------|--------|-----------|----------|-------|----------|
| rxnew | -0.311 | 0.733 | 0.564 | -0.55 | 0.581 |
| logwbc | 1.206 | 3.341 | 0.503 | 2.40 | 0.017 * |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|--------|-----------|------------|-----------|-----------|
| rxnew | 0.733 | 1.365 | 0.243 | 2.21 |
| logwbc | 3.341 | 0.299 | 1.245 | 8.96 |

Concordance= 0.692 (se = 0.085)

Likelihood ratio test= 6.65 on 2 df, p=0.04

Wald test = 6.36 on 2 df, p=0.04

Score (logrank) test = 6.74 on 2 df, p=0.03

The coefficients of treatment look different. Are they statistically different?

```
anderson.coxph.strat.intxn =
coxph(
  formula = surv ~ strata(sex) * (rx + logwbc),
  data = anderson)
```

7. Building Cox Proportional Hazards models

```
anderson.coxph.strat.intxn |> summary()
```

Call:

```
coxph(formula = surv ~ strata(sex) * (rx + logwbc), data = anderson)
```

n= 42, number of events= 30

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|------------------------|--------|-----------|----------|-------|----------|
| rxnew | -0.311 | 0.733 | 0.564 | -0.55 | 0.581 |
| logwbc | 1.206 | 3.341 | 0.503 | 2.40 | 0.017 * |
| strata(sex)male:rxnew | -1.667 | 0.189 | 0.930 | -1.79 | 0.073 . |
| strata(sex)male:logwbc | 0.537 | 1.710 | 0.735 | 0.73 | 0.465 |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|------------------------|-----------|------------|-----------|-----------|
| rxnew | 0.733 | 1.365 | 0.2427 | 2.21 |
| logwbc | 3.341 | 0.299 | 1.2452 | 8.96 |
| strata(sex)male:rxnew | 0.189 | 5.294 | 0.0305 | 1.17 |
| strata(sex)male:logwbc | 1.710 | 0.585 | 0.4048 | 7.23 |

Concordance= 0.797 (se = 0.058)

Likelihood ratio test= 35.8 on 4 df, p=3e-07

Wald test = 21.7 on 4 df, p=2e-04

Score (logrank) test = 33.1 on 4 df, p=1e-06

```
anova(  
  anderson.coxph.strat.intxn,  
  anderson.coxph.strat)
```

7. Building Cox Proportional Hazards models

| loglik | Chisq | Df | Pr(> Chi) |
|--------|-------|----|------------|
| -53.85 | NA | NA | NA |
| -55.73 | 3.766 | 2 | 0.1521 |

We don't have enough evidence to tell the difference between these two models.

7.7.5. Conclusions

- We chose to use a stratified model because of the apparent non-proportionality of the hazard for the sex variable.
- When we fit interactions with the strata variable, we did not get an improved model (via the likelihood ratio test).
- So we use the stratified model with coefficients that are the same across strata.

7.7.6. Another Modeling Approach

- We used an additive model without interactions and saw that we might need to stratify by sex.
- Instead, we could try to improve the model's functional form - maybe the interaction of treatment and sex is real, and after fitting that we might not need separate hazard functions.
- Either approach may work.

```
anderson.coxph.intxn =  
  coxph(  
    formula = surv ~ (rx + logwbc) * sex,  
    data = anderson)  
  
anderson.coxph.intxn |> summary()
```

7. Building Cox Proportional Hazards models

Call:

```
coxph(formula = surv ~ (rx + logwbc) * sex, data = anderson)
```

n= 42, number of events= 30

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|----------------|---------|-----------|----------|-------|----------|
| rxnew | -0.3748 | 0.6874 | 0.5545 | -0.68 | 0.499 |
| logwbc | 1.0637 | 2.8971 | 0.4726 | 2.25 | 0.024 * |
| sexmale | -2.8052 | 0.0605 | 2.0323 | -1.38 | 0.167 |
| rxnew:sexmale | -2.1782 | 0.1132 | 0.9109 | -2.39 | 0.017 * |
| logwbc:sexmale | 1.2303 | 3.4223 | 0.6301 | 1.95 | 0.051 . |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|----------------|-----------|------------|-----------|-----------|
| rxnew | 0.6874 | 1.455 | 0.23185 | 2.038 |
| logwbc | 2.8971 | 0.345 | 1.14730 | 7.315 |
| sexmale | 0.0605 | 16.531 | 0.00113 | 3.248 |
| rxnew:sexmale | 0.1132 | 8.830 | 0.01899 | 0.675 |
| logwbc:sexmale | 3.4223 | 0.292 | 0.99539 | 11.766 |

Concordance= 0.861 (se = 0.036)

Likelihood ratio test= 57 on 5 df, p=5e-11

Wald test = 35.6 on 5 df, p=1e-06

Score (logrank) test = 57.1 on 5 df, p=5e-11

```
cox.zph(anderson.coxph.intxn)
```

| | chisq | df | p |
|--------|-------|----|------|
| rx | 0.136 | 1 | 0.71 |
| logwbc | 1.652 | 1 | 0.20 |
| sex | 1.266 | 1 | 0.26 |
| rx:sex | 0.149 | 1 | 0.70 |

7. Building Cox Proportional Hazards models

```
logwbc:sex 0.102 1 0.75  
GLOBAL      3.747 5 0.59
```

7.8. Time-varying covariates

(adapted from Klein, Moeschberger, et al. (2003), §9.2)

7.8.1. Motivating example: back to the leukemia dataset

```
# load the data:  
data(bmt, package = 'KMsurv')  
bmt |> as_tibble() |> print(n = 5)
```

```
# A tibble: 137 x 22  
  group   t1    t2    d1    d2    d3   ta    da    tc    dc    tp    dp    z1  
  <int> <int> <int> <int> <int> <int> <int> <int> <int> <int> <int> <int> <int>  
1     1  2081  2081     0     0     0    67     1   121     1    13     1    26  
2     1  1602  1602     0     0     0  1602     0   139     1    18     1    21  
3     1  1496  1496     0     0     0  1496     0  307     1    12     1    26  
4     1  1462  1462     0     0     0    70     1    95     1    13     1    17  
5     1  1433  1433     0     0     0  1433     0  236     1    12     1    32  
# i 132 more rows  
# i 9 more variables: z2 <int>, z3 <int>, z4 <int>, z5 <int>, z6 <int>,  
#   z7 <int>, z8 <int>, z9 <int>, z10 <int>
```

This dataset comes from the Copelan et al. (1991) study of allogenic bone marrow transplant therapy for acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL).

7. Building Cox Proportional Hazards models

Outcomes (endpoints)

- The main endpoint is disease-free survival (**t2** and **d3**) for the three risk groups, “ALL”, “AML Low Risk”, and “AML High Risk”.

Possible intermediate events

- graft vs. host disease (**GVHD**), an immunological rejection response to the transplant (bad)
- acute (**AGVHD**)
- chronic (**CGVHD**)
- platelet recovery, a return of platelet count to normal levels (good)

One or the other, both in either order, or neither may occur.

Covariates

- We are interested in possibly using the covariates **z1-z10** to adjust for other factors.
- In addition, the time-varying covariates for acute GVHD, chronic GVHD, and platelet recovery may be useful.

7.8.1.1. Preprocessing

We reformat the data before analysis:

```
# reformat the data:
bmt1 =
  bmt |>
  as_tibble() |>
  mutate(
    id = 1:n(), # will be used to connect multiple records for the same individual
```

7. Building Cox Proportional Hazards models

```
group = group |>
  case_match(
    1 ~ "ALL",
    2 ~ "Low Risk AML",
    3 ~ "High Risk AML") |>
  factor(levels = c("ALL", "Low Risk AML", "High Risk AML")),

`patient age` = z1,

`donor age` = z2,

`patient sex` = z3 |>
  case_match(
    0 ~ "Female",
    1 ~ "Male"),

`donor sex` = z4 |>
  case_match(
    0 ~ "Female",
    1 ~ "Male"),

`Patient CMV Status` = z5 |>
  case_match(
    0 ~ "CMV Negative",
    1 ~ "CMV Positive"),

`Donor CMV Status` = z6 |>
  case_match(
    0 ~ "CMV Negative",
    1 ~ "CMV Positive"),

`Waiting Time to Transplant` = z7,
```

7. Building Cox Proportional Hazards models

```
FAB = z8 |>
  case_match(
    1 ~ "Grade 4 Or 5 (AML only)",
    0 ~ "Other") |>
  factor() |>
  relevel(ref = "Other"),

hospital = z9 |> # `z9` is hospital
  case_match(
    1 ~ "Ohio State University",
    2 ~ "Alferd",
    3 ~ "St. Vincent",
    4 ~ "Hahnemann") |>
  factor() |>
  relevel(ref = "Ohio State University"),

MTX = (z10 == 1) # a prophylatic treatment for GVHD

) |>
select(-(z1:z10)) # don't need these anymore

bmt1 |>
  select(group, id:MTX) |>
  print(n = 10)
```

```
# A tibble: 137 x 12
  group   id `patient age` `donor age` `patient sex` `donor sex`
  <fct> <int>         <int>         <int> <chr>         <chr>
1 ALL     1           26           33 Male         Female
2 ALL     2           21           37 Male         Male
3 ALL     3           26           35 Male         Male
4 ALL     4           17           21 Female        Male
5 ALL     5           32           36 Male         Male
```

7. Building Cox Proportional Hazards models

```
6 ALL      6      22      31 Male      Male
7 ALL      7      20      17 Male      Female
8 ALL      8      22      24 Male      Female
9 ALL      9      18      21 Female    Male
10 ALL     10      24      40 Male      Male
# i 127 more rows
# i 6 more variables: `Patient CMV Status` <chr>, `Donor CMV Status` <chr>,
#   `Waiting Time to Transplant` <int>, FAB <fct>, hospital <fct>, MTX <lgl>
```

7.8.2. Time-Dependent Covariates

- A **time-dependent covariate** (“TDC”) is a covariate whose value changes during the course of the study.
- For variables like age that change in a linear manner with time, we can just use the value at the start.
- But it may be plausible that when and if GVHD occurs, the risk of relapse or death increases, and when and if platelet recovery occurs, the risk decreases.

7.8.3. Analysis in R

- We form a variable `precovery` which is = 0 before platelet recovery and is = 1 after platelet recovery, if it occurs.
- For each subject where platelet recovery occurs, we set up multiple records (lines in the data frame); for example one from $t = 0$ to the time of platelet recovery, and one from that time to relapse, recovery, or death.
- We do the same for acute GVHD and chronic GVHD.
- For each record, the covariates are constant.

7. Building Cox Proportional Hazards models

```
bmt2 = bmt1 |>
  #set up new long-format data set:
  tmerge(bmt1, id = id, tstop = t2) |>

  # the following three steps can be in any order,
  # and will still produce the same result:
  #add aghvd as tdc:
  tmerge(bmt1, id = id, agvhd = tdc(ta)) |>
  #add cghvd as tdc:
  tmerge(bmt1, id = id, cgvhd = tdc(tc)) |>
  #add platelet recovery as tdc:
  tmerge(bmt1, id = id, precovery = tdc(tp))

bmt2 = bmt2 |>
  as_tibble() |>
  mutate(status = as.numeric((tstop == t2) & d3))
# status only = 1 if at end of t2 and not censored
```

Let's see how we've rearranged the first row of the data:

```
bmt1 |>
  dplyr::filter(id == 1) |>
  dplyr::select(id, t1, d1, t2, d2, d3, ta, da, tc, dc, tp, dp)
```

| id | t1 | d1 | t2 | d2 | d3 | ta | da | tc | dc | tp | dp |
|----|------|----|------|----|----|----|----|-----|----|----|----|
| 1 | 2081 | 0 | 2081 | 0 | 0 | 67 | 1 | 121 | 1 | 13 | 1 |

The event times for this individual are:

- $t = 0$ time of transplant
- $tp = 13$ platelet recovery

7. Building Cox Proportional Hazards models

- **ta** = 67 acute GVHD onset
- **tc** = 121 chronic GVHD onset
- **t2** = 2081 end of study, patient not relapsed or dead

After converting the data to long-format, we have:

```
bmt2 |>
  select(
    id,
    tstart,
    tstop,
    agvhd,
    cgvhd,
    precovery,
    status
  ) |>
  dplyr::filter(id == 1)
```

| id | tstart | tstop | agvhd | cgvhd | precovery | status |
|----|--------|-------|-------|-------|-----------|--------|
| 1 | 0 | 13 | 0 | 0 | 0 | 0 |
| 1 | 13 | 67 | 0 | 0 | 1 | 0 |
| 1 | 67 | 121 | 1 | 0 | 1 | 0 |
| 1 | 121 | 2081 | 1 | 1 | 1 | 0 |

Note that **status** could have been 1 on the last row, indicating that relapse or death occurred; since it is false, the participant must have exited the study without experiencing relapse or death (i.e., they were censored).

7.8.4. Event sequences

Let:

7. Building Cox Proportional Hazards models

- A = acute GVHD
- C = chronic GVHD
- P = platelet recovery

Each of the eight possible combinations of A or not-A, with C or not-C, with P or not-P occurs in this data set.

- A always occurs before C, and P always occurs before C, if both occur.
- Thus there are ten event sequences in the data set: None, A, C, P, AC, AP, PA, PC, APC, and PAC.
- In general, there could be as many as $1+3+(3)(2)+6 = 16$ sequences, but our domain knowledge tells us that some are missing: CA, CP, CAP, CPA, PCA, PC, PAC
- Different subjects could have 1, 2, 3, or 4 intervals, depending on which of acute GVHD, chronic GVHD, and/or platelet recovery occurred.
- The final interval for any subject has `status = 1` if the subject relapsed or died at that time; otherwise `status = 0`.
- Any earlier intervals have `status = 0`.
- Even though there might be multiple lines per ID in the dataset, there is never more than one event, so no alterations need be made in the estimation procedures or in the interpretation of the output.
- The function `tmerge` in the `survival` package eases the process of constructing the new long-format dataset.

7.8.5. Model with Time-Fixed Covariates

```
bmt1 =  
  bmt1 |>  
  mutate(surv = Surv(t2,d3))
```

7. Building Cox Proportional Hazards models

```
bmt_coxph_TF = coxph(
  formula = surv ~ group + `patient age` * `donor age` + FAB,
  data = bmt1)
summary(bmt_coxph_TF)
```

Call:

```
coxph(formula = surv ~ group + `patient age` * `donor age` +
      FAB, data = bmt1)
```

n= 137, number of events= 83

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|----------------------------|-----------|-----------|----------|-------|-------------|
| groupLow Risk AML | -1.090648 | 0.335999 | 0.354279 | -3.08 | 0.00208 ** |
| groupHigh Risk AML | -0.403905 | 0.667707 | 0.362777 | -1.11 | 0.26555 |
| `patient age` | -0.081639 | 0.921605 | 0.036107 | -2.26 | 0.02376 * |
| `donor age` | -0.084587 | 0.918892 | 0.030097 | -2.81 | 0.00495 ** |
| FABGrade 4 Or 5 (AML only) | 0.837416 | 2.310388 | 0.278464 | 3.01 | 0.00264 ** |
| `patient age`:`donor age` | 0.003159 | 1.003164 | 0.000951 | 3.32 | 0.00089 *** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|----------------------------|-----------|------------|-----------|-----------|
| groupLow Risk AML | 0.336 | 2.976 | 0.168 | 0.673 |
| groupHigh Risk AML | 0.668 | 1.498 | 0.328 | 1.360 |
| `patient age` | 0.922 | 1.085 | 0.859 | 0.989 |
| `donor age` | 0.919 | 1.088 | 0.866 | 0.975 |
| FABGrade 4 Or 5 (AML only) | 2.310 | 0.433 | 1.339 | 3.988 |
| `patient age`:`donor age` | 1.003 | 0.997 | 1.001 | 1.005 |

Concordance= 0.665 (se = 0.033)

Likelihood ratio test= 32.8 on 6 df, p=1e-05

Wald test = 33 on 6 df, p=1e-05

Score (logrank) test = 35.8 on 6 df, p=3e-06

7. Building Cox Proportional Hazards models

```
drop1(bmt_coxph_TF, test = "Chisq")
```

| | Df | AIC | LRT | Pr(>Chi) |
|-----------------------|----|-------|--------|----------|
| | NA | 725.8 | NA | NA |
| group | 2 | 734.3 | 12.511 | 0.0019 |
| FAB | 1 | 733.0 | 9.216 | 0.0024 |
| patient age:donor age | 1 | 733.3 | 9.514 | 0.0020 |

```
bmt1$mres =
  bmt_coxph_TF |>
  update(. ~ . - `donor age`) |>
  residuals(type="martingale")

bmt1 |>
  ggplot(aes(x = `donor age`, y = mres)) +
  geom_point() +
  geom_smooth(method = "loess", aes(col = "loess")) +
  geom_smooth(method = 'lm', aes(col = "lm")) +
  theme_classic() +
  xlab("Donor age") +
  ylab("Martingale Residuals") +
  guides(col=guide_legend(title = ""))
```

7. Building Cox Proportional Hazards models




Figure 7.17.: Martingale residuals for Donor age

A more complex functional form for donor age seems warranted; left as an exercise for the reader.

Now we will add the time-varying covariates:

```
# add counting process formulation of Surv():
bmt2 =
  bmt2 |>
  mutate(
    surv =
      Surv(
        time = tstart,
        time2 = tstop,
```

7. Building Cox Proportional Hazards models

```
event = status,  
type = "counting"))
```

Let's see how the data looks for patient 15:

```
bmt1 |> dplyr::filter(id == 15) |> dplyr::select(tp, dp, tc,dc, ta, da, FAB, surv, t
```

| tp | dp | tc | dc | ta | da | FAB | surv | t1 | d1 | t2 | d2 | d3 |
|----|----|-----|----|-----|----|-------|------|-----|----|-----|----|----|
| 21 | 1 | 220 | 1 | 418 | 0 | Other | 418 | 418 | 1 | 418 | 0 | 1 |

```
bmt2 |> dplyr::filter(id == 15) |> dplyr::select(id, agvhd, cgvhhd, precovery, surv)
```

| id | agvhd | cgvhhd | precovery | surv |
|----|-------|--------|-----------|------------|
| 15 | 0 | 0 | 0 | (0, 21+] |
| 15 | 0 | 0 | 1 | (21,220+] |
| 15 | 0 | 1 | 1 | (220,418] |

7.8.6. Model with Time-Dependent Covariates

```
bmt_coxph_TV = coxph(  
  formula =  
    surv ~  
    group + `patient age`*`donor age` + FAB + agvhd + cgvhhd + precovery,  
  data = bmt2)  
  
summary(bmt_coxph_TV)
```

7. Building Cox Proportional Hazards models

Call:

```
coxph(formula = surv ~ group + `patient age` * `donor age` +
      FAB + agvhd + cgvhhd + precovery, data = bmt2)
```

n= 341, number of events= 83

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|----------------------------|-----------|-----------|----------|-------|-----------|
| groupLow Risk AML | -1.038514 | 0.353980 | 0.358220 | -2.90 | 0.0037 ** |
| groupHigh Risk AML | -0.380481 | 0.683533 | 0.374867 | -1.01 | 0.3101 |
| `patient age` | -0.073351 | 0.929275 | 0.035956 | -2.04 | 0.0413 * |
| `donor age` | -0.076406 | 0.926440 | 0.030196 | -2.53 | 0.0114 * |
| FABGrade 4 Or 5 (AML only) | 0.805700 | 2.238263 | 0.284273 | 2.83 | 0.0046 ** |
| agvhd | 0.150565 | 1.162491 | 0.306848 | 0.49 | 0.6237 |
| cgvhhd | -0.116136 | 0.890354 | 0.289046 | -0.40 | 0.6878 |
| precovery | -0.941123 | 0.390190 | 0.347861 | -2.71 | 0.0068 ** |
| `patient age`:`donor age` | 0.002895 | 1.002899 | 0.000944 | 3.07 | 0.0022 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|----------------------------|-----------|------------|-----------|-----------|
| groupLow Risk AML | 0.354 | 2.825 | 0.175 | 0.714 |
| groupHigh Risk AML | 0.684 | 1.463 | 0.328 | 1.425 |
| `patient age` | 0.929 | 1.076 | 0.866 | 0.997 |
| `donor age` | 0.926 | 1.079 | 0.873 | 0.983 |
| FABGrade 4 Or 5 (AML only) | 2.238 | 0.447 | 1.282 | 3.907 |
| agvhd | 1.162 | 0.860 | 0.637 | 2.121 |
| cgvhhd | 0.890 | 1.123 | 0.505 | 1.569 |
| precovery | 0.390 | 2.563 | 0.197 | 0.772 |
| `patient age`:`donor age` | 1.003 | 0.997 | 1.001 | 1.005 |

Concordance= 0.702 (se = 0.028)

Likelihood ratio test= 40.3 on 9 df, p=7e-06

Wald test = 42.4 on 9 df, p=3e-06

Score (logrank) test = 47.2 on 9 df, p=4e-07

7. Building Cox Proportional Hazards models

Platelet recovery is highly significant.

Neither acute GVHD (agvhd) nor chronic GVHD (cgvhd) has a statistically significant effect here, nor are they significant in models with the other one removed.

```
update(bmt_coxph_TV, .~-agvhd) |> summary()
```

Call:

```
coxph(formula = surv ~ group + `patient age` + `donor age` +  
      FAB + cgvhd + precovery + `patient age`:`donor age`, data = bmt2)
```

n= 341, number of events= 83

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|----------------------------|-----------|-----------|----------|-------|-----------|
| groupLow Risk AML | -1.049870 | 0.349983 | 0.356727 | -2.94 | 0.0032 ** |
| groupHigh Risk AML | -0.417049 | 0.658988 | 0.365348 | -1.14 | 0.2537 |
| `patient age` | -0.070749 | 0.931696 | 0.035477 | -1.99 | 0.0461 * |
| `donor age` | -0.075693 | 0.927101 | 0.030075 | -2.52 | 0.0118 * |
| FABGrade 4 Or 5 (AML only) | 0.807035 | 2.241253 | 0.283437 | 2.85 | 0.0044 ** |
| cgvhd | -0.095393 | 0.909015 | 0.285979 | -0.33 | 0.7387 |
| precovery | -0.983653 | 0.373942 | 0.338170 | -2.91 | 0.0036 ** |
| `patient age`:`donor age` | 0.002859 | 1.002863 | 0.000936 | 3.05 | 0.0023 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|----------------------------|-----------|------------|-----------|-----------|
| groupLow Risk AML | 0.350 | 2.857 | 0.174 | 0.704 |
| groupHigh Risk AML | 0.659 | 1.517 | 0.322 | 1.349 |
| `patient age` | 0.932 | 1.073 | 0.869 | 0.999 |
| `donor age` | 0.927 | 1.079 | 0.874 | 0.983 |
| FABGrade 4 Or 5 (AML only) | 2.241 | 0.446 | 1.286 | 3.906 |
| cgvhd | 0.909 | 1.100 | 0.519 | 1.592 |
| precovery | 0.374 | 2.674 | 0.193 | 0.726 |

7. Building Cox Proportional Hazards models

```
`patient age`:`donor age`      1.003      0.997      1.001      1.005
```

```
Concordance= 0.701 (se = 0.027 )
Likelihood ratio test= 40 on 8 df, p=3e-06
Wald test              = 42.4 on 8 df, p=1e-06
Score (logrank) test = 47.2 on 8 df, p=1e-07
```

```
update(bmt_coxph_TV, .~-cgvhd) |> summary()
```

Call:

```
coxph(formula = surv ~ group + `patient age` + `donor age` +
      FAB + agvhd + precovery + `patient age`:`donor age`, data = bmt2)
```

```
n= 341, number of events= 83
```

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|----------------------------|-----------|-----------|----------|-------|-----------|
| groupLow Risk AML | -1.019638 | 0.360725 | 0.355311 | -2.87 | 0.0041 ** |
| groupHigh Risk AML | -0.381356 | 0.682935 | 0.374568 | -1.02 | 0.3086 |
| `patient age` | -0.073189 | 0.929426 | 0.035890 | -2.04 | 0.0414 * |
| `donor age` | -0.076753 | 0.926118 | 0.030121 | -2.55 | 0.0108 * |
| FABGrade 4 Or 5 (AML only) | 0.811716 | 2.251769 | 0.284012 | 2.86 | 0.0043 ** |
| agvhd | 0.131621 | 1.140676 | 0.302623 | 0.43 | 0.6636 |
| precovery | -0.946697 | 0.388021 | 0.347265 | -2.73 | 0.0064 ** |
| `patient age`:`donor age` | 0.002904 | 1.002908 | 0.000943 | 3.08 | 0.0021 ** |

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|----------------------------|-----------|------------|-----------|-----------|
| groupLow Risk AML | 0.361 | 2.772 | 0.180 | 0.724 |
| groupHigh Risk AML | 0.683 | 1.464 | 0.328 | 1.423 |
| `patient age` | 0.929 | 1.076 | 0.866 | 0.997 |
| `donor age` | 0.926 | 1.080 | 0.873 | 0.982 |
| FABGrade 4 Or 5 (AML only) | 2.252 | 0.444 | 1.291 | 3.929 |

7. Building Cox Proportional Hazards models

| | | | | |
|---------------------------|-------|-------|-------|-------|
| agvhd | 1.141 | 0.877 | 0.630 | 2.064 |
| precovery | 0.388 | 2.577 | 0.196 | 0.766 |
| `patient age`:`donor age` | 1.003 | 0.997 | 1.001 | 1.005 |

Concordance= 0.701 (se = 0.027)
 Likelihood ratio test= 40.1 on 8 df, p=3e-06
 Wald test = 42.1 on 8 df, p=1e-06
 Score (logrank) test = 47.1 on 8 df, p=1e-07

Let's drop them both:

```
bmt_coxph_TV2 = update(bmt_coxph_TV, . ~ . - agvhd -cgvhd)
bmt_coxph_TV2 |> summary()
```

Call:

```
coxph(formula = surv ~ group + `patient age` + `donor age` +
      FAB + precovery + `patient age`:`donor age`, data = bmt2)
```

n= 341, number of events= 83

| | coef | exp(coef) | se(coef) | z | Pr(> z) | |
|----------------------------|-----------|-----------|----------|-------|----------|----|
| groupLow Risk AML | -1.032520 | 0.356108 | 0.353202 | -2.92 | 0.0035 | ** |
| groupHigh Risk AML | -0.413888 | 0.661075 | 0.365209 | -1.13 | 0.2571 | |
| `patient age` | -0.070965 | 0.931495 | 0.035453 | -2.00 | 0.0453 | * |
| `donor age` | -0.076052 | 0.926768 | 0.030007 | -2.53 | 0.0113 | * |
| FABGrade 4 Or 5 (AML only) | 0.811926 | 2.252242 | 0.283231 | 2.87 | 0.0041 | ** |
| precovery | -0.983505 | 0.373998 | 0.337997 | -2.91 | 0.0036 | ** |
| `patient age`:`donor age` | 0.002872 | 1.002876 | 0.000936 | 3.07 | 0.0021 | ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|-------------------|-----------|------------|-----------|-----------|
| groupLow Risk AML | 0.356 | 2.808 | 0.178 | 0.712 |

7. Building Cox Proportional Hazards models

| | | | | |
|----------------------------|-------|-------|-------|-------|
| groupHigh Risk AML | 0.661 | 1.513 | 0.323 | 1.352 |
| `patient age` | 0.931 | 1.074 | 0.869 | 0.999 |
| `donor age` | 0.927 | 1.079 | 0.874 | 0.983 |
| FABGrade 4 Or 5 (AML only) | 2.252 | 0.444 | 1.293 | 3.924 |
| precovery | 0.374 | 2.674 | 0.193 | 0.725 |
| `patient age`:`donor age` | 1.003 | 0.997 | 1.001 | 1.005 |

Concordance= 0.7 (se = 0.027)

Likelihood ratio test= 39.9 on 7 df, p=1e-06

Wald test = 42.2 on 7 df, p=5e-07

Score (logrank) test = 47.1 on 7 df, p=5e-08

7.9. Recurrent Events

(Adapted from Kleinbaum and Klein, Ch 8)

- Sometimes an appropriate analysis requires consideration of recurrent events.
- A patient with arthritis may have more than one flareup. The same is true of many recurring-remitting diseases.
- In this case, we have more than one line in the data frame, but each line may have an event.
- We have to use a “robust” variance estimator to account for correlation of time-to-events within a patient.

7.9.1. Bladder Cancer Data Set

The bladder cancer dataset from Kleinbaum and Klein contains recurrent event outcome information for eighty-six cancer patients followed for the recurrence of bladder cancer tumor after transurethral surgical excision (Byar and Green 1980). The exposure of interest is the effect of the drug

7. Building Cox Proportional Hazards models

treatment of thiotepa. Control variables are the initial number and initial size of tumors. The data layout is suitable for a counting processes approach.

This drug is still a possible choice for some patients. Another therapeutic choice is Bacillus Calmette-Guerin (BCG), a live bacterium related to cow tuberculosis.

7.9.1.1. Data dictionary

Table 7.15.: Variables in the `bladder` dataset

| Variable | Definition |
|-----------------------|--|
| <code>id</code> | Patient unique ID |
| <code>status</code> | for each time interval: 1 = recurred, 0 = censored |
| <code>interval</code> | 1 = first recurrence, etc. |
| <code>intime</code> | <code>'tstop - tstart</code> (all times in months) |
| <code>tstart</code> | start of interval |
| <code>tstop</code> | end of interval |
| <code>tx</code> | treatment code, 1 = thiotepa |
| <code>num</code> | number of initial tumors |
| <code>size</code> | size of initial tumors (cm) |

- There are 85 patients and 190 lines in the dataset, meaning that many patients have more than one line.
- Patient 1 with 0 observation time was removed.
- Of the 85 patients, 47 had at least one recurrence and 38 had none.
- 18 patients had exactly one recurrence.
- There were up to 4 recurrences in a patient.
- Of the 190 intervals, 112 terminated with a recurrence and 78 were censored.

7. Building Cox Proportional Hazards models

7.9.1.2. Different intervals for the same patient are correlated.

- Is the effective sample size 47 or 112? This might narrow confidence intervals by as much as a factor of $\sqrt{112/47} = 1.54$
- What happens if I have 5 treatment and 5 control values and want to do a t-test and I then duplicate the 10 values as if the sample size was 20? This falsely narrows confidence intervals by a factor of $\sqrt{2} = 1.41$.

```
bladder =  
  paste0(  
    "http://web1.sph.emory.edu/dkleinb/allDatasets",  
    "/surv2datasets/bladder.dta") |>  
  read_dta() |>  
  as_tibble()  
  
bladder = bladder[-1,] #remove subject with 0 observation time  
print(bladder)
```

```
bladder =  
  bladder |>  
  mutate(  
    surv =  
      Surv(  
        time = start,  
        time2 = stop,  
        event = event,  
        type = "counting"))  
  
bladder.cox1 = coxph(  
  formula = surv~tx+num+size,  
  data = bladder)
```

7. Building Cox Proportional Hazards models

```
#results with biased variance-covariance matrix:  
summary(bladder.cox1)
```

Call:

```
coxph(formula = surv ~ tx + num + size, data = bladder)
```

```
n= 190, number of events= 112
```

| | coef | exp(coef) | se(coef) | z | Pr(> z) |
|------|---------|-----------|----------|-------|-------------|
| tx | -0.4116 | 0.6626 | 0.1999 | -2.06 | 0.03947 * |
| num | 0.1637 | 1.1778 | 0.0478 | 3.43 | 0.00061 *** |
| size | -0.0411 | 0.9598 | 0.0703 | -0.58 | 0.55897 |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|------|-----------|------------|-----------|-----------|
| tx | 0.663 | 1.509 | 0.448 | 0.98 |
| num | 1.178 | 0.849 | 1.073 | 1.29 |
| size | 0.960 | 1.042 | 0.836 | 1.10 |

Concordance= 0.624 (se = 0.032)

Likelihood ratio test= 14.7 on 3 df, p=0.002

Wald test = 15.9 on 3 df, p=0.001

Score (logrank) test = 16.2 on 3 df, p=0.001

Note

The likelihood ratio and score tests assume independence of observations within a cluster. The Wald and robust score tests do not.

7. Building Cox Proportional Hazards models

7.9.1.3. adding cluster = id

If we add `cluster= id` to the call to `coxph`, the coefficient estimates don't change, but we get an additional column in the `summary()` output: `robust se`:

```
bladder.cox2 = coxph(
  formula = surv ~ tx + num + size,
  cluster = id,
  data = bladder)

#unbiased though this reduces power:
summary(bladder.cox2)
```

Call:

```
coxph(formula = surv ~ tx + num + size, data = bladder, cluster = id)
```

```
n= 190, number of events= 112
```

| | coef | exp(coef) | se(coef) | robust se | z | Pr(> z) |
|------|---------|-----------|----------|-----------|-------|-----------|
| tx | -0.4116 | 0.6626 | 0.1999 | 0.2488 | -1.65 | 0.0980 . |
| num | 0.1637 | 1.1778 | 0.0478 | 0.0584 | 2.80 | 0.0051 ** |
| size | -0.0411 | 0.9598 | 0.0703 | 0.0742 | -0.55 | 0.5799 |

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|------|-----------|------------|-----------|-----------|
| tx | 0.663 | 1.509 | 0.407 | 1.08 |
| num | 1.178 | 0.849 | 1.050 | 1.32 |
| size | 0.960 | 1.042 | 0.830 | 1.11 |

```
Concordance= 0.624 (se = 0.031 )
```

```
Likelihood ratio test= 14.7 on 3 df, p=0.002
```

7. Building Cox Proportional Hazards models

Wald test = 11.2 on 3 df, p=0.01
Score (logrank) test = 16.2 on 3 df, p=0.001, Robust = 10.8 p=0.01

(Note: the likelihood ratio and score tests assume independence of observations within a cluster, the Wald and robust score tests do not).

robust `se` is larger than `se`, and accounts for the repeated observations from the same individuals:

```
round(bladder.cox2$naive.var, 4)
```

| | | |
|---------|---------|--------|
| 0.0400 | -0.0014 | 0.0000 |
| -0.0014 | 0.0023 | 0.0007 |
| 0.0000 | 0.0007 | 0.0049 |

```
round(bladder.cox2$var, 4)
```

| | | |
|---------|---------|---------|
| 0.0619 | -0.0026 | -0.0004 |
| -0.0026 | 0.0034 | 0.0013 |
| -0.0004 | 0.0013 | 0.0055 |

These are the ratios of correct confidence intervals to naive ones:

```
with(bladder.cox2, diag(var)/diag(naive.var)) |> sqrt()
```

```
[1] 1.244 1.223 1.056
```

We might try dropping the non-significant `size` variable:

7. Building Cox Proportional Hazards models

```
#remove non-significant size variable:  
bladder.cox3 = bladder.cox2 |> update(. ~ . - size)  
summary(bladder.cox3)
```

Call:

```
coxph(formula = surv ~ tx + num, data = bladder, cluster = id)
```

n= 190, number of events= 112

| | coef | exp(coef) | se(coef) | robust se | z | Pr(> z) |
|-----|---------|-----------|----------|-----------|-------|-----------|
| tx | -0.4117 | 0.6625 | 0.2003 | 0.2515 | -1.64 | 0.1017 |
| num | 0.1700 | 1.1853 | 0.0465 | 0.0564 | 3.02 | 0.0026 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

| | exp(coef) | exp(-coef) | lower .95 | upper .95 |
|-----|-----------|------------|-----------|-----------|
| tx | 0.663 | 1.509 | 0.405 | 1.08 |
| num | 1.185 | 0.844 | 1.061 | 1.32 |

Concordance= 0.623 (se = 0.031)

Likelihood ratio test= 14.3 on 2 df, p=8e-04

Wald test = 10.2 on 2 df, p=0.006

Score (logrank) test = 15.8 on 2 df, p=4e-04, Robust = 10.6 p=0.005

(Note: the likelihood ratio and score tests assume independence of observations within a cluster, the Wald and robust score tests do not).

Ways to check PH assumption:

- cloglog
- schoenfeld residuals
- interaction with time

7.10. Age as the time scale

See Canchola et al. (2003).

8. Parametric survival models

Configuring R

Functions from these packages will be used throughout this document:

```
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(dplyr) # manipulate data
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(conflicted) # check for conflicting function definitions
conflicts_prefer(dplyr::filter)
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
knitr::opts_chunk$set(message = FALSE)
pander::panderOptions("table.emphasize.rownames", FALSE)
options('digits' = 4)
```


8.1. Parametric Survival Models

8.1.1. Exponential Distribution

- The exponential distribution is the basic distribution for survival analysis.

$$\begin{aligned}f(t) &= \lambda e^{-\lambda t} \\ \log \{f(t)\} &= \log \{\lambda\} - \lambda t \\ F(t) &= 1 - e^{-\lambda t} \\ S(t) &= e^{-\lambda t} \\ H(t) &= \log \{S(t)\} = -\lambda t \\ h(t) &= \lambda \\ E(T) &= \lambda^{-1}\end{aligned}$$

8.1.2. Weibull Distribution

Using the Kalbfleisch and Prentice (2002) notation:


$$\begin{aligned}f(t) &= \lambda p(\lambda t)^{p-1} e^{-(\lambda t)^p} \\ F(t) &= 1 - e^{-(\lambda t)^p} \\ S(t) &= e^{-(\lambda t)^p} \\ h(t) &= \lambda p(\lambda t)^{p-1} \\ H(t) &= (\lambda t)^p \\ \log \{H(t)\} &= p \log \{\lambda t\} = p \log \{\lambda\} + p \log \{t\} \\ E(T) &= \lambda^{-1} \cdot \Gamma\left(1 + \frac{1}{p}\right)\end{aligned}$$

8. Parametric survival models

i Note

Recall from calculus:

- $\Gamma(t) \stackrel{\text{def}}{=} \int_{u=0}^{\infty} u^{t-1} e^{-u} du$
- $\Gamma(t) = (t-1)!$ for integers $t \in \mathbb{Z}$
- It is implemented by the `gamma()` function in R.



Here are some Weibull density functions, with $\lambda = 1$ and p varying:

```
library(ggplot2)
lambda = 1
ggplot() +
  geom_function(
    aes(col = "0.25"),
    fun = \(x) dweibull(x, shape = 0.25, scale = 1/lambda)) +
```

8. Parametric survival models

```
geom_function(  
  aes(col = "0.5"),  
  fun = \(x) dweibull(x, shape = 0.5, scale = 1/lambda)) +  
geom_function(  
  aes(col = "1"),  
  fun = \(x) dweibull(x, shape = 1, scale = 1/lambda)) +  
geom_function(  
  aes(col = "1.5"),  
  fun = \(x) dweibull(x, shape = 1.5, scale = 1/lambda)) +  
geom_function(  
  aes(col = "2"),  
  fun = \(x) dweibull(x, shape = 2, scale = 1/lambda)) +  
geom_function(  
  aes(col = "5"),  
  fun = \(x) dweibull(x, shape = 5, scale = 1/lambda)) +  
theme_bw() +  
xlim(0, 2.5) +  
ylab("f(t)") +  
theme(axis.title.y = element_text(angle=0)) +  
theme(legend.position="bottom") +  
guides(  
  col =  
    guide_legend(  
      title = "p",  
      label.theme =  
        element_text(  
          size = 12)))
```

8. Parametric survival models




Figure 8.1.: Density functions for Weibull distribution

8.1.2.1. Properties of Weibull hazard functions

- When $p = 1$, the Weibull distribution simplifies to the exponential distribution
- When $p > 1$, the hazard is increasing
- When $p < 1$, the hazard is decreasing

In HW: prove these properties

This distribution provides more flexibility than the exponential.

Here are some Weibull hazard functions, with $\lambda = 1$ and p varying:

8. Parametric survival models

```
library(ggplot2)
library(eha)
lambda = 1

ggplot() +
  geom_function(
    aes(col = "0.25"),
    fun = \(x) hweibull(x, shape = 0.25, scale = 1/lambda)) +
  geom_function(
    aes(col = "0.5"),
    fun = \(x) hweibull(x, shape = 0.5, scale = 1/lambda)) +
  geom_function(
    aes(col = "1"),
    fun = \(x) hweibull(x, shape = 1, scale = 1/lambda)) +
  geom_function(
    aes(col = "1.5"),
    fun = \(x) hweibull(x, shape = 1.5, scale = 1/lambda)) +
  geom_function(
    aes(col = "2"),
    fun = \(x) hweibull(x, shape = 2, scale = 1/lambda)) +
  theme_bw() +
  xlim(0, 2.5) +
  ylab("h(t)") +
  theme(axis.title.y = element_text(angle=0)) +
  theme(legend.position="bottom") +
  guides(
    col =
      guide_legend(
        title = "p",
        label.theme =
          element_text(
            size = 12)))
```

8. Parametric survival models




Figure 8.2.: Hazard functions for Weibull distribution

```
library(ggplot2)
lambda = 1

ggplot() +
  geom_function(
    aes(col = "0.25"),
    fun = \(x) pweibull(lower = FALSE, x, shape = 0.25, scale = 1/lambda)) +
  geom_function(
    aes(col = "0.5"),
    fun = \(x) pweibull(lower = FALSE, x, shape = 0.5, scale = 1/lambda)) +
  geom_function(
    aes(col = "1"),
    fun = \(x) pweibull(lower = FALSE, x, shape = 1, scale = 1/lambda)) +
  geom_function(
```

8. Parametric survival models

```
    aes(col = "1.5"),
    fun = \(x) pweibull(lower = FALSE, x, shape = 1.5, scale = 1/lambda)) +
geom_function(
  aes(col = "2"),
  fun = \(x) pweibull(lower = FALSE, x, shape = 2, scale = 1/lambda)) +
theme_bw() +
xlim(0, 2.5) +
ylab("S(t)") +
theme(axis.title.y = element_text(angle=0)) +
theme(legend.position="bottom") +
guides(
  col =
    guide_legend(
      title = "p",
      label.theme =
        element_text(
          size = 12)))
```

8. Parametric survival models




Figure 8.3.: Survival functions for Weibull distribution

8.1.3. Exponential Regression

For each subject i , define a linear predictor:

$$\begin{aligned}
 \eta(x) &= \beta_0 + (\beta_1 x_1 + \dots + \beta_p x_p) \\
 h(t|x) &= \exp \{ \eta(x) \} \\
 h_0 &\stackrel{\text{def}}{=} h(t|0) \\
 &= \exp \{ \eta(0) \} \\
 &= \exp \{ \beta_0 + (\beta_1 \cdot 0 + \dots + \beta_p \cdot 0) \} \\
 &= \exp \{ \beta_0 + 0 \} \\
 &= \exp \{ \beta_0 \}
 \end{aligned}$$

8. Parametric survival models

We let the linear predictor have a constant term, and when there are no additional predictors the hazard is $\lambda = \exp\{\beta_0\}$. This has a log link as in a generalized linear model. Since the hazard does not depend on t , the hazards are (trivially) proportional.

8.1.4. Accelerated Failure Time

Previously, we assumed the hazards were proportional; that is, the covariates multiplied the baseline hazard function:

$$\begin{aligned} h(T = t|X = x) &\stackrel{\text{def}}{=} p(T = t|X = x, T \geq t) \\ &= h(t|X = 0) \cdot \exp\{\eta(x)\} \\ &= h(t|X = 0) \cdot \theta(x) \\ &= h_0(t) \cdot \theta(x) \end{aligned}$$

and correspondingly,

$$\begin{aligned} H(t|x) &= \theta(x)H_0(t) \\ S(t|x) &= \exp\{-H(t|x)\} \\ &= \exp\{-\theta(x) \cdot H_0(t)\} \\ &= (\exp\{-H_0(t)\})^{\theta(x)} \\ &= (S_0(t))^{\theta(x)} \end{aligned}$$

An alternative modeling assumption would be

$$S(t|X = x) = S_0(t \cdot \theta(x))$$

where $\theta(x) = \exp\{\eta(x)\}$, $\eta(x) = \beta_1 x_1 + \cdots + \beta_p x_p$, and $S_0(t) = P(T \geq t|X = 0)$ is the base survival function.

Then

8. Parametric survival models

$$\begin{aligned}
 E(T|X = x) &= \int_{t=0}^{\infty} S(t|x) dt \\
 &= \int_{t=0}^{\infty} S_0(t \cdot \theta(x)) dt \\
 &= \int_{u=0}^{\infty} S_0(u) du \cdot \theta(x)^{-1} \\
 &= \theta(x)^{-1} \cdot \int_{u=0}^{\infty} S_0(u) du \\
 &= \theta(x)^{-1} \cdot E(T|X = 0)
 \end{aligned}$$

So the mean of T given $X = x$ is the baseline mean divided by $\theta(x) = \exp\{\eta(x)\}$.

This modeling strategy is called an accelerated failure time model, because covariates cause uniform acceleration (or slowing) of failure times.

Additionally:

$$\begin{aligned}
 H(t|x) &= H_0(\theta(x) \cdot t) \\
 h(t|x) &= \theta(x) \cdot h_0(\theta(x) \cdot t)
 \end{aligned}$$

If the base distribution is exponential with parameter λ then

$$\begin{aligned}
 S(t|x) &= \exp\{-\lambda \cdot t\theta(x)\} \\
 &= [\exp\{-\lambda t\}]^{\theta(x)}
 \end{aligned}$$

which is an exponential model with base hazard multiplied by $\theta(x)$, which is also the proportional hazards model.

In terms of the log survival time $Y = \log\{T\}$ the model can be written as

8. Parametric survival models

$$Y = \alpha - \eta + W$$

$$\alpha = -\log \{\lambda\}$$

where W has the extreme value distribution. The estimated parameter λ is the intercept and the other coefficients are those of η , which will be the opposite sign of those for coxph.

For a Weibull distribution, the hazard function and the survival function are

$$h(t) = \lambda p(\lambda t)^{p-1}$$

$$S(t) = e^{-(\lambda t)^p}$$

We can construct a proportional hazards model by using a linear predictor η_i without constant term and letting $\theta_i = e^{\eta_i}$ we have

$$h(t) = \lambda p(\lambda t)^{p-1} \theta_i$$

A distribution with $h(t) = \lambda p(\lambda t)^{p-1} \theta_i$ is a Weibull distribution with parameters $\lambda^* = \lambda \theta_i^{1/p}$ and p so the survival function is

$$S^*(t) = e^{-(\lambda^* t)^p}$$

$$= e^{-(\lambda \theta_i^{1/p} t)^p}$$

$$= S(t \theta_i^{1/p})$$

so this is also an accelerated failure time model.

In terms of the log survival time $Y = \log \{T\}$ the model can be written as

8. Parametric survival models

$$\begin{aligned}Y &= \alpha - \sigma\eta + \sigma W \\ \alpha &= -\log\{\lambda\} \\ \sigma &= 1/p\end{aligned}$$

where W has the extreme value distribution. The estimated parameter λ is the intercept and the other coefficients are those of η , which will be the opposite sign of those for `coxph`.

These AFT models are log-linear, meaning that the linear predictor has a log link. The exponential and the Weibull are the only log-linear models that are simultaneously proportional hazards models. Other parametric distributions can be used for survival regression either as a proportional hazards model or as an accelerated failure time model.

8.1.5. Dataset: Leukemia treatments

Remission survival times on 42 leukemia patients, half on new treatment, half on standard treatment.

This is the same data as the `drug6mp` data from `KMsurv`, but with two other variables and without the pairing.

```
library(haven)
library(survival)
anderson =
  paste0(
    "http://web1.sph.emory.edu/dkleinb/allDatasets",
    "/surv2datasets/anderson.dta") |>
  read_dta() |>
  mutate(
    status = status |>
      case_match(
        1 ~ "relapse",
```

8. Parametric survival models

```
      0 ~ "censored"
    ),
    sex = sex |>
      case_match(
        0 ~ "female",
        1 ~ "male"
      ),

    rx = rx |>
      case_match(
        0 ~ "new",
        1 ~ "standard"
      ),

    surv = Surv(time = survt, event = (status == "relapse"))
  )

print(anderson)
```

8.1.5.1. Cox semi-parametric model

```
anderson.cox0 = coxph(
  formula = surv ~ rx,
  data = anderson)
summary(anderson.cox0)
```

Call:

```
coxph(formula = surv ~ rx, data = anderson)
```

n= 42, number of events= 30

8. Parametric survival models

```

      coef exp(coef) se(coef)      z Pr(>|z|)
rxstandard 1.572      4.817    0.412 3.81 0.00014 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

      exp(coef) exp(-coef) lower .95 upper .95
rxstandard      4.82      0.208      2.15      10.8

Concordance= 0.69 (se = 0.041 )
Likelihood ratio test= 16.4 on 1 df,  p=5e-05
Wald test              = 14.5 on 1 df,  p=1e-04
Score (logrank) test = 17.2 on 1 df,  p=3e-05

```

8.1.5.2. Weibull parametric model

```

anderson.weib <- survreg(
  formula = surv ~ rx,
  data = anderson,
  dist = "weibull")
summary(anderson.weib)

```

```

Call:
survreg(formula = surv ~ rx, data = anderson, dist = "weibull")

      Value Std. Error      z      p
(Intercept)  3.516      0.252 13.96 < 2e-16
rxstandard   -1.267      0.311 -4.08 4.5e-05
Log(scale)   -0.312      0.147 -2.12  0.034

Scale= 0.732

```

8. Parametric survival models

Weibull distribution

Loglik(model)= -106.6 Loglik(intercept only)= -116.4

Chisq= 19.65 on 1 degrees of freedom, p= 9.3e-06

Number of Newton-Raphson Iterations: 5

n= 42

8.1.5.3. Exponential parametric model

```
anderson.exp <- survreg(  
  formula = surv ~ rx,  
  data = anderson,  
  dist = "exp")  
summary(anderson.exp)
```

Call:

```
survreg(formula = surv ~ rx, data = anderson, dist = "exp")
```

| | Value | Std. Error | z | p |
|-------------|--------|------------|-------|---------|
| (Intercept) | 3.686 | 0.333 | 11.06 | < 2e-16 |
| rxstandard | -1.527 | 0.398 | -3.83 | 0.00013 |

Scale fixed at 1

Exponential distribution

Loglik(model)= -108.5 Loglik(intercept only)= -116.8

Chisq= 16.49 on 1 degrees of freedom, p= 4.9e-05


Number of Newton-Raphson Iterations: 4

n= 42

8. Parametric survival models

8.1.5.4. Diagnostic - complementary log-log survival plot

```
library(survminer)
survfit(
  formula = surv ~ rx,
  data = anderson) |>
ggsurvplot(fun = "cloglog")
```



If the cloglog plot is linear, then a Weibull model may be ok.

8.2. Combining left-truncation and interval-censoring

From [<https://stat.ethz.ch/pipermail/r-help/2015-August/431733.html>]:

8. *Parametric survival models*

coxph does left truncation but not left (or interval) censoring
survreg does interval censoring but not left truncation (or time
dependent covariates).

- Terry Therneau, August 31, 2015

References

- Canchola, Alison J, Susan L Stewart, Leslie Bernstein, Dee W West, Ronald K Ross, Dennis Deapen, Richard Pinder, et al. 2003. “Cox Regression Using Different Time-Scales.” *Western Users of SAS Software*. https://www.lexjansen.com/wuss/2003/DataAnalysis/i-cox_time_scales.pdf.
- Casella, George, and Roger Berger. 2002. *Statistical Inference*. 2nd ed. Cengage Learning. <https://www.cengage.com/c/statistical-inference-2e-casella-berger/9780534243128/>.
- Chatterjee, Samprit, and Ali S Hadi. 2015. *Regression Analysis by Example*. John Wiley & Sons.
- Dobson, Annette J, and Adrian G Barnett. 2018. *An Introduction to Generalized Linear Models*. 4th ed. CRC press. <https://doi.org/10.1201/9781315182780>.
- Dunn, Peter K, Gordon K Smyth, et al. 2018. *Generalized Linear Models with Examples in r*. Vol. 53. Springer. <https://link.springer.com/book/10.1007/978-1-4419-0118-7>.
- Fieller, Nick. 2016. *Basics of Matrix Algebra for Statistics with r*. Chapman; Hall/CRC. <https://doi.org/10.1201/9781315370200>.
- Grambsch, Patricia M, and Terry M Therneau. 1994. “Proportional Hazards Tests and Diagnostics Based on Weighted Residuals.” *Biometrika* 81 (3): 515–26. <https://doi.org/10.1093/biomet/81.3.515>.
- Klein, John P, Melvin L Moeschberger, et al. 2003. *Survival Analysis: Techniques for Censored and Truncated Data*. Vol. 1230. Springer. <https://link.springer.com/book/10.1007/b97377>.
- Kutner, Michael H, Christopher J Nachtsheim, John Neter, and William Li. 2005. *Applied Linear Statistical Models*. McGraw-Hill.

References

- Lawrance, Rachael, Evgeny Degtyarev, Philip Griffiths, Peter Trask, Helen Lau, Denise D'Alessio, Ingolf Griebisch, Gudrun Wallenstein, Kim Cocks, and Kaspar Rufibach. 2020. "What Is an Estimand, and How Does It Relate to Quantifying the Effect of Treatment on Patient-Reported Quality of Life Outcomes in Clinical Trials?" *Journal of Patient-Reported Outcomes* 4 (1): 1–8.
- McLachlan, Geoffrey J, and Thriyambakam Krishnan. 2007. *The EM Algorithm and Extensions*. 2nd ed. John Wiley & Sons. <https://doi.org/10.1002/9780470191613>.
- Pohl, Moritz, Lukas Baumann, Rouven Behnisch, Marietta Kirchner, Johannes Krisam, and Anja Sander. 2021. "Estimands—A Basic Element for Clinical Trials." *Deutsches Ärzteblatt International* 118 (51–52): 883–88. <https://doi.org/10.3238/arztebl.m2021.0373>.
- Van Buuren, Stef. 2018. *Flexible Imputation of Missing Data*. CRC press. <https://stefvanbuuren.name/fimd/>.
- Vittinghoff, Eric, David V Glidden, Stephen C Shiboski, and Charles E McCulloch. 2012. *Regression Methods in Biostatistics: Linear, Logistic, Survival, and Repeated Measures Models*. 2nd ed. Springer. <https://doi.org/10.1007/978-1-4614-1353-0>.

A. Probability

A.1. Random variables

A.1.1. Types of random variables

Definition A.1 (binary variable). A **binary variable** is a random variable which has only two possible values in its range.

Examples of binary outcomes include:

- exposure (exposed vs unexposed)
- disease (diseased vs healthy)
- recovery (recovered vs unrecovered)
- relapse (relapse vs remission)
- return to hospital (returned vs not)
- vital status (dead vs alive)

A.2. Characteristics of probability distributions

Definition A.2 (Density function). The density function $f(t)$ for a random variable T at value t can be defined as the derivative of the cumulative probability function $P(T \leq t)$; that is:

$$f(t) \stackrel{\text{def}}{=} \frac{d}{dt} \Pr(T \leq t)$$

A. Probability

Definition A.3 (Hazard function). The hazard function for a random variable T at value t is the conditional density of T at t , given $T \geq t$; that is:

$$h(t) \stackrel{\text{def}}{=} p(T = t | T \geq t)$$

If T represents the time at which an event occurs, then $h(t)$ is the probability that the event occurs at time t , given that it has not occurred prior to time t .

Definition A.4 (Expectation, expected value). The *expectation* or *expected value* of a random variable X , denoted $\mathbb{E}[X]$, $\mu(X)$, or μ_X , is its mean:

$$\mathbb{E}[X] = \int_{x \in \mathcal{R}(X)} x \cdot p(X = x) dx$$

(c.f. https://en.wikipedia.org/wiki/Expected_value)

Definition A.5 (Variance). The variance of a random variable X is the expectation of the squared difference between X and $\mathbb{E}[X]$; that is:

$$\text{Var}(X) \stackrel{\text{def}}{=} \mathbb{E}[(X - \mathbb{E}[X])^2]$$

Theorem A.1 (Alternative expression for variance).

$$\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$$

Proof. By linearity of expectation, we have:

A. Probability

$$\begin{aligned}
 \text{Var}(X) &\stackrel{\text{def}}{=} \mathbb{E}[(X - \mathbb{E}[X])^2] \\
 &= \mathbb{E}[X^2 - 2X\mathbb{E}[X] + (\mathbb{E}[X])^2] \\
 &= \mathbb{E}[X^2] - \mathbb{E}[2X\mathbb{E}[X]] + \mathbb{E}[(\mathbb{E}[X])^2] \\
 &= \mathbb{E}[X^2] - 2\mathbb{E}[X]\mathbb{E}[X] + (\mathbb{E}[X])^2 \\
 &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2
 \end{aligned}$$

□

Definition A.6 (Precision). The **precision** of a random variable X , often denoted $\tau(X)$, τ_X , or shorthanded as τ , is the inverse of that random variable's variance; that is:

$$\tau(X) \stackrel{\text{def}}{=} (\text{Var}(X))^{-1}$$

Definition A.7 (Standard deviation). The standard deviation of a random variable X is the square-root of the variance of X :

$$\text{SD}(X) \stackrel{\text{def}}{=} \sqrt{\text{Var}(X)}$$

A.3. Key probability distributions

Definition A.8 (Bernoulli distribution). The **Bernoulli distribution** family for a random variable X is defined as:

$$\begin{aligned}
 \Pr(X = x) &= \mathbb{1}_{x \in \{1\}} \pi^x (1 - \pi)^{1-x} \\
 &= \begin{cases} \pi, & x = 1 \\ 1 - \pi, & x = 0 \end{cases}
 \end{aligned}$$

B. Estimation

B.1. Probabilitistic models

Definition B.1 (Scientific models). **Scientific models** are attempts to describe *physical conditions or changes* that occur in the world and universe around us.

Example B.1 (Scientific models in epidemiology). Epidemiologists typically study *biological conditions and changes*, such as the spread of infectious diseases through populations, or the effects of environmental factors on individuals.

B.1.1. Statistical analysis of scientific models

When we perform statistical analyses, we use data to help us choose between models - specifically, to determine which models best explain that data.

However, physical processes do not produce data on their own. Data is only produced when scientists implement an *observation process* (i.e., a *scientific study*), which is distinct from the underlying *physical process*. In some cases, the observation process and the physical process interact with each other. This phenomenon is called the “observer effect”¹.

¹https://en.wikipedia.org/wiki/Observer_effect

B. Estimation

In order to learn about the physical processes we are ultimately interested in, we often need to make special considerations for the observation process that produced the data which we are analyzing. In particular, if some of the planned observations in the study design were not completed, we will likely need to account for the incompleteness of the resulting data set in our analysis. If we are not sure why some observations are incomplete, we may need to model the observation process in addition to the physical process we were originally interested in. For example, if some participants in a study dropped out part-way through the study, we may need investigate why those participants dropped out, as opposed to other participants who completed the study.

These kinds of *missing data* issues are outside of the scope of this course; see Van Buuren (2018) for more details.

B.2. Estimands, estimates, and estimators

B.2.1. Estimands

Definition B.2 (Estimand). An **estimand** is an unknown quantity whose value we want to know (Pohl et al. 2021; Lawrance et al. 2020).

Example B.2 (Mean height of students). If we are trying to determine the mean height of students at our school, then the population mean is our estimand.

In statistical contexts, most estimands are parameters of probabilistic models, or functions of model parameters.

Notation for estimands

Model parameters and other estimands are often symbolized using lower-case Greek letters: $\alpha, \beta, \gamma, \delta$, etc.

B.2.2. Estimates

Definition B.3 (Estimate/estimated value). In statistics, an **estimate** or **estimated value** is an informed guess of an estimand's value, based on observed data.

Example B.3 (Mean height of students). Suppose we measure the heights of 50 random students from our school, and the sample mean was 175cm. We might use 175cm as an *estimate* of the population mean.

B.2.3. Estimators

Definition B.4 (Estimator). An **estimator** is a function $\hat{\theta}(x_1, \dots, x_n)$ that transforms data x_1, \dots, x_n into an estimate.

i Estimators are random variables

When estimators are applied to random variables, the estimators are also random variables.

i Notation for estimators

Estimators are often symbolized by placing a $\hat{}$ (“hat”) symbol on top of the corresponding estimand; for example, $\hat{\theta}$.

Usually, their dependence on the data is implicit:

$$\hat{\theta} \stackrel{\text{def}}{=} \hat{\theta}(x_1, \dots, x_n)$$

Example B.4 (Mean height of students). If we want to estimate the mean height of students at our university, which we will represent as μ , we might measure the heights of $n = 50$ randomly sampled students as random variables X_1, \dots, X_n . Then we could use the function

B. Estimation

$$\hat{\mu}(X_1, \dots, X_n) = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{\text{def}}{=} \bar{X}$$

as an *estimator* to produce an *estimate* $\hat{\mu} = \bar{x}$ of μ .

Another estimator would be just the height of the first student sampled:

$$\hat{\mu}^{(2)}(X_1, \dots, X_n) = X_1$$

A third possible estimator would be the mean of all sampled students' heights, except for the two most extreme; that is, if we re-order the observations $X_{(1)} = \min_{i \in 1:n} X_i$, $X_{(2)} = \min_{i \in \{1:n\} - \arg X_{(1)}} X_i$, ..., $X_{(n)} = \max_{i \in 1:n} X_i$, then we could define the estimator:

$$\hat{\mu}^{(3)}(X_1, \dots, X_n) = \frac{1}{n} \sum_{i=2}^{n-1} X_{(i)}$$

Which of these estimators is best? It depends on how we evaluate them (see Section B.3 below).

B.2.4. Contrasting estimands, estimates, and estimators

It's helpful to keep in mind the mathematical type of each estimation concept:

- estimands are numbers (or vector of numbers)
- estimates are also numbers (or vectors)
- estimators are functions of random variables, so they are also random variables

B.3. Accuracy of estimators

B.3.1. Accuracy

To determine which estimator is best, we need to define *best*. “Accuracy” is usually most important; easy computation is usually secondary.

Definition B.5 (Accuracy). The **accuracy** of an estimator for a given estimand does not have a consensus formal definition, but all of the usual candidates are related to the distributions of the *errors* made by the resulting estimates.

B.3.2. Error

Definition B.6 (Error). The **error** of an estimate $\hat{\theta}$, often denoted $\epsilon(\hat{\theta})$, is the difference between the estimate and its estimand θ ; that is:

$$\epsilon(\hat{\theta}) \stackrel{\text{def}}{=} \hat{\theta} - \theta$$

Some frequently-used measures of accuracy include:

B.3.3. Mean squared error

Definition B.7 (Mean squared error). The **mean squared error** of an estimator $\hat{\theta}$, denoted $\text{MSE}(\hat{\theta})$, is the expectation of the square of the error²:

$$\text{MSE}(\hat{\theta}) \stackrel{\text{def}}{=} \mathbb{E}[(\epsilon(\hat{\theta}))^2]$$

²I might sometimes switch the order of x , θ ; this is unintentional and not meaningful.

B.3.4. Mean absolute error

Definition B.8 (Mean absolute error). The **mean absolute error** of an estimator is the expectation of the absolute value of the error:

$$\text{MAE}(\hat{\theta}) \stackrel{\text{def}}{=} \mathbb{E} [|\epsilon(\hat{\theta})|]$$

B.3.5. Bias

Definition B.9 (Bias). The **bias** of an estimator $\hat{\theta}$ for an estimand θ is the expected value of the error:

$$\text{Bias}(\hat{\theta}) \stackrel{\text{def}}{=} \mathbb{E} [\epsilon(\hat{\theta})] \tag{B.1}$$

Theorem B.1 (Bias equals Expectation minus Truth).

$$\text{Bias}(\hat{\theta}) = \mathbb{E} [\hat{\theta}] - \theta$$

Proof.

$$\begin{aligned} \text{Bias}(\hat{\theta}) &\stackrel{\text{def}}{=} \mathbb{E} [\epsilon(\hat{\theta})] \\ &= \mathbb{E} [\hat{\theta} - \theta] \\ &= \mathbb{E} [\hat{\theta}] - \mathbb{E} [\theta] \\ &= \mathbb{E} [\hat{\theta}] - \theta \end{aligned}$$

The third equality is by the linearity of expectation. □

B. Estimation

Theorem B.2 (Mean Squared Error equals Bias Squared plus Variance).
For any one-dimensional estimator $\hat{\theta}$:

$$MSE(\hat{\theta}) = \left(Bias(\hat{\theta}) \right)^2 + Var(\hat{\theta}) \quad (B.2)$$

Proof. Let's start by expanding each term of the right-hand side:

$$\begin{aligned} \left(Bias(\hat{\theta}) \right)^2 &= \left(\mathbb{E}[\hat{\theta}] - \theta \right)^2 \\ &= \left(\mathbb{E}[\hat{\theta}] \right)^2 - 2\mathbb{E}[\hat{\theta}] \theta + \theta^2 \end{aligned}$$

$$Var(\hat{\theta}) = \mathbb{E}[\hat{\theta}^2] - \left(\mathbb{E}[\hat{\theta}] \right)^2$$

Now, add them together and simplify:

$$\begin{aligned} \left(Bias(\hat{\theta}) \right)^2 + Var(\hat{\theta}) &= \left(\mathbb{E}[\hat{\theta}] \right)^2 - 2\mathbb{E}[\hat{\theta}] \theta + \theta^2 + \mathbb{E}[\hat{\theta}^2] - \left(\mathbb{E}[\hat{\theta}] \right)^2 \\ &= \mathbb{E}[\hat{\theta}^2] - 2\mathbb{E}[\hat{\theta}] \theta + \theta^2 \end{aligned}$$

Now let's expand the left-hand side to reach the same expression:

$$\begin{aligned} MSE(\hat{\theta}) &= \mathbb{E}[(\epsilon(\hat{\theta}))^2] \\ &= \mathbb{E}[(\hat{\theta} - \theta)^2] \\ &= \mathbb{E}[\hat{\theta}^2 - 2\hat{\theta}\theta - \theta^2] \\ &= \mathbb{E}[\hat{\theta}^2] - \mathbb{E}[2\hat{\theta}\theta] + \mathbb{E}[\theta^2] \\ &= \mathbb{E}[\hat{\theta}^2] - 2\mathbb{E}[\hat{\theta}] \theta + \theta^2 \end{aligned}$$

B. Estimation

$\text{MSE}(\hat{\theta})$ and $(\text{Bias}(\hat{\theta}))^2 + \text{Var}(\hat{\theta})$ both equal $\mathbb{E}[\hat{\theta}^2] - 2\mathbb{E}[\hat{\theta}]\theta + \theta^2$. Equality is transitive, so $\text{MSE}(\hat{\theta})$ and $(\text{Bias}(\hat{\theta}))^2 + \text{Var}(\hat{\theta})$ are equal to each other:

$$\text{MSE}(\hat{\theta}) = (\text{Bias}(\hat{\theta}))^2 + \text{Var}(\hat{\theta})$$

□

B.3.5.1. Unbiased estimators

Definition B.10 (unbiased estimator). An estimator $\hat{\theta}$ is **unbiased** if $\text{Bias}(\hat{\theta}) = 0$.

Theorem B.3 (properties of unbiased estimators). *If $\hat{\theta}$ is unbiased, then:*

$$\mathbb{E}[\hat{\theta}] = \theta \tag{B.3}$$

$$\text{MSE}(\hat{\theta}) = \text{Var}(\hat{\theta}) \tag{B.4}$$

Proof. If $\hat{\theta}$ is unbiased, then:

Equation B.3:

$$\text{Bias}(\hat{\theta}) = 0$$

$$\mathbb{E}[\hat{\theta}] - \theta = 0$$

$$\mathbb{E}[\hat{\theta}] = \theta$$

Equation B.4:

B. Estimation

$$\begin{aligned}\text{MSE}(\hat{\theta}) &\stackrel{\text{def}}{=} \mathbb{E} \left[\left(\epsilon(\hat{\theta}) \right)^2 \right] \\ &= \mathbb{E} \left[\left(\hat{\theta} - \theta \right)^2 \right] \\ &= \mathbb{E} \left[\left(\hat{\theta} - \mathbb{E}[\hat{\theta}] \right)^2 \right] \\ &\stackrel{\text{def}}{=} \text{Var}(\hat{\theta})\end{aligned}$$

(Alternative proof of Equation B.4) We could have started from Theorem B.2 instead:

$$\begin{aligned}\text{MSE}(\hat{\theta}) &= \left(\text{Bias}(\hat{\theta}) \right)^2 + \text{Var}(\hat{\theta}) \\ &= (0)^2 + \text{Var}(\hat{\theta}) \\ &= 0 + \text{Var}(\hat{\theta}) \\ &= \text{Var}(\hat{\theta})\end{aligned}$$

□

B.3.6. Standard error

Definition B.11 (Standard error). The **standard error** of an estimator $\hat{\theta}$ is just the **standard deviation** of $\hat{\theta}$; that is:

$$\text{SE}(\hat{\theta}) \stackrel{\text{def}}{=} \text{SD}(\hat{\theta})$$

B. Estimation

“Standard error” is a confusing concept in a few ways. First of all, it isn’t even defined as a characteristic of the **error**, $\epsilon(\hat{\theta})$! Moreover, it is just a synonym for standard deviation, so it seems like a redundant concept. However, standard errors help us construct p-values and confidence intervals, so they come up a lot - often enough to give them their own name.

We can relate standard error to actual error, by rearranging the result from Theorem **B.2**:

$$\begin{aligned}\text{Var}(\hat{\theta}) &= \text{Var}(\hat{\theta} - \theta) \\ &= \text{Var}(\epsilon(\hat{\theta}))\end{aligned}$$

So the variance of the estimator is equal to the variance of the error, and the standard error is equal to the standard deviation of the error:

$$\text{SE}(\hat{\theta}) = \text{SD}(\epsilon(\hat{\theta}))$$

Corollary B.1 (Standard error squared equals MSE minus squared bias).
standard error is what is left over of MSE after bias is removed:

$$\left(\text{SE}(\hat{\theta})\right)^2 = \text{MSE}(\hat{\theta}) - \left(\text{Bias}(\hat{\theta})\right)^2$$

Proof.

$$\begin{aligned}\text{MSE}(\hat{\theta}) &= \left(\text{Bias}(\hat{\theta})\right)^2 + \text{Var}(\hat{\theta}) \\ \therefore \text{Var}(\hat{\theta}) &= \text{MSE}(\hat{\theta}) - \left(\text{Bias}(\hat{\theta})\right)^2 \\ \therefore \left(\text{SE}(\hat{\theta})\right)^2 &= \text{MSE}(\hat{\theta}) - \left(\text{Bias}(\hat{\theta})\right)^2\end{aligned}$$

□

B. Estimation

Corollary B.2 (For unbiased estimators, SE = RMSE). If $\mathbb{E} [\epsilon(\hat{\theta})] = 0$, then:

$$SE(\hat{\theta}) = \sqrt{MSE(\hat{\theta})}$$

(this result is equivalent to Equation [B.4](#))

C. Introduction to Maximum Likelihood Inference

These notes are derived primarily from Dobson and Barnett (2018) (mostly chapters 1-5).

Some material was also taken from McLachlan and Krishnan (2007) and Casella and Berger (2002).

C.1. Introduction

Definition C.1 (Likelihood). Let \tilde{x} be a dataset with corresponding random variable \tilde{X} . Let $p_{\Theta}(\tilde{X})$ be a probability model for the distribution of \tilde{X} with parameter vector Θ .

Then the **likelihood** of parameter value θ (for model $p_{\Theta}(\tilde{X})$) is the *joint probability* of \tilde{x} given $\Theta = \theta$:

$$\begin{aligned}\mathcal{L}(\theta) &\stackrel{\text{def}}{=} p_{\theta}(\tilde{X}) \\ &= p_{\theta}(X_1 = x_1, \dots, X_n = x_n)\end{aligned}$$

The likelihood is a function that takes θ (and implicitly, \tilde{X}) as inputs and outputs a single number, the joint probability of \tilde{X} for model $p_{\Theta}(\tilde{X})$ with $\Theta = \theta$.

Definition C.2 (Maximum likelihood estimate). The **maximum likelihood estimate** of a parameter vector θ is the value of θ that maximizes the likelihood:

$$\hat{\theta}_{\text{ML}} = \arg \max_{\theta} \mathcal{L}(\theta)$$

C.2. Example: Maximum likelihood for Tropical Cyclones in Australia

The `cyclones` dataset in the `dobson` package (Table C.1) records the number of tropical cyclones in Northeastern Australia during 13 November-to-April cyclone seasons (more details in Dobson and Barnett (2018) §1.6.5 and `help(cyclones, package = "dobson")`). Figure C.1 graphs the number of cyclones (y-axis) by season (x-axis). Let's use Y_i to represent these counts, where i is an indexing variable for the seasons and Y_i is the number of cyclones in season i .

Suppose we want to learn about how many cyclones to expect per season.

```
library(dobson)
library(dplyr)
data(cyclones)
library(pander)
pander(cyclones |> relocate(season, .before = everything()))
```

Table C.1.: Number of tropical cyclones during a season from November to April in Northeastern Australia

| season | years | number |
|--------|--------|--------|
| 1 | 1956/7 | 6 |
| 2 | 1957/8 | 5 |

C. Introduction to Maximum Likelihood Inference

Table C.1.: Number of tropical cyclones during a season from November to April in Northeastern Australia

| season | years | number |
|--------|---------|--------|
| 3 | 1958/9 | 4 |
| 4 | 1959/60 | 6 |
| 5 | 1960/1 | 6 |
| 6 | 1961/2 | 3 |
| 7 | 1962/3 | 12 |
| 8 | 1963/4 | 7 |
| 9 | 1964/5 | 4 |
| 10 | 1965/6 | 2 |
| 11 | 1966/7 | 6 |
| 12 | 1967/8 | 7 |
| 13 | 1968/9 | 4 |

```
library(ggplot2)
library(dplyr)
cyclones |>
  mutate(years = years |> factor(level = years)) |>
  ggplot(
    aes(x = years,
         y = number,
         group = 1)) +
  geom_point() +
  geom_line() +
  xlab("Season") +
  ylab("Number of cyclones") +
  theme_bw() +
  expand_limits(y = 0) +
  theme(axis.text.x = element_text(vjust = .5, angle = 45))
```




Figure C.1.: Number of tropical cyclones per season in northeastern Australia, 1956-1969

There's no obvious correlation between adjacent seasons, so let's assume that each season is independent of the others.

Let's also assume that they are identically distributed; let's denote this distribution as $P(Y = y)$ (note that there's no index i in this expression, since we are assuming the Y_i s are identically distributed). We can visualize the distribution using a bar plot (Figure C.2). Table C.2 provides summary statistics.

```
cyclones |>
  ggplot() +
  geom_histogram(aes(x = number)) +
  expand_limits(x = 0) +
```

C. Introduction to Maximum Likelihood Inference

```
theme_bw() +  
xlab("Number of cyclones") +  
ylab("Count (number of seasons)")
```




Figure C.2.: Bar plot of cyclones per season

```
cyclones |> table1::table1(x = ~ number)
```

We want to estimate $P(Y = y)$; that is, $P(Y = y)$ is our estimand.

We could estimate $P(Y = y)$ for each value of y in $0 : \infty$ separately (“nonparametrically”) using the fraction of our data with $Y_i = y$, but then we would be estimating an infinitely large set of parameters, and we would have low precision. We will probably do better with a parametric model.

C. Introduction to Maximum Likelihood Inference

Table C.2.: Summary statistics for `cyclones` data

| | Overall |
|-------------------|-------------------|
| | (N=13) |
| number | |
| Mean (SD) | 5.54 (2.47) |
| Median [Min, Max] | 6.00 [2.00, 12.0] |

Exercise C.1. What parametric probability distribution family might we use to model this empirical distribution?

Solution. Let's use the Poisson. The Poisson distribution is appropriate for this data, because the data are counts that could theoretically take any integer value (discrete) in the range $0 : \infty$. Visually, the plot of our data closely resembles a Poisson or binomial distribution. Since cyclones do not have an “upper limit” on the number of events we could potentially observe in one season, the Poisson distribution is more appropriate than the binomial.

Exercise C.2. Write down the Poisson distribution's probability mass function.

Solution.

$$P(Y = y) = \frac{\lambda^y e^{-\lambda}}{y!}$$

Now, we can estimate the parameter λ for this distribution using maximum likelihood estimation.

What is the likelihood?

C. Introduction to Maximum Likelihood Inference

Exercise C.3. Write down the likelihood (probability mass function or probability density function) of a single observation x , according to your model.

Solution.

$$\begin{aligned}\mathcal{L}(\lambda; x) &= p(X = x | \Lambda = \lambda) \\ &= \frac{\lambda^x e^{-\lambda}}{x!}\end{aligned}$$

Exercise C.4. Write down the vector of parameters in your model.

Solution. There is only one parameter, λ :

$$\theta = (\lambda)$$

Exercise C.5. Write down the population mean and variance of a single observation from your chosen probability model, as a function of the parameters (extra credit - derive them).

Solution.

- Population mean: $E[X] = \lambda$
- Population variance: $\text{Var}(X) = \lambda$

Exercise C.6. Write down the likelihood of the full dataset.

Solution.

$$\begin{aligned}\mathcal{L}(\lambda; \tilde{x}) &= P(\tilde{X} = \tilde{x}) \\ &= P(X_1 = x_1, X_2 = x_2, \dots, X_{13} = x_{13}) \\ &= \prod_{i=1}^{13} P(X_i = x_i) \\ &= \prod_{i=1}^{13} \frac{\lambda^{x_i} e^{-\lambda}}{x_i!}\end{aligned}$$

C. Introduction to Maximum Likelihood Inference

Exercise C.7. Graph the likelihood as a function of λ .

Solution.

```
lik = function(lambda, y = cyclones$number, n = length(y))
{
  lambda^sum(y) * exp(-n*lambda) / prod(factorial(y))
}

library(ggplot2)
ggplot() +
  geom_function(fun = lik, n = 1001) +
  xlim(min(cyclones$number), max(cyclones$number)) +
  theme_bw() +
  ylab("likelihood") +
  xlab('lambda')
```

C. Introduction to Maximum Likelihood Inference




Figure C.3.: Likelihood of Dobson cyclone data

Exercise C.8. Write down the log-likelihood of the full dataset.

C. Introduction to Maximum Likelihood Inference

Solution.

$$\begin{aligned}\ell(\lambda; \tilde{x}) &= \log \{ \mathcal{L}(\lambda; \tilde{x}) \} \\ &= \log \left\{ \prod_{i=1}^n \frac{\lambda^{x_i} e^{-\lambda}}{x_i!} \right\} \\ &= \sum_{i=1}^n \log \left\{ \frac{\lambda^{x_i} e^{-\lambda}}{x_i!} \right\} \\ &= \sum_{i=1}^n \log \{ \lambda^{x_i} \} + \log \{ e^{-\lambda} \} - \log \{ x_i! \} \\ &= \sum_{i=1}^n x_i \log \{ \lambda \} - \lambda - \log \{ x_i! \} \\ &= \sum_{i=1}^n x_i \log \{ \lambda \} - \sum_{i=1}^n \lambda - \sum_{i=1}^n \log \{ x_i! \} \\ &= \sum_{i=1}^n x_i \log \{ \lambda \} - n\lambda - \sum_{i=1}^n \log \{ x_i! \}\end{aligned}$$

Exercise C.9. Graph the log-likelihood as a function of λ .

Solution.

```
loglik = function(lambda, y = cyclones$number, n = length(y))
{
  sum(y) * log(lambda) - n*lambda - sum(log(factorial(y)))
}

ll_plot = ggplot() +
  geom_function(fun = loglik, n = 1001) +
  xlim(min(cyclones$number), max(cyclones$number)) +
  theme_bw() +
  ylab("log-likelihood") +
  xlab('lambda')
ll_plot
```




Figure C.4.: log-likelihood of Dobson cyclone data

Exercise C.10. Derive the score function for the dataset.

Solution. The score function is the first derivative(s) of the log-likelihood:

C. Introduction to Maximum Likelihood Inference

$$\begin{aligned}\ell'(\lambda; \tilde{x}) &= \frac{\partial}{\partial \lambda} \sum_{i=1}^n x_i \log \{\lambda\} - n\lambda - \sum_{i=1}^n \log \{x_i!\} \\ &= \frac{\partial}{\partial \lambda} \sum_{i=1}^n x_i \log \{\lambda\} - \frac{\partial}{\partial \lambda} n\lambda - \frac{\partial}{\partial \lambda} \sum_{i=1}^n \log \{x_i!\} \\ &= \sum_{i=1}^n x_i \frac{\partial}{\partial \lambda} \log \{\lambda\} - n \frac{\partial}{\partial \lambda} \lambda - \sum_{i=1}^n \frac{\partial}{\partial \lambda} \log \{x_i!\} \\ &= \sum_{i=1}^n x_i \frac{1}{\lambda} - n - 0 \\ &= \frac{1}{\lambda} \sum_{i=1}^n x_i - n\end{aligned}$$

Exercise C.11. Graph the score function.

Solution.

```
score = function(lambda, y = cyclones$number, n = length(y))
{
  sum(y)/lambda - n
}

ggplot() +
  geom_function(fun = score, n = 1001) +
  xlim(min(cyclones$number), max(cyclones$number)) +
  theme_bw() +
  ylab("log-likelihood") +
  xlab('lambda') +
  geom_hline(yintercept = 0, col = 'red')
```




Figure C.5.: score function of Dobson cyclone data

Exercise C.12. Derive the Hessian matrix.

Solution. The Hessian function for an iid sample is the 2nd derivative(s) of the log-likelihood:

$$\begin{aligned}
 \ell''(\lambda; \tilde{x}) &= \frac{\partial}{\partial \lambda} \left(\frac{1}{\lambda} \sum_{i=1}^n x_i - n \right) \\
 &= \frac{\partial}{\partial \lambda} \frac{1}{\lambda} \sum_{i=1}^n x_i - \frac{\partial}{\partial \lambda} n \\
 &= -\frac{1}{\lambda^2} \sum_{i=1}^n x_i \\
 &= -\frac{1}{\lambda^2} n \bar{x}
 \end{aligned}$$

C. Introduction to Maximum Likelihood Inference

Exercise C.13. Graph the Hessian.

Solution.

```
hessian = function(lambda, y = cyclones$number, n = length(y))
{
  -sum(y)/lambda^2
}

ggplot() +
  geom_function(fun = hessian, n = 1001) +
  xlim(min(cyclones$number), max(cyclones$number)) +
  theme_bw() +
  ylab("log-likelihood") +
  xlab('lambda') +
  geom_hline(yintercept = 0, col = 'red')
```




Figure C.6.: Hessian function of Dobson cyclone data

Exercise C.14. Write the score equation (estimating equation).

Solution.

$$\ell'(\lambda; \tilde{x}) = 0$$

Exercise C.15. Solve the estimating equation for λ :

C. Introduction to Maximum Likelihood Inference

Solution.

$$\begin{aligned} 0 &= \frac{1}{\lambda} \sum_{i=1}^n x_i - n \\ n &= \frac{1}{\lambda} \sum_{i=1}^n x_i \\ n\lambda &= \sum_{i=1}^n x_i \\ \lambda &= \frac{1}{n} \sum_{i=1}^n x_i \\ &= \bar{x} \end{aligned}$$

Let's call this solution of the estimating equation $\tilde{\lambda}$ for now:

$$\tilde{\lambda} \stackrel{\text{def}}{=} \bar{x}$$

Exercise C.16. Confirm that the Hessian $\ell''(\lambda; \tilde{x})$ is negative when evaluated at $\tilde{\lambda}$.

Solution.

$$\begin{aligned} \ell''(\tilde{\lambda}; \tilde{x}) &= -\frac{1}{\tilde{\lambda}^2} n\bar{x} \\ &= -\frac{1}{\bar{x}^2} n\bar{x} \\ &= -\frac{n}{\bar{x}} \\ &< 0 \end{aligned}$$

Exercise C.17. Find the MLE of λ .

Solution. Since $\ell''(\tilde{\lambda}; \tilde{x}) < 0$, $\tilde{\lambda}$ is at least a local maximizer of the likelihood function $\mathcal{L}(\lambda)$. Since there is only one solution to the estimating equation

C. Introduction to Maximum Likelihood Inference

and the Hessian is negative definite everywhere, $\tilde{\lambda}$ must also be the global maximizer of $\mathcal{L}(\lambda; \tilde{x})$:

```
mle = mean(cyclones$number)
```

$$\hat{\lambda}_{\text{ML}} = \bar{x} = 5.53846154$$

Exercise C.18. Graph the log-likelihood with the MLE superimposed.

Solution.

```
library(dplyr)

mle_data = tibble(x = mle, y = loglik(mle))
ll_plot + geom_point(data = mle_data, aes(x = x, y = y), col = 'red')
```




Figure C.7.: log-likelihood of Dobson cyclone data with MLE

```
obs_inf = function(lambda, y = cyclones$number, n = length(y))  
{  
  sum(y)/lambda^2  
}
```

C.3. Maximum likelihood inference for univariate Gaussian models

Suppose $X_1, \dots, X_n \sim_{iid} N(\mu, \sigma^2)$. Let $X = (X_1, \dots, X_n)^\top$ be these random variables in vector format. Let x_i and x denote the corresponding observed data. Let $\theta = (\mu, \sigma^2)^\top$ be the vector of parameters. Let Θ denote the parameters as a random vector.

C. Introduction to Maximum Likelihood Inference

Then the log-likelihood $\ell \stackrel{\text{def}}{=} \ell(X; \theta) \stackrel{\text{def}}{=} p(X = x \mid \Theta = \theta)$ is:

$$\begin{aligned}\ell &\propto -\frac{n}{2} \log \{\sigma^2\} - \frac{1}{2} \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^2} \\ &= -\frac{n}{2} \log \{\sigma^2\} - \frac{1}{2\sigma^2} \sum_{i=1}^n x_i^2 - 2x_i\mu + \mu^2\end{aligned}$$

C.3.1. MLE of μ

Then:

$$\begin{aligned}\frac{d\ell}{d\mu} &= -\frac{1}{2} \sum_{i=1}^n \frac{-2(x_i - \mu)}{\sigma^2} \\ &= \frac{1}{\sigma^2} \left[\left(\sum_{i=1}^n x_i \right) - n\mu \right]\end{aligned}$$

If $\frac{d\ell}{d\mu} = 0$, then $\mu = \bar{x} \stackrel{\text{def}}{=} \frac{1}{n} \sum_{i=1}^n x_i$.

$$\frac{d^2\ell}{(d\mu)^2} = \frac{-n}{\sigma^2} < 0$$

So $\hat{\mu}_{ML} = \bar{x}$.

C.3.2. MLE of σ^2

💡 Reparametrizing the Gaussian distribution

When solving for $\hat{\sigma}_{ML}$, you can treat σ^2 as an atomic variable (don't differentiate with respect to σ or things get messy). In fact, you can replace σ^2 with $1/\tau$ and differentiate with respect to τ instead, and the process might be even easier.

$$\begin{aligned}\frac{d\ell}{d\sigma^2} &= \frac{d}{d\sigma^2} \left(-\frac{n}{2} \log \{\sigma^2\} - \frac{1}{2} \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^2} \right) \\ &= -\frac{n}{2} (\sigma^2)^{-1} + \frac{1}{2} (\sigma^2)^{-2} \sum_{i=1}^n (x_i - \mu)^2\end{aligned}$$

If $\frac{d\ell}{d\sigma^2} = 0$, then:

$$\begin{aligned}\frac{n}{2} (\sigma^2)^{-1} &= \frac{1}{2} (\sigma^2)^{-2} \sum_{i=1}^n (x_i - \mu)^2 \\ \sigma^2 &= \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2\end{aligned}$$

We plug in $\hat{\mu}_{ML} = \bar{x}$ to maximize globally (a technique called profiling):

$$\sigma_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

Now:

C. Introduction to Maximum Likelihood Inference

$$\begin{aligned}
 \frac{d^2 \ell}{(d\sigma^2)^2} &= \frac{d}{d\sigma^2} \left\{ -\frac{n}{2} (\sigma^2)^{-1} + \frac{1}{2} (\sigma^2)^{-2} \sum_{i=1}^n (x_i - \mu)^2 \right\} \\
 &= \left\{ -\frac{n}{2} \frac{d}{d\sigma^2} (\sigma^2)^{-1} + \frac{1}{2} \frac{d}{d\sigma^2} (\sigma^2)^{-2} \sum_{i=1}^n (x_i - \mu)^2 \right\} \\
 &= \left\{ \frac{n}{2} (\sigma^2)^{-2} - (\sigma^2)^{-3} \sum_{i=1}^n (x_i - \mu)^2 \right\} \\
 &= (\sigma^2)^{-2} \left\{ \frac{n}{2} - (\sigma^2)^{-1} \sum_{i=1}^n (x_i - \mu)^2 \right\}
 \end{aligned}$$

Evaluated at $\mu = \bar{x}, \sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$, we have:

$$\begin{aligned}
 \frac{d^2 \ell}{(d\sigma^2)^2} &= (\hat{\sigma}^2)^{-2} \left\{ \frac{n}{2} - (\hat{\sigma}^2)^{-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right\} \\
 &= (\hat{\sigma}^2)^{-2} \left\{ \frac{n}{2} - (\hat{\sigma}^2)^{-1} n \hat{\sigma}^2 \right\} \\
 &= (\hat{\sigma}^2)^{-2} \left\{ \frac{n}{2} - n \right\} \\
 &= (\hat{\sigma}^2)^{-2} n \left\{ \frac{1}{2} - 1 \right\} \\
 &= (\hat{\sigma}^2)^{-2} n \left(-\frac{1}{2} \right) < 0
 \end{aligned}$$

Finally, we have:

C. Introduction to Maximum Likelihood Inference

$$\begin{aligned}
\frac{d^2\ell}{d\mu d\sigma^2} &= \frac{d}{d\mu} \left\{ -\frac{n}{2} (\sigma^2)^{-1} + \frac{1}{2} (\sigma^2)^{-2} \sum_{i=1}^n (x_i - \mu)^2 \right\} \\
&= \frac{1}{2} (\sigma^2)^{-2} \frac{d}{d\mu} \sum_{i=1}^n (x_i - \mu)^2 \\
&= \frac{1}{2} (\sigma^2)^{-2} \sum_{i=1}^n -2(x_i - \mu) \\
&= -(\sigma^2)^{-2} \sum_{i=1}^n (x_i - \mu)
\end{aligned}$$

Evaluated at $\mu = \hat{\mu} = \bar{x}$, $\sigma^2 = \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$, we have:

$$\frac{d^2\ell}{d\mu d\sigma^2} = -(\hat{\sigma}^2)^{-2} (n\bar{x} - n\bar{x}) = 0$$

C.3.3. Covariance matrix of MLEs

C.3.3.1. The score function

Let θ be the vector of all parameters; here, $\theta = (\mu, \sigma^2)^\top$.

Let $\ell'(x, \theta) \stackrel{\text{def}}{=} \frac{d}{d\theta} \ell(x, \theta) = \begin{pmatrix} \frac{d}{d\mu} \ell(\theta; x) \\ \frac{d}{d\sigma^2} \ell(\theta; x) \end{pmatrix} = \begin{pmatrix} \ell'_\mu(\theta; x) \\ \ell'_{\sigma^2}(\theta; x) \end{pmatrix}$.

$\ell'(x, \theta)$ is the function we set equal to 0 and solve to find the MLE:

$$\hat{\theta}_{ML} = \{\theta : \ell'(x, \theta) = 0\}$$

The function $\ell'(x, \theta)$ is so central that it has its own name, the “score” or “gradient” function. Statisticians also often skip writing the arguments

(x, θ) , so $\ell' \stackrel{\text{def}}{=} \ell'(x, \theta)$.¹ Some statisticians use U or S instead of ℓ' . I prefer ℓ' . Why use up extra letters?

C.3.3.2. The (Fisher) (expected) information matrix

The variance of $\ell'(x, \theta)$, $\text{Cov}\{\ell'(x, \theta)\}$, is also very important; we call it the “expected information matrix”, “Fisher information matrix”, or just “information matrix”, and we represent it using the symbol \mathfrak{I} (`\frakturI` in Unicode, `\mathfrak{I}` in LaTeX).

Review of variances and covariances

Variances and covariances of one-dimensional random variables

For a one-dimensional random variables X ,

$$\text{Var}(X) \stackrel{\text{def}}{=} \text{E}[(X - \text{E}[X])^2] = \text{E}[X^2] - (\text{E}[X])^2$$

For any two-dimensional random variables, X, Y :

$$\text{Cov}(X, Y) = \text{E}[(X - \text{E}[X])(Y - \text{E}[Y])] = \text{E}[XY] - \text{E}[X]\text{E}[Y]$$

Therefore, $\text{Var}(X) = \text{Cov}(X, X) = \text{E}[XX] - \text{E}[X]\text{E}[X] = \text{E}[X^2] - (\text{E}[X])^2$

$$\mathfrak{I} \stackrel{\text{def}}{=} \mathfrak{I}(\theta) \stackrel{\text{def}}{=} \text{Cov}(\ell' | \theta) = \text{E}[\ell' \ell'^\top] - \text{E}[\ell'] \text{E}[\ell']^\top$$

Sometimes we write $\text{Cov}(X) \stackrel{\text{def}}{=} \text{Cov}(X, X) = \text{Var}(X)$.

¹I might sometimes switch the order of x, θ ; this is unintentional and not meaningful.

Variances and covariances of $p \times 1$ random vectors

Now, for a $p \times 1$ dimensional random vector X ,

$$\begin{aligned}
 \text{Var}(X) &\stackrel{\text{def}}{=} \text{Cov}(X) \\
 &\stackrel{\text{def}}{=} E \left[(X - E[X])^\top (X - E[X]) \right] \\
 &= E \left[(X^\top - E[X]^\top) (X - E[X]) \right] \\
 &= E \left[X^\top X - E[X]^\top X - X^\top E[X] + E[X]^\top E[X] \right] \\
 &= E \left[X^\top X \right] - E[X]^\top E[X] - E[X]^\top E[X] + E[X]^\top E[X] \\
 &= E \left[X^\top X \right] - 2E[X]^\top E[X] + E[X]^\top E[X] \\
 &= E \left[X^\top X \right] - E[X]^\top E[X]
 \end{aligned}$$

The elements of \mathfrak{J} are:

$$\left\{ \mathfrak{J}_{ij} \stackrel{\text{def}}{=} \text{Cov}(\ell'_i, \ell'_j) = E[\ell'_i \ell'_j] - E[\ell'_i] E[\ell'_j] \right\}$$

In our motivating example, $i, j \in \{\mu, \sigma^2\}$. Here,

$$\begin{aligned}
 E[\ell'] &\stackrel{\text{def}}{=} \int_{x \in \mathfrak{R}(x)} \ell'(x, \theta) p(X = x | \theta) dx \\
 &= \int_{x \in \mathfrak{R}(x)} \left(\frac{d}{d\theta} \log \{p(X = x | \theta)\} \right) p(X = x | \theta) dx \\
 &= \int_{x \in \mathfrak{R}(x)} \frac{\frac{d}{d\theta} p(X = x | \theta)}{p(X = x | \theta)} p(X = x | \theta) dx \\
 &= \int_{x \in \mathfrak{R}(x)} \frac{d}{d\theta} p(X = x | \theta) dx
 \end{aligned}$$

And similarly

C. Introduction to Maximum Likelihood Inference

$$\mathbb{E} \left[\ell' \ell'^{\top} \right] \stackrel{\text{def}}{=} \int_{x \in R(x)} \ell'(x, \theta) \ell'(x, \theta)^{\top} p(X = x \mid \theta) dx$$

Note that $\mathbb{E} [\ell']$ and $\mathbb{E} [\ell' \ell'^{\top}]$ are functions of θ but not of x ; the expectation operator removed x .

Also note that for most of the distributions you are familiar with (including Gaussian, binomial, Poisson, exponential),

$$\mathbb{E} [\ell'] = 0$$

So

$$\mathfrak{J} = \mathbb{E} [\ell' \ell'^{\top}]$$

Moreover, for those distributions (called the “exponential family”), we have:

$$\mathfrak{J} = -\mathbb{E} [\ell''] = \mathbb{E} [-\ell'']$$

(see Dobson and Barnett 4e, §3.17), where

$$\ell'' \stackrel{\text{def}}{=} \frac{d}{d\theta} \ell'(x, \theta)^{\top} = \frac{d}{d\theta} \frac{d}{d\theta^{\top}} \ell(x, \theta)$$

is the $p \times p$ matrix whose elements are:

$$\ell''_{ij} \stackrel{\text{def}}{=} \frac{d}{d\theta_i} \frac{d}{d\theta_j} \log \{p(X = x \mid \theta)\}$$

ℓ'' could be called the “Hessian” of the log-likelihood function.

C. Introduction to Maximum Likelihood Inference

Sometimes, we use $I(\theta; x) \stackrel{\text{def}}{=} -\ell''$ (note the standard-font “I” here). $I(\theta; x)$ is the observed information, precision, or concentration matrix (Negative Hessian).

! Key point

The asymptotics of MLEs gives us $\hat{\theta}_{ML} \sim N(\theta, \mathcal{I}^{-1}(\theta))$, approximately, for large sample sizes.

We can estimate $\mathcal{I}^{-1}(\theta)$ by working out $-\text{E}[\ell'']$ or $\text{E}[\ell' \ell'^\top]$ and plugging in $\hat{\theta}_{ML}$, but sometimes we instead use $I(\hat{\theta}_{ML}; x)$ for convenience; there are some cases where it’s provably better according to some criteria (Efron & Hinkley 1978).

Note that later, when we are trying to find MLEs for likelihoods that we can’t easily differentiate, we will “hill-climb” using the Newton-Raphson algorithm:

$$\begin{aligned}\hat{\theta} &\leftarrow \hat{\theta} + [I(\hat{\theta}, y)]^{-1} \ell'(y, \hat{\theta}) \\ &= \hat{\theta} - [\ell''(y, \hat{\theta})]^{-1} \ell'(y, \hat{\theta})\end{aligned}$$

Here, for computational simplicity, we will sometimes use $\mathcal{I}^{-1}(\theta)$ in place of $I(\hat{\theta}, y)$; doing so is called “Fisher scoring” or the “method of scoring”. Note that this is the opposite of the substitution that we are making for estimating the variance of the MLE; this time we should technically use the observed information but we use the expected information instead.

There’s also an “empirical information matrix” (see McLachlan and Krishnan 2007).

$$I_e(\theta, y) = \sum_{i=1}^n \ell'_i \ell'_i{}^\top - \frac{1}{n} \ell' \ell'^\top$$

C. Introduction to Maximum Likelihood Inference

where ℓ_i is the log-likelihood of the i th observation. Note that $\ell' = \sum_{i=1}^n \ell'_i$.

$\frac{1}{n}I_e(\theta, y)$ is the sample equivalent of

$$\mathfrak{J} \stackrel{\text{def}}{=} \mathfrak{J}(\theta) \stackrel{\text{def}}{=} \text{Cov}(\ell' | \theta) = \text{E} \left[\ell' \ell'^{\top} \right] - \text{E} \left[\ell' \right] \text{E} \left[\ell' \right]^{\top}$$

$$\left\{ \mathfrak{J}_{jk} \stackrel{\text{def}}{=} \text{Cov}(\ell'_j, \ell'_k) = \text{E} \left[\ell'_j \ell'_k \right] - \text{E} \left[\ell'_j \right] \text{E} \left[\ell'_k \right] \right\}$$

$I_e(\theta, y)$ is sometimes computationally easier to compute for Newton-Raphson-type maximization algorithms.

Back to our Gaussian example:

$$I = \begin{bmatrix} \frac{n}{\hat{\sigma}^2} & 0 \\ 0 & (\hat{\sigma}^2)^{-2} n \left(-\frac{1}{2}\right) \end{bmatrix} = \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix}$$

So:

$$I^{-1} = \frac{1}{ad} \begin{bmatrix} d & 0 \\ 0 & a \end{bmatrix} = \begin{bmatrix} \frac{1}{a} & 0 \\ 0 & \frac{1}{d} \end{bmatrix}$$

$$I^{-1} = \begin{bmatrix} \frac{\hat{\sigma}^2}{n} & 0 \\ 0 & \frac{2(\hat{\sigma}^2)^2}{n} \end{bmatrix}$$

See Casella and Berger (2002) p322, example 7.2.12.

To prove it's a maximum, need:

- $\ell' = 0$
- At least one diagonal element of \mathfrak{l}'' is negative.
- Determinant of \mathfrak{l}'' is positive.

C.3.4. Confidence intervals for MLEs

C.3.5. p-values and hypothesis tests for MLEs

C.3.6. Likelihood ratio tests for MLEs

[We haven't gone over this yet]

log(likelihood ratio) tests (c.f. Dobson and Barnett 2018, sec. 5.7):

$$2(\ell - \ell_0) \sim \chi^2(p - q)$$

See also <https://online.stat.psu.edu/stat504/book/export/html/657>

C.3.7. Prediction intervals for MLEs


$$\bar{X} \in \left[\hat{\mu} \pm z_{1-\alpha/2} \frac{\sigma}{m} \right]$$

Where m is the sample size of the new data to be predicted (typically 1, except for binary outcomes, where it needs to be bigger for prediction intervals to make sense)

D. Common Mistakes

D.1. Parameters versus random variables

The parameters of a probability distribution shouldn't involve the random variables being modeled:

 This is wrong

$$\begin{aligned} X &\sim \text{Pois}(\lambda) \\ \hat{\lambda}_{ML} &\rightarrow_D N(\bar{X}, \lambda/n) \end{aligned}$$

Solution.

$$\hat{\lambda}_{ML} \rightarrow_D N(\lambda, \lambda/n)$$

Expectations are means, not sums, despite the similarity of Σ and E . Really, we should use μ instead of E .

E. Notation

Table E.1.: Notation used in this book

| symbol | meaning | LaTeX |
|-----------------|---------------------------|------------------------------------|
| \neg | not | <code>\neg</code> |
| \forall | all | <code>\forall</code> |
| \exists | some | <code>\exists</code> |
| \cup | union, “or” | <code>\cup</code> |
| \cap | intersection, “and” | <code>\cap</code> |
| $ $ | given, conditional on | <code>\mid</code> , <code> </code> |
| \sum | sum | <code>\sum</code> |
| \prod | product | <code>\prod</code> |
| μ | mean | <code>\mu</code> |
| $\mathbb{E}[X]$ | expectation of X | <code>\text{\texttt{E}}[X]</code> |

Index

estimand, 379
estimate, 380
estimated value, 380
estimator, 380
expectation, 376
expected value, 376