

Homological Algebra

Luiz Alberto

September 24, 2026

Contents

1	Some notes from 2026 - class	2
1.1	Derived functors	2
2	Exercises	7
2.1	List 1	7
2.2	List 2	14

Chapter 1

Some notes from 2026 - class

1.1 Derived functors

- (Homotopy) We define a equivalence relation on $\text{Hom}(C_\bullet, D_\bullet)$ as follows: Let $f, g : (C_\bullet, d) \rightarrow (D_\bullet, d)$ be two morphisms of complexes, then we say that $f \sim g$ if there exist $h_n : C_n \rightarrow D_{n+1}$, $\forall n$ such that the diagram below commutes.

$$\begin{array}{ccccccc}
 C_{n-1} & \xleftarrow{d_n} & C_n & \xleftarrow{d_{n+1}} & C_{n+1} & \xleftarrow{\quad} & \cdots \\
 \downarrow h_{n-1} & \searrow & \downarrow h_n & \searrow & \downarrow h_{n+1} & \searrow & \\
 D_{n-1} & \xleftarrow{d_n} & D_n & \xleftarrow{d_{n+1}} & D_{n+1} & \xleftarrow{\quad} & \cdots
 \end{array}$$

We write $h \circ d + d \circ h = g - f$. Observe that d decreases the index and h increases the index. So $(h \circ d)_n = h_{n-1} \circ d_n$ and $(d \circ h)_n = d_{n+1} \circ h_n$.

Definition 1.1.1. We say that $f_\bullet : C_\bullet \rightarrow D_\bullet$ is a Homotopy equivalence if there exists $g_\bullet : D_\bullet \rightarrow C_\bullet$ such that $g_\bullet \circ f_\bullet \sim \text{Id}_{C_\bullet}$ and $f_\bullet \circ g_\bullet \sim \text{Id}_{D_\bullet}$.

Notation 1.1.2. Normally we will avoid to write $(C_\bullet, d)$ for complexes and adopt the minimalistic notation by just writing “ C ” instead. Same with morphism of complexes.

Proposition 1.1.3. Let $f, g : C \rightarrow D$, $f \sim g$, then $H_n(f) = H_n(g)$, $\forall n$.

Proof: Remember that we define $H_n(f) : H_n(C) \rightarrow H_n(D) : [x] \mapsto [f(x)]$. But since $g - f = hd + dh$, we get that $g(x) - f(x) = dh(x)$, $x \in \text{Ker}(d : C_n \rightarrow C_{n-1})$ and $[f(x)] = [g(x)]$. $\square$

Corollary 1.1.4. If $f : C \rightarrow D$ is Homotopy equivalence then $H_n(C) \cong H_n(D)$.

Proof: By hypothesis, there exists $g : D \rightarrow C$ such that $f \circ g \sim \text{Id}_D$, $g \circ f \sim \text{Id}_C$. Therefore $H_n(f) \circ H_n(g) = H_n(\text{Id}_D) = \text{Id}_{H_n(D)}$ and etc. $\square$

- (Injective resolution) Here $\mathcal{C}$ denotes an abelian category if not stated otherwise.

Definition 1.1.5. We say that $I \in \mathcal{C}$ is injective if $\text{Hom}_{\mathcal{C}}(-, I) : \mathcal{C} \rightarrow \underline{\text{Ab}}$ is an exact functor. This is equivalent to say that whenever we have a monomorphism $i : A \rightarrow B$ and a morphism $f : A \rightarrow I$, then there exists $\tilde{f} : B \rightarrow I$.

$$\begin{array}{ccc} A & \xrightarrow{i} & B \\ f \downarrow & \swarrow \tilde{f} & \\ I & & \end{array}$$

Definition 1.1.6. Let $A \in \mathcal{C}$, then an injective resolution for A is a complex

$$I^\bullet = (0 \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \dots) \quad (1.1)$$

such that

- (1) I^n is injective;
- (2) The augmented complex $\tilde{I}^\bullet = (0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow \dots)$ is acyclic.

Definition 1.1.7. We say that $\mathcal{C}$ has enough injectives if for any $A \in \mathcal{C}$ there exists an injective object $I \in \mathcal{C}$ and a monomorphism $f : A \rightarrow I$.

Proposition 1.1.8. If $\mathcal{C}$ has enough injectives, then any $A \in \mathcal{C}$ has a injective resolution.

Proof: Draw the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{\epsilon} & I^0 & \xrightarrow{\quad \quad \quad} & I^1 \dots \\ & & & & \searrow & & \swarrow \\ & & & & & I^0/\epsilon(A) = Q^0 & \end{array}$$

and repeat

$$\begin{array}{ccc} I^1 & \xrightarrow{\quad \quad \quad} & I^2 \dots \\ & \searrow & \swarrow \\ & & I^1/Q^0 = Q^1 \end{array}$$

etc...

□

- (Derived functors) Assume that $\mathcal{A}$ has enough injectives. If $F : \mathcal{A} \rightarrow \mathcal{B}$ is a left exact functor we define

$$R^n F(A) = H^n(F(I^\bullet)) \quad (1.2)$$

where $I^\bullet$ is some injective resolution of A .

Remark 1.1.9. We have

$$\begin{aligned}
R^0 F(A) &= H^0(F(I^\bullet)) \\
&= H^0(0 \rightarrow F(I^0) \rightarrow F(I^1) \rightarrow \dots) \\
&= \text{Ker}(F(I^0) \rightarrow F(I^1)) / \text{Im}(0 \rightarrow F(I^0)) \\
&= F(A)
\end{aligned} \tag{1.3}$$

Here we used that F is left exact, so $0 \rightarrow F(A) \rightarrow F(I^0) \rightarrow F(I^1)$ is exact.

Notation 1.1.10. From now on, we will adopt the following convention: We will draw arrows away from left to right. So, index decreases from left to right in homology and increases from left to right in cohomology.

$$\begin{aligned}
(\text{Homology}) \quad & \dots \rightarrow C_2 \rightarrow C_1 \rightarrow C_0 \rightarrow 0 \\
(\text{Cohomology}) \quad & 0 \rightarrow C^0 \rightarrow C^1 \rightarrow C^2 \rightarrow \dots
\end{aligned} \tag{1.4}$$

Lemma 1.1.11. Let $f : A \rightarrow B$ be a morphism and take injective resolutions $A \rightarrow I^\bullet, B \rightarrow J^\bullet$. Then there exists lifting $f^\bullet : I^\bullet \rightarrow J^\bullet$, i.e., $\{f^n\}_{n \geq 0}$ such that

$$\begin{array}{ccccccc}
0 & \longrightarrow & A & \xrightarrow{\eta} & I^0 & \xrightarrow{d} & I^1 \xrightarrow{d} \dots \\
& & \downarrow f & & \downarrow f^0 & & \downarrow f^1 \\
0 & \longrightarrow & B & \xrightarrow{\eta} & J^0 & \xrightarrow{d} & J^1 \xrightarrow{d} \dots
\end{array}$$

Proof: First we define f^0 by the diagram below.

$$\begin{array}{ccc}
A & \xrightarrow{\eta} & I^0 \\
& \searrow \eta \circ f & \downarrow \exists f^0 \\
& & J^0
\end{array}$$

For the second block we construct the diagram

$$\begin{array}{ccccc}
I^0 & & & & I^1 \\
\downarrow & \searrow & & \swarrow & \downarrow \\
& & Q^0 = I^0 / \text{Ker}(d^0) & & \\
& & \vdots & & \\
J^0 & \xrightarrow{\quad} & & \xrightarrow{\quad} & J^1 \\
& \searrow & & \swarrow & \\
& & R^0 = J^0 / \text{Ker}(d^0) & &
\end{array}$$

The dashed morphism is defined by $Q^0 \rightarrow R^0 : [x] \mapsto [f^0(x)]$. It's well defined, because if $x - x' \in \text{Ker}(d^0) = \text{Im}(\eta)$ implies that $x - x' = \eta(a)$. So $f^0(x) - f^0(x') = f^0\eta(a) = \eta f(a)$ and $d\eta = 0$. Now we define f^1 from the diagram

$$\begin{array}{ccc} & & I^1 \\ & \nearrow & \downarrow \exists f^1 \\ Q^0 & & J^1 \\ & \searrow & \nearrow \\ & & R^0 \end{array}$$

and etc...

□

Lemma 1.1.12. Any two liftings are homotopic.

Proof: We want to find h such that makes the diagram below commutes.

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{\eta} & I^0 & \xrightarrow{d} & I^1 \xrightarrow{d} \dots \\ & \searrow 0 & \downarrow 0 & \swarrow h^0 & \downarrow h^0 - g^0 & \swarrow h^1 & \downarrow f^1 - g^1 \\ 0 & \longrightarrow & B & \xrightarrow{\eta} & J^0 & \xrightarrow{d} & J^1 \xrightarrow{d} \dots \end{array}$$

For $n = 0$, we need $h^1 \circ d^0 + d^1 \circ h^0 = g^0 - f^0$. It's clear that $h^0 = 0$ makes the first block commutes, then $h^1 \circ d^0 = g^0 - f^0$. However, we have

$$\begin{array}{ccc} & Q^0 = I^0 / \text{Ker}(d^0) & \\ \nearrow & & \searrow \\ I^0 & \xrightarrow{\quad} & I^1 \\ \downarrow & \nwarrow & \downarrow \\ J^0 & \xleftarrow{\quad} & \end{array}$$

As before, we define $Q^0 \rightarrow J^0 : x + \text{Ker}(d^0) \mapsto f^0(x) - g^0(x)$. Now it's well defined because if $x \in \text{Ker}(d^0) = \text{Im}(\eta)$, we get $f^0(x) - g^0(x) = (f^0 - g^0)\eta(a) = 0$. □

Theorem 1.1.13. (1) $RF^n(A)$ independent of choice of resolution $I^\bullet$;
(2) $f : A \rightarrow B$ induces $R^n F(A) \rightarrow R^n F(B)$;
(3) If $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is a SES, then there exists a long exact sequence:

$$\begin{array}{ccccccc}
& & & \cdots & \longleftarrow & & \\
& & & & & \nearrow & \\
& & R^1 F(A) & \longrightarrow & R^1 F(B) & \longrightarrow & R^1 F(C) \\
& & & \nwarrow & & \searrow & \\
& 0 & \longrightarrow & F(A) & \longrightarrow & F(B) & \longrightarrow & F(C)
\end{array}$$

Proof: Let $A \in \mathcal{C}$ and take two injective resolutions $I^\bullet, J^\bullet$ of A . Then the identity morphism $A \rightarrow A$ lifts to morphisms $f : I^\bullet \rightarrow J^\bullet, g : J^\bullet \rightarrow I^\bullet$. In particular $fg : I^\bullet \rightarrow I^\bullet$ is a complex morphism, so it must be homotopic to the identity morphism, i.e., $fg \sim \text{Id}_{I^\bullet} \Rightarrow F(f) \circ F(g) \sim \text{Id}_{F(I^\bullet)}$. In a similar way we see that $F(g) \circ F(f) \sim \text{Id}_{F(J^\bullet)}$, so $H^n(F(I^\bullet)) \cong H^n(F(J^\bullet))$. The second one also follows directly from the lemmas above.

TODO: Prove (3)

- (Projectives)
- (Group cohomology)
- (Lie algebra cohomology)

Chapter 2

Exercises

2.1 List 1

Problem 1. State the definition of monomorphism, and prove that an injective function of sets is a monomorphism in $\mathbf{Set}$.

Solution: A monomorphism $f \in \mathbf{Hom}_{\mathcal{C}}(B, C)$ is a morphism that is left cancelative in the sense that for any object A and morphisms $g, h \in \mathbf{Hom}_{\mathcal{C}}(A, B)$, we have:

$$fg = fh \Rightarrow g = h. \quad (2.1)$$

Now, if $\mathcal{C} = \mathbf{Set}$, the equality $fg = fh$ implies $f(g(a)) = f(h(a))$, $\forall a \in A$ and if f injective, by definition, we have $g(a) = h(a)$, $\forall a \in A$, i.e., $g = h$.

Problem 2. Prove that $h : \mathbb{Z} \hookrightarrow \mathbb{Q}$ is an epimorphism in $\mathbf{Ring}$.

Solution: Let $f, g : \mathbb{Q} \rightarrow R$ be two ring morphisms such that $fh = gh$, i.e., $f(n) = g(n)$, $\forall n \in \mathbb{Z}$. Here, we assume that a morphism in $\mathbf{Ring}$ send unity to unity, so $f(1) = g(1) = 1_R$. Furthermore, since $\mathbb{Q}$ is a field, $f : \mathbb{Q} \hookrightarrow R$ and $f(\mathbb{Q}^\times) \subset R^\times$. In particular $m1_R \in R^\times$. Now we have

$$m1_R(f(\frac{1}{m}) - g(\frac{1}{m})) = 0 \Rightarrow f(\frac{1}{m}) = g(\frac{1}{m}). \quad (2.2)$$

This proves that $f = g$.

Problem 3. Write the proof that any two terminal objects in a category $\mathcal{C}$ are isomorphic to each other.

Solution: Let T, T' be two terminal objects in $\mathcal{C}$. Let $f \in \text{Hom}_{\mathcal{C}}(T, T')$ and $T' \in \text{Hom}_{\mathcal{C}}(T', T)$, then $f \circ g \in \text{Hom}_{\mathcal{C}}(T, T) = \{\mathbb{1}_T\}$. Therefore $fg = \mathbb{1}_T$ and in a similar way we see that $gf = \mathbb{1}_{T'}$. This proves that f is an isomorphism.

Problem 4. State an "explicit" definition of the free product $G * H$ of groups G and H . Prove that the free product is a coproduct in Grp .

Solution: Let $A = G \amalg H$, we define

$$G * H = \coprod_{n \geq 1} A^n / \sim \quad (2.3)$$

where we the equivalence relation is defined by

$$\begin{aligned} (a_1, \dots, a_i, a_{i+1}, \dots, a_n) &\sim (a_1, \dots, a_i a_{i+1}, \dots, a_n) \text{ if } a_1, a_2 \in G \text{ or } a_1, a_2 \in H \\ (a_1, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_n) &\sim (a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n) \text{ if } a_i = 1_G \text{ or } a_i = 1_H \end{aligned} \quad (2.4)$$

The multiplication is defined by

$$[(a_1, \dots, a_n)] * [(b_1, \dots, b_m)] = [(a_1, \dots, a_n, b_1, \dots, b_m)]. \quad (2.5)$$

By the equivalence relations defined above, the multiplication rule is clearly well defined.

We see that $(G * H, *)$ is a group. The associativity and existence of unity is clear and the inverse operation is defined by

$$[(a_1, a_2, \dots, a_n)]^{-1} = [(a_n^{-1}, \dots, a_2^{-1}, a_1^{-1})]. \quad (2.6)$$

Now we define morphisms

$$i_1 : G \rightarrow G * H : g \mapsto [(g)], \quad i_2 : H \rightarrow G * H : h \mapsto [(h)]. \quad (2.7)$$

We claim that $(G * H, i_1, i_2)$ is a coproduct of G and H . Suppose that we have morphisms $f_1 : G \rightarrow K, f_2 : H \rightarrow K$, then we define a morphism

$$f : G * H \rightarrow K : [(a_1, \dots, a_n)] \mapsto f_{j_1}(a_1) \cdots f_{j_n}(a_n) \quad (2.8)$$


where we let $j_k = 1$ (resp. 2) if $a_k \in G$ (resp. H). Furthermore, we see that $f i_j = f_j$, $j = 1, 2$. For uniqueness, we observe that if $h : G * H \rightarrow K$ is a morphism, such that $h i_j = f_j$, $j = 1, 2$, then $h([(g)]) = f_1(g)$ and $h([(h)]) = f_2(h)$. So

$$h([(a_1, \dots, a_n)]) = h([(a_1)] * \cdots * [(a_n)]) = h([(a_1)]) \cdots h([(a_n)]) \quad (2.9)$$


and we see that $h([(a_k)]) = f_{j_k}(a_k)$ as defined above, therefore $h = f$.

Problem 5. Let $\mathcal{C}$ be a pre-additive category and $X, Y \in \mathcal{C}$. Show that if a product $X \times Y$ of X and Y exists, then $X \times Y$ is also a coproduct of X and Y .

Solution: A product $X \times Y$ comes with projections $p_1 : X \times Y \rightarrow X$ and $p_2 : X \times Y \rightarrow Y$ such that the following universal property is satisfied.



So we define $i_1 : X \rightarrow X \times Y$ and $i_2 : Y \rightarrow X \times Y$ using this universal property:




So we have the following properties

$$\begin{cases} 1_X = p_1 i_1, & p_2 i_1 = 0_{XY} \\ 1_Y = p_2 i_2, & p_1 i_2 = 0_{YX} \end{cases} \quad (2.10)$$


Now, we claim that

$$i_1 p_1 + i_2 p_2 = 1_{X \times Y} \quad (*) \quad (2.11)$$

which can be seen by the diagram below and the universal property of the product.



Now, suppose that we have morphisms $f_1 : X \rightarrow Z, f_2 : Y \rightarrow Z$. Then if we define $h = f_1 p_1 + f_2 p_2$, the diagram below commutes.




Furthermore, relations (2.10) and equation (*) guarantees that this is the unique expression possible for h that makes the above diagram commutes.

Problem 6. Show that existence of binary products implies existence of all finite n -ary products for $n \geq 2$.

Solution: By a finite n -ary product ($n \geq 2$), we mean an object $X = \prod_{i=1}^n X_i$ with morphisms $p_i : X \rightarrow X_i$ satisfying the following universal property. Take a collection of morphisms $f_i : Y \rightarrow X_i$, ($i = 1, \dots, n$), then there exist an unique morphism $f : Y \rightarrow X$ such that $p_i f = f_i$, $\forall i$. By induction, assume that there exist all finite $(n-1)$ -ary products and let $X' = \prod_{i=1}^{n-1} X_i$. We want to show that $X = X' \times X_n$ satisfies the universal property for a n -ary product. The projections are obtained as follows: we have the projection $p_n : X \rightarrow X_n$ and for $i < n$ we compose the projection $q : X \rightarrow X'$ with the i th projection $p'_i : X' \rightarrow X_i$ to get $p_i : X \rightarrow X_i$.

Now assume we have morphisms $f_i : Y \rightarrow X_i$, ($i = 1, \dots, n$). Then, by the universal property of the finite $(n-1)$ -ary product, there exists a (unique) morphism $f' : Y \rightarrow X'$ such that $p'_i f' = f_i$, $i < n$. Now we have the diagram:



Now we observe that

$$p_i h = (p'_i q) h = p'_i (q h) = p'_i f' = f_i, \quad i < n. \quad (2.12)$$

This show existence of h . Now, for uniqueness, suppose that h' is such that $p_i h' = f_i$, $\forall i$. Then, for $i < n$ we have $p'_i (h' q) = f_i$. So by the universal property of the finite $(n-1)$ -ary product X' , we must have $h' q = f'$. Then, by applying the universal property of the product $X = X' \times X_n$, we must have $h' = h$.

Problem 7. Consider a composition

$$A \longrightarrow B \longrightarrow C$$

in an abelian category, in which $g \circ f = 0$. Using only universal properties, show that there is a canonical morphism from $\text{im}(f)$ to $\text{ker}(g)$. [Recall $\text{im}(f) = \text{ker}(\text{coker}(f))$ by definition.]

Solution: From the universal property of the cokernel we have the diagram.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow g \\ & \text{Coker}(f) & \\ \downarrow & \nearrow & \downarrow \\ 0 & & C \end{array}$$

(Note: In the original diagram, there is a diagonal arrow from A to $\text{Coker}(f)$ and a dashed arrow from $\text{Coker}(f)$ to C labeled $\exists! \tilde{g}$. The arrow from B to $\text{Coker}(f)$ is labeled p .)

Now, by the universal property of the kernel, we have:

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & & \\ \downarrow & \searrow \exists! \tilde{f} & \nearrow i & \searrow g & \\ & \text{Ker}(p) = \text{Im}(f) & & & C \\ \downarrow & & \downarrow p & \nearrow \tilde{g} & \\ 0 & \xrightarrow{\quad} & \text{coker}(f) & & \end{array}$$

Finally, by applying the universal property of the kernel again, we have:

$$\begin{array}{ccc} \text{Im}(f) & \xrightarrow{i} & B \\ \downarrow & \searrow \exists! \gamma & \downarrow g \\ & \text{Ker}(g) & \\ \downarrow & \nearrow & \downarrow \\ 0 & & C \end{array}$$

as desired.

Problem 8. Let $G : \mathbf{Ab} \rightarrow \mathbf{Set}$ be the forgetful functor, and let $F : \mathbf{Set} \rightarrow \mathbf{Ab}$ be the functor of forming a free abelian group. Show that F is left adjoint to G .

Solution: We have the map

$$\text{Hom}_{\mathbf{Ab}}(F(X), Y) \rightarrow \text{Hom}_{\mathbf{Set}}(X, G(Y)) : f \mapsto f|_X \quad (2.13)$$

Here, we let $F(X) = \{f : X \rightarrow \mathbb{Z} : f(x) \neq 0 \text{ for a finite number of } x \in X\}$ and denote $\hat{x} \in F(X)$ for the map defined by $\hat{x}(y) = \delta_{xy}$. So we see that the set theoretic map $\iota : X \rightarrow F(X) : x \rightarrow \hat{x}$ is an injection and this justifies the notation $f|_X$ used in (*). Moreover, show that this map is an isomorphism is equivalent to show that we have the following universal property: For any set map of sets $f : X \rightarrow G(Y)$, there exists a unique morphism $\hat{f} : F(X) \rightarrow Y$ such that $\hat{f}\iota = f$. So we have no choice but define $\hat{f}(\hat{x}) = f(x)$ and extend to $F(X)$ by $\mathbb{Z}$ -linearity, this defines $\hat{f}$ uniquely.

Problem 9. Compute $(\mathbb{Z}/5) \otimes_{\mathbb{Z}} (\mathbb{Z}/6)$ and $(\mathbb{Z}/p^2) \otimes_{\mathbb{Z}} (\mathbb{Z}/p)$ where p is a prime.

Solution: First we claim that $(\mathbb{Z}/5) \otimes_{\mathbb{Z}} (\mathbb{Z}/6) = (0)$. In fact, if $a \otimes b \in (\mathbb{Z}/5) \otimes_{\mathbb{Z}} (\mathbb{Z}/6)$, then since $6^2 \equiv 1 \pmod{5}$, we have

$$a \otimes b = a \cdot 6^2 \otimes b = a \cdot 6 \otimes 6 \cdot b = a \cdot 6 \otimes 0 = 0. \quad (2.14)$$

For the second group, we claim that $(\mathbb{Z}/p^2) \otimes_{\mathbb{Z}} (\mathbb{Z}/p) \cong \mathbb{Z}/p$. To show this, we show that $\mathbb{Z}/p$ satisfies the universal property of the tensor product (here we will use the notation of problem 10). Let $\varphi_0 : \mathbb{Z}/p^2 \times \mathbb{Z}/p \rightarrow \mathbb{Z}/p : ([m], [n]) \mapsto [mn]$, it's a well defined balanced map, i.e., $\varphi_0 \in \text{Bal}_{A,B}(\mathbb{Z}/p)$. In particular $\mathbb{Z}/p$ is generated by the image of φ_0 . Now, let $f \in \text{Bal}_{A,B}(X)$, we want to define $\tilde{f}$ in such a way the following diagram commutes.

$$\begin{array}{ccc} \mathbb{Z}/p^2 \times \mathbb{Z}/p & \xrightarrow{\varphi_0} & \mathbb{Z}/p \\ f \downarrow & \swarrow \tilde{f} & \\ X & & \end{array}$$

So we set $\tilde{f}([n]) = f(1, [n])$, which is well defined while makes the diagram commutes.

Problem 10. Let R be a unital ring. In order to formulate the definition of tensor product of R -modules, we introduce a definition:

definition 0.1. Let A be a right R -module and B a left R -module, and X an abelian group. A balanced map $f \in \text{Bal}_{A,B}(X)$ is a function $f : A \times B \rightarrow X$ such that

$$f(a_1 + a_2, b) = f(a_1, b) + f(a_2, b) \quad \text{for all } a_1, a_2 \in A \text{ and } b \in B$$

$$f(a, b_1 + b_2) = f(a, b_1) + f(a, b_2) \quad \text{for all } a \in A \text{ and } b_1, b_2 \in B$$

$$f(ar, b) = f(a, rb) \quad \text{for all } a \in A, b \in B \text{ and } r \in R.$$

Now a tensor product of a right R -module A with a left R -module B is an abelian group $A \otimes_R B$ together with a $\varphi_0 \in \text{Bal}_{A,B}(A \otimes_R B)$, such that for any balanced map f as below, there exists a unique linear map $\tilde{f}$ making the diagram commute.

$$\begin{array}{ccc} A \times B & \xrightarrow{\varphi_0} & A \otimes_R B \\ & \searrow f & \swarrow \tilde{f} \\ & X & \end{array}$$

Verify that

$$A \otimes_R B \cong Q = \frac{A \otimes_{\mathbb{Z}} B}{\langle ar \otimes b - a \otimes rb \mid a \in A, b \in B, r \in R \rangle}.$$

Solution: To prove the asserted isomorphism, we show that the quotient of abelian groups Q satisfies the universal property of the tensor product $A \otimes_R B$. Let $f \in \text{Bal}_{A,B}(X)$, then f is balanced by viewing A, B as abelian groups. By the universal property of $A \otimes_{\mathbb{Z}} B$, there exist maps $\hat{\varphi}_0$ and $\hat{f}$, such that:

$$\begin{array}{ccc} A \times B & \xrightarrow{\hat{\varphi}_0} & A \otimes_{\mathbb{Z}} B \\ & \searrow f & \swarrow \hat{f} \\ & X & \end{array}$$

Now, since $f \in \text{Bal}_{A,B}(X)$, we have

$$\text{Ker}(\hat{f}) \supset \langle ar \otimes b - a \otimes rb \mid a \in A, b \in B, r \in R \rangle. \quad (2.15)$$

Then, there exists $\tilde{f}$ such that the following diagram commutes:

$$\begin{array}{ccccc} A \times B & \xrightarrow{\hat{\varphi}_0} & A \otimes_{\mathbb{Z}} B & \xrightarrow{p} & Q \\ & \searrow f & \swarrow \hat{f} & & \uparrow \tilde{f} \\ & X & & & \end{array}$$

Now, by construction, the quotient Q is generated by the image of the map $\varphi_0 = p\hat{\varphi}_0$ and since $f = \tilde{f}\varphi_0$ we see that $\tilde{f}$ is uniquely determined by f . Therefore (Q, φ_0) satisfies the universal property for the tensor product $A \otimes_R B$ and $Q \cong A \otimes_R B$.

2.2 List 2

Let A be an abelian group, and consider the action

$$A \times A \longrightarrow A$$

given by addition in the group. Does this action make A into an A -module? Justify your answer.

The homologies $H_n(G, A)$ of a group G with coefficients in the G -module A are the derived functors of the coinvariants functor

$$(-)_G = \mathbb{Z} \otimes_{\mathbb{Z}G} (-).$$

(where $\mathbb{Z}$ is the trivial G -module). Using a convenient projective resolution of $\mathbb{Z}$, or otherwise, compute group homology $H_n(G, A)$ where G is the cyclic group $\mathbb{Z}/n$ and A is a trivial G -module.

The Curry adjunction: Let $A \in \mathbf{Set}$ be a fixed set, and let $L : \mathbf{Set} \rightarrow \mathbf{Set}$ and $R : \mathbf{Set} \rightarrow \mathbf{Set}$ be the functors

$$L(X) = X \times A \quad \text{and} \quad R(X) = \text{Hom}_{\mathbf{Set}}(A, X).$$

Prove that L is left adjoint to R , i.e., construct a natural isomorphism

$$\text{curry} : \text{Hom}_{\mathbf{Set}}(X \times A, Y) \longrightarrow \text{Hom}_{\mathbf{Set}}(X, \text{Hom}_{\mathbf{Set}}(A, Y)).$$

Prove the following lemma from class.

Lemma 0.1. If $L : \mathcal{A} \rightarrow \mathcal{B}$ is a functor with right adjoint R , and

$$\eta : \text{Id} \Rightarrow R \circ L \quad \text{and} \quad \varepsilon : L \circ R \Rightarrow \text{Id}$$

are as in class (see Corollary 3.4.9). Then

$$\varepsilon_{L(X)} \circ L(\eta_X) = \text{Id}_{L(X)}.$$

Hint: There are several equivalent definitions of adjunction. The one I have given is in terms of a natural isomorphism

$$\beta_{X,Y} : \text{Hom}(L(X), Y) \longrightarrow \text{Hom}(X, R(Y)).$$

There are other definitions, in particular one that is in terms of two natural transformations

$$\eta : \text{Id}_{\mathcal{A}} \Rightarrow R \circ L, \quad \varepsilon : L \circ R \Rightarrow \text{Id}_{\mathcal{B}}.$$

One of the axioms in this second definition is

$$\begin{array}{ccc} L & \xrightarrow{L(\eta)} & LRL \\ & \searrow \text{Id}_L & \downarrow \varepsilon_L \\ & & L \end{array}$$

The identity to be proved follows fairly easily from this axiom. So the exercise, really, is to understand this second definition of adjoint functors, and why it follows from the first.

Deduce the classical version of Hilbert 90 from the more general cohomological statement, using the “convenient complex” for $H^*(\mathbb{Z}/n, -)$ that we met in class:

Theorem 0.2. Let L/K be a Galois extension, and suppose that

$$G = \text{Gal}(L/K)$$

is cyclic,

$$G = \langle \sigma \rangle \cong \mathbb{Z}/n.$$

Then for every $\alpha \in L$ of norm $N(\alpha) = 1$, there exists $\beta \in L$ such that

$$\alpha = \frac{\sigma(\beta)}{\beta}.$$

Let G be a group and $N \subset G$ a normal subgroup. If A is a G -module, verify that the invariants A^N carries a natural action of the quotient group G/N . Thus we may consider the functor

$$(-)^N : G\text{-mod} \longrightarrow (G/N)\text{-mod}.$$

Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a left exact functor, and $E : \mathcal{B} \rightarrow \mathcal{C}$ an exact functor. Assume $\mathcal{A}$ has enough injectives. Prove that

$$[R^n(E \circ F)](A) \cong [E \circ R^n F](A)$$

for all $A \in \mathcal{A}$. Deduce that there exists a natural action of G/N on the cohomology

$$H^*(N, A)$$

(where G, N, A as in Exercise 6).

Let G be a group and $N \subset G$ a normal subgroup, and A a G -module. Consider the bar complex, whose terms are

$$C^p(N, A) = \text{Maps}(N^p \rightarrow A).$$

Check that

$$[g \cdot \varphi](n_1, \dots, n_p) = g\varphi(g^{-1}n_1g, \dots, g^{-1}n_pg),$$

for all $p \in \mathbb{Z}_+$, defines a G -action, and a G/N -action, on

$$H^*(N, A).$$

We continue in the context of the previous exercise. But now G is the symmetric group S_3 and $N \cong \mathbb{Z}/3$ is the normal subgroup generated by a 3-cycle. Consider $\mathbb{Z}$ equipped with the trivial G -module structure. Determine the G/N -module structure of the

$$H^k(N, \mathbb{Z}).$$

Optional challenge question Show that the G/N -actions of Exercises 7 and 8, coincide.