

# Discrete Algebraic Structures

WiSe 2025/2026

Prof. Dr. Antoine Wiehe  
Research Group for Theoretical Computer Science



Exam: 10.03.

- What a relation is
- How to draw a relation  $R \subseteq A \times B$
- How to compute the composition and transpose of binary relations
- The basic properties  
(anti)reflexivity, (anti)symmetry, transitivity
- How to compute the closures

Relations we encounter in “real life” are often of one of two types:

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:


- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company

Relations we encounter in “real life” are often of one of two types:

$$\mathcal{R} \subseteq A \times A$$

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




SameCompany  $\subseteq$  People  $\times$  People

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$



Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company




$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Things can be ordered:

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:

- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company



$\text{SameCompany} \subseteq \text{People} \times \text{People}$


Things can be ordered:

- $x$  appears before  $y$  in the dictionary
- The string  $x$  is not longer than the string  $y$
- $x$  is taller than  $y$
- $y$  is a multiple of  $x$

Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:


- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company



$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Things can be ordered:

- $x$  appears before  $y$  in the dictionary
- The string  $x$  is not longer than the string  $y$
- $x$  is taller than  $y$
- $y$  is a multiple of  $x$






Relations we encounter in “real life” are often of one of two types:

Things can share a **common property**:



- $x = y$ : share the same value
- $x, y$  are numbers that end with the same digit
- $x, y$  are in the same family
- $x, y$  work in the same company



$\text{SameCompany} \subseteq \text{People} \times \text{People}$

Things can be ordered:

- $x$  appears before  $y$  in the dictionary
- The string  $x$  is not longer than the string  $y$
- $x$  is taller than  $y$
- $y$  is a multiple of  $x$



**Definition.** A relation  $R \subseteq A \times A$  is:

- **reflexive**: if for all  $a \in A$ ,  $(a, a) \in R$
- **antireflexive**: if for all  $a \in A$ ,  $(a, a) \notin R$
- **symmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$ , then  $(b, a) \in R$
- **antisymmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$  and  $a \neq b$  then  $(b, a) \notin R$
- **transitive**: if for all  $a, b, c \in A$ , if  $(a, b) \in R$  and  $(b, c) \in R$  then  $(a, c) \in R$ .

**Definition.** A relation  $R \subseteq A \times A$  is:

- **reflexive**: if for all  $a \in A$ ,  $(a, a) \in R$
- **antireflexive**: if for all  $a \in A$ ,  $(a, a) \notin R$
- **symmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$ , then  $(b, a) \in R$
- **antisymmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$  and  $a \neq b$  then  $(b, a) \notin R$
- **transitive**: if for all  $a, b, c \in A$ , if  $(a, b) \in R$  and  $(b, c) \in R$  then  $(a, c) \in R$ .

	Reflexive	Antireflexive	Symmetric	Antisymmetric	Transitive
Equivalence relation					
Order					
Strict order					

**Definition.** A relation  $R \subseteq A \times A$  is:

- **reflexive**: if for all  $a \in A$ ,  $(a, a) \in R$
- **antireflexive**: if for all  $a \in A$ ,  $(a, a) \notin R$
- **symmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$ , then  $(b, a) \in R$
- **antisymmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$  and  $a \neq b$  then  $(b, a) \notin R$
- **transitive**: if for all  $a, b, c \in A$ , if  $(a, b) \in R$  and  $(b, c) \in R$  then  $(a, c) \in R$ .

	Reflexive	Antireflexive	Symmetric	Antisymmetric	Transitive
Equivalence relation	✓		✓		✓
Order	✓			✓	✓
Strict order		✓		✓	✓

**Definition.** A relation  $R \subseteq A \times A$  is:

- **reflexive**: if for all  $a \in A$ ,  $(a, a) \in R$
- **antireflexive**: if for all  $a \in A$ ,  $(a, a) \notin R$
- **symmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$ , then  $(b, a) \in R$
- **antisymmetric**: if for all  $a, b \in A$ , if  $(a, b) \in R$  and  $a \neq b$  then  $(b, a) \notin R$
- **transitive**: if for all  $a, b, c \in A$ , if  $(a, b) \in R$  and  $(b, c) \in R$  then  $(a, c) \in R$ .

	Reflexive	Antireflexive	Symmetric	Antisymmetric	Transitive
Equivalence relation	✓		✓		✓
Order	✓			✓	✓
Strict order		✓		✓	✓
Quasi-order	✓				✓

# Equivalence relations

**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$   $\{0, +, -\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$   $\{\text{even, odd}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$   $\mathbb{N}_0$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$



**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$   $\{0, +, -\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$   $\{\text{even, odd}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$   $\mathbb{N}_0$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$

We use symbols like  $\sim, \simeq, \equiv$  for relations that are equivalence relations

We then write  $a \simeq b$  instead of  $(a, b) \in \simeq$


**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$   $\{0, +, -\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$   $\{\text{even}, \text{odd}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$   $\mathbb{N}_0$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$

We use symbols like  $\sim, \simeq, \equiv$  for relations that are equivalence relations

We then write  $a \simeq b$  instead of  $(a, b) \in \simeq$




**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$   $\{0, +, -\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$   $\{\text{even}, \text{odd}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$   $\mathbb{N}_0$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$

We use symbols like  $\sim, \simeq, \equiv$  for relations that are equivalence relations

We then write  $a \simeq b$  instead of  $(a, b) \in \simeq$




**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$   $\{0, +, -\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$   $\{\text{even}, \text{odd}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$   $\mathbb{N}_0$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$

We use symbols like  $\sim, \simeq, \equiv$  for relations that are equivalence relations

We then write  $a \simeq b$  instead of  $(a, b) \in \simeq$




**Definition.** A relation  $R \subseteq A \times A$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$   $\{0, +, -\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$   $\{\text{even}, \text{odd}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$   $\mathbb{N}_0$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$

We use symbols like  $\sim, \simeq, \equiv$  for relations that are equivalence relations

We then write  $a \simeq b$  instead of  $(a, b) \in \simeq$




**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is


$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$

**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$



0



+

$x \sim y$  if  
 $x, y$  have same  
sign

$$[-1]_{\sim} = \{-2, -3, -4, -1, \dots\}$$


$$[0]_{\sim} = \{0\}$$

$$[1]_{\sim} = \{1, 2, 3, 4, \dots\}$$

$$[-2]_{\sim} = \{-1, -2, -3, -4, \dots\}$$

**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$



Define  $a \sim b$  on  $\mathbb{N}_0$  as “ $a$  and  $b$  have the same parity”.

What is  $[0]_{\sim}$ ?

~~$\emptyset$~~    
  ~~$\{0\}$~~    
  ~~$\{1\}$~~    
  $\{0, 2, 4, \dots\}$    
  $\{0, 2, 4, \dots\}$







**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$

**Theorem.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a, b \in A$ . The following are equivalent:

1.  $a \sim b$
2.  $[a]_{\sim} = [b]_{\sim}$
3.  $[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$



**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$

**Theorem.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a, b \in A$ . The following are equivalent:


1.  $a \sim b$
2.  $[a]_{\sim} = [b]_{\sim}$
3.  $[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$

Theoretically, one needs to prove **6 implications**:

$$\varphi_1 \wedge \varphi_2 \wedge \dots \wedge \varphi_6$$

III

$$\varphi_1 \wedge \varphi_2 \wedge \varphi_3$$




**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$

**Theorem.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a, b \in A$ . The following are equivalent:

1.  $a \sim b$
2.  $[a]_{\sim} = [b]_{\sim}$
3.  $[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$

Theoretically, one needs to prove **6 implications**:



By **logical equivalence**, it suffices to prove **3 implications**!

**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$

**Theorem.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a, b \in A$ . The following are equivalent:

1.  $a \sim b$
2.  $[a]_{\sim} = [b]_{\sim}$
3.  $[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$

Theoretically, one needs to prove **6 implications**:

Hyp:  $a \sim b$   
 $\text{ff}$ :  $\sim$  is an eq. relation

- $x \in [a]_{\sim}$  so in fact  $x \sim a$
- $x \sim b$  by transitivity of  $\sim$

$$a \sim b$$



$$[a]_{\sim} = [b]_{\sim}$$

Goal:  $x \in [b]_{\sim}$

By **logical equivalence**, it suffices to prove **3 implications**!

**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$

**Theorem.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a, b \in A$ . The following are equivalent:

1.  $a \sim b$
2.  $[a]_{\sim} = [b]_{\sim}$
3.  $[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$

Theoretically, one needs to prove **6 implications**:

$$a \in [a]_{\sim} \neq \emptyset$$

$$[a]_{\sim} \cap [b]_{\sim} \neq \emptyset \quad \Longleftarrow \quad [a]_{\sim} = [b]_{\sim}$$

By **logical equivalence**, it suffices to prove **3 implications**!


**Definition.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a \in A$ . The **equivalence class** of  $a$  is

$$[a]_{\sim} = \{b \in A \mid a \sim b\}$$


**Theorem.** Let  $\sim$  be an equivalence relation on  $A$ , and let  $a, b \in A$ . The following are equivalent:

1.  $a \sim b$
2.  $[a]_{\sim} = [b]_{\sim}$
3.  $[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$

Theoretically, one needs to prove **6 implications**:



$$[a]_{\sim} \cap [b]_{\sim} \neq \emptyset$$



By **logical equivalence**, it suffices to prove **3 implications**!


**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\forall S \in P \ S \neq \emptyset$$

$$\forall a \in A \exists S \in P : a \in S$$

$$\forall S, T \in P (S \neq T \Rightarrow S \cap T = \emptyset)$$




$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\}$   
is a partition of  $\mathbb{Z}$

**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\begin{aligned} \forall S \in P \quad S &\neq \emptyset \\ \forall a \in A \exists S \in P : a &\in S \\ \forall S, T \in P (S \neq T \Rightarrow S \cap T &= \emptyset) \end{aligned}$$



$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\}$   
is a partition of  $\mathbb{Z}$

$f$ : equivalence relation  $\sim \mapsto$  partition  $\{[a]_\sim \mid a \in A\}$




**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\forall S \in P \ S \neq \emptyset$$

$$\forall a \in A \exists S \in P : a \in S$$

$$\forall S, T \in P (S \neq T \Rightarrow S \cap T = \emptyset)$$




$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\}$   
is a partition of  $\mathbb{Z}$

$f$ : equivalence relation  $\sim \mapsto$  partition  $\{[a]_\sim \mid a \in A\}$

**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\begin{aligned} \forall S \in P \quad S &\neq \emptyset \\ \forall a \in A \exists S \in P : a &\in S \\ \forall S, T \in P (S \neq T \Rightarrow S \cap T &= \emptyset) \end{aligned}$$



$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\}$   
is a partition of  $\mathbb{Z}$


$f$ : equivalence relation  $\sim \mapsto$  partition  $\{[a]_\sim \mid a \in A\}$

$g$ : partition  $P \mapsto \{(a, b) \in A^2 \mid a, b \text{ belong to the same part}\}$

**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\begin{aligned} &\forall S \in P \quad S \neq \emptyset \\ &\forall a \in A \exists S \in P : a \in S \\ &\forall S, T \in P (S \neq T \Rightarrow S \cap T = \emptyset) \end{aligned}$$



$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\}$   
is a partition of  $\mathbb{Z}$


$f$ : equivalence relation  $\sim \mapsto$  partition  $\{[a]_\sim \mid a \in A\}$

$g$ : partition  $P \mapsto \{(a, b) \in A^2 \mid a, b \text{ belong to the same part}\}$

**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\begin{aligned} \forall S \in P \quad S &\neq \emptyset \\ \forall a \in A \exists S \in P : a &\in S \\ \forall S, T \in P (S \neq T \Rightarrow S \cap T &= \emptyset) \end{aligned}$$



$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\}$   
is a partition of  $\mathbb{Z}$

$f$ : equivalence relation  $\sim \mapsto$  partition  $\{[a]_{\sim} \mid a \in A\}$

$g$ : partition  $P \mapsto \{(a, b) \in A^2 \mid a, b \text{ belong to the same part}\}$

these two functions are inverse of each other!


$$(g \circ f)(\sim) = \sim$$

$$(f \circ g)(P) = P$$

**Definition.** Let  $A$  be a set. A **partition** of  $A$  is a set  $P$  of **subsets of  $A$** , called the **parts**, such that:

- the parts are not empty
- every  $a \in A$  belongs to one of the parts of  $P$
- distinct parts in  $P$  are disjoint.

$$\begin{aligned} \forall S \in P \quad S &\neq \emptyset \\ \forall a \in A \exists S \in P : a &\in S \\ \forall S, T \in P (S \neq T \Rightarrow S \cap T &= \emptyset) \end{aligned}$$



$P = \{\mathbb{Z}_-, \{0\}, \mathbb{Z}_+\} = \mathbb{Z}/\sim$   
is a partition of  $\mathbb{Z}$

We have proved:

**Theorem.** There exists a bijection between the set of all equivalence relations on  $A$ , and the set of all partitions of  $A$ .

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$

Partitions


$\{0, +, -\}$

$\{\text{even}, \text{odd}\}$

$\mathbb{N}_0$

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$




$\{0, +, -\}$   
 $\{\text{even, odd}\}$   
 $\mathbb{N}_0$

partition associated  
with the equivalence  
relation

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$



$\{0, +, -\}$   
 $\{\text{even, odd}\}$   
 $\mathbb{N}_0$


partition associated  
with the equivalence  
relation

“thing” shared by the  
elements that are  
equivalent



**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$




partition associated  
with the equivalence  
relation

“thing” shared by the  
elements that are  
equivalent

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$



partition associated  
with the equivalence  
relation

“thing” shared by the  
elements that are  
equivalent

**Notation.** We write  $A/\sim$  for the partition of  $A$  given by the equivalence classes of  $\sim$ .


$$A/\sim = \{[a]_\sim \mid a \in A\}$$

Partitions and quotients will appear many many times in your studies, for example:

- modular arithmetic, Lagrange’s theorem (later this semester): **RSA** cryptosystem
- quotient of vector spaces,  $V/W$  (Math 2 or 3): **persistent homology**, recognizing patterns in big data
- automata theory (2nd semester)

**Examples.** All of these are equivalence relations:

- $\{(x, y) \in \mathbb{Z}^2 \mid x, y \text{ have the same sign}\}$
- $\{(x, y) \in (\mathbb{N}_0)^2 \mid x, y \text{ have the same parity}\}$
- $\{(x, y) \in \text{People}^2 \mid x, y \text{ have the same height}\}$
- $\{(\varphi, \psi) \in \text{PropositionalFormulas}^2 \mid \varphi \equiv \psi\}$
- $\{(x, y) \in A^2 \mid x = y\}$
- $A^2$



partition associated  
with the equivalence  
relation

“thing” shared by the  
elements that are  
equivalent

**Notation.** We write  $A/\sim$  for the partition of  $A$  given by the equivalence classes of  $\sim$ .

$$A/\sim = \{[a]_\sim \mid a \in A\}$$

Partitions and quotients will appear many many times in your studies, for example:

- modular arithmetic, Lagrange’s theorem (later this semester): **RSA** cryptosystem
- quotient of vector spaces,  $V/W$  (Math 2 or 3): **persistent homology**, recognizing patterns in big data
- automata theory (2nd semester)

**Intuition:** forget some differences between elements,  
only remember some specific aspects

Orders

- Generalize properties of  $(\mathbb{N}, <)$  and  $(\mathbb{Q}, <)$  to other sets
- Having an order can help to design good algorithms (binary search)
- Many problems (in math, in CS, natural sciences) are solved by computing minimal elements for some order

- Generalize properties of  $(\mathbb{N}, <)$  and  $(\mathbb{Q}, <)$  to other sets
- Having an order can help to design good algorithms (binary search)
- Many problems (in math, in CS, natural sciences) are solved by computing minimal elements for some order

**Question.** What properties does the relation  $\leq$  have (on  $\mathbb{N}, \mathbb{Z}, \dots$ )?

- Reflexive ✓
- Antireflexive ✗
- Symmetric ✗
- Antisymmetric ✓
- Transitive ✓



all the loops  
Q  
Q  
② • For all  $x$ ,  $x \leq x$

• For all  $x, y$ :  $x \leq y \Rightarrow y \leq x$

• For all  $x, y$ :  $(x \leq y \wedge y \leq x) \Rightarrow x = y$

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Notation:  $(A, R)$  is a **poset**.



**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Examples.** The following relations are all orders:

- $\{(n, m) \in \mathbb{N}^2 \mid n \leq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \geq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \text{ divides } m\}$
- $\{(S, T) \in \mathcal{P}(A)^2 \mid S \subseteq T\}$
- $\{(s, t) \in \text{BinaryStrings}^2 \mid s \text{ is a prefix of } t\}$



Notation:  $(A, R)$  is a **poset**.

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Examples.** The following relations are all orders:

- $\{(n, m) \in \mathbb{N}^2 \mid n \leq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \geq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \text{ divides } m\}$
- $\{(S, T) \in \mathcal{P}(A)^2 \mid S \subseteq T\}$
- $\{(s, t) \in \text{BinaryStrings}^2 \mid s \text{ is a prefix of } t\}$

Notation:  $(A, R)$  is a **poset**.




**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Examples.** The following relations are all orders:

- $\{(n, m) \in \mathbb{N}^2 \mid n \leq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \geq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \text{ divides } m\}$
- $\{(S, T) \in \mathcal{P}(A)^2 \mid S \subseteq T\}$
- $\{(s, t) \in \text{BinaryStrings}^2 \mid s \text{ is a prefix of } t\}$

Notation:  $(A, R)$  is a **poset**.








**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Examples.** The following relations are all orders:




- $\{(n, m) \in \mathbb{N}^2 \mid n \leq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \geq m\}$
- $\{(n, m) \in \mathbb{N}^2 \mid n \text{ divides } m\}$
- $\{(S, T) \in \mathcal{P}(A)^2 \mid S \subseteq T\}$
- $\{(s, t) \in \text{BinaryStrings}^2 \mid s \text{ is a prefix of } t\}$

Notation:  $(A, R)$  is a **poset**.






**Question.** How much information is actually necessary to “remember” the order we are drawing?






**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It’s reflexive, so we know the loops are there



**Question.** How much information is actually necessary to “remember” the order we are drawing?




- It’s reflexive, so we know the loops are there



**Question.** How much information is actually necessary to “remember” the order we are drawing?




- It’s reflexive, so we know the loops are there






**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It's reflexive, so we know the loops are there
- It's transitive, so some edges are implied by others




**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It's reflexive, so we know the loops are there
- It's transitive, so some edges are implied by others





**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It’s reflexive, so we know the loops are there
- It’s transitive, so some edges are implied by others




**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It's reflexive, so we know the loops are there
- It's transitive, so some edges are implied by others
- The increasing direction is already shown by going up, so arrows are not necessary




**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It’s reflexive, so we know the loops are there
- It’s transitive, so some edges are implied by others
- The increasing direction is already shown by going up, so arrows are not necessary



**Question.** How much information is actually necessary to “remember” the order we are drawing?




- It's reflexive, so we know the loops are there
- It's transitive, so some edges are implied by others
- The increasing direction is already shown by going up, so arrows are not necessary



**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It’s reflexive, so we know the loops are there
- It’s transitive, so some edges are implied by others
- The increasing direction is already shown by going up, so arrows are not necessary

These are the **Hasse diagrams** of the orders.



**Question.** How much information is actually necessary to “remember” the order we are drawing?

- It's reflexive, so we know the loops are there
- It's transitive, so some edges are implied by others
- The increasing direction is already shown by going up, so arrows are not necessary

These are the **Hasse diagrams** of the orders.

In general: consider an order  $\leq$  on  $A$  and  $a, b \in A$  distinct with  $a \leq b$

We say  $b$  **covers**  $a$  if there is nothing between  $a$  and  $b$

$$\forall c \in A (a \leq c \leq b \Rightarrow c \in \{a, b\})$$

**Definition.** If  $A$  is finite, then a **Hasse diagram** of  $\leq$  is a drawing of  $\leq_{\text{cover}} = \{(a, b) \in A^2 \mid b \text{ covers } a\}$ , where if  $a \leq_{\text{cover}} b$ , then  $b$  is above  $a$ .



We say  $b$  **covers**  $a$  if there is nothing between  $a$  and  $b$

$$\forall c \in A (a \leq c \leq b \Rightarrow c \in \{a, b\})$$

**Definition.** If  $A$  is finite, then a **Hasse diagram** of  $\leq$  is a drawing of  $\leq_{\text{cover}} = \{(a, b) \in A^2 \mid b \text{ covers } a\}$ , where if  $a \leq_{\text{cover}} b$ , then  $b$  is above  $a$ .

For infinite orders, sometimes it is not possible to draw a Hasse diagram:  
in  $\mathbb{Q}$  with the usual ordering, does 1 cover 0?



We say  $b$  **covers**  $a$  if there is nothing between  $a$  and  $b$

$$\forall c \in A (a \leq c \leq b \Rightarrow c \in \{a, b\})$$

**Definition.** If  $A$  is finite, then a **Hasse diagram** of  $\leq$  is a drawing of  $\leq_{\text{cover}} = \{(a, b) \in A^2 \mid b \text{ covers } a\}$ , where if  $a \leq_{\text{cover}} b$ , then  $b$  is above  $a$ .

For infinite orders, sometimes it is not possible to draw a Hasse diagram:

in  $\mathbb{Q}$  with the usual ordering, does 1 cover 0?

In this case,  $\leq_{\text{cover}}$  is the empty set!

We say  $b$  **covers**  $a$  if there is nothing between  $a$  and  $b$

$$\forall c \in A (a \leq c \leq b \Rightarrow c \in \{a, b\})$$

**Definition.** If  $A$  is finite, then a **Hasse diagram** of  $\leq$  is a drawing of  $\leq_{\text{cover}} = \{(a, b) \in A^2 \mid b \text{ covers } a\}$ , where if  $a \leq_{\text{cover}} b$ , then  $b$  is above  $a$ .


For infinite orders, sometimes it is not possible to draw a Hasse diagram:

in  $\mathbb{Q}$  with the usual ordering, does 1 cover 0?

In this case,  $\leq_{\text{cover}}$  is the empty set!

Let  $A = \{n \in \mathbb{N} \mid n \text{ divides } 12\}$ , and let  $R = \{(n, m) \in A^2 \mid n \text{ divides } m\}$ .  
Draw the Hasse diagram of  $R$ .

$$A = \{1, 2, 3, 4, 6, 12\}$$



**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.




**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Question.** Is  $<$  an order?

- Reflexive?
- Antisymmetric?
- Transitive?




**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Question.** Is  $<$  an order?

- Reflexive? 
- Antisymmetric? 
- Transitive? 

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

**Question.** Is  $<$  an order?

- Reflexive? 
- Antisymmetric? 
- Transitive? 

**Definition.** A relation is a **strict order** if it is **antireflexive**, antisymmetric, and transitive.

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Consider the relation  $R$  defined by  $\{(\varphi, \psi) \mid \text{if } \varphi \text{ is true, then } \psi \text{ is true}\}$

- $(p, p \vee q) \in R$
- $((p \Rightarrow q) \wedge p, q) \in R$
- $(p \wedge q, q \wedge p) \in R$
- $(p, p \wedge q) \notin R$




**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Consider the relation  $R$  defined by  $\{(\varphi, \psi) \mid \text{if } \varphi \text{ is true, then } \psi \text{ is true}\}$

- $(p, p \vee q) \in R$
- $((p \Rightarrow q) \wedge p, q) \in R$
- $(p \wedge q, q \wedge p) \in R$
- $(p, p \wedge q) \notin R$

**Question.** Is  $R$  an order?

- Reflexive?
- Antisymmetric?
- Transitive?






**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Consider the relation  $R$  defined by  $\{(\varphi, \psi) \mid \text{if } \varphi \text{ is true, then } \psi \text{ is true}\}$

- $(p, p \vee q) \in R$
- $((p \Rightarrow q) \wedge p, q) \in R$
- $(p \wedge q, q \wedge p) \in R$
- $(p, p \wedge q) \notin R$

**Definition.** A relation is a **quasiorder** if it is reflexive and transitive.

**Question.** Is  $R$  an order?

- Reflexive? 
- Antisymmetric? 
- Transitive? 

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.


Consider the relation  $R$  defined by  $\{(\varphi, \psi) \mid \text{if } \varphi \text{ is true, then } \psi \text{ is true}\}$

- $(p, p \vee q) \in R$
- $((p \Rightarrow q) \wedge p, q) \in R$
- $(p \wedge q, q \wedge p) \in R$
- $(p, p \wedge q) \notin R$

**Definition.** A relation is a **quasiorder** if it is reflexive and transitive.

**Question.** Is  $R$  an order?

- Reflexive? ✓
- Antisymmetric? ✗
- Transitive? ✓



(loops omitted for readability)




**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.


Consider the relation  $R$  defined by  $\{(\varphi, \psi) \mid \text{if } \varphi \text{ is true, then } \psi \text{ is true}\}$

- $(p, p \vee q) \in R$
- $((p \Rightarrow q) \wedge p, q) \in R$
- $(p \wedge q, q \wedge p) \in R$
- $(p, p \wedge q) \notin R$


**Definition.** A relation is a **quasiorder** if it is reflexive and transitive.

**Question.** Is  $R$  an order?

- Reflexive? 
- Antisymmetric? 
- Transitive? 



$\rightsquigarrow$



(Hasse diagram after contracting the dotted bubbles into single elements)

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Consider  $R$  defined by  $\{(x, y) \in \mathbb{N}^2 \mid x \leq y\}$ .

**Question.** Is  $R$  an order?

- Reflexive?
- Antisymmetric?
- Transitive?

**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Consider  $R$  defined by  $\{(x, y) \in \mathbb{N}^2 \mid x \leq y\}$ .

**Question.** Is  $R$  an order?

- Reflexive? ✓
- Antisymmetric? ✓
- Transitive? ✓

**Definition.** We say an order is **linear** if for every  $a, b \in A$ , we have  $a \leq b$  or  $b \leq a$ .

In other words: any two elements can be compared using the order.

The Hasse diagram (if it exists) looks like a line.



**Definition.** A relation  $R \subseteq A \times A$  is an **order** if it is reflexive, antisymmetric, and transitive.

Consider  $R$  defined by  $\{(x, y) \in \mathbb{N}^2 \mid x \leq y\}$ .

**Question.** Is  $R$  an order?


- Reflexive? ✓
- Antisymmetric? ✓
- Transitive? ✓

**Definition.** We say an order is **linear** if for every  $a, b \in A$ , we have  $a \leq b$  or  $b \leq a$ .

In other words: any two elements can be compared using the order.  
The Hasse diagram (if it exists) looks like a line.

Which of the following orders are linear?

- $\leq$  on  $\mathbb{Q}$
- $\preceq_{\text{prefix}}$  on binary strings
- $\leq$  on  $\mathbb{R}$
- $\subseteq$  on  $\mathcal{P}(A)$






**Definition.** Let  $\leq$  be an order on  $A$  and  $a \in A$ .

We say that  $a$  is **minimal** if there is no  $b \in A \setminus \{a\}$  such that  $b \leq a$ .






**Definition.** Let  $\leq$  be an order on  $A$  and  $a \in A$ .

We say that  $a$  is **minimal** if there is no  $b \in A \setminus \{a\}$  such that  $b \leq a$ .



**Definition.** Let  $\leq$  be an order on  $A$  and  $a \in A$ .




We say that  $a$  is **minimal** if there is no  $b \in A \setminus \{a\}$  such that  $b \leq a$ .



Many problems in IT/science can be rephrased as finding a minimal element in a given order.  
Being able to formalize the order you care about allows you to use known algorithms/methods.

**Definition.** Let  $\leq$  be an order on  $A$  and  $a \in A$ .

We say that  $a$  is **minimal** if there is no  $b \in A \setminus \{a\}$  such that  $b \leq a$ .






Many problems in IT/science can be rephrased as finding a minimal element in a given order. Being able to formalize the order you care about allows you to use known algorithms/methods.

**Theorem.** Every non-empty finite order has a minimal element.

**Definition.** Let  $\leq$  be an order on  $A$  and  $a \in A$ .

We say that  $a$  is **minimal** if there is no  $b \in A \setminus \{a\}$  such that  $b \leq a$ .



Many problems in IT/science can be rephrased as finding a minimal element in a given order. Being able to formalize the order you care about allows you to use known algorithms/methods.

**Theorem.** Every non-empty finite order has a minimal element.

Proof by contradiction!

**Definition.** Let  $\leq$  be an order on  $A$ ,  $S \subseteq A$ , and  $a \in A$ . We say:





- $a$  is a **lower bound** of  $S$  if for all  $s \in S$ , we have  $a \leq s$
- $a$  is a **greatest lower bound** of  $S$  if it is a lower bound and for every lower bound  $b$  of  $S$ , we have  $b \leq a$ .

We write  $a \wedge b$  for the greatest lower bound of  $\{a, b\}$ , if it exists.

**Definition.** Let  $\leq$  be an order on  $A$ ,  $S \subseteq A$ , and  $a \in A$ . We say:

- $a$  is a **lower bound** of  $S$  if for all  $s \in S$ , we have  $a \leq s$
- $a$  is a **greatest lower bound** of  $S$  if it is a lower bound and for every lower bound  $b$  of  $S$ , we have  $b \leq a$ .

We write  $a \wedge b$  for the greatest lower bound of  $\{a, b\}$ , if it exists.




**Definition.** Let  $\leq$  be an order on  $A$ ,  $S \subseteq A$ , and  $a \in A$ . We say:

- $a$  is a **lower bound** of  $S$  if for all  $s \in S$ , we have  $a \leq s$
- $a$  is a **greatest lower bound** of  $S$  if it is a lower bound and for every lower bound  $b$  of  $S$ , we have  $b \leq a$ .

We write  $a \wedge b$  for the greatest lower bound of  $\{a, b\}$ , if it exists.





Let  $A = \{n \in \mathbb{N} \mid n \text{ divides } 12\}$ , and let  $R = \{(n, m) \in A^2 \mid n \text{ divides } m\}$ .  
What is  $4 \wedge 3$ ?



**Definition.** Let  $\leq$  be an order on  $A$ ,  $S \subseteq A$ , and  $a \in A$ . We say:

- $a$  is an **upper bound** of  $S$  if for all  $s \in S$ , we have  $s \leq a$
- $a$  is a **lowest upper bound** of  $S$  if it is an upper bound and for every upper bound  $b$  of  $S$ , we have  $a \leq b$ .

We write  $a \vee b$  for the lowest upper bound of  $\{a, b\}$ , if it exists.






**Definition.** Let  $\leq$  be an order on  $A$ ,  $S \subseteq A$ , and  $a \in A$ . We say:

- $a$  is an **upper bound** of  $S$  if for all  $s \in S$ , we have  $s \leq a$
- $a$  is a **lowest upper bound** of  $S$  if it is an upper bound and for every upper bound  $b$  of  $S$ , we have  $a \leq b$ .

We write  $a \vee b$  for the lowest upper bound of  $\{a, b\}$ , if it exists.

Let  $A = \{n \in \mathbb{N} \mid n \text{ divides } 12\}$ , and let  $R = \{(n, m) \in A^2 \mid n \text{ divides } m\}$ .  
What is  $2 \vee 3$ ?



- How to recognize an equivalence relation, an order
- Compute equivalence classes
- Compute  $A/\sim$
- Draw a Hasse diagram
- Recognize minimal/maximal elements
- Recognize lower/upper bounds

	Reflexive	Antireflexive	Symmetric	Antisymmetric	Transitive
Equivalence relation	✓		✓		✓
Order	✓			✓	✓
Strict order		✓		✓	✓
Quasi-order	✓				✓