

Discrete Algebraic Structures

WiSe 2025/2026

Prof. Dr. Antoine Wiehe
Research Group for Theoretical Computer Science



- Registration for the exam is now open (until December 14)
Register on **TUNE**
- Also register for the **Studienleistung** if not done already (and if you can)
- Exam on **March 10** (whole day, in groups of ~ 100)

- Registration for the exam is now open (until December 14)
Register on **TUNE**
- Also register for the **Studienleistung** if not done already (and if you can)
- Exam on **March 10** (whole day, in groups of ~ 100)

Exam has 2 parts:

- **Part 1** (around 100 points): mostly multiple choice questions + numerical questions
- Multiple choice questions: **negative points** for wrong answers

DO NOT TRY TO GUESS

- Registration for the exam is now open (until December 14)
Register on **TUNE**
- Also register for the **Studienleistung** if not done already (and if you can)
- Exam on **March 10** (whole day, in groups of ~ 100)

Exam has 2 parts:

- **Part 1** (around 100 points): mostly multiple choice questions + numerical questions
- Multiple choice questions: **negative points** for wrong answers

DO NOT TRY TO GUESS

A ☐ ☒ ☐ ☐ $A \cap (B \cup C) = A \cup (B \cap C)$ -1.67 Punkte

B ☐ ☒ ☐ ☐ $A \setminus (A \setminus B) = A \cap B$ -1.67 Punkte

C ☐ ☒ ☐ ☐ $A \cap (B \cup C) = (A \cup B) \cap (A \cup C)$ -1.67 Punkte

D ☐ ☒ ☐ ☐ $\emptyset \in A$ 1.67 Punkte

E ☐ ☒ ☐ ☐ $A \setminus (B \setminus C) = C \setminus B$ 1.67 Punkte

F ☐ ☒ ☐ ☐ $\emptyset \subseteq A$ 0.00 Punkte

- Registration for the exam is now open (until December 14)
Register on **TUNE**
- Also register for the **Studienleistung** if not done already (and if you can)
- Exam on **March 10** (whole day, in groups of ~ 100)

Exam has 2 parts:

- **Part 1** (around 100 points): mostly multiple choice questions + numerical questions
- Multiple choice questions: **negative points** for wrong answers
DO NOT TRY TO GUESS
- Also some questions with computations (just enter a number)

- Registration for the exam is now open (until December 14)
Register on **TUNE**
- Also register for the **Studienleistung** if not done already (and if you can)
- Exam on **March 10** (whole day, in groups of ~ 100)

Exam has 2 parts:

- **Part 1** (around 100 points): mostly multiple choice questions + numerical questions
- Multiple choice questions: **negative points** for wrong answers
DO NOT TRY TO GUESS
- Also some questions with computations (just enter a number)
- **Part 2** (around 100 points): **proofs**

- Registration for the exam is now open (until December 14)
Register on **TUNE**
- Also register for the **Studienleistung** if not done already (and if you can)
- Exam on **March 10** (whole day, in groups of ~ 100)

Exam has 2 parts:

- **Part 1** (around 100 points): mostly multiple choice questions + numerical questions
- Multiple choice questions: **negative points** for wrong answers
DO NOT TRY TO GUESS
- Also some questions with computations (just enter a number)
- **Part 2** (around 100 points): **proofs**

Strategy to not fail:

- Focus on Part 1: enough points to get a 2.0-2.3 if done perfectly
- Part 2 only for those of you who hope to get 1.x
Do not attempt if you are not confident with writing proofs (points are not easy to get)

- Countable sets, uncountable sets
- Countable sets = what can be represented exactly on a computer
- Combinatorial proofs as a way to prove equalities/inequalities about numbers using functions

$$\text{Injection } A \rightarrow B \quad \leftrightarrow \quad |A| \leq |B|$$

$$\text{Surjection } A \rightarrow B \quad \leftrightarrow \quad |A| \geq |B|$$

$$\text{Bijection } A \rightarrow B \quad \leftrightarrow \quad |A| = |B|$$

- Drawing a tuple/(multi)set with/without replacement

n = size of the set we are drawing from

k = number of draws

	Order matters	Order does not matter
Replacement	n^k	$\binom{n+k-1}{n-1}$
No replacement	$n^{\underline{k}}$	$\frac{n!}{k!(n-k)!}$

3 “advanced” proof techniques for counting:

- Double counting
- The inclusion-exclusion principle
- Application: counting partitions and **surjective** functions
- The **pigeonhole principle** and **Ramsey's theorem**

3 “advanced” proof techniques for counting:

- Double counting
- The inclusion-exclusion principle
- Application: counting partitions and **surjective** functions
- The **pigeonhole principle** and **Ramsey's theorem**


Talk about **asymptotic growth** of counting functions

Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?

Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?




Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?

Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$




Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?

Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$




Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?

Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$




Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?

Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.




Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?

Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b
- Size of R_a ?



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b
- Size of R_a ? $\deg(a)$
- $R = \bigcup_{a \in A} R_a$



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b
- Size of R_a ? $\deg(a)$
- $R = \bigcup_{a \in A} R_a$
- $R_a \cap R_{a'} = \emptyset$ if $a \neq a'$



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b
- Size of R_a ? $\deg(a)$
- $R = \bigcup_{a \in A} R_a$
- $R_a \cap R_{a'} = \emptyset$ if $a \neq a'$
- By the union rule, $|R| = \sum_{a \in A} \deg(a)$. □



Idea:

- we know if $f: A \rightarrow B$ is bijective, then $|A| = |B|$
- what can we do with **relations** instead of functions?


Definition. Let $R \subseteq A \times B$, $a \in A$, $b \in B$.

- $\deg(a) = |\{b \in B \mid (a, b) \in R\}|$
- $\deg(b) = |\{a \in A \mid (a, b) \in R\}|$

Theorem. $\sum_{a \in A} \deg(a) = |R| = \sum_{b \in B} \deg(b)$.

Proof. Theorem has a $\Sigma \rightsquigarrow$ partition R into subsets

- R_a : all edges “touching” a
- R_b : all edges “touching” b
- Size of R_a ? $\deg(a)$
- $R = \bigcup_{a \in A} R_a$
- $R_a \cap R_{a'} = \emptyset$ if $a \neq a'$
- By the union rule, $|R| = \sum_{a \in A} \deg(a)$. □



When to use this?

If:

- The degrees are known
- $|A|$ is known


Then we can know $|B|$.

We have seen: $\binom{n}{k} = \frac{n^{\underline{k}}}{k!} = \frac{n!}{k!(n-k)!}$

We have seen: $\binom{n}{k} = \frac{n^{\underline{k}}}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$





We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$

$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$





We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$

$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$





(For the example: $k = 3$)

We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$
- Define $R \subseteq A \times B$:
 $R = \{((a_1, \dots, a_k), T) \in A \times B \mid \{a_1, \dots, a_k\} = T\}$

$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$





(For the example: $k = 3$)

We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$
- Define $R \subseteq A \times B$:
 $R = \{((a_1, \dots, a_k), T) \in A \times B \mid \{a_1, \dots, a_k\} = T\}$
- degree of elements of A ?

$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$



(For the example: $k = 3$)

We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:



- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$
- Define $R \subseteq A \times B$:
 $R = \{((a_1, \dots, a_k), T) \in A \times B \mid \{a_1, \dots, a_k\} = T\}$
- degree of elements of A ?

What is $\deg(T)$ for every $T \in B$?

- 1
- $n!$
- $2k$
- $k!$



$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$





(For the example: $k = 3$)

We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$

Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$
- Define $R \subseteq A \times B$:
 $R = \{((a_1, \dots, a_k), T) \in A \times B \mid \{a_1, \dots, a_k\} = T\}$
- degree of elements of A ?
- degree of elements of B is $k!$

$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$



(For the example: $k = 3$)

We have seen: $\binom{n}{k} = \frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$



Proof using double counting:

- Want to know: $|\mathcal{P}_k(\{1, \dots, n\})|$
- Define $A = \{(a_1, \dots, a_k) \in \{1, \dots, n\}^k \mid \forall i \neq j : a_i \neq a_j\}$
- Define $R \subseteq A \times B$:
 $R = \{((a_1, \dots, a_k), T) \in A \times B \mid \{a_1, \dots, a_k\} = T\}$
- degree of elements of A ?
- degree of elements of B is $k!$

By double-counting:

$$\sum_{a \in A} \deg(a) = \sum_{T \in B} \deg(T)$$

$$A = \{(a_1, \dots, a_k) \in \dots^k \mid \forall i \neq j : a_i \neq a_j\}$$



(For the example: $k = 3$)


Union rule: $|A \cup B| =$

Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?

Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$


How to compute $|A \cup B \cup C|$?



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?


$$|A| + |B| + |C|$$



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?


$$|A| + |B| + |C|$$



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?


$$|A| + |B| + |C|$$



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?


$$|A| + |B| + |C|$$



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?


$$|A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C|$$



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?


$$|A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C| = |A \cup B \cup C|$$



Union rule: $|A \cup B| = |A| + |B| - |A \cap B|$

How to compute $|A \cup B \cup C|$?

$$|A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C| = |A \cup B \cup C|$$



This is called the **inclusion-exclusion principle**.

Ideally, one would like to be able to count more complicated expressions like $|A \cup B \cup C \cup D|$

Ideally, one would like to be able to count more complicated expressions like $|A \cup B \cup C \cup D|$

Theorem. If A_1, \dots, A_n are finite sets, then

$$|A_1 \cup \dots \cup A_n| = \sum_{r=1}^n (-1)^{r+1} \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} |A_{i_1} \cap \dots \cap A_{i_r}|$$

Ideally, one would like to be able to count more complicated expressions like $|A \cup B \cup C \cup D|$

Theorem. If A_1, \dots, A_n are finite sets, then

$$|A_1 \cup \dots \cup A_n| = \sum_{r=1}^n (-1)^{r+1} \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} |A_{i_1} \cap \dots \cap A_{i_r}|$$

sign alternates between $+$ and $-$

“Loop” over all subsets $\{i_1, \dots, i_r\}$
of $\{1, \dots, n\}$ with k elements

Ideally, one would like to be able to count more complicated expressions like $|A \cup B \cup C \cup D|$

Theorem. If A_1, \dots, A_n are finite sets, then

$$|A_1 \cup \dots \cup A_n| = \sum_{r=1}^n (-1)^{r+1} \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} |A_{i_1} \cap \dots \cap A_{i_r}|$$

sign alternates between $+$ and $-$

Proof: by induction on $n \geq 2$, base case is the union rule

“Loop” over all subsets $\{i_1, \dots, i_r\}$
of $\{1, \dots, n\}$ with k elements

Ideally, one would like to be able to count more complicated expressions like $|A \cup B \cup C \cup D|$

Theorem. If A_1, \dots, A_n are finite sets, then

$$|A_1 \cup \dots \cup A_n| = \sum_{r=1}^n (-1)^{r+1} \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} |A_{i_1} \cap \dots \cap A_{i_r}|$$

sign alternates between $+$ and $-$

Proof: by induction on $n \geq 2$, base case is the union rule

“Loop” over all subsets $\{i_1, \dots, i_r\}$
of $\{1, \dots, n\}$ with k elements

In some special cases, all the intersections $|A_{i_1} \cap \dots \cap A_{i_r}|$ have the same size N_r , we then get:

Theorem. Let A_1, \dots, A_n be finite sets and $N_1, \dots, N_n \in \mathbb{N}$ be such that for all $r \in \{1, \dots, n\}$ and all distinct $i_1, \dots, i_r \in \{1, \dots, n\}$, the intersection $A_{i_1} \cap \dots \cap A_{i_r}$ has size N_r . Then

$$|A_1 \cup \dots \cup A_n| = \sum_{r=1}^n (-1)^{r+1} \binom{n}{r} N_r$$

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$

Naive attempt:

```
from itertools import combinations

smallPrimes = primes(100) # compute all primes up to 100
finalSet = set()
for (p1,p2,p3,p4) in combinations(smallPrimes, 4):
    :

print(len(finalSet))
```

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$

Naive attempt:

```
from itertools import combinations

smallPrimes = primes(100) # compute all primes up to 100
finalSet = set()
for (p1,p2,p3,p4) in combinations(smallPrimes,4):

    print(len(finalSet))
```

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$

Naive attempt:

```
from itertools import combinations

smallPrimes = primes(100) # compute all primes up to 100
finalSet = set()
for (p1,p2,p3,p4) in combinations(smallPrimes,4):
    # take all multiples of p1*p2*p3*p4 that are smaller than 10^16
    for k in range(10**16 / (p1*p2*p3*p4)):
        finalSet = finalSet | {p1*p2*p3*p4 * k}
print(len(finalSet)) # outputs the correct number
```

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| \stackrel{?}{=} \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}|$$

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| \stackrel{?}{=} \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}|$$

How many times did we count $2 \times 3 \times 5 \times 7 \times 11$?

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| \stackrel{?}{=} \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}| \quad \text{---} \quad \sum_{p_1, p_2, p_3, p_4, p_5 \text{ distinct}} |D_{p_1 p_2 p_3 p_4 p_5}|$$

(A blue circle with -4 is around the first sum, and a blue arrow points from a blue circle with 5 in the text below to it.)

How many times did we count $2 \times 3 \times 5 \times 7 \times 11$? $\binom{5}{4} = 5$ times

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| \stackrel{?}{=} \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}| \quad \text{---} \quad \sum_{p_1, p_2, p_3, p_4, p_5 \text{ distinct}} |D_{p_1 p_2 p_3 p_4 p_5}|$$

(A blue circle highlights the -4 in the first sum, and a blue arrow points from a circled 5 in the text below to it.)

How many times did we count $2 \times 3 \times 5 \times 7 \times 11$? $\binom{5}{4} = 5$ times

How many times did we count $2 \times 3 \times 5 \times 7 \times 11 \times 13$?

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| \stackrel{?}{=} \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}| \quad \text{---} \quad \sum_{p_1, p_2, p_3, p_4, p_5 \text{ distinct}} |D_{p_1 p_2 p_3 p_4 p_5}|$$

(Note: A blue circle highlights the -4 in the first sum, and a blue arrow points from a blue circle around the 5 in the second sum to the -4.)

How many times did we count $2 \times 3 \times 5 \times 7 \times 11$? $\binom{5}{4} = 5$ times

How many times did we count $2 \times 3 \times 5 \times 7 \times 11 \times 13$? $\binom{6}{4} - 4\binom{6}{5} = -9$ times

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| = \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}| \quad \text{--4--} \quad \sum_{p_1, p_2, p_3, p_4, p_5 \text{ distinct}} |D_{p_1 p_2 p_3 p_4 p_5}| \quad +10 \quad \sum_{p_1, \dots, p_6 \text{ distinct}} |D_{p_1 \dots p_6}| - \dots$$

How many times did we count $2 \times 3 \times 5 \times 7 \times 11$? $\binom{5}{4} = 5$ times

How many times did we count $2 \times 3 \times 5 \times 7 \times 11 \times 13$? $\binom{6}{4} - 4\binom{6}{5} = -9$ times

(From projecteuler.net)

At Least Four Distinct Prime Factors Less Than 100

Problem 268



It can be verified that there are 23 positive integers less than 1000 that are divisible by at least four distinct primes less than 100.

Find how many positive integers less than 10^{16} are divisible by at least four distinct primes less than 100.

Setup: $S = \{n \in \{1, \dots, 10^{16}\} \mid n \text{ is divisible by four distinct primes less than } 100\}$
 $S = \bigcup_{p_1, p_2, p_3, p_4 \text{ distinct primes}} D_{p_1 p_2 p_3 p_4}$ where $D_k = \text{set of multiples of } k \text{ smaller than } 10^{16}$
 $|D_k| = 10^{16}/k$

$$|S| = \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}| \quad \text{--4--} \quad \sum_{p_1, p_2, p_3, p_4, p_5 \text{ distinct}} |D_{p_1 p_2 p_3 p_4 p_5}| \quad +10 \quad \sum_{p_1, \dots, p_6 \text{ distinct}} |D_{p_1 \dots p_6}| - \dots$$

How many times did we count $2 \times 3 \times 5 \times 7 \times 11$? $\binom{5}{4} = 5$ times

How many times did we count $2 \times 3 \times 5 \times 7 \times 11 \times 13$? $\binom{6}{4} - 4\binom{6}{5} = -9$ times

How many times did we count $p_1 \times p_2 \times \dots \times p_r$? $1 - (-1)^r \binom{r-1}{3}$ times

$$|S| = \sum_{p_1, p_2, p_3, p_4 \text{ distinct}} |D_{p_1 p_2 p_3 p_4}| \quad \textcircled{-4} \sum_{p_1, \dots, p_5 \text{ distinct}} |D_{p_1 p_2 p_3 p_4 p_5}| + \dots + \boxed{(-1)^r \binom{r-1}{3}} \sum_{p_1, \dots, p_r \text{ distinct}} |D_{p_1 \dots p_r}|$$

```

from itertools import combinations # iterator on subsets
from math import comb # gives the binomial coefficients

smallPrimes = primes(100)
result = 0
N = 10**16
for r in range(4, len(smallPrimes)+1):
    sum_for_fixed_r = 0
    for combination_of_primes in combinations(smallPrimes, r):
        product = 1
        for p in combination_of_primes:
            product *= p
        sum_for_fixed_r += N//product
    result += (-1)**r * comb(r-1, 3) * sum_for_fixed_r
print(result)

```


What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{\underline{|A|}}$ **injective** functions $f: A \rightarrow B$

What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$

Ideas:

- (1) If we know the number of functions that are **not** surjective, we know the number of surjective functions
- (2) The set of **not surjective functions** is the union of simple sets

What we have seen so far: if A, B are finite, then there are:


- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$

Ideas:

(1) If we know the number of functions that are **not** surjective, we know the number of surjective functions

(2) The set of **not surjective functions** is the union of simple sets

$$\text{Let } F_i = \{f: A \rightarrow B \mid b_i \notin f[A]\}$$



A function in F_1

What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$

Ideas:


(1) If we know the number of functions that are **not** surjective, we know the number of surjective functions

(2) The set of **not surjective functions** is the union of simple sets

$$\text{Let } F_i = \{f: A \rightarrow B \mid b_i \notin f[A]\}$$

Suppose $|A| = 8$ and $|B| = 4$.
What is $|F_i|$?

- 4
- 4^7
- 3^8
- 3^7



A function in F_1

What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$


Ideas:

(1) If we know the number of functions that are **not** surjective, we know the number of surjective functions

(2) The set of **not surjective functions** is the union of simple sets

$$\text{Let } F_i = \{f: A \rightarrow B \mid b_i \notin f[A]\}$$

$$|F_i \cap F_j| = \quad \text{if } i < j$$



A function in $F_1 \cap F_2$?

What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$

Ideas:


(1) If we know the number of functions that are **not** surjective, we know the number of surjective functions

(2) The set of **not surjective functions** is the union of simple sets

$$\text{Let } F_i = \{f: A \rightarrow B \mid b_i \notin f[A]\}$$

$$|F_i \cap F_j| = (|B| - 2)^{|A|} \text{ if } i < j$$

$$|F_i \cap F_j \cap F_k| = \quad \text{if } i < j < k$$



A function in $F_1 \cap F_2 \cap F_3$?

What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$

Ideas:

(1) If we know the number of functions that are **not** surjective, we know the number of surjective functions


(2) The set of **not surjective functions** is the union of simple sets

$$\text{Let } F_i = \{f: A \rightarrow B \mid b_i \notin f[A]\}$$

$$|F_i \cap F_j| = (|B| - 2)^{|A|} \text{ if } i < j$$

$$|F_i \cap F_j \cap F_k| = (|B| - 3)^{|A|} \text{ if } i < j < k$$

$$|F_{i_1} \cap \dots \cap F_{i_r}| = \quad \text{if } i_1 < \dots < i_r$$



What we have seen so far: if A, B are finite, then there are:

- $|B|^{|A|}$ functions $f: A \rightarrow B$
- $|B|^{|A|}$ **injective** functions $f: A \rightarrow B$

Ideas:

(1) If we know the number of functions that are **not** surjective, we know the number of surjective functions

(2) The set of **not surjective functions** is the union of simple sets

$$\text{Let } F_i = \{f: A \rightarrow B \mid b_i \notin f[A]\}$$


$$|F_i \cap F_j| = (|B| - 2)^{|A|} \text{ if } i < j$$

$$|F_i \cap F_j \cap F_k| = (|B| - 3)^{|A|} \text{ if } i < j < k$$

$$|F_{i_1} \cap \dots \cap F_{i_r}| = (|B| - r)^{|A|} \text{ if } i_1 < \dots < i_r$$


Theorem. The number of surjective functions $A \rightarrow B$ is given by

$$\sum_{r=0}^{|B|} (-1)^r \binom{|B|}{r} (|B| - r)^{|A|}$$




Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function




Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function



Clustering is an important problem in data science/machine learning


Goal: find a clustering of the points that minimizes an objective function



Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function

```
S = set_of_clusterings(nb_clusters=5)
for clustering in S:
    # check if this clustering
    # is good enough
```




Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function

```
S = set_of_clusterings(nb_clusters=5)
for clustering in S:
    # check if this clustering
    # is good enough
```

Clustering \leftrightarrow partition of the set
cluster \leftrightarrow part




How many partitions with k parts of a set with n elements are there?

Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function

```
S = set_of_clusterings(nb_clusters=5)
for clustering in S:
    # check if this clustering
    # is good enough
```

Clustering \leftrightarrow partition of the set \leftrightarrow eq. relation
cluster \leftrightarrow part \leftrightarrow eq. class



How many partitions with k parts of a set with n elements are there?


Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function

```
S = set_of_clusterings(nb_clusters=5)
for clustering in S:
    # check if this clustering
    # is good enough
```

Clustering \leftrightarrow partition of the set \leftrightarrow eq. relation
 cluster \leftrightarrow part \leftrightarrow eq. class

How many partitions with k parts of a set with n elements are there?




Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function


```
S = set_of_clusterings(nb_clusters=5)
for clustering in S:
    # check if this clustering
    # is good enough
```

Clustering \leftrightarrow partition of the set \leftrightarrow eq. relation
 cluster \leftrightarrow part \leftrightarrow eq. class




How many partitions with k parts of a set with n elements are there?

partitions of $\{1, \dots, n\}$
 in k parts



surjective
 functions




Clustering is an important problem in data science/machine learning

Goal: find a clustering of the points that minimizes an objective function

```
S = set_of_clusterings(nb_clusters=5)
for clustering in S:
    # check if this clustering
    # is good enough
```

Clustering \leftrightarrow partition of the set \leftrightarrow eq. relation
 cluster \leftrightarrow part \leftrightarrow eq. class




How many partitions with k parts of a set with n elements are there?

Theorem. The number of **partitions** of $\{1, \dots, n\}$ into **k parts** is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

$n = 39, k = 5 \rightsquigarrow 1817478286796884221993319200$ partitions into k parts



Usually, to prove $\exists x \varphi$ one must give a concrete x that works

Usually, to prove $\exists x \varphi$ one must give a concrete x that works

Sometimes, **counting** is enough:

- Pigeonhole principle
- Ramsey theory
- Probabilistic method

Usually, to prove $\exists x \varphi$ one must give a concrete x that works

Sometimes, **counting** is enough:

- Pigeonhole principle
- Ramsey theory
- Probabilistic method

Theorem. Suppose $f: A \rightarrow B$ is a function and $|A| > |B|$.
Then **there exist** distinct $x, y \in A$ such that $f(x) = f(y)$.

Usually, to prove $\exists x \varphi$ one must give a concrete x that works

Sometimes, **counting** is enough:

- Pigeonhole principle
- Ramsey theory
- Probabilistic method

Theorem. Suppose $f: A \rightarrow B$ is a function and $|A| > |B|$.
Then **there exist** distinct $x, y \in A$ such that $f(x) = f(y)$.

In fact if A is much bigger than B , one can obtain more elements like this:

Theorem (Pigeonhole principle). Suppose $f: A \rightarrow B$ is a function and $|A| > k|B|$ for some $k \in \mathbb{N}$.
Then there exists distinct $x_1, \dots, x_{k+1} \in A$ such that $f(x_i) = f(x_j)$ for all $i, j \in \{1, \dots, k+1\}$.

Usually, to prove $\exists x \varphi$ one must give a concrete x that works

Sometimes, **counting** is enough:


- Pigeonhole principle
- Ramsey theory
- Probabilistic method

Theorem. Suppose $f: A \rightarrow B$ is a function and $|A| > |B|$. Then **there exist** distinct $x, y \in A$ such that $f(x) = f(y)$.

In fact if A is much bigger than B , one can obtain more elements like this:

Theorem (Pigeonhole principle). Suppose $f: A \rightarrow B$ is a function and $|A| > k|B|$ for some $k \in \mathbb{N}$. Then there exists distinct $x_1, \dots, x_{k+1} \in A$ such that $f(x_i) = f(x_j)$ for all $i, j \in \{1, \dots, k+1\}$.

Example. For any 9 numbers a_1, \dots, a_9 from $\{1, \dots, 16\}$, two numbers are at distance exactly 2.



Theorem. Let $r \in \mathbb{N}$. There exists $n \in \mathbb{N}$ such that for every symmetric relation $R \subseteq A^2$, if $|A| \geq n$, there exist $a_1, \dots, a_r \in A$ distinct such that either:

- $(a_i, a_j) \in R$ for all $i \neq j$.
- or $(a_i, a_j) \notin R$ for all $i \neq j$.

Theorem. Let $r \in \mathbb{N}$. There exists $n \in \mathbb{N}$ such that for every symmetric relation $R \subseteq A^2$, if $|A| \geq n$, there exist $a_1, \dots, a_r \in A$ distinct such that either:




- $(a_i, a_j) \in R$ for all $i \neq j$.
- or $(a_i, a_j) \notin R$ for all $i \neq j$.

Example. We choose $r = 3$. Ramsey's Theorem gives us $n = 6$

Theorem. Let $r \in \mathbb{N}$. There exists $n \in \mathbb{N}$ such that for every symmetric relation $R \subseteq A^2$, if $|A| \geq n$, there exist $a_1, \dots, a_r \in A$ distinct such that either:

- $(a_i, a_j) \in R$ for all $i \neq j$.
- or $(a_i, a_j) \notin R$ for all $i \neq j$.




Example. We choose $r = 3$. Ramsey's Theorem gives us $n = 6$



Theorem. Let $r \in \mathbb{N}$. There exists $n \in \mathbb{N}$ such that for every symmetric relation $R \subseteq A^2$, if $|A| \geq n$, there exist $a_1, \dots, a_r \in A$ distinct such that either:

- $(a_i, a_j) \in R$ for all $i \neq j$.
- or $(a_i, a_j) \notin R$ for all $i \neq j$.

Example. We choose $r = 3$. Ramsey's Theorem gives us $n = 6$



For $r = 4$, $n = 18$ is enough

For $r = 5$, nobody knows what n is!

Theorem. The number of **partitions** of $\{1, \dots, n\}$ into **k parts** is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?

Theorem. The number of **partitions** of $\{1, \dots, n\}$ into **k parts** is


$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?

Theorem. The number of partitions of $\{1, \dots, n\}$ into k parts is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?



Theorem. The number of **partitions** of $\{1, \dots, n\}$ into **k parts** is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?

Definition. Let $f, g: \mathbb{N} \rightarrow \mathbb{N}$. We say **$f \in O(g)$** if there are $C, n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $f(n) \leq C \cdot g(n)$.

Theorem. The number of **partitions** of $\{1, \dots, n\}$ into **k parts** is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?

Definition. Let $f, g: \mathbb{N} \rightarrow \mathbb{N}$. We say **$f \in O(g)$** if there are $C, n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $f(n) \leq C \cdot g(n)$.

Exercise. Prove that the relation $\{(f, g) \mid f \in O(g)\}$ is a **quasiorder**.

Theorem. The number of partitions of $\{1, \dots, n\}$ into k parts is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?

Definition. Let $f, g: \mathbb{N} \rightarrow \mathbb{N}$. We say $f \in O(g)$ if there are $C, n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $f(n) \leq C \cdot g(n)$.

Exercise. Prove that the relation $\{(f, g) \mid f \in O(g)\}$ is a **quasiorder**.

	Order matters	Order does not matter
Replacement	n^k	$\binom{n+k-1}{n-1}$
No replacement	$n^{\underline{k}}$	$\frac{n!}{k!(n-k)!}$

Number of surjective functions:
(from a set of size n to a set of size k)

$$\sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Number of partitions:
(of a set of size n into k parts)

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Theorem. The number of **partitions** of $\{1, \dots, n\}$ into **k parts** is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?

Definition. Let $f, g: \mathbb{N} \rightarrow \mathbb{N}$. We say **$f \in O(g)$** if there are $C, n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $f(n) \leq C \cdot g(n)$.

Exercise. Prove that the relation $\{(f, g) \mid f \in O(g)\}$ is a **quasiorder**.

	Order matters	Order does not matter
Replacement	n^k	$\binom{n+k-1}{n-1} \in O(n^k)$
No replacement	$n^{\underline{k}} \in O(n^k)$	$\frac{n!}{k!(n-k)!} \in O(n^k)$

Number of surjective functions:
(from a set of size n to a set of size k)

$$O\left(\sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n\right) \ni k^n$$

Number of partitions:
(of a set of size n into k parts)

$$O\left(\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n\right) \ni k^n$$

Theorem. The number of partitions of $\{1, \dots, n\}$ into k parts is

$$\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n$$

Any idea how big this is?
How about k^n ?

Definition. Let $f, g: \mathbb{N} \rightarrow \mathbb{N}$. We say $f \in O(g)$ if there are $C, n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $f(n) \leq C \cdot g(n)$.

Exercise. Prove that the relation $\{(f, g) \mid f \in O(g)\}$ is a **quasiorder**.

	Order matters	Order does not matter
Replacement	n^k	$\binom{n+k-1}{n-1} \in O(n^k)$
No replacement	$n^{\underline{k}} \in O(n^k)$	$\frac{n!}{k!(n-k)!} \in O(n^k)$

Number of surjective functions:
(from a set of size n to a set of size k)

$$O\left(\sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n\right) \ni k^n$$

Number of partitions:
(of a set of size n into k parts)

$$O\left(\frac{1}{k!} \sum_{r=0}^k (-1)^r \binom{k}{r} (k-r)^n\right) \ni k^n$$

Take away message:
 $O(n^k)$ is “ok” for algorithms if k small
Don’t ever try $O(k^n)$, even if k is small

- How to use double counting
- The inclusion-exclusion principle for 2 and 3 sets
- How to apply the pigeonhole principle