

Prep Course Mathematics

Lecture 6 – Vectors and systems of linear equations

Christian Seifert (Deutsch), Jens-Peter M. Zemke (English)



Content

1. Vectors

- ▶ Vectors in geometry
- ▶ Basic arithmetic
- ▶ Linear combinations


2. Systems of linear equations

- ▶ Method of substitution
- ▶ Method of equalization
- ▶ Solving by graphing
- ▶ Method of elimination

Vectors

Vectors in two dimensions (2D)

Plane: two coordinate axes (e.g. x - and y -axis) intersecting in the zero-point (origin)



Example: the vector


$$\mathbf{v} := \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

starts at an arbitrary point and goes 2 steps in the direction of the x -axis and 3 steps in the direction of the y -axis.

A vector in this plane is an arrow of which only the elongation in the direction of the x - and y -axes is known.

Vectors in two dimensions (2D)

Plane: two coordinate axes (e.g. x - and y -axis) intersecting in the zero-point (origin)



Example: the vector


$$\mathbf{v} := \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

starts at an arbitrary point and goes 2 steps in the direction of the x -axis and 3 steps in the direction of the y -axis.

A vector in this plane is an arrow of which only the elongation in the direction of the x - and y -axes is known.

Vectors in three dimensions (3D)

Space: three coordinate axes (e.g., x_1 -, x_2 - and x_3 -axis) intersecting in the zero-point (origin)



Example: the vector

$$\mathbf{v} := \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$$

goes 1 step in the direction of the x_1 -axis,
3 steps in the direction of the x_2 -axis and
2 steps in the direction of the x_3 -axis.

A vector in this space is an arrow of which only the elongation in the direction of the x_1 -, x_2 - and x_3 -axes is known.

Vectors in n dimensions (nD)

Vectors in n dimensions

Let $n \in \mathbb{N}$. Every object \mathbf{v} of the form

$$\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \quad \text{with} \quad v_1, v_2, \dots, v_n \in \mathbb{R}$$

is called a real **vector**.

Vectors in n dimensions (nD)

Vectors in n dimensions

Let $n \in \mathbb{N}$. Every object \mathbf{v} of the form


$$\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \quad \text{with} \quad v_1, v_2, \dots, v_n \in \mathbb{R}$$

is called a real **vector**. The set of all such vectors is denoted by \mathbb{R}^n . The entries v_1, v_2, \dots, v_n are called **components**.

Notes:


- ▶ *Today we mainly consider the case $n \in \{2, 3\}$. However, in Linear Algebra we will often work with larger $n \in \mathbb{N}$.*
- ▶ *In different lectures you may see different notation for vectors: \mathbf{v} , \vec{v} , \underline{v} etc.*

Exercise



Exercise: Which of the given vectors are equal? Give their representations with coordinates.

Exercise



Exercise: Which of the given vectors are equal? Give their representations with coordinates.

Solution:


$$\mathbf{c} = \mathbf{k} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}, \quad \mathbf{b} = \mathbf{i} = \mathbf{l} = \begin{pmatrix} 1 \\ -1.5 \end{pmatrix}.$$

Points and position vectors

A point P is given as a *tuple* (v_1, v_2, \dots, v_n) , e.g., $P = (1, 3, 2)$.

Definition (position vector)

Let O be the origin (of the coordinate system). If P is a point, then the vector going from O to P is called the **position vector** of P . We often identify points with their position vectors.



Length of a vector

Length of a vector

Consider any vector

$$\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \in \mathbb{R}^n.$$

The **length** of this \mathbf{v} is given by

$$\|\mathbf{v}\| := \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}.$$

Length of a vector

Length of a vector

Consider any vector

$$\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \in \mathbb{R}^n.$$

The **length** of this \mathbf{v} is given by

$$\|\mathbf{v}\| := \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}.$$


Example: the length of the vector

$$\mathbf{v} := \begin{pmatrix} 4 \\ -3 \end{pmatrix}$$

is

$$\|\mathbf{v}\| = \sqrt{4^2 + (-3)^2} = \sqrt{25} = 5.$$

See the following picture:



Addition/subtraction of vectors

Addition/subtraction of vectors in \mathbb{R}^n

The addition/subtraction of two vectors in \mathbb{R}^n is done componentwise.

Given two vectors

$$\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \in \mathbb{R}^n, \quad \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{pmatrix} \in \mathbb{R}^n,$$

we define

$$\mathbf{v} + \mathbf{w} := \begin{pmatrix} v_1 + w_1 \\ v_2 + w_2 \\ \vdots \\ v_n + w_n \end{pmatrix}, \quad \mathbf{v} - \mathbf{w} := \begin{pmatrix} v_1 - w_1 \\ v_2 - w_2 \\ \vdots \\ v_n - w_n \end{pmatrix}.$$

Addition/subtraction of vectors

Examples:

$$(2) + (5) = (7),$$

$$\begin{pmatrix} 2 \\ 1 \\ 2 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix},$$

$$\begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} + \begin{pmatrix} -v_1 \\ \vdots \\ -v_n \end{pmatrix} =$$

$$\begin{pmatrix} 2 \\ -3 \end{pmatrix} + \begin{pmatrix} -5 \\ 6 \end{pmatrix} = \begin{pmatrix} -3 \\ 3 \end{pmatrix},$$

$$\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} =$$

$$\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} =$$

Addition/subtraction of vectors

Examples:

$$(2) + (5) = (7),$$

$$\begin{pmatrix} 2 \\ 1 \\ 2 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix},$$

$$\begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} + \begin{pmatrix} -v_1 \\ \vdots \\ -v_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix},$$

$$\begin{pmatrix} 2 \\ -3 \end{pmatrix} + \begin{pmatrix} -5 \\ 6 \end{pmatrix} = \begin{pmatrix} -3 \\ 3 \end{pmatrix},$$


$$\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix},$$

$$\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}.$$

Vectors between two points

Vectors between two points

Let A and B be two points. The vector \mathbf{v}_{AB} going from A to B can be obtained by subtracting the position vector of A from the position vector of B .


$$\mathbf{v}_{AB} = \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \\ \vdots \end{pmatrix}$$


$A = (a_1, a_2, \dots)$ $B = (b_1, b_2, \dots)$

Vectors between two points

Vectors between two points

Let A and B be two points. The vector \mathbf{v}_{AB} going from A to B can be obtained by subtracting the position vector of A from the position vector of B .


$$\mathbf{v}_{AB} = \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \\ \vdots \end{pmatrix}$$

$A = (a_1, a_2, \dots)$ $B = (b_1, b_2, \dots)$


Example: The vector going from $A := (2, 4, -6)$ to $B := (3, -1, 9)$ is

$$\mathbf{v}_{AB} = \begin{pmatrix} 3 \\ -1 \\ 9 \end{pmatrix} - \begin{pmatrix} 2 \\ 4 \\ -6 \end{pmatrix} = \begin{pmatrix} 3 - 2 \\ -1 - 4 \\ 9 + 6 \end{pmatrix} = \begin{pmatrix} 1 \\ -5 \\ 15 \end{pmatrix}.$$


Addition/subtraction of vectors

Addition of vectors:

The addition of vectors corresponds to a composition of movements that are described by the vectors.



Example:




Exercise

Of a parallelogram in space with corner points A, B, C, D three are given as follows:

$$A := (2, 3, 4),$$

$$B := (0, 7, 0),$$

$$D := (-2, 6, 4).$$



- (i) Determine the vectors \mathbf{v}_{AB} and \mathbf{v}_{AD} which lead from A to B and from A to D .
- (ii) Which coordinates does the point C have?
- (iii) Determine the circumference U of the parallelogram.
- (iv) Determine the length d_{AC} of the diagonal AC .


Solution

- (i) Determine the vectors \mathbf{v}_{AB} and \mathbf{v}_{AD} .

$$A := (2, 3, 4),$$

$$B := (0, 7, 0),$$

$$D := (-2, 6, 4).$$



$$\mathbf{v}_{AB} = \begin{pmatrix} 0 \\ 7 \\ 0 \end{pmatrix} - \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} -2 \\ 4 \\ -4 \end{pmatrix} = \mathbf{v}_{DC}, \quad \mathbf{v}_{AD} = \begin{pmatrix} -2 \\ 6 \\ 4 \end{pmatrix} - \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} -4 \\ 3 \\ 0 \end{pmatrix} = \mathbf{v}_{BC}.$$


Solution

(ii) Which coordinates does the point C have?

$$A := (2, 3, 4),$$

$$B := (0, 7, 0),$$

$$D := (-2, 6, 4).$$



$$\mathbf{v}_{OC} = \mathbf{v}_{OD} + \mathbf{v}_{DC} = \begin{pmatrix} -2 \\ 6 \\ 4 \end{pmatrix} + \begin{pmatrix} -2 \\ 4 \\ -4 \end{pmatrix} = \begin{pmatrix} -4 \\ 10 \\ 0 \end{pmatrix}, \quad C = (-4, 10, 0).$$


Solution

- (iii) Determine the circumference U of the parallelogram.

$$A := (2, 3, 4),$$

$$B := (0, 7, 0),$$

$$D := (-2, 6, 4).$$



$$\|\mathbf{v}_{AB}\| = \left\| \begin{pmatrix} -2 \\ 4 \\ -4 \end{pmatrix} \right\| = 6, \quad \|\mathbf{v}_{AD}\| = \left\| \begin{pmatrix} -4 \\ 3 \\ 0 \end{pmatrix} \right\| = 5, \quad U = 2(\|\mathbf{v}_{AB}\| + \|\mathbf{v}_{AD}\|) = 22.$$


Solution

- (iv) Determine the length d_{AC} of the diagonal AC .

$$A := (2, 3, 4),$$

$$B := (0, 7, 0),$$

$$D := (-2, 6, 4).$$



$$\mathbf{v}_{AC} = \mathbf{v}_{AB} + \mathbf{v}_{BC} = \begin{pmatrix} -2 \\ 4 \\ -4 \end{pmatrix} + \begin{pmatrix} -4 \\ 3 \\ 0 \end{pmatrix} = \begin{pmatrix} -6 \\ 7 \\ -4 \end{pmatrix}, \quad d_{AC} = \|\mathbf{v}_{AC}\| = \sqrt{36 + 49 + 16} = \sqrt{101}.$$

Scalar multiplication of vectors

Scalar multiplication of vectors in \mathbb{R}^n

The scalar multiplication of vectors in \mathbb{R}^n is done componentwise.

Given a real number (scalar) $\lambda \in \mathbb{R}$ and a vector

$$\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \in \mathbb{R}^n,$$


we define

$$\lambda \cdot \mathbf{v} := \begin{pmatrix} \lambda \cdot v_1 \\ \lambda \cdot v_2 \\ \vdots \\ \lambda \cdot v_n \end{pmatrix} \in \mathbb{R}^n.$$


Scalar multiplication of vectors

Examples: Consider the vector $\mathbf{v} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$. We determine and draw the following three vectors:


(i) $-1 \cdot \mathbf{v}$



(ii) $2 \cdot \mathbf{v}$




(iii) $\frac{1}{2} \cdot \mathbf{v}$



Scalar multiplication of vectors


Examples: Consider the vector $\mathbf{v} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$. We determine and draw the following three vectors:

(i) $-1 \cdot \mathbf{v} = \begin{pmatrix} -1 \\ -2 \end{pmatrix}$




change of direction

(ii) $2 \cdot \mathbf{v} = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$



elongation


(iii) $\frac{1}{2} \cdot \mathbf{v} = \begin{pmatrix} 0.5 \\ 1 \end{pmatrix}$



shortening

Scalar multiplication of vectors

scalar multiplication of vectors: the scalar multiplication of a vector with a constant α corresponds to a stretching by the factor $|\alpha|$, while the direction is switched if α is negative.



Examples

$$3 \cdot \begin{pmatrix} 13 \\ -12 \end{pmatrix} - 2 \cdot \begin{pmatrix} 5 \\ -4 \end{pmatrix} = \begin{pmatrix} 39 \\ -36 \end{pmatrix} - \begin{pmatrix} 10 \\ -8 \end{pmatrix} = \begin{pmatrix} 29 \\ -28 \end{pmatrix},$$

$$2 \cdot \left[\begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} + \begin{pmatrix} 0 \\ 5 \\ -5 \end{pmatrix} \right] - \begin{pmatrix} 6 \\ -2 \\ 2 \end{pmatrix} = 2 \cdot \begin{pmatrix} 3 \\ 7 \\ -6 \end{pmatrix} - \begin{pmatrix} 6 \\ -2 \\ 2 \end{pmatrix} = \begin{pmatrix} 6 \\ 14 \\ -12 \end{pmatrix} - \begin{pmatrix} 6 \\ -2 \\ 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 16 \\ -14 \end{pmatrix},$$

$$-1 \cdot \begin{pmatrix} 1 \\ 4 \\ 7 \end{pmatrix} + 2 \cdot \begin{pmatrix} 2 \\ 5 \\ 8 \end{pmatrix} - 1 \cdot \begin{pmatrix} 3 \\ 6 \\ 9 \end{pmatrix} = \begin{pmatrix} -1 \\ -4 \\ -7 \end{pmatrix} + \begin{pmatrix} 4 \\ 10 \\ 16 \end{pmatrix} + \begin{pmatrix} -3 \\ -6 \\ -9 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Rules of calculation of vector addition and scalar multiplication

Rules of calculation in \mathbb{R}^n

Let $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ be vectors and $\alpha, \beta \in \mathbb{R}$ be scalars. Then the following hold:

- (i) Commutativity: $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$
- (ii) Associativity: $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
- (iii) Distributivity I: $\alpha \cdot (\mathbf{u} + \mathbf{v}) = \alpha \cdot \mathbf{u} + \alpha \cdot \mathbf{v}$
- (iv) Distributivity II: $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v}$

Exercises

(a) Calculate one solution of

$$(i) \quad 4 \cdot \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix} - 5 \cdot \begin{pmatrix} -3 \\ 8 \\ -2 \end{pmatrix}, \quad (ii) \quad \begin{pmatrix} 0 \\ 3 \\ -5 \end{pmatrix} - \left[\begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix} - \begin{pmatrix} -1 \\ 1 \\ -2 \end{pmatrix} \right].$$

(b) Find a vector \mathbf{x} which satisfies one of the given equations:

$$(i) \quad 2\mathbf{x} - 3 \cdot \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} = 4\mathbf{x} + \begin{pmatrix} 1 \\ 7 \\ 2 \end{pmatrix}, \quad (ii) \quad 2 \cdot \left[\begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} - \mathbf{x} \right] = \mathbf{x} + \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}.$$

(c) Find scalars $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ such that the following equation is true:

$$\alpha_1 \cdot \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} + \alpha_2 \cdot \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix} + \alpha_3 \cdot \begin{pmatrix} 4 \\ -2 \\ -2 \end{pmatrix} = \begin{pmatrix} 3 \\ -7 \\ -1 \end{pmatrix}.$$

Solutions

(a) Calculate

$$(i) \quad 4 \cdot \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix} - 5 \cdot \begin{pmatrix} -3 \\ 8 \\ -2 \end{pmatrix}, \quad (ii) \quad \begin{pmatrix} 0 \\ 3 \\ -5 \end{pmatrix} - \left[\begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix} - \begin{pmatrix} -1 \\ 1 \\ -2 \end{pmatrix} \right].$$

Solution:

$$\begin{aligned} & 4 \cdot \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix} - 5 \cdot \begin{pmatrix} -3 \\ 8 \\ -2 \end{pmatrix} \\ &= \begin{pmatrix} 12 \\ 0 \\ -8 \end{pmatrix} + \begin{pmatrix} 15 \\ -40 \\ 10 \end{pmatrix} = \begin{pmatrix} 27 \\ -40 \\ 2 \end{pmatrix}. \end{aligned}$$

$$\begin{aligned} & \begin{pmatrix} 0 \\ 3 \\ -5 \end{pmatrix} - \left[\begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix} - \begin{pmatrix} -1 \\ 1 \\ -2 \end{pmatrix} \right] \\ &= \begin{pmatrix} 0 \\ 3 \\ -5 \end{pmatrix} - \begin{pmatrix} 3 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} -3 \\ 3 \\ -4 \end{pmatrix}. \end{aligned}$$

Solutions

(b) Find a vector \mathbf{x} which satisfies the given equation:

$$(i) \quad 2\mathbf{x} - 3 \cdot \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} = 4\mathbf{x} + \begin{pmatrix} 1 \\ 7 \\ 2 \end{pmatrix}, \quad (ii) \quad 2 \cdot \left[\begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} - \mathbf{x} \right] = \mathbf{x} + \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}.$$

Solution:

$$\begin{aligned} 2\mathbf{x} + \begin{pmatrix} -3 \\ 3 \\ 0 \end{pmatrix} &= 4\mathbf{x} + \begin{pmatrix} 1 \\ 7 \\ 2 \end{pmatrix} \\ \begin{pmatrix} -3 \\ 3 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ 7 \\ 2 \end{pmatrix} &= \begin{pmatrix} -4 \\ -4 \\ -2 \end{pmatrix} = 2\mathbf{x} \\ \begin{pmatrix} -2 \\ -2 \\ -1 \end{pmatrix} &= \mathbf{x}. \end{aligned}$$

$$\begin{aligned} \begin{pmatrix} 6 \\ 2 \\ 2 \end{pmatrix} - 2\mathbf{x} &= \mathbf{x} + \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} \\ \begin{pmatrix} 6 \\ 2 \\ 2 \end{pmatrix} - \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} &= \begin{pmatrix} 6 \\ -1 \\ 1 \end{pmatrix} = 3\mathbf{x} \\ \begin{pmatrix} 2 \\ -1/3 \\ 1/3 \end{pmatrix} &= \mathbf{x}. \end{aligned}$$

Solutions

(c) Find scalars $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ such that the following equation is true:

$$\alpha_1 \cdot \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} + \alpha_2 \cdot \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix} + \alpha_3 \cdot \begin{pmatrix} 4 \\ -2 \\ -2 \end{pmatrix} = \begin{pmatrix} 3 \\ -7 \\ -1 \end{pmatrix}.$$

Solution:

$$\begin{array}{lcl} \text{I:} & 2\alpha_1 + 1\alpha_2 + 4\alpha_3 & = 3 \\ \text{II:} & 3\alpha_2 - 2\alpha_3 & = -7 \\ \text{III:} & -2\alpha_3 & = -1 \end{array}$$

$$\text{III: } -2\alpha_3 = -1 \rightsquigarrow \boxed{\alpha_3 = 1/2}$$

$$\text{II: } 3\alpha_2 = -7 + 2\alpha_3 = -7 + 1 = -6 \rightsquigarrow \boxed{\alpha_2 = -2}$$

$$\text{I: } 2\alpha_1 = 3 - 4\alpha_3 - 1\alpha_2 = 3 - 2 + 2 = 3 \rightsquigarrow \boxed{\alpha_1 = 3/2}$$

Linear combination

Linear combination

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ and $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ be given. Then we call the vector

$$\mathbf{v} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_k \mathbf{v}_k$$

a **linear combination** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. We also say that \mathbf{v} can be generated by $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$.

Linear combination

Linear combination

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ and $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ be given. Then we call the vector

$$\mathbf{v} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_k \mathbf{v}_k$$

a **linear combination** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. We also say that \mathbf{v} can be generated by $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$.

Examples:


- ▶ $\mathbf{a} + \mathbf{b} = 1 \cdot \mathbf{a} + 1 \cdot \mathbf{b}$ is a linear combination of \mathbf{a} and \mathbf{b} .
- ▶ $\mathbf{a} = 1 \cdot \mathbf{a} + 0 \cdot \mathbf{b}$ is a linear combination of \mathbf{a} and \mathbf{b} .
- ▶ $\mathbf{b} = 0 \cdot \mathbf{a} + 1 \cdot \mathbf{b}$ is a linear combination of \mathbf{a} and \mathbf{b} .
- ▶ $\begin{pmatrix} 16 \\ 2 \\ 3 \end{pmatrix} = 2 \cdot \begin{pmatrix} 3 \\ 1 \\ -1 \end{pmatrix} + 5 \cdot \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$ is a linear combination of $\begin{pmatrix} 3 \\ 1 \\ -1 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$.
- ▶ The zero vector in \mathbb{R}^n is a linear combination of every tuple of vectors in \mathbb{R}^n .

Linear combination

Example:

Every point (position vector) $\mathbf{x} \in \mathbb{R}^3$ lying in the (x_1, x_2) -plane can be generated by the vectors


$$\mathbf{e}_1 := \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{e}_2 := \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}.$$



Example

Every point (position vector) $\mathbf{x} \in \mathbb{R}^2$ can be generated by the vectors

$$\mathbf{v}_1 := \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \quad \mathbf{v}_2 := \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$



Span

Span

Consider vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$. The set of all linear combinations of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ is called **span** (or linear hull) of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. Shortly:



$$\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k) := \{ \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_k \mathbf{v}_k : \lambda_1, \dots, \lambda_k \in \mathbb{R} \}.$$

Span

Span

Consider vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$. The set of all linear combinations of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ is called **span** (or linear hull) of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. Shortly:

$$\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k) := \{ \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_k \mathbf{v}_k : \lambda_1, \dots, \lambda_k \in \mathbb{R} \}.$$





Span

Span



Consider vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$. The set of all linear combinations of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ is called **span** (or linear hull) of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. Shortly:

$$\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k) := \{ \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_k \mathbf{v}_k : \lambda_1, \dots, \lambda_k \in \mathbb{R} \}.$$





Exercise

Sketch the following spans:







$\text{Span}\left(\begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}\right)$



$\text{Span}\left(\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}\right)$





Exercise

Sketch the following spans:



Exercise

Sketch the following spans:



Linear combination \rightarrow system of linear equations

The following question appears frequently in Linear Algebra: “Given a fixed vector \mathbf{v} , can it be written as **linear combination** of some other given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots$?”

$$\begin{pmatrix} 8 \\ 17 \\ 0 \end{pmatrix} = \boxed{} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + \boxed{} \begin{pmatrix} -3 \\ -5 \\ 5 \end{pmatrix} + \boxed{} \begin{pmatrix} 5 \\ 9 \\ -4 \end{pmatrix}$$

Linear combination \rightarrow system of linear equations

The following question appears frequently in Linear Algebra: “Given a fixed vector \mathbf{v} , can it be written as **linear combination** of some other given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots$?”

$$\begin{pmatrix} 8 \\ 17 \\ 0 \end{pmatrix} = \boxed{x_1} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + \boxed{x_2} \begin{pmatrix} -3 \\ -5 \\ 5 \end{pmatrix} + \boxed{x_3} \begin{pmatrix} 5 \\ 9 \\ -4 \end{pmatrix}$$

Linear combination \rightarrow system of linear equations

The following question appears frequently in Linear Algebra: “Given a fixed vector \mathbf{v} , can it be written as **linear combination** of some other given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots$?”

$$\begin{pmatrix} 8 \\ 17 \\ 0 \end{pmatrix} = \boxed{x_1} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + \boxed{x_2} \begin{pmatrix} -3 \\ -5 \\ 5 \end{pmatrix} + \boxed{x_3} \begin{pmatrix} 5 \\ 9 \\ -4 \end{pmatrix}$$

This corresponds to asking whether a certain **system of linear equations** has a solution:

$$\begin{array}{rcrcrcrcrcl} 8 & = & 1x_1 & - & 3x_2 & + & 5x_3 & & \\ 17 & = & 2x_1 & - & 5x_2 & + & 9x_3 & & \\ 0 & = & -1x_1 & + & 5x_2 & - & 4x_3 & & \end{array}$$

Systems of linear equations

System of linear equations

System of linear equations

Let $m, n \in \mathbb{N}$. A **system of linear equations** (LES) in the variables x_1, x_2, \dots, x_n is of the form

$$\begin{aligned}a_{1,1} x_1 + a_{1,2} x_2 + \cdots + a_{1,n} x_n &= b_1, \\a_{2,1} x_1 + a_{2,2} x_2 + \cdots + a_{2,n} x_n &= b_2, \\&\vdots \\a_{m,1} x_1 + a_{m,2} x_2 + \cdots + a_{m,n} x_n &= b_m.\end{aligned}$$

with $a_{i,j}$ and b_i being (usually real) numbers.

System of linear equations

System of linear equations

Let $m, n \in \mathbb{N}$. A **system of linear equations** (LES) in the variables x_1, x_2, \dots, x_n is of the form

$$\begin{aligned}a_{1,1} x_1 + a_{1,2} x_2 + \cdots + a_{1,n} x_n &= b_1, \\a_{2,1} x_1 + a_{2,2} x_2 + \cdots + a_{2,n} x_n &= b_2, \\&\vdots \\a_{m,1} x_1 + a_{m,2} x_2 + \cdots + a_{m,n} x_n &= b_m.\end{aligned}$$

with $a_{i,j}$ and b_i being (usually real) numbers. An assignment of values for x_1, \dots, x_n such that all equations are satisfied is called a **solution** of this system of equations. Such a solution is written as a vector.

System of linear equations

Example: the system

$$\begin{array}{rrcrcl} 1x_1 & - & 3x_2 & + & 5x_3 & = & 8 \\ 2x_1 & - & 5x_2 & + & 9x_3 & = & 17 \\ -1x_1 & + & 5x_2 & - & 4x_3 & = & 0 \end{array}$$

is a system of linear equations with 3 equations and 3 variables.

System of linear equations

Example: the system

$$\begin{array}{rrcrcl} 1x_1 & - & 3x_2 & + & 5x_3 & = & 8 \\ 2x_1 & - & 5x_2 & + & 9x_3 & = & 17 \\ -1x_1 & + & 5x_2 & - & 4x_3 & = & 0 \end{array}$$

is a system of linear equations with 3 equations and 3 variables.

A solution is given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 7 \\ 3 \\ 2 \end{pmatrix}.$$

System of linear equations

Example: the system

$$\begin{array}{rrcrcl} 1x_1 & - & 3x_2 & + & 5x_3 & = & 8 \\ 2x_1 & - & 5x_2 & + & 9x_3 & = & 17 \\ -1x_1 & + & 5x_2 & - & 4x_3 & = & 0 \end{array}$$

is a system of linear equations with 3 equations and 3 variables.

A solution is given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 7 \\ 3 \\ 2 \end{pmatrix}.$$

Check:

$$7 \cdot \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + 3 \cdot \begin{pmatrix} -3 \\ -5 \\ 5 \end{pmatrix} + 2 \cdot \begin{pmatrix} 5 \\ 9 \\ -4 \end{pmatrix} = \begin{pmatrix} 7 \\ 14 \\ -7 \end{pmatrix} + \begin{pmatrix} -9 \\ -15 \\ 15 \end{pmatrix} + \begin{pmatrix} 10 \\ 18 \\ -8 \end{pmatrix} = \begin{pmatrix} 8 \\ 17 \\ 0 \end{pmatrix}.$$

Method of equalization

Method of equalization

In the method of equalization we isolate the same variable in 2 equations and use the obtained expressions as left- and right-hand side. This way we obtain an equation which does not use one of the given variables.

Method of equalization

Method of equalization

In the method of equalization we isolate the same variable in 2 equations and use the obtained expressions as left- and right-hand side. This way we obtain an equation which does not use one of the given variables.

Example: We solve the following system with the method of equalization:

$$\begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ -4x_1 & + & 2x_2 & = & -9 \end{array}$$

Method of equalization

Method of equalization

In the method of equalization we isolate the same variable in 2 equations and use the obtained expressions as left- and right-hand side. This way we obtain an equation which does not use one of the given variables.

Example: We solve the following system with the method of equalization:

$$\begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ -4x_1 & + & 2x_2 & = & -9 \end{array}$$

Computation:

$$\begin{array}{rclcl} 2x_1 = 15 + 4x_2 & \Rightarrow & x_1 = 15/2 + 2 \cdot x_2 & \Rightarrow & \frac{15}{2} + 2x_2 = \frac{9}{4} + \frac{1}{2}x_2, \\ -4x_1 = -9 - 2x_2 & \Rightarrow & x_1 = 9/4 + 1/2 \cdot x_2 & \Rightarrow & \end{array}$$

thus,

$$\left(2 - \frac{1}{2}\right)x_2 = \frac{3}{2}x_2 = \frac{9}{4} - \frac{15}{2} = -\frac{21}{4}, \quad x_2 = -\frac{7}{2}, \quad x_1 = \frac{15}{2} + 2x_2 = \frac{15}{2} - \frac{14}{2} = \frac{1}{2}.$$

Method of substitution

Method of substitution

In the method of substitution we isolate a variable in an equation and replace it in the other equations. By this we obtain equations which do not use this variable.

Method of substitution

Method of substitution

In the method of substitution we isolate a variable in an equation and replace it in the other equations. By this we obtain equations which do not use this variable.

Example: we solve the following system with the method of substitution:

$$\begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ -4x_1 & + & 2x_2 & = & -9 \end{array}$$

Method of substitution

Method of substitution

In the method of substitution we isolate a variable in an equation and replace it in the other equations. By this we obtain equations which do not use this variable.

Example: we solve the following system with the method of substitution:

$$\begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ -4x_1 & + & 2x_2 & = & -9 \end{array}$$

Computation like before:

$$2x_1 = 15 + 4x_2 \quad \Rightarrow \quad x_1 = \frac{15}{2} + 2x_2.$$

Substitution of first variable in second equation:


$$-4 \left(\frac{15}{2} + 2x_2 \right) + 2x_2 = -30 - 8x_2 + 2x_2 = -9 \quad \Rightarrow \quad -6x_2 = 21 \quad \Rightarrow \quad x_2 = -\frac{7}{2}.$$

First variable like before: $x_1 = 1/2$.

Solving by graphing

Solving by graphing (2 variables)

Consider a system of linear equations with 2 variables. Then each equation can be represented by a line in a 2-dimensional coordinate system. The set of solutions then is represented by the intersection of all lines.




Solving by graphing

Solving by graphing (2 variables)

Consider a system of linear equations with 2 variables. Then each equation can be represented by a line in a 2-dimensional coordinate system. The set of solutions then is represented by the intersection of all lines.

Example: we solve the following system by graphing:

$$\begin{array}{rclcl} 4x_1 & - & 2x_2 & = & 6 \\ 2x_1 & + & x_2 & = & 5 \end{array}$$



Solving by graphing

Solving by graphing (2 variables)


Consider a system of linear equations with 2 variables. Then each equation can be represented by a line in a 2-dimensional coordinate system. The set of solutions then is represented by the intersection of all lines.

Example: we solve the following system by graphing:

$$\begin{array}{rclcl} 4x_1 & - & 2x_2 & = & 6 \\ 2x_1 & + & x_2 & = & 5 \end{array}$$

Solve for x_2 :

$$\begin{aligned} 2x_2 &= -6 + 4x_1, \\ x_2 &= 5 - 2x_1. \end{aligned}$$



Solving by graphing

Solving by graphing (2 variables)

Consider a system of linear equations with 2 variables. Then each equation can be represented by a line in a 2-dimensional coordinate system. The set of solutions then is represented by the intersection of all lines.


Example: we solve the following system by graphing:

$$\begin{array}{rclcl} 4x_1 & - & 2x_2 & = & 6 \\ 2x_1 & + & x_2 & = & 5 \end{array}$$

Solve for x_2 :

$$x_2 = -3 + 2x_1,$$

$$x_2 = 5 - 2x_1.$$



Solving by graphing

Solving by graphing (2 variables)

Consider a system of linear equations with 2 variables. Then each equation can be represented by a line in a 2-dimensional coordinate system. The set of solutions then is represented by the intersection of all lines.


Example: we solve the following system by graphing:

$$\begin{array}{rclcl} 4x_1 & - & 2x_2 & = & 6 \\ 2x_1 & + & x_2 & = & 5 \end{array}$$

Solve for x_2 . Sketch lines & intersection:

$$x_2 = -3 + 2x_1,$$

$$x_2 = 5 - 2x_1.$$



Solving by graphing

Solving by graphing (2 variables)


Consider a system of linear equations with 2 variables. Then each equation can be represented by a line in a 2-dimensional coordinate system. The set of solutions then is represented by the intersection of all lines.

Example: we solve the following system by graphing:

$$\begin{array}{rclcl} 4x_1 & - & 2x_2 & = & 6 \\ 2x_1 & + & x_2 & = & 5 \end{array}$$


Solve for x_2 . Sketch lines & intersection:

$$\begin{array}{l} x_2 = -3 + 2x_1, \\ x_2 = 5 - 2x_1, \end{array} \quad \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$




Solvability: number of solutions


For any lines there exist three kinds of intersections:



intersection point
(one solution)




parallel
(no solution)




identical
(infinitely many solutions)

Solvability: number of solutions


For any lines there exist three kinds of intersections:



intersection point
(one solution)



parallel
(no solution)



identical
(infinitely many solutions)

In general the following is true:

Solvability of systems of linear equations

Every system of linear equations has either (i) exactly one solution or (ii) no solution or (iii) infinitely many solutions.

Method of elimination

Method of elimination

In the method of elimination two equations (or multiples of them) are added/subtracted such that at least one variable is eliminated.

Method of elimination

Method of elimination

In the method of elimination two equations (or multiples of them) are added/subtracted such that at least one variable is eliminated.

Example 1: we solve the following system with the method of elimination:

$$\begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ -4x_1 & + & 2x_2 & = & -9 \end{array}$$

Method of elimination

Method of elimination

In the method of elimination two equations (or multiples of them) are added/subtracted such that at least one variable is eliminated.

Example 1: we solve the following system with the method of elimination:

$$\begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ -4x_1 & + & 2x_2 & = & -9 \end{array} \quad \Rightarrow \quad \begin{array}{rclcl} 2x_1 & - & 4x_2 & = & 15 \\ & & -6x_2 & = & 21 \end{array}$$

Solution:

- ▶ We subtract the first equation $-2 = \frac{-4}{2}$ times from the second, changing the second equation.
- ▶ We solve the new second equation:

$$x_2 = -\frac{21}{6} = -\frac{7}{2}.$$

- ▶ We utilize the first equation:

$$2x_1 = 15 + 4x_2 = 15 - 14 = 1 \quad \Rightarrow \quad x_1 = \frac{1}{2}, \quad \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1/2 \\ -7/2 \end{pmatrix}.$$

Method of elimination

Method of elimination

In the method of elimination two equations (or multiples of them) are added/subtracted such that at least one variable is eliminated.

Example 2: we solve the following system with the method of elimination:

$$\begin{array}{rclcl} 4x_1 & - & 5x_2 & = & 2 \\ -8x_1 & + & 10x_2 & = & -4 \end{array}$$

Method of elimination

Method of elimination

In the method of elimination two equations (or multiples of them) are added/subtracted such that at least one variable is eliminated.

Example 2: we solve the following system with the method of elimination:

$$\begin{array}{rclcl} 4x_1 & - & 5x_2 & = & 2 \\ -8x_1 & + & 10x_2 & = & -4 \end{array} \Rightarrow \begin{array}{rclcl} 4x_1 & - & 5x_2 & = & 2 \\ & & 0x_2 & = & 0 \end{array}$$

Solution:

- ▶ We subtract the first equation $-2 = \frac{-8}{4}$ times from the second, changing the second equation.
- ▶ The new second equation is always satisfied:

$$0 \cdot x_2 = 0 \Rightarrow x_2 \in \mathbb{R}.$$

- ▶ The first equation yields as solution a line:

$$x_1 = \frac{1}{2} + \frac{5}{4}x_2 \Rightarrow \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} + \frac{5}{4}x_2 \\ x_2 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ 0 \end{pmatrix} + x_2 \cdot \begin{pmatrix} \frac{5}{4} \\ 1 \end{pmatrix}, \quad x_2 \in \mathbb{R}.$$

Exercises

Solve the following systems with the method of elimination:

$$(a) \quad \begin{array}{rclcl} 1x_1 & - & 4x_2 & = & 1 \\ 2x_1 & - & 2x_2 & = & 3 \end{array}$$

$$(b) \quad \begin{array}{rclcl} -3x_1 & + & 2x_2 & = & 5 \\ 9x_1 & - & 6x_2 & = & 13 \end{array}$$

Solution of part (a):

Exercises

Solve the following systems with the method of elimination:

$$(a) \quad \begin{array}{rclcl} 1x_1 & - & 4x_2 & = & 1 \\ 2x_1 & - & 2x_2 & = & 3 \end{array}$$

$$(b) \quad \begin{array}{rclcl} -3x_1 & + & 2x_2 & = & 5 \\ 9x_1 & - & 6x_2 & = & 13 \end{array}$$

Solution of part (a):

► Compute $II - 2I$:

$$\begin{array}{rclcl} 1x_1 & - & 4x_2 & = & 1 \\ 2x_1 & - & 2x_2 & = & 3 \end{array} \Rightarrow \begin{array}{rclcl} 1x_1 & - & 4x_2 & = & 1 \\ & & 6x_2 & = & 1 \end{array}$$

► Solve second new equation:

$$6x_2 = 1 \Rightarrow x_2 = \frac{1}{6}.$$

► Insert into first equation:

$$x_1 = 1 + 4x_2 = 1 + \frac{4}{6} = \frac{10}{6} = \frac{5}{3} \quad \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} \frac{5}{3} \\ \frac{1}{6} \end{pmatrix}.$$

Exercises

Solve the following systems with the method of elimination:

$$(a) \quad \begin{array}{rclcl} 1x_1 & - & 4x_2 & = & 1 \\ 2x_1 & - & 2x_2 & = & 3 \end{array}$$

$$(b) \quad \begin{array}{rclcl} -3x_1 & + & 2x_2 & = & 5 \\ 9x_1 & - & 6x_2 & = & 13 \end{array}$$

Solution of part (b):

Exercises

Solve the following systems with the method of elimination:

$$(a) \quad \begin{array}{rclcl} 1x_1 & - & 4x_2 & = & 1 \\ 2x_1 & - & 2x_2 & = & 3 \end{array}$$

$$(b) \quad \begin{array}{rclcl} -3x_1 & + & 2x_2 & = & 5 \\ 9x_1 & - & 6x_2 & = & 13 \end{array}$$

Solution of part (b):

- Compute $II + 3I$:

$$\begin{array}{rclcl} -3x_1 & + & 2x_2 & = & 5 \\ 9x_1 & - & 6x_2 & = & 13 \end{array} \Rightarrow \begin{array}{rclcl} -3x_1 & + & 2x_2 & = & 5 \\ 0x_2 & = & 28 & & \end{array}$$

- Solve second new equation:

$$0x_2 = 28 \Rightarrow \text{not solvable.}$$

- The linear system is not solvable.

Solving bigger systems of linear equations

Observations

- ▶ The set of solutions does not change if the ordering of the equations is changed.
- ▶ The set of solutions does not change if one equation is subtracted (multiple times) from the other equations.

Solving bigger systems of linear equations

Observations

- ▶ The set of solutions does not change if the ordering of the equations is changed.
- ▶ The set of solutions does not change if one equation is subtracted (multiple times) from the other equations.

Recipe for calculations:

- ▶ Bring the system to a “triangle form” or “row echelon form”.
- ▶ Afterwards determine the set of solutions.

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right) \begin{array}{l} \text{II} - 2 \cdot \text{I} \\ \text{III} + 1 \cdot \text{I} \end{array} \rightsquigarrow \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 4 & 6 & -10 \end{array} \right)$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 4 & 6 & -10 \end{array} \right) \text{ III} - 4 \cdot \text{II} \rightsquigarrow \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 2 & -2 \end{array} \right)$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 2 & -2 \end{array} \right)$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 2 & -2 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -3 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ -2 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ \end{pmatrix}.$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 2 & -2 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -3 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ -2 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ -1 \end{pmatrix}.$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 2 & -2 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -3 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ -2 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2 + 1 = -1 \\ -1 \\ -1 \end{pmatrix}.$$

Example

We determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 3x_3 & = & 3 \\ 2x_1 & + & 7x_2 & - & 5x_3 & = & 4 \\ -1x_1 & + & 1x_2 & + & 9x_3 & = & -13 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 2 & 7 & -5 & 4 \\ -1 & 1 & 9 & -13 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -3 & 3 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 2 & -2 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -3 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ -2 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 + 3 - 3 = 3 \\ -2 + 1 = -1 \\ -1 \end{pmatrix}.$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right) \begin{array}{l} \text{II} - 3 \cdot \text{I} \\ \text{III} - 2 \cdot \text{I} \end{array} \rightsquigarrow \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & -4 & 0 & 4 \end{array} \right)$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & -4 & 0 & 4 \end{array} \right) \text{ III} + 2 \cdot \text{II} \quad \rightsquigarrow \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -2 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 5 \\ -4 \\ -4 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ \end{pmatrix}.$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -2 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 5 \\ -4 \\ -4 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ -2 \end{pmatrix}.$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -2 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 5 \\ -4 \\ -4 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} (-4 + 2)/2 = -1 \\ -2 \\ -2 \end{pmatrix}.$$

Exercise

Determine the unique solution of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & + & 3x_2 & - & 2x_3 & = & 5 \\ 3x_1 & + & 11x_2 & - & 5x_3 & = & 11 \\ 2x_1 & + & 2x_2 & - & 4x_3 & = & 14 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 11 & -5 & 11 \\ 2 & 2 & -4 & 14 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 2 & 1 & -4 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

- Solve from below:

$$\left(\begin{array}{ccc} 1 & 3 & -2 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 5 \\ -4 \\ -4 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 5 + 3 - 4 = 4 \\ (-4 + 2)/2 = -1 \\ -2 \end{pmatrix}.$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right) \begin{array}{l} \text{II} - 1 \cdot \text{I} \\ \text{III} + 2 \cdot \text{I} \end{array} \rightsquigarrow \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & -3 & 6 & -9 \end{array} \right)$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & -3 & 6 & -9 \end{array} \right) \text{ III} + 3 \cdot \text{II} \rightsquigarrow \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

- Solve from below:

$$\begin{pmatrix} 1 & -5 & 4 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ \end{pmatrix}$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

- Solve from below: ($x_3 \in \mathbb{R}$)

$$\begin{pmatrix} 1 & -5 & 4 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ x_3 \end{pmatrix}$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

- Solve from below: ($x_3 \in \mathbb{R}$)

$$\begin{pmatrix} 1 & -5 & 4 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 + 2x_3 \\ x_3 \end{pmatrix}$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

- Solve from below: ($x_3 \in \mathbb{R}$, $-2 + 5(3 + 2x_3) - 4x_3 = 13 + 6x_3$)

$$\begin{pmatrix} 1 & -5 & 4 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 13 + 6x_3 \\ 3 + 2x_3 \\ x_3 \end{pmatrix}$$

An example with infinitely many solutions

We determine the solutions of the following system of linear equations:

$$\begin{array}{rrcrcl} 1x_1 & - & 5x_2 & + & 4x_3 & = & -2 \\ 1x_1 & - & 4x_2 & + & 2x_3 & = & 1 \\ -2x_1 & + & 7x_2 & - & 2x_3 & = & -5 \end{array} \quad \text{short,} \quad \left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 1 & -4 & 2 & 1 \\ -2 & 7 & -2 & -5 \end{array} \right).$$

Solution:

- Compute upper triangular form:

$$\left(\begin{array}{ccc|c} 1 & -5 & 4 & -2 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

- Solve from below: $(x_3 \in \mathbb{R}, -2 + 5(3 + 2x_3) - 4x_3 = 13 + 6x_3)$

$$\begin{pmatrix} 1 & -5 & 4 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 13 + 6x_3 \\ 3 + 2x_3 \\ x_3 \end{pmatrix} = \begin{pmatrix} 13 \\ 3 \\ 0 \end{pmatrix} + x_3 \cdot \begin{pmatrix} 6 \\ 2 \\ 1 \end{pmatrix}.$$