

Slides are available on my website.
<http://lukaseigentler.github.io>

Variation in resource quality causes maintenance of individual variation in a competitive trait

Behavior, Evolution & Emergence (BEE'26)


18 June 2026

Lukas Eigentler (University of Warwick)
joint work with

Klaus Reinhold (Bielefeld), David W. Kikuchi (Oregon State), Valeria Giunta (Swansea)

Individual variation

- Evolution does not necessarily lead to identical individuals.
- Individual variation is ubiquitous in many populations.
- This includes **genetic variation**.
- Potential causes: temporal/spatial environmental variability, negative frequency-dependent selection, ...



Bluegill size differences¹

¹ (Yokogawa, K.: *Ichthyological Research* 60.1 [2012])

Individual variation in competitive traits

- Traits determining intraspecific competitiveness often vary within populations.
- Examples: armaments, body size, sensory capabilities to detect food, ...
- **Investment into competitiveness typically bears a cost.**
- Trade off between ability to attain resources and resource investment into reproduction.




Source: Wikimedia Commons

Previous work

- Previous work mainly considered a small finite number of genotypes.

Previous work

- Previous work mainly considered a small finite number of genotypes.
- Notable exception: Assuming a continuous competitive trait in a population competing for two different resource quality levels found either dimorphisms or repeated cycles of “arms races” depending on the difference between the two resource quality levels.²




Source: Wikimedia Commons

(Baldauf, S. A., Engqvist, L. and Weissing, F. J.: *Nat Commun* 5.1 [2014])

Previous work

- Previous work mainly considered a small finite number of genotypes.
- Notable exception: Assuming a continuous competitive trait in a population competing for two different resource quality levels found either dimorphisms or repeated cycles of “arms races” depending on the difference between the two resource quality levels.²
- What if resource qualities also vary continuously?



Source: Wikimedia Commons


(Baldauf, S. A., Engqvist, L. and Weissing, F. J.: *Nat Commun* 5.1 [2014])

Individual based model



- Large population of fixed size N .


Individual based model



- Large population of fixed size N .
- Individual $i \in \{1, \dots, N\}$ has competitive trait $c_i \geq 0$.


Individual based model

- Large population of fixed size N .
- Individual $i \in \{1, \dots, N\}$ has competitive trait $c_i \geq 0$.
- N resources of quality $d \leq r_j \leq 1, j \in \{1, \dots, N\}$ drawn from a continuous distribution. d is the ratio of the worst to the best resource.




Individual based model

- Large population of fixed size N .
- Individual $i \in \{1, \dots, N\}$ has competitive trait $c_i \geq 0$.
- N resources of quality $d \leq r_j \leq 1, j \in \{1, \dots, N\}$ drawn from a continuous distribution. d is the ratio of the worst to the best resource.
- Best competitor attains best resource, and so on.



Individual based model


- Large population of fixed size N .
- Individual $i \in \{1, \dots, N\}$ has competitive trait $c_i \geq 0$.
- N resources of quality $d \leq r_j \leq 1, j \in \{1, \dots, N\}$ drawn from a continuous distribution. d is the ratio of the worst to the best resource.
- Best competitor attains best resource, and so on.
- Reproductive fitness is $W_i = r_i(1 - c_i)$.



Reproductive fitness: $W_i = r_i(1 - c_i)$


Individual based model

- Large population of fixed size N .
- Individual $i \in \{1, \dots, N\}$ has competitive trait $c_i \geq 0$.
- N resources of quality $d \leq r_j \leq 1, j \in \{1, \dots, N\}$ drawn from a continuous distribution. **d is the ratio of the worst to the best resource.**
- Best competitor attains best resource, and so on.
- Reproductive fitness is $W_i = r_i(1 - c_i)$.
- Next generation is created through draw with replacements with weights W_i and mutations.





Resource distributions

- A: linearly increasing
- B: linearly increasing and decreasing
- C: uniform
- D: linearly decreasing
- E: half-normal
- F: power law




Arms races vs. stable polymorphisms

- Dynamics either represent **arms races** or **stable polymorphisms**.
- Arms race: mean competitive trait oscillates with little individual variation at a time.
If all individuals have $c = c^*$, then $c = c^* + \varepsilon > c^*$ has higher fitness.
If c^* is high, low competitive traits can invade; in particular, $c_1 = 0$ (fitness $W = (1 - c_1)d = d$) has larger fitness than $c_2 > 1 - d$ because $W = (1 - c_2)r < dr \leq d$.
- Stable polymorphism: mean competitive trait remains constant over time; large individual variation at all times.




Worst-to-best resource ratio

- The ratio d of the worst to the best resource determines mean investment into competition.
- Arms races occur when d is small.
- Period of arms race cycles increases as d decreases; unclear behaviour for $d \rightarrow 0$.
- No arms races for power law distribution.
- Stable mean trait values occur when d is large.
- The threshold $d = d^*$ depends on the resource distribution.




Worst-to-best resource ratio

- Little individual variation occurs during arms races.
- Stable mean trait values are always associated with stable polymorphisms.
- Individual variation is close to maximum (uniform distribution with $0 \leq c \leq 1 - d$).




Special case: power law

- Can show analytically that the power law with pdf $g(r) = \frac{d}{(1-d)r^2}$ leads to uniform trait distribution.




Special case: power law

- Can show analytically that the power law with pdf $g(r) = \frac{d}{(1-d)r^2}$ leads to uniform trait distribution.
- look for trait distribution in which all values $0 \leq c \leq 1 - d$ attain equal fitness
 $W = (1 - c)r(c) = d \Leftrightarrow r(c) = \frac{d}{1-c}.$



Special case: power law

- Can show analytically that the power law with pdf $g(r) = \frac{d}{(1-d)r^2}$ leads to uniform trait distribution.
- look for trait distribution in which all values $0 \leq c \leq 1 - d$ attain equal fitness
 $W = (1 - c)r(c) = d \Leftrightarrow r(c) = \frac{d}{1-c}$.
- Hierarchical resource assignment means $P(r < r^*) = P(c < c^*)$, where $r^* = \frac{d}{1-c^*}$.




Special case: power law

- Can show analytically that the power law with pdf $g(r) = \frac{d}{(1-d)r^2}$ leads to uniform trait distribution.
- look for trait distribution in which all values $0 \leq c \leq 1 - d$ attain equal fitness
 $W = (1 - c)r(c) = d \Leftrightarrow r(c) = \frac{d}{1-c}$.
- Hierarchical resource assignment means $P(r < r^*) = P(c < c^*)$, where $r^* = \frac{d}{1-c^*}$.
Therefore, the cdf of the trait distribution is

$$P(c < c^*) = P(r < r^*) = \frac{r^* - d}{(1 - d)r^*} = \frac{c^*}{1 - d}.$$

$\Rightarrow$ uniform distribution.




Conclusions – IBM

- Continuous individual variation in a competitive trait is a consequence of intraspecific competition for resources with continuously varying quality.
- The ratio between worst and best quality resources determines (i) mean investment into competition, and (ii) whether arms races with little individual variation or stable polymorphisms occur.
- The distribution of resource quality is important.
- Resource distributions with more low-quality resources lead to stable polymorphisms in more cases.

What next? Nonlocal PDE!

- Main limitation: IBMs are hard to analyse.
- PDE models allow for bifurcation analysis.
- Formal derivation of a PDE from the IBM is (we think) not possible.



A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.

A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.
- Let $u(t, c)$ be the size of the population with trait value $c \geq 0$ at time $t \geq 0$.

A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.
- Let $u(t, c)$ be the size of the population with trait value $c \geq 0$ at time $t \geq 0$.
- Fixed population size $\Rightarrow \int_0^\infty u(t, c)dc \equiv 1$ fixed, i.e. u is a pdf.

A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.
- Let $u(t, c)$ be the size of the population with trait value $c \geq 0$ at time $t \geq 0$.
- Fixed population size $\Rightarrow \int_0^\infty u(t, c)dc \equiv 1$ fixed, i.e. u is a pdf.
- Evolution equation:

$$\frac{\partial u}{\partial t} = [\text{mutation}] + [\text{selection}].$$

A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.
- Let $u(t, c)$ be the size of the population with trait value $c \geq 0$ at time $t \geq 0$.
- Fixed population size $\Rightarrow \int_0^\infty u(t, c)dc \equiv 1$ fixed, i.e. u is a pdf.
- Evolution equation:

$$\frac{\partial u}{\partial t} = [\text{mutation}] + [\text{selection}].$$

- Mutation: mutation in a single gene will have very small impact on trait value $\Rightarrow$ diffusion.

A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.
- Let $u(t, c)$ be the size of the population with trait value $c \geq 0$ at time $t \geq 0$.
- Fixed population size $\Rightarrow \int_0^\infty u(t, c)dc \equiv 1$ fixed, i.e. u is a pdf.
- Evolution equation:

$$\frac{\partial u}{\partial t} = [\text{mutation}] + [\text{selection}].$$

- Mutation: mutation in a single gene will have very small impact on trait value $\Rightarrow$ diffusion.
- Selection: net growth of population with trait c .

A phenomenological PDE

- Idea: derive a phenomenological PDE using the same assumptions as for the IBM.
- Let $u(t, c)$ be the size of the population with trait value $c \geq 0$ at time $t \geq 0$.
- Fixed population size $\Rightarrow \int_0^\infty u(t, c)dc \equiv 1$ fixed, i.e. u is a pdf.
- Evolution equation:

$$\frac{\partial u}{\partial t} = [\text{mutation}] + [\text{selection}].$$

- Mutation: mutation in a single gene will have very small impact on trait value $\Rightarrow$ diffusion.
- Selection: net growth of population with trait c .

- $$\frac{\partial u(t, c)}{\partial t} = \frac{\partial^2 u(t, c)}{\partial c^2} + \lambda(t, c)u(t, c).$$

Growth rate

- Growth rate

$$\lambda(t, c) = \underbrace{W(t, c)}_{\text{fitness of phenotype } c} - \underbrace{\overline{W}(t)}_{\text{average fitness}} .$$

Growth rate

- Growth rate

$$\lambda(t, c) = \underbrace{W(t, c)}_{\text{fitness of phenotype } c} - \underbrace{\overline{W}(t)}_{\text{average fitness}} .$$

- Average fitness

$$\overline{W}(t) = \int_0^\infty W(t, c) u(t, c) dc.$$

Growth rate

- Growth rate

$$\lambda(t, c) = \underbrace{W(t, c)}_{\text{fitness of phenotype } c} - \underbrace{\overline{W}(t)}_{\text{average fitness}} .$$

- Average fitness

$$\overline{W}(t) = \int_0^\infty W(t, c) u(t, c) dc .$$

- Fitness of phenotype c

$$W(t, c) = \underbrace{(1 - c)}_{\text{proportion left for reproduction}} \cdot \underbrace{r(t, c)}_{\text{resource quality attained}} .$$

Resource quality

- For $0 \leq d < 1$, let the pdf of the resource distribution be $g : [d, 1] \rightarrow [0, \infty)$, such that the inverse $G^{-1} : [0, 1] \rightarrow [d, 1]$ of its cdf $G : [d, 1] \rightarrow [0, 1]$ exists.

Resource quality

- For $0 \leq d < 1$, let the pdf of the resource distribution be $g : [d, 1] \rightarrow [0, \infty)$, such that the inverse $G^{-1} : [0, 1] \rightarrow [d, 1]$ of its cdf $G : [d, 1] \rightarrow [0, 1]$ exists.
- Denote by $U(t, c) = \int_0^c u(t, \tilde{c}) d\tilde{c}$ the cdf of the population with respect to the trait c .

Resource quality

- For $0 \leq d < 1$, let the pdf of the resource distribution be $g : [d, 1] \rightarrow [0, \infty)$, such that the inverse $G^{-1} : [0, 1] \rightarrow [d, 1]$ of its cdf $G : [d, 1] \rightarrow [0, 1]$ exists.
- Denote by $U(t, c) = \int_0^c u(t, \tilde{c}) d\tilde{c}$ the cdf of the population with respect to the trait c .
- Hierarchical resource assignment means that for any fixed trait value c^* , the attained resource quality r^* is defined implicitly by

$$G(r^*) = U(t, c^*).$$

Resource quality

- For $0 \leq d < 1$, let the pdf of the resource distribution be $g : [d, 1] \rightarrow [0, \infty)$, such that the inverse $G^{-1} : [0, 1] \rightarrow [d, 1]$ of its cdf $G : [d, 1] \rightarrow [0, 1]$ exists.
- Denote by $U(t, c) = \int_0^c u(t, \tilde{c}) d\tilde{c}$ the cdf of the population with respect to the trait c .
- Hierarchical resource assignment means that for any fixed trait value c^* , the attained resource quality r^* is defined implicitly by

$$G(r^*) = U(t, c^*).$$

•


$$\Rightarrow r(t, c) = G^{-1}(U(t, c)).$$

Full nonlocal model

$$\frac{\partial u(t, c)}{\partial t} = \frac{\partial^2 u(t, c)}{\partial c^2} + \left[(1 - c)G^{-1} \left(\int_0^c u(t, \tilde{c}) d\tilde{c} \right) - \int_0^\infty (1 - c)G^{-1} \left(\int_0^c u(t, \tilde{c}) d\tilde{c} \right) u(t, c) dc \right] u(t, c).$$


Numerical simulations

- Results qualitatively the same as for IBM.
- Note lack of oscillations for small d .




1d numerical bifurcation analysis **work in progress**

Tool: discretise trait space $\Rightarrow$ continuation of large ODE system using AUTO-07p.



From high d to low d : constant sol $\rightarrow$ onset of periodic sol via Hopf bifurcation $\rightarrow$ termination of periodic sol at either Hopf bifurcation or homoclinic-fold bifurcation $\rightarrow$ constant sol.

2d numerical bifurcation analysis: m - d **work in progress**




Periodic sol require sufficiently low mutation rate. Onset at low d occurs at Hopf bifurcation for “large” m ; at homoclinic-fold bifurcation at “small” m .

2d bif analysis: generalised linear distribution **work in progress**

Distributions A, D are special cases of general distribution with pdf

$$g(r; \alpha) = \alpha r + \beta, \quad d \leq r \leq 1, \quad -\frac{2}{(1-d)^2} \leq \alpha \leq \frac{2}{(1-d)^2}$$

⇒ Continuation in α to study dynamics for this generalised distribution.



Analytical results **work in progress** – Valeria Giunta

We can show analytically that

- the existence of lower and upper bounds for steady state solutions.
- trait distributions are monomodal with peak at $c = 0$ for sufficiently large d .
- $d > \bar{d}$ is a sufficient condition for constant solutions to be stable.

Conclusions – PDE

- Numerical bifurcation analysis of PDE model is possible.
- Pattern onset at Hopf bifurcation (high/low d) and homoclinic-fold bifurcation (low d).
- Can perform 2-parameter continuation of these bifurcations.
- Low mutation rates required for periodic “arms race” solutions.
- Can find codimension 2 bifurcation, such as Bogdanov-Takens.
- Can perform continuation in a parameter describing the resource quality distribution to understand how different distributions affect results.

Open questions

- What are the most realistic resource quality distributions?
- How to include more realism, such as spatial dynamics or sexual reproduction?
- Are there better tools for numerical bifurcation analysis of nonlocal PDEs (BifurcationKit, pde2path)?
- Can we formally derive a continuum model from the IBM?

References

Slides are available on my website.

<http://lukaseigentler.github.io>

- [1] Reinhold, K., Eigentler, L. and Kikuchi, D. W.: 'Evolution of individual variation in a competitive trait: a theoretical analysis'. *J. Evolution. Biol.* 37.5 (2024), pp. 538–547.