

# Space Physics Project , FYS3610

Sigve Harang, Lukas Powalla

November 29, 2015

# Contents

<b>1 Theory part</b>	<b>3</b>
1.1 The Dungey cycle - a steady state picture . . . . .	3
1.2 Auroral substorm . . . . .	4
1.2.1 groth phase . . . . .	4
1.2.2 Expansion phase . . . . .	4
1.2.3 recovery phase . . . . .	4
1.3 Currents . . . . .	4
1.3.1 Earth's current system . . . . .	5
1.4 Twin cell convection . . . . .	5
<b>2 Observations</b>	<b>6</b>
2.1 ACE . . . . .	6
2.2 Groundbasedmagnetometer . . . . .	8
2.3 SuperDARN . . . . .	9
2.4 AMPERE . . . . .	10
2.5 Svalbard All-Sky Imager . . . . .	11
<b>3 discussion</b>	<b>13</b>
<b>4 conclusion</b>	<b>14</b>

# Introduction


In this project we are going to analyse the ionospheric and auroral dynamics and its relation to the solar wind conditions. One of the most relevant physical mechanism is magnetic reconnection. We will also investigate in near earth current flows. Currents and magnetic fields are connected with each other through Maxwell's equations and influence each other. Finally, we will compare our observations to theory. We will investigate in a specific timeslot of 6 hours from the 6. January 2011 from 18:00 o'clock to the 07. January 2011 00:00 o'clock.

## 1 Theory part

In order to understand the movements and changes in the system of the earth and the sun, we want to introduce some basic concepts of space physics. A really important steady state picture of what is going on with respect to the earth's magnetic field, the interaction of the earth's magnetic field with the IMF and the consequences of the earth's environment (such as current flows or aurora) is the Dungey cycle. (the Dungey cycle is only valid when the IMF is southward orientated) The Dungey cycle is as already mentioned a steady state picture. This means that the magnetic flux added to the open magnetic field lines through day side reconnection is equal to the magnetic flux subtracting from the open magnetic field lines through night side reconnection. However, this cycle is in reality in general not a steady state picture. We observe different phases such as adding magnetic flux (growing of the polar cap), some kind of explosive expansion (subtraction of magnetic flux from the polar cap) and a recovery phase. This phenomena is called a "Auroral substorm". In addition to that, we can extract information about what is going on with the system earth-sun by looking at different data sets containing information about the magnetic field on earth, the solar wind bulk speed, the auroral intensity, the current density and further data from different physical measurements. Based on this measurements we will interpret the received data and relate it to the theoretical predictions and the models we introduced in the theoretical part.

### 1.1 The Dungey cycle - a steady state picture

Figure 1: Dungey cycle; THIS IS PARTLY WRONG!!!!



If the magnetic field is northward orientated ( $B_z > 0$ ) nothing happens at the boundary of the terrestrial magnetic field and the magnetic field of the sun/ the solar wind. IMF can not mix with the terrestrial magnetic field because everything is frozen in. Since the terrestrial magnetic field and the IMF are not anti-parallel orientated, magnetic reconnection does not occur and hence, everything is and stays frozen in. However, if the IMF is southward orientated ( $B_z < 0$ ), we do have anti-parallel orientated magnetic field at the boundary of terrestrial magnetic field and the IMF (at the magnetopause). Therefore, magnetic reconnection occurs at the magnetopause, which causes a cyclic reconnection pattern at day/night-side. If we assume a steady state picture, we call the cycle, which happens for southward IMF, the dungey cycle. The different parts of the cycle can be described as follows:

- 1 opposite polarity field lines reconnect at magnetopause (if IMF southward orientated)
- 2 magnetic field lines are still connected to the solar wind. (frozen in again) Therefore, the magnetic field line is draged along with the solar wind in the direction of the tale. (magnetic tension force)

- 3 magnetic flux is added to the tail and compresses the plasma sheet (magnetic pressure increases, plasma sheet gets thinner and thinner)
- 4 magnetic reconnection occurs in the tail.
- 5 reconnected magnetic field lines return to the day side (imagination of "rubber bands" help to understand this)

## 1.2 Auroral substorm

As already mentioned in the introduction, the assumption of a steady state picture is not always a appropriate description of what happens in reality. To specify this, the rate of adding magnetic flux and the rate of subtracting magnetic flux from the polar cap are in general not equal. However, after studying data from different measurements, Akasofu could state out in 1964 different phases of a cycle so called:

- 1 growth phase
- 2 expansion phase
- 3 recovery phase

### 1.2.1 growth phase

The day side reconnection dominates and the aurora moves to lower latitudes. The auroral activity increases in the cusp and open magnetic flux is added to the polar cap. The magnetic flux, which is added due to the day side reconnection makes the tail radius increase.(because the field lines are draged tail-wards with the IMF )


### 1.2.2 Expansion phase

Sudden onset of night-side reconnection at the NENL. Sudden brightening of the night side aurora. Poleward and westward movement of the aurora (on average). It closes open magnetic flux.

### 1.2.3 recovery phase

The night side reconnection continues at smaller rates and the auroral display quietens. Furthermore, the polar cap contracts and the magnetosphere returns to pro growth phase "equilibrium".

Figure 2: Auroral substorm



## 1.3 Currents

When we are dealing with the interaction of the earth's magnetic field with the IMF, it seems to be reasonable to investigate into how the currents close to earth behave. The reason for this is due to the connection of magnetic fields

and flowing currents. We know from Maxwell's equations that flowing electric currents cause a magnetic field. In particular, this relation is given by Ampere's law (which follows from Maxwell's equations):

$$\nabla \times \vec{B} = \mu_0 \cdot \vec{J} \quad (1)$$

Therefore, we can derive the current flow by looking at the magnetic field. (in particular by looking at the curl of the magnetic field) (see measurement AMPERE)

In general, we can express the current flow in a given system with given magnetic field and electric field as follows:

$$\vec{J} = en(\vec{v}_i - \vec{v}_e) = \sigma_p \vec{E}'_{\perp} + \sigma_H \frac{\vec{E}'_{\perp} \times \vec{B}}{B} + \sigma_{\parallel} \vec{E}_{\parallel} \quad (2)$$

Here,  $\sigma_p$  is the Pedersen conductivity,  $\sigma_H$  is the Hall conductivity and  $\sigma_{\parallel}$  is a conductivity, which belongs to a current parallel to the electric field. We notice that the different currents caused by different conductivities flow in orthogonal directions with respect to each other. The orientation of  $\vec{E}$  means parallel or orthogonal to the magnetic field. A discrete expression of the conductivities will not be needed in this report.

$$\vec{E} = \vec{E}_{\perp} + \vec{E}_{\parallel} \quad (3)$$

$$\vec{E}'_{\perp} = \vec{E}_{\perp} + \vec{v}_n \times \vec{B} \quad (4)$$

$$\vec{E}'_{\perp} \approx \vec{E}_{\perp} \quad (5)$$

### 1.3.1 Earth's current system

The magnetic field of the terrestrial magnetosphere is produced by superposition of magnetic fields from a variety of sources. [6][p.400] Near the earth's surface, the biggest contribution comes from the currents within the earth's liquid core. However, electric currents like the solar dynamo, the equatorial electrojet, the convection electrojets and the sub storm electrojets play also an important role. [6][p.400]. We want to give a short overview about the current system of the earth. We want to limit us on currents, which can actually be measured by the measurements used in this report. The currents, which flow near the polar cap can be described as follows: On the polar cap, we find Hall currents, Pedersen currents and Field-aligned currents. The Electron Hall currents flow at lower altitudes than the Pedersen currents. The flow of the Hall currents is also associated with the so called twin cell convection. The ionospheric Pedersen currents are connected to the Field aligned currents (FAC). The FAC are on the one hand side closed with the magnetopause current and on the other hand side with the ring current. This field aligned currents are discovered by a Norwegian explorer called Kristian Birkeland in 1908. All current systems need to be closed, which is the case. The current flow on the polar cap can be seen in figure 3.




Figure 3: electric currents on the polar cap[1]

### 1.4 Twin cell convection

If we look at the Dungey cycle ( $B_z < 0$ ), we can figure out the movement of the open/closed field lines at the polar caps, which gives us the movement of the particles at the polar cap. Let us first explain, what leads to the so called "Twin-cell"-convection. If we have southward IMF, the orientation of the IMF and the earth's magnetic field are opposite to each other. Like discussed in chapter ??, magnetic reconnection on the day side of the earth can occur. The magnetic reconnection at the day side adds open magnetic field lines to the polar cap. In other words, day side reconnection adds magnetic flux to the polar cap. The deformation of the usually round open/closed-field boundary

leads to a movement, which tries to "smooth out" the little bump in the boundary. Magnetic field lines naturally want themselves to orientate in a straight line.

Let us now look at the night side. If we look at the night side events, we know that magnetic reconnection occurs in the tail.(dungey cycle) Reasons for this happening are for example that through day side reconnection, magnetic flux is added to the tail, the magnetic pressure increases and leads in the end to magnetic reconnection in the tail. The magnetic reconnection to the tail closes open magnetic field lines. This means that the polar cap shrinks a bit on the night side. In other words, magnetic flux is subtracted at the night side of the earth from the polar cap. The subtraction leads to a little dint in the polar cap. The magnetic field lines tend to "smooth out" the little dint, which leads to a corresponding movement.

In total, we expect now a movement, which is well known as the twin cell convection. The electric potential on the poles will give us the movement of the charged particles. The currents move along potential lines. The equipotential lines are therefore the trajectories of the charged particles. The currents are called "Hall"-currents. The hall-currents can be found in an altitude of approximately 110 km. (electro-jets)

## 2 Observations

All in all, we use five different sources and data to draw an image of what is going on with the interaction of the IMF with the earth's magnetic field. We will just shortly list the used methods in this reports. A detailed description of each of the methods can be found in the separated chapters. First of all, we use a satellite called ACE (Advanced composition explorer) in order to get knowledge about the magnetic field of the IMF at earth's position, which tells us whether day side magnetic reconnection occurs or not . We also use Groundbasedmagnetometer to get data from the magnetic field on earth. Depressions in the magnetic field can finally be used to detect auroral sub storms. Moreover, we want to look at data from SuperDARN in order to know the movement of charged particles in the ionosphere. The data from AMPERE gives rise to Region 1 and Region 2 currents. Finally, the All-sky-Camera at Svalbard helps us to detect sub storms.

### 2.1 ACE

In this section, we want to discuss the data from the ACE. (Advanced Composition Explorer) The ACE satellite orbits the Lagrangian point  $L_1$ .






Figure 4: position of ACE [2]

This Lagrangian point sits exactly between the earth and the sun.(see figure 4) The solar wind particles move radially outwards from the sun. Therefore, the data from ACE can be used to describe the IMF at earth's position when we take the time into account, which is needed from the solar wind to travel from the position of the ACE to earth. If we take a look at the position of ACE and the solar-wind speed at the position of ACE, we can calculate the time the IMF needs to travel from ACE to earth.

Finally, we know what the IMF looks like at earth's position at a given time on earth. This is important because we know from the chapter ??, that the  $B_z$  tells us something about magnetic reconnection on the day side. If  $B_z > 0$  (or northward IMF orientation), there is no magnetic reconnection on the day side. There is no open magnetic flux added to the polar cap. However, if the  $B_z$  component is negative (or southward orientation of IMF) magnetic reconnection on the day side can occur and the dungey cycle kicks in. It is therefore very important to get data from the  $B_z$  component of the IMF. The other components of the IMF can be used at a later point to tell something about the specific shape of the twin-cell convection. In figures 5a and 6, we find data from the solarwindspeed and the ACE position. Both together will be used to determine the time delay of the IMF. The coordinate system GSE (Geocentric Solar Ecliptic). If we assume a average velocity of the solar wind (see 5a  $V_x$ ) of  $400 \frac{m}{s}$  and a average distance of  $1.4802 \cdot$



(a) Solarwind data



(b) magnetic field measured at the ACE

Figure 5: ACE data




Figure 6: ACE distance from earth

$10^6$ km from ACE to earth, we get a average time delay of 1 hour and.(range between 55 minutes and 70 minutes) This is the time, the solar wind needs to travel from ACE to earth.

From 18 o'clock to approximately 20:15 o'clock, the IMF at ACEs' position is slightly positive. We can see that the  $B_z$ -component of the IMF at the position of ACE turns negative at approximately 20:15 o'clock. (to  $B_z \approx -12 nT$ ) From there on, the IMF stays almost all the time negative until 23:45 o'clock. At 21:20 o'clock, the  $B_z$  component turned for a short time to relatively high positive values.

## 2.2 Groundbasedmagnetometer

The Groundbasedmagnetometer we use is a collection of 32 station on the surface of the earth measuring the magnetic field. (by the use of magnetometers) The collection is called IMAGE (International Monitor for Auroral Geomagnetic Effects). We have chosen a night side event, which means that we have chosen a time when the All-sky-camera in Svalbard is on the night side. If we have a look at our data, we should be aware of indications concerning auroral sub storms. During a sub storm, the westward electro-jets increase. The currents cause according to Ampere's rule a depression in the north-south component of the magnetic field on earth. After the depression, the magnitude of the north-south component slowly increases to its original value. Observing this depression is therefore a first hint of a sub storm. Let us now have a closer look at the data.




Figure 7: Data from Groundbasedmagnetometer, different components

In figure 7, you can see data from IMAGE. The figures 7a, 7b and 7c show different orthogonal components of the magnetic field. In figure 7a, you can see the south- north component of the magnetic field, which is the important component for us. The vertical axis of the plots show the time and the different horizontal lines are the magnetic field components measured at different stations on earth. On the right side, you can see abbreviations of the different stations. Note, that they are ordered concerning their latitude.

The first thing, which strikes the eye is that the magnetic field component is mostly constant from 18:00 o'clock until approximately 22:00 o'clock. At approximately 22:00 o'clock (depends on the latitude of the station), we can see fluctuation of the magnetic field component and then a sudden depression of the magnetic field component. The first minimum of the measured magnetic field can be seen between 22:30 and 23:00 o'clock. This is a hint to a sub storm.(see chapter 3 ) We can also see a poleward moving of the depression especially for the 8 most northward stations (KEV to NAL). We can also see that the stations MEK to TAR don't measure a depression of the magnetic field. The stations KEV to OIJ measure the depression approximately at the same time.

## 2.3 SuperDARN

SuperDARN stands for Super Dual Auroral Radar Network and this network consists of more than 30 low-power HF radars that look into Earth's upper atmosphere. (from mid-latitudes to polar regions) The radars can among others be used to measure the motion of charged particles in the ionosphere via the Doppler shift effect. (scattering of radiation) The motion of the plasma is interesting for us because we know what we expect from the orientation of the IMF is southward. The twin cell convection is therefore related to the dungey cycle. How can we visualize the motion of the hall currents? Unfortunately, we have not a complete data coverage all over the polar cap. Therefore, it seems to be useful to search for another way to visualize the motion of the plasma. The velocity of the hall currents is:  $\vec{v} = \frac{\vec{E} \times \vec{B}}{B^2}$  and the formula for the electric potential is:  $\vec{E} = -\vec{\nabla}\phi$ . It now turns out that lines of equal potential are perpendicular to the electric field, which is also the case for the hall currents. It can be shown that indeed the electric potential can describe the motion of the hall currents. The currents move then on equal potential lines. We use the data of the Radars (motion of plasma) to make a expansion in spherical harmonics for the electric potential. The plotting of the potential can then be used to describe the plasma flow.




Figure 8: data from Superdarn for different times

The plot of the electric field on the polar cap is shown in figure 8 for different times. All figure you can qualitatively see two regions. One region with a positive electric potential at 3 o'clock and one region with a negative electric potential at 9 o'clock. The orientation of there "cells" is qualitatively the same. (two cells are separated by a line, which goes from 12 o'clock to 6 o'clock) (look at figure 8) The strength of the electric potential variates a lot with time. We can compare the strength by comparing the contour lines of the potential.

From 18 o'clock to 21 o'clock, we have approximately the same strength of the electric potential. At this time, we also have approximately the same magnetic open flux through the polar cap.  $\phi_{pc} = 31 kV$  to  $\phi_{pc} = 41 kV$ . The size

of the polar cap stays the same in this time interval. We can also see that the area, where the electric potential is significantly different from 0 is fairly small and located at high latitudes. (above 70° north)(see figures 8a to 8d)

At 22 o'clock, we can see a huge difference to the previous time step. The strength of the electric potential increased. The area of significantly from 0 different electric potential increased too to about 60° latitude. The open magnetic flux through the polar cap is now  $\phi_{pc} = 76 kV$ . (see figure 8e)

At 23 o'clock on the 6th of January 2011, The area of significantly from 0 different electric potential reaches the maximum from all the shown time frames. (expanded almost to 50° latitude) The strength of the electric field stays approximately the same as well as the open magnetic flux through the polar cap. (see figure 8f)

At 00 o'clock on the 7th of January 2011, the properties described at 23 o'clock stay the same. However, we see on the top of figure 8g a small positive electric area with small electric field, which we couldn't observe at 23 o'clock.

At 01 o'clock, we observe a less stronger electric field than in the previous time step. We also observe some perturbations referred to the "two-cell-observation" from the previous steps.

## 2.4 AMPERE

AMPERE (Active Magnetosphere and Planetary Electrodynamics Response Experiment) is a earth observing system, which can be used to measure near-realtime magnetic field by 66 commercial satellites. The data from the magnetic field collected by AMPERE can be used to improve our understanding of space weather. In our project, we want to use AMPERE to determine the Birkeland currents. The Birkeland currents are field aligned currents. As described in the theory part, flowing electric currents cause a curl of the magnetic field. If there are such FAC, we should also see a curl in the magnetic field measured by the satellites. We don't want to have a look at the explicit magnetic field data, but directly at the simulated current flow based on the measured magnetic field of AMPERE. This means that we expect to see the Region 1 currents at high latitudes and the Region 2 currents at lower latitude. In addition to that, we are interested to see how the currents change with respect to time.




Figure 9: this figure shows the measured magnetic field vectors by the satellites at 23 o'clock of the 6th January and the field align currents in blue and red

In figure 9, we can see the measured magnetic field by the satellites visualized through little arrows. We convince ourselves that a curl in the magnetic field can be related to a area with blue or red colour, which belong to field aligned currents. Red colour means that the field aligned currents flow away from the surface of the earth and blue towards the earth.

With this pre-knowledge, we can now have a closer look at the data. From 18 o'clock (see figure 10a) until 20 o'clock(see figure 10c), we don't see significant currents. At 21 o'clock (see figure 10d), we see a small increase of the currents indicated by a bit more intensive red and blue colour. However, it is hard to derive a structure of the current-flow. At 22 o'clock (see figure 10e), the current flow is a bit more intense than one hour before. We can now see four different areas. A inner circle consisting of a current coming down from space to earth on the right half on the circle and of currents flowing into space the left part of the circle. The outer circle consists of a current coming down from space to earth on the left half on the circle and of currents flowing into space the right part of the circle. This basic structure can also be seen at 23 o'clock and at 00 o'clock.(see figure 10f and 10g) The currents reach their maximum at approximately 23 o'clock. From then on, the currents decrease. At 02 o'clock (see figure 10i), we can say that there aren't any significant currents any more.




Figure 10: Data from AMPERE at different times

## 2.5 Svalbard All-Sky Imager

The Svalbard All-Sky Imager measures the intensity of two different wavelength all over the sky. Every pixel represents a small part of the sky. We look at two different wavelength ( $\lambda_1 = 630\text{nm}$ ,  $\lambda_2 = 557.7\text{nm}$ ). The reason for this is that we want to have a look at the most important constituents of the northern lights.  $\lambda_1$  belongs to the relaxation from the 1D orbital to the 3p orbital of atomic oxygen and  $\lambda_2$  belongs to the relaxation from the 1s orbital to the 1D orbital. We have different images for each of this wavelength. (compare reference [6][p.469f] ) in order to visualize the time-development of the intensity of this light, it is convenient to use a so called keogram. A keogram takes a slice of each image at each time step and adds the different slices together in order to have a time evolution of exact one longitude. Longitudinal movements are easily to detect in keograms, however for the exact latitudinal movements one should check the full images. In the images, red areas represent relatively high intensity of the specific wavelength and black relatively low intensity. We should keep in mind that the intensities are rescaled for every picture in order to have a better visualization.

The figure 11 shows a general overview of the whole day. (06.01.2011) Figure 11a shows the overview of the whole day for  $\lambda_2$  and 11b shows the overview of the whole day for  $\lambda_1$ . The interval from 00 o'clock to 18 o'clock is not interesting for us. We can see that from 18 o'clock to approximately 22 o'clock, the intensity on both images are really low; whereas from 22 o'clock to 00 o'clock (7th January) we can see that we have high intensity over all longitudes

represented by the red colour.




Figure 11: Overview over the Svalbard-Imagers

The figure 11a, we can see keograms of each a hour of time. In figure 12a to 12d, we don't see anything special. The red area at latitudes below 20 degrees can be associated with trees. At 22:30 o'clock, we can see first a movement of a area of high intensity of light from north to south and at 22:45, we see a higher intensity of the light up to 76.5 degree latitude. (see figure 12e) At 23 o'clock, we can see a lot of areas with high light-intensity. The longest time interval of high light intensity all over the latitudes is at approximately 23:30 o'clock and lasts about 5-10 minutes. At 23:45 o'clock, the intensity is low and at 23:55, higher intensities are again measured at high latitudes.




Figure 12: Ketograms from the SvalbardImager for  $\lambda = 5577 \cdot 10^{-10} \text{m}$ , different times

The Svalbard imager for the wavelength of  $\lambda = 630\text{nm}$  shows qualitatively the same as the imager for  $\lambda = 577.7\text{nm}$ . This can be seen in figure 13. The figures 13a to 13f show each intervals of 1 hour.




Figure 13: Ketograms from the SvalbardImager for  $\lambda = 6300 \cdot 10^{-10}$  m, different times

### 3 discussion

In this section, we want to explain our observations. We will also try to connect our observation with what we expect from theory. It is also very interesting to see, whether the data from the different measurements can be interpreted in a sense-full, physical way.

The dunhey cycle, auroral sub storms and the twin-cell connection are all dependent on the IMF. Especially the sign of the  $B_z$ -component is really important because it determines, whether magnetic reconnection occurs or not. (compare 1.1) Therefore, it makes sense to look first at the data from ACE. If we assume that the solar wind needs on average 1 hour to travel from ACE to earth, we can say that the  $B_z$ -component is positive from 19 o'clock to 21:15 o'clock at earth's position. At approximately 21:15, the negative-turn of  $B_z$  should reach the earth. While the  $B_z$  component is negative, we expect that magnetic reconnection on day side kicks in. When magnetic reconnection on day side occurs, open magnetic flux is added to the polar cap. If magnetic reconnection on night side doesn't take place, the polar cap should grow. Southward orientation of the IMF is needed in order to have a so called groth phase of a sub storm. On average, we observed a southward orientation of the magnetic field until 01:45 o'clock of the 7th of January 2011. However, we can see a sudden peak in the  $B_z$  component at 22:20 o'clock at earth's place.

Next, we want to know whether we have any hints to a sub storm. A good method to get knowledge about whether a sub storm takes place or not is to look at the data from the ground-based-magnetometers. A sub storm leads in general to westward electro-jets, which lead to a magnetic field according to Amperes rule. In the end, if a sub storm kicks in, this leads to a depression of the south-north component of the magnetic field measured on earth. We now have a look at the data from 2.4. As described there, we can see a depression of the south-nord component of the magnetic field at approximately 22 o'clock. The depression reaches a minimum at approximately 22:30 o'clock. (depending on the station) This depression is a strong hint to a sub storm kicking in between 22 o'clock and 23 o'clock.

Furthermore, we want to have a look at the data from SuperDARN. (see chapter 2.3) We notice, that the shape of the electric potential fits to the theoretical predictions from theory. We can see two cells, which can be characterised through a region with negative electric potential (blue) and a region with positive electric potential (red). In the region with negative electric potential, the charged particles move clockwise and in the positive region the particles move anti-clockwise. Also the orientation of the two cells fits. The direction to the sun is in the figures 8 at 12 o'clock. (referring to a watch for defining direction with 12 o'clock on top) The twin cells should be separated by a line from approximately 6 o'clock to 12 o'clock.(directions) This depends also on the components of the IMF. The orientation of the twin cells seems to be right.

Looking at the time development of the electric field, we can now connect the increase of the electric potential from 21 to 22 o'clock with other physical phenomena. From section 2.1, we know that the negative turn of the  $B_z$  component of the IMF is at about 21:45 o'clock on earth. A southward IMF results in magnetic reconnection on day side. Magnetic reconnection on day side is directly couplet to the twin cell convection. Twin cell convection increases. This is exactly what we can see in the data. From the Groundbasedmagnetometer, we got a strong hint of an auroral sub storm kicking in between 22:00 o'clock and 23:00 o'clock. Sub storms decrease the total magnetic flux in the polar cap. We can see that from 22 o'clock to 23 o'clock, that  $\phi_{pc}$  stays approximately constant. All in all, we can say that the theory matches the experimental data concerning this measurement.

We want now to discuss the data from the Measurement AMPERE.(observations described in chapter 2.4) The curl of the magnetic field measured parallel to the surface of the earth determines the current flow perpendicular to the surface to the earth. The qualitatively development of the measurend currents from section 2.4 ca be explained through...

Finally, let us look at the Svalbard-imager. The first significant development in the imager can be seen at approximately 22:35 o'clock. A small brightened area moves equator-ward and westward and at 22:45 o'clock, we can see an increase of intensity of both wavelength for lower latitudes. This seems to be the first sub storm detected in this night. The time of the sub storm seems to be reasonable. In the time before the substorn, we know from the data from ACE that the  $B_z$  component on the IMF was negative from 22:15 o'clock on. In this time, energy could be stored in the magnetic tail of the earth by adding magnetic flux. From reference [6][p.419, Fig. 13.14.], we know that "sudden changes in the solar-wind dynamic pressure and, more frequently, northward turnings of the IMF can trigger the sudden unloading of stored energy". At approximately 22:20, the IMF- $B_z$ -component on earth turns suddenly northward and southward again. It is possible that this sudden change triggered the onset on the sub storm. In the time after 23:00 o'clock, we can see several timeslots of each 5-10 minutes, where we measure high intensity of lightening. One peaks around 23:15 o'clock, one peaks at 23:30 and the last one peaks about 0:00 o'clock. In between, we an see significant less intensity. This can be interpreted as several sub storms.

## 4 conclusion

## References

- [1] [http://www.windows2universe.org/earth/Magnetosphere/tour/tour\\_earth\\_magnetosphere\\_07.html&edu=high](http://www.windows2universe.org/earth/Magnetosphere/tour/tour_earth_magnetosphere_07.html&edu=high), 27.11.2015
- [2] [http://www.nasa.gov/mission\\_pages/sunearth/news/ace-15th.html](http://www.nasa.gov/mission_pages/sunearth/news/ace-15th.html), 27.11.2015
- [3] Gerd W. Prölss: Physics of the Earth's Space Environment (An Introduction). Bonn dec 2003
- [4] Sandholt, P.E., C.J. Farrugia, J. Moen, Ø. Noraberg, B. Lybekk, T. Sten, and T.L. Hansen, A classification of dayside auroral forms and activities as a function of IMF orientation, *J. Geophys. Res.*, 103, 23,325-23,345, 1998.
- [5] Ruohoniemi, J. M., R. A. Greenwald, Dependencies of high-latitude plasma convection: Consideration of interplanetary magnetic field, seasonal and universal time factors in statistical patterns, *J. Geophys. Res.*, 110, A09204, doi:10.1029/2004JA010815, 2005.
- [6] Kievelson and Russel, Introduction to Space Physics, Chapters 9, 13, and 14