

Space Physics Project , FYS3610

Sigve Harang, Lukas Powalla

November 17, 2015

Contents

1 Theory part	3
1.1 The Dungey cycle - a steady state picture	3
1.2 Auroral substorm	4
1.2.1 groth phase	4
1.2.2 Expansion phase	4
1.2.3 recovery phase	4
1.3 currents	4
1.4 Twin cell convection	5
1.5 Earth- Corotation and Dungey-cycle	5
2 Observations	5
2.1 ACE	5
2.2 AMPERE	5
2.3 Groundbasedmagnetometer	5
2.4 Superdarn	5
2.5 Svalbardimager	5
3 discussion	5
4 conclusion	5

Introduction


In this project we are going to analyse the ionospheric and auroral dynamics and its relation to the solar wind conditions. Furthermore, we will connect our observations to the current systems as well as the auroral activity and polar cap. We will investigate in a specific timeslot of 24 hours.

1 Theory part

In order to understand the movements and changes in the system of the earth and the sun, we want to introduce some basic concepts of space physics. A really important steady state picture of what is going on with respect to the earth's magnetic field, the interaction of the earth's magnetic field with the IMF and the consequences of the earth's environment (such as current flows or aurora) is the Dungey cycle. (only when IMF is southward orientated) However, the dungey cycle is as already mentioned a steady state picture. (This means for example that the magnetic flux added to the open magnetic field lines through day side reconnection is equal to the magnetic flux subtracting from the open magnetic field lines through night side reconnection) However, this cycle is in reality not a steady state picture. There are different phases of adding magnetic flux (growing of the polar cap), some kind of explosive expansion (subtraction of magnetic flux from the polar cap) and a recovery phase. This phenomena is called a "Auroral substorm". In addition to that, we can extract information about what is going on with the system earth-sun by looking at different data sets containing information about the magnetic field on earth, the solar wind bulk speed, the auroral intensity, the current density and further data from different physical measurements. Based on this measurements we will interpret the received data and relate it to the theoretical predictions and the models we introduced in the theoretical part.

1.1 The Dungey cycle - a steady state picture

Figure 1: Dungey cycle



If the magnetic field is northward orientated ($B_z > 0$) nothing happens at the boundary of the terrestrial magnetic field and the magnetic field of the sun/ the solar wind. IMF can not mix with the terrestrial magnetic field because everything is frozen in. Since the terrestrial magnetic field and the IMF are not anti-parallel orientated, magnetic reconnection does not occur and hence, everything is and stays frozen in. However, if the IMF is southward orientated ($B_z < 0$), we do have anti-parallel orientated magnetic field at the boundary of terrestrial magnetic field and the IMF (at the magnetopause). Therefore, magnetic reconnection occurs at the magnetopause, which causes a cyclic reconnection pattern at day/night-side. If we assume a steady state picture, we call the cycle, which happens for southward IMF the dungey cycle. The different parts of the cycle can be described as follows:

- 1 opposite polarity field lines reconnect at magnetopause (if IMF southward orientated)
- 2 magnetic field lines are still connected to the solar wind. (frozen in again) Therefore, the magnetic field line is draged along with the solar wind in the direction of the tale. (magnetic tension force)
- 3 magnetic flux is added to the tail and compresses the plasma sheet (magnetic pressure increases, plasma sheet gets thinner and thinner)

4 magnetic reconnection occurs in the tail.

5 reconnected magnetic field lines return to the day side (imagination of "rubber bands" help to understand this)

1.2 Auroral substorm

As already mentioned in the introduction, the assumption of a steady state picture is not an appropriate description of what happens in reality. To specify this, the rate of adding magnetic flux and the rate of subtracting magnetic flux from the polar cap are not equal. However, after studying data from different measurements, Akasofu could state out in 1964 different phases of a cycle so called:

- 1 groth phase
- 2 expansion phase
- 3 recovery phase

1.2.1 groth phase

The day side reconnection dominates and the aurora moves to lower latitudes. The auroral activity increases in the cusp and open magnetic flux is added to the polar cap. The magnetic flux, which is added due to the day side reconnection makes the tail radius increase.(because the field lines are draged tail-wards with the IMF)


1.2.2 Expansion phase

Sudden onset of night-side reconnection at the NENL. Sudden brightening of the night side aurora. Poleward and westward movement of the aurora (on average). It closes open magnetic flux.

1.2.3 recovery phase

The night side reconnection continues at smaller rates and the auroral display quietens. Furthermore, the polar cap contracts and the magnetosphere returns to pro groth phase "equilibrium".

Figure 2: Auroral substorm



1.3 currents

In general, we can express the current flow in a given system with given magnetic field and electric field as follows:

$$\vec{j} = en(\vec{v}_i - \vec{v}_e) = \sigma_p \vec{E}'_{\perp} + \sigma_H \frac{\vec{E}'_{\perp} \times \vec{B}}{B} + \sigma_{\parallel} \vec{E}_{\parallel} \quad (1)$$

Here, σ_p is the Pedersen conductivity, σ_H is the Hall conductivity and σ_{\parallel} is a conductivity, which belongs to a current parallel to the electric field. We notice that the different currents caused by different conductivities flow in orthogonal

directions with respect to each other. The orientation of \vec{E} means parallel or orthogonal to the magnetic field. A discrete expression of the conductivities will not be needed in this report.

$$\vec{E} = \vec{E}_\perp + \vec{E}_\parallel \quad (2)$$

$$\vec{E}'_\perp = \vec{E}_\perp + \vec{v}_n \times \vec{B} \quad (3)$$

$$\vec{E}'_\perp \approx \vec{E}_\perp \quad (4)$$

The Electron Hall currents flow at lower altitudes than the Pedersen currents. The ionospheric Pedersen currents are connected to the Field aligned currents (FAC). The FAC are on the one hand side closed with the magnetopause current and on the other hand side with the ring current. All current systems need to be closed, which is the case.

1.4 Twin cell convection

If we look at the Dungey cycle ($B_z < 0$), we can figure out the movement of the open/closed field lines at the polar caps, which gives us the movement of the particles at the polar cap. Let us first explain, what leads to the so called "Twin-cell"-convection. If we have southward IMF, the orientation of the IMF and the earth's magnetic field are opposite to each other. Like discussed in chapter 1.1, magnetic reconnection on the day side of the earth can occur. The magnetic reconnection at the day side adds open magnetic field lines to the polar cap. In other words, day side reconnection adds magnetic flux to the polar cap. The deformation of the usually round open/closed-field boundary leads to a movement, which tries to "smooth out" the little bump in the boundary. Magnetic field lines naturally want themselves to orientate in a straight line.

Let us now look at the night side. If we look at the night side events, we know that magnetic reconnection occurs in the tail.(dungey cycle) Reasons for this happening are for example that through day side reconnection, magnetic flux is added to the tail, the magnetic pressure increases and leads in the end to magnetic reconnection in the tail. The magnetic reconnection to the tail closes open magnetic field lines. This means that the polar cap shrinks a bit on the night side. In other words, magnetic flux is subtracted at the night side of the earth from the polar cap. The subtraction leads to a little dint in the polar cap. The magnetic field lines tend to "smooth out" the little dint, which leads to a corresponding movement.

In total, we expect now a movement, which is well known as the twin cell convection. The electric potential on the poles will give us the movement of the charged particles. The currents move along potential lines. The equipotential lines are therefore the trajectories of the charged particles. The currents are called "Hall"-currents. The hall-currents can be found in an altitude of approximately 110 km. (electro-jets)

1.5 Earth- Corotation and Dungey-cycle

2 Observations

2.1 ACE

2.2 AMPERE

2.3 Groundbasedmagnetometer

2.4 Superdarn

2.5 Svalbardimager

3 discussion

4 conclusion






Figure 3: Data from AMPERE at different times




(a) Groundbasedmagnetometer x- (b) Groundbasedmagnetometer y- (c) Groundbasedmagnetometer z- direction direction direction

Figure 4: Data from Groundbasedmagnetometer, different components



(a) Superdarn at 18:00 o'clock

Figure 5: data from Superdarn for different times



(a) SvalbardImager for 5577 Å

(b) SvalbardImager for 6300 Å

Figure 6: Overview over the Svalbard-Imagers




Figure 7: Ketograms from the SvalbardImager for $\lambda = 5577 \cdot 10^{-10}$ m, different times




Figure 8: Ketograms from the SvalbardImager for $\lambda = 6300 \cdot 10^{-10}$ m, different times