

Thesis

Lukas Prader

10th September 2023

Contents

1	Introduction	3
2	Methods	4
2.1	Datasets	4
2.2	Data preparation	4
2.3	Model building	4
2.4	Analysis	5
3	Background	5
3.1	Invasion theory	5
3.2	Harmonia axyridis	5

1 Introduction

(citations missing)

The topic of invasive species has become more and more important, even more so with recent changes in climate and habitats due to human influence. In order to deal with invasive species and the impact they can have on existing ecosystems, species distribution models (SDMs) have been used to predict the potential habitat and with that the threat of an emerging invasive species. Especially time-partitioned models can provide more insight into the process of invasion. Ensemble models are often used in applications where models are projected, since aspects of different modelling approaches can be combined to hopefully gain a more complete model of the species distribution.

There are certain challenges when using SDMs in invasive research though. For example, the invading species is usually not currently in equilibrium with its environment, leading to underestimations of the final niche occupied. A similar difficulty is modelling the invaded range with data from a species native range since the native niche can be quite different from the realized invaded niche. Nonetheless, SDMs are frequently used in trying to predict the spread of invasive species, which is why it is important to gain a better understanding of invasion processes and an SDM's ability to accurately represent it.

Harmonia axyridis, also known as the harlequin or Asian lady beetle, is an invasive species already established in non-native habitats all around the world. The invasion process has already been studied extensively and there have been SDMs modelling the invaded range in Europe as well, though mostly on a national scale.

One goal of this thesis is to look into the limitations of models built early in the invasion process of a species. By iterating over the years of the invasion, model performance can be evaluated with consideration to the current state of invasion. Computing the occupied niche separately for each year also provides more insight into the invasion process and might help to understand modelling limitations for certain years.

2 Methods

(citations missing)

2.1 Datasets

In order to also have some kind of influence of the human nature of first introduction, land cover data was used from the Copernicus Land cover Classification dataset with yearly resolution starting from 2002 up to 2020.

All traditional 19 bioclim variables were obtained from the CHELSA V2.1 climatologies datasets too, using the 1981-2010 time frame for all years from 2002 to 2010 as well as the MPI-ESM 1.2 ssp370 scenario 2011-2040 for all years from 2011 to 2022. For occurrence data, all global occurrences of *Harmonia axyridis* were downloaded from the GBIF database.

2.2 Data preparation

All bioclim and land cover layers were resampled to a matching resolution of 30 arc seconds and cropped to two spatial extents, Europe and the presumed native range referencing (Orlova-Bienkowskaja, Ukrainsky & Brown, 2015). The presence-only points from GBIF were subset to the afore mentioned spatial extents and then checked for missing values for latitude, longitude, year or coordinate uncertainty. No occurrences after 2022 were used, also no points with a coordinate uncertainty larger than 1 km. In Europe, the initial cut off year for presences was 1991, since this is the year of invasion according to the EASIN website. Afterwards, using the library ‘CoordinateCleaner’, all remaining data points were again checked for common errors or biases in the respective subset. To prepare the data for modelling, pseudoabsences were generated for each year. For this, the subset of a specific year was taken and used to generate pseudo-absences limited to a radius around the occurrence points. For each presence point, a buffer circle was generated with a maximum distance of 18 km. This buffer circle also excluded any area closer than 1 km to any presence point of any year to limit contradiction in environmental space.

2.3 Model building

For each year, the following Models were computed: General Linear (GLM), General Additive (GAM), Boosted Regression Trees (BRT) and Maximum Entropy (MAXENT). A model for a specific year always included all points from past years as well. The iterative models that were built use only data points from Europe, though there was one model created only with native occurrences and predicted for each year in Europe. For all used occurrence points prior to 2002 and after 2020, the land cover data of 2002 or 2020 was used as a substitute.

Variance inflation factors were used to select the variables used for model building. For this, a GLM was computed only using Europe data from 2002.

This model included all 19 bioclim variables and their squared values as separate variables. For land cover variables, a PCA was computed from the 2002 data. PCA axes were included in the model until a cut off of 90% of explained variance was reached. Variance inflation factors were computed for this GLM and the variable with the highest VIF was dropped until none of the remaining variables had a VIF greater than 10. Quadratic versions of bioclim variables were always dropped before their linear counterparts, even if their VIF was lower.

2.4 Analysis

All SDM models of each year were tested for their accuracy on predicting the occurrences of the following year and the final year of 2022. For each year, the occupied niche was also computed by running a PCA analysis on the bioclim variables. The niche can then be visualized by plotting a dynamic occurrence density grid for the first two PCA axes (Broennimann et al. 2011).

3 Background

This section aims to give sufficient background on *Harmonia axyridis* and the theory necessary to understand this thesis.

3.1 Invasion theory

basic theory of how invasions work and maybe a bit on sdms in that area

3.2 *Harmonia axyridis*

Harmonia axyridis, also known as the Harlequin ladybird or multicoloured Asian lady beetle, is of the family of the Coccinellidae and has its native origin in Asia [1]. At first widely introduced as a control species against pest aphids, *H. axyridis* has turned out to be a highly invasive species reaching an almost global distribution [2]. In America, the species was introduced as early as 1916 (California) and in 1988, first populations outside intended release were found [3]. Usage of *H. axyridis* for biological control in Europe dates back as far as 1990 (France) [4]. First invasive occurrences were confirmed in multiple countries during the early 2000s, including Germany (2000), Belgium (2001), the Netherlands (2002) and the United Kingdom (2003) [1]. The first confirmation in Austria, where it was never used for biological control, was 2006 [5]. It has been shown that all established invasive populations outside of North America have their origin in the first established population in eastern North America, with the European populations being significantly influenced by the used biocontrol strain [6].

The impact of *H. axyridis* on invaded areas is diverse. In some contexts, the ladybird has been shown to have a negative impact on the diversity and abundance of native ladybird species [1]. Many studies show intra guild predation and direct interspecific competition in favour of *H. axyridis* [7]. It has also been shown that the species feeds on a variety of damaged fruit crops, for

example grapes, apples, stone fruit and berry crops [8]. The aggregating behaviour of *H. axyridis*, mostly as a strategy for overwintering, is also a cause of disturbance, since private homes and facilities are invaded by large amounts of beetles at a given time [9].

References

- [1] H. E. Roy et al. “The harlequin ladybird, *Harmonia axyridis*: global perspectives on invasion history and ecology”. In: *Biological invasions* 18 (2016), pp. 997–1044. DOI: 10.1007/s10530-016-1077-6.
- [2] P. Brown et al. “*Harmonia axyridis* in Europe: spread and distribution of a non-native coccinellid”. In: *From biological control to invasion: the ladybird *Harmonia axyridis* as a model species* (2008), pp. 5–21. DOI: 10.1007/978-1-4020-6939-0_2.
- [3] J. B. Chapin, V. Brou et al. “*Harmonia axyridis* (Pallas), the third species of the genus to be found in the United States (Coleoptera: Coccinellidae)”. In: *Proc. Entomol. Soc. Wash* 93.3 (1991), pp. 630–635. URL: <https://www.biodiversitylibrary.org/partpdf/55539>.
- [4] J. Coutanceau. “*Harmonia axyridis* (Pallas, 1773): Une coccinelle asiatique introduite, acclimatée et en extension en France”. In: *Bulletin de la Société Entomologique de France* 111 (Jan. 2006), pp. 395–401. DOI: 10.3406/bsef.2006.16343.
- [5] W. Rabitsch and R. Schuh. “First record of the multicoloured Asian ladybird *Harmonia axyridis* (Pallas, 1773) in Austria”. In: *Beiträge zur Entomofaunistik* 7 (2006), pp. 161–164. URL: https://www.zobodat.at/pdf/BEEF_7_0161-0164.pdf.
- [6] E. Lombaert et al. “Bridgehead effect in the worldwide invasion of the bio-control harlequin ladybird”. In: *PloS one* 5.3 (2010), e9743. DOI: 10.1371/journal.pone.0009743.
- [7] J. K. Pell et al. “Intraguild predation involving *Harmonia axyridis*: a review of current knowledge and future perspectives”. In: *BioControl* 53 (2008), pp. 147–168. DOI: 10.1007/978-1-4020-6939-0_10.
- [8] R. Koch et al. “Phytophagous preferences of the multicolored Asian lady beetle (Coleoptera: Coccinellidae) for autumn-ripening fruit”. In: *Journal of Economic Entomology* 97.2 (2004), pp. 539–544. DOI: 10.1093/jee/97.2.539.
- [9] C. Nalepa, G. Kennedy and C. Brownie. “Role of visual contrast in the alighting behavior of *Harmonia axyridis* (Coleoptera: Coccinellidae) at overwintering sites”. In: *Environmental Entomology* 34.2 (2005), pp. 425–431. DOI: 10.1603/0046-225X-34.2.425.

Appendix