



name: <unnamed>
 log: C:\Users\chopikwi\Dropbox\PushButtonReproduction.smcl
 log type: smcl
 opened on: 22 Jun 2022, 17:41:09

```
1 . do "C:\Users\chopikwi\Downloads\nyhan-reifler-causal-corrections.do"
2 . *****
3 . *Replication code for *
4 . *Displacing Misinformation about Events: *
5 . *An Experimental Test of Causal Corrections *
6 . *Brendan Nyhan and Jason Reifler *
7 . *Forthcoming, JEPS *
8 . *****
9 .
10 . clear all
11 .
12 . cd "C:\Users\chopikwi\Downloads"
    C:\Users\chopikwi\Downloads
13 .
14 . use "causal-replication.dta", clear
15 .
16 . /*Table 2 - note lincom results correspond to significance test table in append
    > ix*/
17 . reg swensenfav innuendo denial causal [aweight=aw]
    (sum of wgt is 3.9441e+03)
```

Source	SS	df	MS	Number of obs =	987
Model	73.4211912	3	24.4737304	F(3, 983) =	19.67
Residual	1222.83522	983	1.24398293	Prob > F =	0.0000
				R-squared =	0.0566
				Adj R-squared =	0.0538
				Root MSE =	1.1153
Total	1296.25642	986	1.31466168		

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
innuendo	-.5839232	.100686	-5.80	0.000	-.7815073 -.3863391
denial	-.3894682	.1007865	-3.86	0.000	-.5872497 -.1916866
causal	.0753454	.1010347	0.75	0.456	-.1229231 .2736139
_cons	2.752434	.071879	38.29	0.000	2.61138 2.893488

```
18 .
19 . lincom denial-innuendo

( 1)  - innuendo + denial = 0
```

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
(1)	.194455	.0998118	1.95	0.052	-.0014138 .3903238

20 . lincom causal-innuendo

(1) - innuendo + causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.6592686	.1000624	6.59	0.000	.4629081	.8556291

21 . lincom causal-denial

(1) - denial + causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.4648136	.1001636	4.64	0.000	.2682544	.6613727

22 .

23 . svy: reg swensenfav innuendo denial causal
(running regress on estimation sample)

Survey: Linear regression

Number of strata	=	1	Number of obs	=	987
Number of PSUs	=	987	Population size	=	980.27809
			Design df	=	986
			F(3, 984)	=	3.98
			Prob > F	=	0.0078
			R-squared	=	0.0307

swensenfav	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
innuendo	-.5228951	.1731961	-3.02	0.003	-.8627703	-.1830198
denial	-.4097696	.1490654	-2.75	0.006	-.7022916	-.1172476
causal	-.1983303	.1578649	-1.26	0.209	-.50812	.1114594
_cons	2.855356	.1207248	23.65	0.000	2.618449	3.092263

24 .

25 . lincom denial-innuendo

(1) - innuendo + denial = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.1131255	.1518829	0.74	0.457	-.1849255	.4111765

26 . lincom causal-innuendo

(1) - innuendo + causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.3245648	.160528	2.02	0.043	.009549	.6395806

27 . lincom causal-denial

(1) - denial + causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.2114393	.1341375	1.58	0.115	-.0517885	.4746671

28 .
29 . reg acceptedbribes innuendo denial causal [aweight=aw]
(sum of wgt is 3.9414e+03)

Source	SS	df	MS	Number of obs = 986		
Model	106.101485	3	35.3671616	F(3, 982) =	31.72	
Residual	1094.80676	982	1.1148745	Prob > F =	0.0000	
				R-squared =	0.0884	
				Adj R-squared =	0.0856	
Total	1200.90824	985	1.21919619	Root MSE =	1.0559	

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
innuendo	.7845069	.095376	8.23	0.000	.5973426	.9716712
denial	.5506573	.095568	5.76	0.000	.3631162	.7381984
causal	.0693211	.0954347	0.73	0.468	-.1179583	.2566005
_cons	2.86127	.0679831	42.09	0.000	2.727861	2.994679

30 .
31 . lincom denial-innuendo

(1) - innuendo + denial = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.2338497	.0947969	-2.47	0.014	-.4198774	-.0478219

32 . lincom causal-innuendo

(1) - innuendo + causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.7151858	.0946624	-7.56	0.000	-.9009497	-.5294219

33 . lincom causal-denial

(1) - denial + causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.4813362	.0948559	-5.07	0.000	-.6674797	-.2951926

```
34 .
35 . svy: reg acceptedbribes innuendo denial causal
    (running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	1	Number of obs	=	986
Number of PSUs	=	986	Population size	=	985.14713
			Design df	=	985
			F(3, 983)	=	8.14
			Prob > F	=	0.0000
			R-squared	=	0.0607

acceptedbr~s	Linearized		t	P> t	[95% Conf. Interval]	
	Coef.	Std. Err.				
innuendo	.668764	.1573979	4.25	0.000	.3598904	.9776377
denial	.5179593	.1477819	3.50	0.000	.2279557	.8079629
causal	.1596657	.1491155	1.07	0.285	-.1329549	.4522864
_cons	2.942776	.1094129	26.90	0.000	2.728067	3.157485

```
36 .
37 . lincom denial-innuendo
```

(1) - innuendo + denial = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.1508047	.1505696	-1.00	0.317	-.4462788	.1446694

```
38 . lincom causal-innuendo
```

(1) - innuendo + causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.5090983	.1518787	-3.35	0.001	-.8071414	-.2110552

```
39 . lincom causal-denial
```

(1) - denial + causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.3582936	.1418893	-2.53	0.012	-.6367336	-.0798536

```
40 .
```

```
41 . reg resigninvest denial causal [aweight=aw]
    (sum of wgt is 2.9802e+03)
```

Source	SS	df	MS	Number of obs = 764		
Model	43.3852407	2	21.6926204	F(2, 761) =	38.95	
Residual	423.8789	761	.557002497	Prob > F =	0.0000	
				R-squared =	0.0928	
				Adj R-squared =	0.0905	
Total	467.264141	763	.612403854	Root MSE =	.74633	

resigninvest	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
denial	-.2460515	.0661224	-3.72	0.000	-.3758555	-.1162475
causal	-.5812725	.0661139	-8.79	0.000	-.7110598	-.4514852
_cons	3.455409	.0467082	73.98	0.000	3.363717	3.547101

```
42 .
43 . lincom causal-denial
```

(1) - **denial** + **causal** = 0

resigninvest	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.335221	.0661809	-5.07	0.000	-.4651398	-.2053021

```
44 .
45 . svy: reg resigninvest denial causal
    (running regress on estimation sample)
```

Survey: Linear regression

Number of strata =	1	Number of obs =	764
Number of PSUs =	764	Population size =	756.43179
		Design df =	763
		F(2, 762) =	7.18
		Prob > F =	0.0008
		R-squared =	0.0521

resigninvest	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
denial	-.2582214	.0976405	-2.64	0.008	-.4498974	-.0665455
causal	-.4205997	.113207	-3.72	0.000	-.6428339	-.1983655
_cons	3.402598	.0772646	44.04	0.000	3.250922	3.554274

```
46 .
47 . lincom causal-denial
```

(1) - **denial** + **causal** = 0

resigninvest	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.1623783	.1020287	-1.59	0.112	-.3626686	.0379121

```

48 .
49 . /*Figures 1-3*/
50 .
51 . preserve

52 .
53 . collapse (mean) swensenfav resigninvest acceptedbribes (semean) swensenfavse=sw
> ensenfav resigninvestse=resigninvest acceptedbribesse=acceptedbribes [aweight=a
> w],by(swensenrand)

54 .
55 . foreach var of varlist swensenfav resigninvest acceptedbribes {
    2. gen `var'1l=`var'-(1.96*`var'se)
    3. gen `var'ul=`var'+(1.96*`var'se)
    4. }
    (1 missing value generated)
    (1 missing value generated)

56 .
57 . twoway (scatter swensenfav swensenrand, ytitle("") mcolor(black) graphregion(fc
> olor(white) ifcolor(none)) plotregion(fcolor(none) lcolor(white) ifcolor(none)
> ilcolor(none)) scheme(s2mono) yscale(r(1 3.25)) xlabel(1 `"' "No reason" "given
> " "' 2 `"' "Innuendo" "only" "' 3 `"' "Innuendo +" "allegation denial" "' 4 `"'
> "Innuendo +" "denial + causal" "' ,labsize(*.8)) ylabel(1 `"' "Very" "unfavorab
> le" "' 2 `"' "Somewhat" "unfavorable" "' 3 `"' "Slightly" "unfavorable" "' ,nogr
> id labsize(*.8) angle(0)) yscale(r(1(1)3)) xscale(r(.75 4.25)) xtick(none) xmti
> ck(none) xtitle("Reason for resignation",margin(medsmall) size(*.8))) (rspike s
> wensenfavul swensenfav1l swensenrand,lcolor(black) legend(off))

58 . graph export "fig1.eps",replace
    (note: file fig1.eps not found)
    (file fig1.eps written in EPS format)

59 .
60 . twoway (scatter acceptedbribes swensenrand, ytitle("") mcolor(black) graphregi
> on(fcolor(white) ifcolor(none)) plotregion(fcolor(none) lcolor(white) ifcolor(n
> one) ilcolor(none)) scheme(s2mono) yscale(r(1 3.25)) xlabel(1 `"' "No reason" "
> given" "' 2 `"' "Innuendo" "only" "' 3 `"' "Innuendo +" "allegation denial" "'
> 4 `"' "Innuendo +" "denial + causal" "' ,labsize(*.8)) ylabel(1 `"' "Not at all"
> "likely" "' 2 `"' "A little" "likely" "' 3 `"' "Moderately" "likely" "' 4 `"' "
> Very" "likely" "' ,nogrid labsize(*.8) angle(0)) yscale(r(1(1)3)) xscale(r(.75
> 4.25)) xtick(none) xmtick(none) xtitle("Reason for resignation",margin(medsmall
> ) size(*.8))) (rspike acceptedbribesul acceptedbribes1l swensenrand,lcolor(blac
> k) legend(off))

61 . graph export "fig2.eps",replace
    (note: file fig2.eps not found)
    (file fig2.eps written in EPS format)

62 .
63 . drop if swensenrand==1
    (1 observation deleted)

64 .
65 . twoway (scatter resigninvest swensenrand, ytitle("") mcolor(black) graphregion(
> fcolor(white) ifcolor(none)) plotregion(fcolor(none) lcolor(white) ifcolor(none)
> ) ilcolor(none)) scheme(s2mono) yscale(r(1 3.25)) xlabel(2 `"' "Innuendo" "only
> " "' 3 `"' "Innuendo +" "allegation denial" "' 4 `"' "Innuendo +" "denial + cau
> sal" "' ,labsize(*.8)) ylabel(1 `"' "Not likely" "at all" "' 2 `"' "Not very" "l
> ikely" "' 3 `"' "Somewhat" "likely" "' 4 `"' "Very" "likely" "' ,nogrid labsize(
> *.8) angle(0)) yscale(r(1(1)3)) xscale(r(1.75 4.25)) xtick(none) xmtick(none) x
> title("Reason for resignation",margin(medsmall) size(*.8))) (rspike resigninves
> tul resigninvest1l swensenrand,lcolor(black) legend(off))

```

```

66 . graph export "fig3.eps",replace
    (note: file fig3.eps not found)
    (file fig3.eps written in EPS format)

67 .
68 . restore

69 .
70 . /*Appendix*/
71 .
72 . /*balance tests - unweighted*/
73 . /*note the putexcel command requires Stata 13*/
74 . putexcel B1=("Control") C1=("Innuendo") D1=("Denial") E1=("Causal") F1=("Total"
    > ) using jepsbalance, replace
    file jepsbalance.xlsx saved

75 .
76 . gen fakewt=1 /*fake survey weights*/

77 . svyset [pweight=fakewt] /*more convenient for output of proportions*/

    pweight: fakewt
    VCE: linearized
    Single unit: missing
    Strata 1: <one>
    SU 1: <observations>
    FPC 1: <zero>

78 .
79 . /*estimated - don't have exact birthdate*/
80 . gen age=2012-pp_birthy

81 .
82 . gen agecat4=.
    (1000 missing values generated)

83 . replace agecat4=1 if age>=18 & age<=29
    (83 real changes made)

84 . replace agecat4=2 if age>=30 & age<=44
    (141 real changes made)

85 . replace agecat4=3 if age>=45 & age<=59
    (440 real changes made)

86 . replace agecat4=4 if age>=60 & age!=.
    (336 real changes made)

87 .
88 . gen control=(swensenrand==1)

89 .
90 . /*age marginals*/
91 . putexcel A22=("Age") A23=("18-29") A24=("30-44") A25=("45-59") A26=("60+") usin
    > g jepsbalance, modify
    file jepsbalance.xlsx saved

```

```
92 .
93 . tab agecat4 swensenrand, chi
```

agecat4	swensenrand				Total
	1	2	3	4	
1	27	14	21	21	83
2	34	37	27	43	141
3	101	99	114	126	440
4	69	95	84	88	336
Total	231	245	246	278	1,000

Pearson chi2(9) = 12.0650 Pr = 0.210

```
94 . svy: tab agecat4 swensenrand
(running tabulate on estimation sample)
```

```
Number of strata   =          1          Number of obs       =       1000
Number of PSUs     =       1000          Population size      =       1000
                                   Design df          =         999
```

agecat4	swensenrand				Total
	1	2	3	4	
1	.027	.014	.021	.021	.083
2	.034	.037	.027	.043	.141
3	.101	.099	.114	.126	.44
4	.069	.095	.084	.088	.336
Total	.231	.245	.246	.278	1

Key: cell proportions

```
Pearson:
Uncorrected chi2( 9) = 12.0650
Design-based F( 9, 8991) = 1.3392 P = 0.2105
```

```
95 .
96 . svy, subpop(control): tab agecat4
(running tabulate on estimation sample)
```

```
Number of strata   =          1          Number of obs       =       1000
Number of PSUs     =       1000          Population size      =       1000
                                   Subpop. no. of obs    =        231
                                   Subpop. size           =        231
                                   Design df              =         999
```

agecat4	proportions
1	.1169
2	.1472
3	.4372
4	.2987
Total	1

Key: proportions = cell proportions

```

97 . matrix AgeControl = e(b) '
98 . putexcel B23=matrix(AgeControl) using jepsbalance, modify
   file jepsbalance.xlsx saved
99 .
100 . svy,subpop(innuendo): tab agecat4
    (running tabulate on estimation sample)

```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	245
			Subpop. size	=	245
			Design df	=	999

agecat4	proportions
1	.0571
2	.151
3	.4041
4	.3878
Total	1

Key: proportions = **cell proportions**

```

101 . matrix AgeDanger = e(b) '
102 . putexcel C23=matrix(AgeDanger) using jepsbalance, modify
    file jepsbalance.xlsx saved
103 .
104 . svy,subpop(denial): tab agecat4
    (running tabulate on estimation sample)

```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	246
			Subpop. size	=	246
			Design df	=	999

agecat4	proportions
1	.0854
2	.1098
3	.4634
4	.3415
Total	1

Key: proportions = **cell proportions**

```

105 . matrix AgeCorrection = e(b) '

```

```
106 . putexcel D23=matrix(AgeCorrection) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
107 .
108 . svy, subpop(causal): tab agecat4
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	278
			Subpop. size	=	278
			Design df	=	999

agecat4	proportions
1	.0755
2	.1547
3	.4532
4	.3165
Total	1

Key: proportions = **cell proportions**

```
109 . matrix AgeCorrection2 = e(b) '
```

```
110 . putexcel E23=matrix(AgeCorrection2) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
111 .
112 . svy: tab agecat4
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Design df	=	999

agecat4	proportions
1	.083
2	.141
3	.44
4	.336
Total	1

Key: proportions = **cell proportions**

```
113 . matrix AgeTotal = e(b) '
```

```
114 . putexcel F23=matrix(AgeTotal) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
115 .
116 . /*gender marginals*/
117 . putexcel A4=("Gender") A5=("Male") A6=("Female") using jepsbalance, modify
file jepsbalance.xlsx saved
```

```
118 .
119 . tab pp_gender swensenrand, chi
```

Profile - Gender	swensenrand				Total
	1	2	3	4	
Male	119	104	112	131	466
Female	112	141	134	147	534
Total	231	245	246	278	1,000

Pearson chi2(3) = **4.0831** Pr = **0.253**

```
120 . svy: tab pp_gender swensenrand
(running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Design df	=	999

Profile - Gender	swensenrand				Total
	1	2	3	4	
Male	.119	.104	.112	.131	.466
Female	.112	.141	.134	.147	.534
Total	.231	.245	.246	.278	1

Key: **cell proportions**

Pearson:
 Uncorrected chi2(3) = **4.0831**
 Design-based F(3.00, 2997.00)= **1.3597** P = **0.2533**

```
121 .
122 . svy, subpop(control): tab pp_gender
(running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	231
			Subpop. size	=	231
			Design df	=	999

Profile - Gender	proportions
Male	.5152
Female	.4848
Total	1

Key: proportions = **cell proportions**

```

123 . matrix GenderControl = e(b) '
124 . putexcel B5=matrix(GenderControl) using jepsbalance, modify
    file jepsbalance.xlsx saved
125 .
126 . svy,subpop(innuendo): tab pp_gender
    (running tabulate on estimation sample)

```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	245
			Subpop. size	=	245
			Design df	=	999

Profile - Gender	proportions
Male	.4245
Female	.5755
Total	1

Key: proportions = cell proportions

```

127 . matrix GenderDanger = e(b) '
128 . putexcel C5=matrix(GenderDanger) using jepsbalance, modify
    file jepsbalance.xlsx saved
129 .
130 . svy,subpop(denial): tab pp_gender
    (running tabulate on estimation sample)

```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	246
			Subpop. size	=	246
			Design df	=	999

Profile - Gender	proportions
Male	.4553
Female	.5447
Total	1

Key: proportions = cell proportions

```

131 . matrix GenderCorrection = e(b) '
132 . putexcel D5=matrix(GenderCorrection) using jepsbalance, modify
    file jepsbalance.xlsx saved

```

```
133 .
134 . svy, subpop(causal): tab pp_gender
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	278
			Subpop. size	=	278
			Design df	=	999

Profile - Gender	proportions
Male	.4712
Female	.5288
Total	1

Key: proportions = **cell proportions**

```
135 . matrix GenderCorrection2 = e(b) '
136 . putexcel E5=matrix(GenderCorrection2) using jepsbalance, modify
    file jepsbalance.xlsx saved
137 .
138 . svy: tab pp_gender
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Design df	=	999

Profile - Gender	proportions
Male	.466
Female	.534
Total	1

Key: proportions = **cell proportions**

```
139 . matrix GenderTotal = e(b) '
140 . putexcel F5=matrix(GenderTotal) using jepsbalance, modify
    file jepsbalance.xlsx saved
141 .
142 . /*education marginals*/
143 . putexcel A8=("Education") A9=("High School or less") A10=("Some college") A11=(
    > "College grad") A12=("Post-grad") using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```

144 .
145 . gen neweduc=pp_educ

146 . recode neweduc (2=1) (3=2) (4=2) (5=3) (6=4)
      (neweduc: 975 changes made)

147 .
148 . tab pp_educ swensenrand, chi

```

Profile - Education	swensenrand				Total
	1	2	3	4	
No HS	7	6	8	4	25
High school graduate	68	78	72	78	296
Some college	61	56	65	71	253
2-year	20	26	20	22	88
4-year	53	58	50	60	221
Post-grad	22	21	31	43	117
Total	231	245	246	278	1,000

Pearson chi2(15) = 12.0933 Pr = 0.672

```

149 . svy: tab pp_educ swensenrand
      (running tabulate on estimation sample)

```

```

Number of strata   =          1          Number of obs       =       1000
Number of PSUs     =       1000          Population size      =       1000
                                   Design df          =          999

```

Profile - Education	swensenrand				Total
	1	2	3	4	
No HS	.007	.006	.008	.004	.025
High sch	.068	.078	.072	.078	.296
Some col	.061	.056	.065	.071	.253
2-year	.02	.026	.02	.022	.088
4-year	.053	.058	.05	.06	.221
Post-gra	.022	.021	.031	.043	.117
Total	.231	.245	.246	.278	1

Key: cell proportions

```

Pearson:
  Uncorrected chi2( 15) = 12.0933
  Design-based F( 15.00, 14985.00)= 0.8054 P = 0.6728

```

```

150 .
151 . tab neweduc swensenrand, chi

```

neweduc	swensenrand				Total
	1	2	3	4	
1	75	84	80	82	321
2	81	82	85	93	341
3	53	58	50	60	221
4	22	21	31	43	117
Total	231	245	246	278	1,000

Pearson chi2(9) = 8.3770 Pr = 0.497

```
152 . svy: tab neweduc swensenrand
      (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Design df	=	999

neweduc	swensenrand				
	1	2	3	4	Total
1	.075	.084	.08	.082	.321
2	.081	.082	.085	.093	.341
3	.053	.058	.05	.06	.221
4	.022	.021	.031	.043	.117
Total	.231	.245	.246	.278	1

Key: **cell proportions**

Pearson:

Uncorrected	chi2(9)	=	8.3770	
Design-based	F(9, 8991)	=	0.9299	P = 0.4975

```
153 .  
154 . svy, subpop(control): tab neweduc  
(running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	231
			Subpop. size	=	231
			Design df	=	999

neweduc	proportions
1	.3247
2	.3506
3	.2294
4	.0952
Total	1

Key: proportions = cell proportions

```
155 . matrix EducControl = e(b)'
```

```
156 . putexcel B9=matrix(EducControl) using jepsbalance, modify
file jepsbalance.xlsx saved
```

```
157 .
158 . svy, subpop(innuendo): tab neweduc
(running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	245
			Subpop. size	=	245
			Design df	=	999

neweduc	proportions
1	.3429
2	.3347
3	.2367
4	.0857
Total	1

Key: proportions = cell proportions

159 . matrix EducDanger = e(b) '

160 . putexcel C9=matrix(EducDanger) using jepsbalance, modify
file jepsbalance.xlsx saved

161 .

162 . svy,subpop(denial): tab neweduc
(running tabulate on estimation sample)

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	246
			Subpop. size	=	246
			Design df	=	999

neweduc	proportions
1	.3252
2	.3455
3	.2033
4	.126
Total	1

Key: proportions = cell proportions

163 . matrix EducCorrection = e(b) '

164 . putexcel D9=matrix(EducCorrection) using jepsbalance, modify
file jepsbalance.xlsx saved

165 .

166 . svy,subpop(causal): tab neweduc
(running tabulate on estimation sample)

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	278
			Subpop. size	=	278
			Design df	=	999

neweduc	proportions
1	.295
2	.3345
3	.2158
4	.1547
Total	1

Key: proportions = cell proportions

```
167 . matrix EducCorrection2 = e(b) '
168 . putexcel E9=matrix(EducCorrection2) using jepsbalance, modify
    file jepsbalance.xlsx saved
169 .
170 . svy: tab neweduc
    (running tabulate on estimation sample)
```

```
Number of strata   =           1           Number of obs       =       1000
Number of PSUs     =       1000           Population size      =       1000
                                           Design df           =        999
```

neweduc	proportions
1	.321
2	.341
3	.221
4	.117
Total	1

Key: proportions = **cell proportions**

```
171 . matrix EducTotal = e(b) '
172 . putexcel F9=matrix(EducTotal) using jepsbalance, modify
    file jepsbalance.xlsx saved
173 .
174 . /*race marginals*/
175 . gen newrace=pp_race
176 . recode newrace (5=4) (6=4) (7=4)
    (newrace: 61 changes made)
177 .
178 . putexcel A14=("Race/ethnicity") A15=("White") A16=("Black") A17=("Hispanic") A1
    > 8=("Other") using jepsbalance, modify
    file jepsbalance.xlsx saved
179 .
180 . tab pp_race swensenrand, chi
```

Profile - Race	swensenrand				Total
	1	2	3	4	
White	174	191	180	209	754
Black	25	23	23	31	102
Hispanic	15	18	20	15	68
Asian	3	3	4	5	15
Native American	2	4	4	3	13
Mixed	8	2	5	8	23
Other	4	4	10	7	25
Total	231	245	246	278	1,000

Pearson chi2(18) = 11.7337 Pr = 0.861

```
181 . svy: tab pp_race swensenrand
(running tabulate on estimation sample)
```

```
Number of strata   =           1           Number of obs   =       1000
Number of PSUs     =       1000           Population size   =       1000
                                   Design df       =           999
```

Profile - Race	swensenrand				Total
	1	2	3	4	
White	.174	.191	.18	.209	.754
Black	.025	.023	.023	.031	.102
Hispanic	.015	.018	.02	.015	.068
Asian	.003	.003	.004	.005	.015
Native A	.002	.004	.004	.003	.013
Mixed	.008	.002	.005	.008	.023
Other	.004	.004	.01	.007	.025
Total	.231	.245	.246	.278	1

Key: **cell proportions**

```
Pearson:
Uncorrected  chi2( 18)           =    11.7337
Design-based F( 18.00, 17982.00)=    0.6512   P = 0.8612
```

```
182 .
183 . tab newrace swensenrand, chi
```

newrace	swensenrand				Total
	1	2	3	4	
1	174	191	180	209	754
2	25	23	23	31	102
3	15	18	20	15	68
4	17	13	23	23	76
Total	231	245	246	278	1,000

```
Pearson chi2( 9) =    5.5006   Pr = 0.789
```

```
184 . svy: tab newrace swensenrand
(running tabulate on estimation sample)
```

```
Number of strata   =           1           Number of obs   =       1000
Number of PSUs     =       1000           Population size   =       1000
                                   Design df       =           999
```

newrace	swensenrand				Total
	1	2	3	4	
1	.174	.191	.18	.209	.754
2	.025	.023	.023	.031	.102
3	.015	.018	.02	.015	.068
4	.017	.013	.023	.023	.076
Total	.231	.245	.246	.278	1

Key: **cell proportions**

```
Pearson:
Uncorrected  chi2( 9)           =    5.5006
Design-based F( 9.00, 8991.00)=    0.6106   P = 0.7891
```

```
185 .
186 . svy, subpop(control): tab newrace
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	231
			Subpop. size	=	231
			Design df	=	999

newrace	proportions
1	.7532
2	.1082
3	.0649
4	.0736
Total	1

Key: proportions = **cell proportions**

```
187 . matrix RaceControl = e(b) '
```

```
188 . putexcel B15=matrix(RaceControl) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
189 .
190 . svy, subpop(innuendo): tab newrace
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	245
			Subpop. size	=	245
			Design df	=	999

newrace	proportions
1	.7796
2	.0939
3	.0735
4	.0531
Total	1

Key: proportions = **cell proportions**

```
191 . matrix RaceDanger = e(b) '
```

```
192 . putexcel C15=matrix(RaceDanger) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
193 .
```

```
194 . svy, subpop(denial): tab newrace
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	246
			Subpop. size	=	246
			Design df	=	999

newrace	proportions
1	.7317
2	.0935
3	.0813
4	.0935
Total	1

Key: proportions = cell proportions

```
195 . matrix RaceCorrection = e(b) '
```

```
196 . putexcel D15=matrix(RaceCorrection) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
197 .
```

```
198 . svy, subpop(causal): tab newrace
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Subpop. no. of obs	=	278
			Subpop. size	=	278
			Design df	=	999

newrace	proportions
1	.7518
2	.1115
3	.054
4	.0827
Total	1

Key: proportions = cell proportions

```
199 . matrix RaceCorrection2 = e(b) '
```

```
200 . putexcel E15=matrix(RaceCorrection2) using jepsbalance, modify
    file jepsbalance.xlsx saved
```

```
201 .
```

```
202 . svy: tab newrace
    (running tabulate on estimation sample)
```

Number of strata	=	1	Number of obs	=	1000
Number of PSUs	=	1000	Population size	=	1000
			Design df	=	999

newrace	proportions
1	.754
2	.102
3	.068
4	.076
Total	1

Key: proportions = cell proportions

203 . matrix RaceTotal = e(b)'

204 . putexcel F15=matrix(RaceTotal) using jepsbalance, modify
file jepsbalance.xlsx saved

205 .

206 . /*ordered probit results*/

207 .

208 . oprobit swensenfav innuendo denial causal [aweight=aw]

(sum of wgt is 3.9441e+03)

Iteration 0: log likelihood = -1474.7046

Iteration 1: log likelihood = -1445.26

Iteration 2: log likelihood = -1445.2554

Iteration 3: log likelihood = -1445.2554

Ordered probit regression

Number of obs = 987

LR chi2(3) = 58.90

Prob > chi2 = 0.0000

Pseudo R2 = 0.0200

Log likelihood = -1445.2554

swensenfav	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
innuendo	-.574654	.0960842	-5.98	0.000	-.7629756	-.3863325
denial	-.3841301	.0952463	-4.03	0.000	-.5708094	-.1974507
causal	.0459583	.094494	0.49	0.627	-.1392464	.2311631
/cut1	-1.06395	.0763423			-1.213578	-.9143214
/cut2	-.2181641	.0708799			-.3570862	-.079242
/cut3	.7512049	.0733655			.6074111	.8949987
/cut4	1.401776	.0862095			1.232809	1.570744
/cut5	2.094072	.1271028			1.844955	2.343189

209 .

210 . lincom denial-innuendo

(1) - [swensenfav]innuendo + [swensenfav]denial = 0

swensenfav	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	.190524	.0952539	2.00	0.045	.0038297	.3772182

211 . lincom causal-innuendo

(1) - [swensenfav]innuendo + [swensenfav]causal = 0

swensenfav	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	.6206124	.095704	6.48	0.000	.4330359	.8081888

212 . lincom causal-denial

(1) - [swensenfav]denial + [swensenfav]causal = 0

swensenfav	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	.4300884	.094851	4.53	0.000	.2441838	.615993

213 .

214 . svy: oprobit swensenfav innuendo denial causal
(running oprobit on estimation sample)

Survey: Ordered probit regression

Number of strata	=	1	Number of obs	=	987
Number of PSUs	=	987	Population size	=	987
			Design df	=	986
			F(3, 984)	=	18.06
			Prob > F	=	0.0000

swensenfav	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
innuendo	-.5711644	.0943605	-6.05	0.000	-.7563349	-.385994
denial	-.3798074	.0932185	-4.07	0.000	-.562737	-.1968779
causal	.0312641	.0899189	0.35	0.728	-.1451904	.2077185
/cut1	-1.053315	.0725627	-14.52	0.000	-1.19571	-.9109203
/cut2	-.2173536	.0666615	-3.26	0.001	-.3481684	-.0865388
/cut3	.7531182	.0678114	11.11	0.000	.6200469	.8861896
/cut4	1.396552	.0761784	18.33	0.000	1.247061	1.546042
/cut5	2.07787	.1179546	17.62	0.000	1.846399	2.309341

215 .

216 . lincom denial-innuendo

(1) - [swensenfav]innuendo + [swensenfav]denial = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.191357	.0984453	1.94	0.052	-.0018294	.3845435

217 . lincom causal-innuendo

(1) - [swensenfav]innuendo + [swensenfav]causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.6024285	.0984385	6.12	0.000	.4092554	.7956016

218 . lincom causal-denial

(1) - [swensenfav]denial + [swensenfav]causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.4110715	.0970218	4.24	0.000	.2206785	.6014645

219 .

220 . oprobit acceptedbribes innuendo denial causal [aweight=aw]

(sum of wgt is 3.9414e+03)
Iteration 0: log likelihood = -1458.9113
Iteration 1: log likelihood = -1414.8777
Iteration 2: log likelihood = -1414.8611
Iteration 3: log likelihood = -1414.8611

Ordered probit regression

Number of obs = 986
LR chi2(3) = 88.10
Prob > chi2 = 0.0000
Pseudo R2 = 0.0302

Log likelihood = -1414.8611

acceptedbribes	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
innuendo	.776435	.0964726	8.05	0.000	.5873521	.9655179
denial	.5364651	.0955094	5.62	0.000	.3492702	.7236601
causal	.0836056	.0946095	0.88	0.377	-.1018256	.2690369
/cut1	-1.19551	.0825283			-1.357262	-1.033757
/cut2	-.2949245	.0715228			-.4351066	-.1547424
/cut3	.5082236	.072069			.366971	.6494762
/cut4	1.622064	.0835497			1.458309	1.785818

221 .

222 . lincom denial-innuendo

(1) - [acceptedbribes]innuendo + [acceptedbribes]denial = 0

acceptedbr~s	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	-.2399698	.0946655	-2.53	0.011	-.4255108	-.0544289

223 . lincom causal-innuendo

(1) - [acceptedbribes]innuendo + [acceptedbribes]causal = 0

acceptedbr~s	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	-.6928294	.0956554	-7.24	0.000	-.8803105	-.5053482

224 . lincom causal-denial

(1) - [acceptedbribes]denial + [acceptedbribes]causal = 0

acceptedbr~s	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	-.4528595	.0948091	-4.78	0.000	-.638682	-.2670371

225 .

226 . svy: oprobit acceptedbribes innuendo denial causal
(running oprobit on estimation sample)

Survey: Ordered probit regression

Number of strata	=	1	Number of obs	=	986
Number of PSUs	=	986	Population size	=	986
			Design df	=	985
			F(3 , 983)	=	28.77
			Prob > F	=	0.0000

acceptedbribes	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
innuendo	.7742062	.0941625	8.22	0.000	.589424	.9589883
denial	.5339922	.0932866	5.72	0.000	.3509289	.7170555
causal	.099282	.0974736	1.02	0.309	-.0919978	.2905617
/cut1	-1.184781	.0835541	-14.18	0.000	-1.348745	-1.020817
/cut2	-.2872679	.0717806	-4.00	0.000	-.4281283	-.1464075
/cut3	.512992	.0717101	7.15	0.000	.3722699	.6537141
/cut4	1.61116	.0836572	19.26	0.000	1.446994	1.775327

227 .

228 . lincom denial-innuendo

(1) - [acceptedbribes]innuendo + [acceptedbribes]denial = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.240214	.0899578	-2.67	0.008	-.4167449	-.063683

229 . lincom causal-innuendo

(1) - [acceptedbribes]innuendo + [acceptedbribes]causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.6749242	.0974239	-6.93	0.000	-.8661064	-.483742

230 . lincom causal-denial

(1) - [acceptedbribes]denial + [acceptedbribes]causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.4347102	.0960961	-4.52	0.000	-.6232869	-.2461336

231 .

232 . oprobit resigninvest denial causal [aweight=aw]

(sum of wgt is 2.9802e+03)
Iteration 0: log likelihood = **-844.04756**
Iteration 1: log likelihood = **-808.44627**
Iteration 2: log likelihood = **-808.40597**
Iteration 3: log likelihood = **-808.40597**

Ordered probit regression

Number of obs = **764**
LR chi2(2) = **71.28**
Prob > chi2 = **0.0000**
Pseudo R2 = **0.0422**

Log likelihood = **-808.40597**

resigninvest	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
denial	-.4051632	.1011984	-4.00	0.000	-.6035084	-.206818
causal	-.8529667	.1020236	-8.36	0.000	-1.052929	-.6530041
/cut1	-2.358818	.1158121			-2.585806	-2.131831
/cut2	-1.460528	.0872103			-1.631457	-1.289598
/cut3	-.0842087	.0742481			-.2297323	.0613149

233 .

234 . lincom causal-denial

(1) - [resigninvest]denial + [resigninvest]causal = 0

resigninvest	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
(1)	-.4478035	.0983262	-4.55	0.000	-.6405192	-.2550877

```
235 .
236 . svy: oprobit resigninvest denial causal
      (running oprobit on estimation sample)
```

Survey: Ordered probit regression

```
Number of strata   =           1           Number of obs       =          764
Number of PSUs     =          764           Population size      =          764
                                           Design df           =          763
                                           F(           2,       762) =          33.76
                                           Prob > F             =          0.0000
```

resigninvest	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
denial	-.4031662	.0980295	-4.11	0.000	-.5956058	-.2107267
causal	-.8420202	.1024705	-8.22	0.000	-1.043178	-.6408627
/cut1	-2.343823	.1128273	-20.77	0.000	-2.565311	-2.122334
/cut2	-1.448461	.0834922	-17.35	0.000	-1.612363	-1.28456
/cut3	-.0864461	.0738371	-1.17	0.242	-.2313941	.0585019

```
237 .
238 . lincom causal-denial
```

(1) - [resigninvest]denial + [resigninvest]causal = 0

resigninvest	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.438854	.0972905	-4.51	0.000	-.6298429	-.2478651

```
239 .
240 . /*unweighted results*/
241 .
242 . reg swensenfav innuendo denial causal
```

Source	SS	df	MS	Number of obs =	987
Model	71.2144156	3	23.7381385	F(3, 983) =	18.90
Residual	1234.87677	983	1.25623273	Prob > F =	0.0000
				R-squared =	0.0545
				Adj R-squared =	0.0516
Total	1306.09119	986	1.32463609	Root MSE =	1.1208

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
innuendo	-.581532	.1037343	-5.61	0.000	-.7850982	-.3779658
denial	-.3856136	.1037343	-3.72	0.000	-.5891798	-.1820474
causal	.0613401	.1010847	0.61	0.544	-.1370266	.2597067
_cons	2.748879	.0750555	36.62	0.000	2.601591	2.896166

243 .
244 . lincom denial-innuendo

(1) - innuendo + denial = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.1959184	.1012668	1.93	0.053	-.0028056	.3946423

245 . lincom causal-innuendo

(1) - innuendo + causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.642872	.0985509	6.52	0.000	.4494778	.8362663

246 . lincom causal-denial

(1) - denial + causal = 0

swensenfav	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	.4469537	.0985509	4.54	0.000	.2535594	.6403479

247 .
248 . reg acceptedbribes innuendo denial causal

Source	SS	df	MS	Number of obs = 986		
Model	102.994993	3	34.3316645	F(3, 982) =	30.45	
Residual	1107.12671	982	1.12742028	Prob > F =	0.0000	
Total	1210.1217	985	1.22854995	R-squared =	0.0851	
				Adj R-squared =	0.0823	
				Root MSE =	1.0618	

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
innuendo	.7869404	.098368	8.00	0.000	.5939046	.9799761
denial	.5509052	.0984646	5.59	0.000	.3576799	.7441305
causal	.0855267	.0956062	0.89	0.371	-.1020893	.2731428
_cons	2.856502	.0711034	40.17	0.000	2.71697	2.996034

249 .
250 . lincom denial-innuendo

(1) - innuendo + denial = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.2360352	.0962297	-2.45	0.014	-.4248748	-.0471956

251 . lincom causal-innuendo

(1) - innuendo + causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.7014136	.0933029	-7.52	0.000	-.8845096	-.5183176

252 . lincom causal-denial

(1) - denial + causal = 0

acceptedbr~s	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.4653784	.0934047	-4.98	0.000	-.6486743	-.2820826

253 .

254 . reg resigninvest denial causal

Source	SS	df	MS	Number of obs = 764		
Model	43.8991173	2	21.9495587	F(2, 761) =	38.61	
Residual	432.655856	761	.568535948	Prob > F =	0.0000	
				R-squared =	0.0921	
				Adj R-squared =	0.0897	
Total	476.554974	763	.624580569	Root MSE =	.75401	

resigninvest	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
denial	-.2459016	.0682652	-3.60	0.000	-.379912	-.1118913
causal	-.5781064	.0662569	-8.73	0.000	-.7081745	-.4480384
_cons	3.454918	.0482708	71.57	0.000	3.360158	3.549678

255 .

256 . lincom causal-denial

(1) - denial + causal = 0

resigninvest	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.3322048	.0662569	-5.01	0.000	-.4622728	-.2021368

257 .

258 . /*significance tests: see lincom results for Table 2 above*/

259 .

260 . /*test for recall diffs*/

261 . gen recall=(GSU213==3)+(GSU214==2)+(GSU215==3)

```

262 .
263 . preserve

264 .
265 . drop if swensenrand==1
    (231 observations deleted)

266 .
267 . reg recall denial causal [aweight=aw]
    (sum of wgt is 3.0000e+03)

```

Source	SS	df	MS	Number of obs =	769
Model	1.44331156	2	.721655779	F(2, 766) =	0.99
Residual	560.533629	766	.73176714	Prob > F =	0.3735
				R-squared =	0.0026
				Adj R-squared =	-0.0000
Total	561.976941	768	.731740809	Root MSE =	.85543

recall	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
denial	.0941114	.0755612	1.25	0.213	-.0542202	.242443
causal	.0045917	.0755612	0.06	0.952	-.1437399	.1529233
_cons	2.212666	.0534298	41.41	0.000	2.10778	2.317553

```

268 .
269 . lincom causal-denial

    ( 1)  - denial + causal = 0

```

recall	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.0895197	.0755612	-1.18	0.236	-.2378513	.0588119

```

270 .
271 . svy: reg recall denial causal
    (running regress on estimation sample)

```

Survey: Linear regression

Number of strata	=	1	Number of obs	=	769
Number of PSUs	=	769	Population size	=	769
			Design df	=	768
			F(2, 767)	=	0.98
			Prob > F	=	0.3748
			R-squared	=	0.0024

recall	Linearized Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
denial	.0926332	.0750609	1.23	0.218	-.0547157	.239982
causal	.0035824	.0761647	0.05	0.962	-.1459332	.1530981
_cons	2.212245	.0537147	41.19	0.000	2.1068	2.31769

```
272 .
273 . lincom causal-denial
```

```
( 1)  - denial + causal = 0
```

recall	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
(1)	-.0890507	.0752639	-1.18	0.237	-.236798	.0586966

```
274 .
275 . restore

276 .
    end of do-file

277 . save "causal-replication.dta", replace
    file causal-replication.dta saved

278 . exit, clear
```