

Homework 1

CSC 445-01: Theory of Computation

Matthew Mabrey, Luke Kurlandski

February 8, 2021

Question I

Question II

Given a relation on the set of all nonempty subsets of N tell whether it is reflexive, symmetric, or transitive.

a

R is defined by: ARB if and only if $A \subseteq B$

- Reflexive because the relation over any given element subset relates every element to itself. In our case $X \subseteq X$ holds true since a set will always be a subset of itself.
- Not symmetric because an element X being a subset of the element Y only guarantees that Y contains all of X 's elements but not vice versa. So every $XRY \neq YRX$.
- Transitive because an element X can only be a subset of Y if it contains all of X 's elements. Then for Y to be a subset of Z it must contain all of Y 's elements. Thus guaranteeing that all of X 's elements are contained within Z therefore making $X \subseteq Z$ and the relation $XRY \wedge YRZ \rightarrow XRZ$ hold true.

b

R is defined by: ARB if and only if $A \cap B \neq \emptyset$

- Reflexive because $X \cap X$ will never be an empty set since the parent set only contains nonempty subsets of N and so X intersecting with itself will always contain at least one element.
- Symmetric because the relation in this case does not care about order since if the intersection of two sets $X \cap Y$ results in a nonempty set then $Y \cap X$ will also yield the same result set.
- Not transitive because while $X \cap Y$ may result in a nonempty set based on some subset of shared elements, the intersection $Y \cap Z$ could result in a nonempty set based on a different subset of shared elements. Therefore the transitive relation $XRY \wedge YRZ \rightarrow XRZ$ would not always hold true.

c

R is defined by: ARB if and only if $1 \in A \cap B$

- Reflexive because if the element X contains 1 then the intersection X with itself will always contain 1.
- Symmetric because the order of the sets in the intersection $X \cap Y$ doesn't matter and if the result set contains 1 then the intersection $Y \cap X$ set will also contain 1.
- Transitive because if the result set of $X \cap Y$ contains 1 and the result set of $Y \cap Z$ also contains 1 then all 3 sets must contain 1 and so $X \cap Z$ will also contain one and the transitive relation holds true.

Question III

We will prove

$$P(n): \sum_{i=1}^n \frac{1}{i(i+1)} = \frac{n}{n+1} \quad \left| \quad n \geq 0, n \in \mathbb{Z} \right.$$

Base case:

$$\begin{aligned} P(0) &: \sum_{i=1}^0 \frac{1}{i(i+1)} = \frac{n}{n+1} \\ &: \sum_{i=1}^0 \frac{1}{i(i+1)} = \frac{0}{0+1} \\ &: 0 = 0 \\ &: \text{TRUE} \end{aligned}$$

Inductive Hypothesis:

$$P(k): \sum_{i=1}^k \frac{1}{i(i+1)} = \frac{k}{k+1} \quad \left| \quad k \geq 0, k \in \mathbb{Z} \right.$$

Induction:

$$\begin{aligned} P(k+1): \frac{k+1}{k+2} &= \sum_{i=1}^{k+1} \frac{1}{i(i+1)} \\ &= \sum_{i=1}^k \frac{1}{i(i+1)} + \frac{1}{(k+1)((k+1)+1)} \\ &= \frac{k}{k+1} + \frac{1}{(k+1)(k+2)} \\ &= \frac{k(k+2)+1}{(k+1)(k+2)} \\ &= \frac{(k+1)^2}{(k+1)(k+2)} \\ \frac{k+1}{(k+2)} &= \frac{k+1}{(k+2)} \\ P(k+1): &\text{TRUE} \end{aligned}$$

Conclusion:

$$\left(P(0) \wedge \left(P(k) \rightarrow P(k+1) \right) \right) \rightarrow P(n)$$

Question IV

Truth Table

p	q	$(p \rightarrow q) \wedge (p \rightarrow \neg q)$	$p \wedge (p \rightarrow q)$	$(p \rightarrow q) \wedge (\neg p \rightarrow q)$	$p \iff (p \iff q)$	$q \wedge (p \rightarrow q)$
T	T	F	T	T	T	T
T	F	F	F	F	F	F
F	T	T	F	T	F	T
F	F	T	F	F	F	F

Equivalence

<i>Problem</i>	<i>Original</i>	<i>Simplified</i>
a	$(p \rightarrow q) \wedge (p \rightarrow \neg q)$	$\neg p$
b	$p \wedge (p \rightarrow q)$	$p \wedge q$
c	$(p \rightarrow q) \wedge (\neg p \rightarrow q)$	q
d	$p \iff (p \iff q)$	$p \wedge q$
e	$q \wedge (p \rightarrow q)$	q

Question V

Given $P = \emptyset, Q = \{\epsilon\}, R = \{aba, bcb, cac\}$, and $S = \{aba, abc, abb\}$

0.1 1

$P \subset Q$ is correct because \emptyset is a proper subset of all sets (besides itself) since all elements contained in P are contained within Q , while $P \neq Q$.

0.2 2

$R \cap S = \{aba\}$ because \cap is the intersection of both sets and the only element both sets share is aba .

0.3 3

$R \triangle S = \{bcb, cac, abc, abb\}$ because \triangle is the disjoint union of both sets. In other words, the set of elements from both sets that are not shared between them. Thus the resulting set is every element besides the shared element aba from the previous problem.

0.4 4

$||R|| = 3$ because the cardinality of a set is the number of elements contained within the set.

Question VI

Given $P = (x \rightarrow y) \wedge (y \rightarrow z)$ where x, y , and z are boolean variables

0.5 1

The value of P is false when $x, z = \text{true}$ and $y = \text{false}$ because the first implication $(x \rightarrow y)$ in the *AND* statement evaluates to false. This is because the hypothesis in the implication is true while the conclusion is false, which is the only case where the implication will evaluate to false.

0.6 2

The value of P is again false when $y = \text{true}$ and $x, z = \text{false}$ because the second implication $(y \rightarrow z)$ in the *AND* statement evaluates to false. The same reasoning as above applies where the hypothesis in the implication is true, while the conclusion is false, causing the implication to be false.

Question VII

0.7 1


Let the 4-state finite automaton that accepts the words over $\{a, b\}$ that end with aba be formally defined by

$$M = (\{q_0, q_1, q_2, q_3\}, \{a, b\}, \delta, q_0, \{q_3\}),$$

where δ is given by

	a	b
q_0	q_1	q_0
q_1	q_1	q_2
q_2	q_3	q_0
q_3	q_1	q_2

The state diagram then constructed from this definition is thus



0.8 2

Let the 6-state finite automaton that accepts the words over $\{0, 1\}$ that contain either 000 or 111 be formally defined by

$$M = (\{q_0, q_1, q_2, q_3, q_4, q_5\}, \{0, 1\}, \delta, q_0, \{q_5\}),$$

where δ is given by

	0	1
q_0	q_1	q_0
q_1	q_1	q_2
q_2	q_3	q_0
q_3	q_1	q_2
q_4	q_1	q_2
q_5	q_1	q_2

The state diagram then constructed from this definition is thus

