

Hydrologic modeling at Luke

"Process models and academic curiosity
for sustainable use of boreal ecosystems"

- Samuli Launiainen, Antti-Jussi Kieloaho, Kersti Haahti, Mingfu Guan, Aura Salmivaara

Krycklan model work



© Natural Resources Institute Finland





Why carbon sink increases at Hyytiälä Fluxnet site?



- Two decades of eddy-covariance data at Hyytiälä SMEAR II –site
- Increasing annual CO₂ sink that is mostly due to enhanced productivity
 - NEE: from -180 to ~-200 g C m⁻² a⁻¹ (+53%)
 - GPP: from ~1000 to 1200 g C m⁻² a⁻¹ (+19%)
- Also annual ecosystem ET has increased (+22%)
- Plausible explanations for increased productivity?
 - CO₂ fertilization
 - Increasing T & longer growing season
 - Stand development
 - Footprint changes



APES: Atmosphere-Plant Exchange simulator



APES:

- Multi-layer, multi-species SVAT-model
- CO₂-H₂O-energy
- Sub-daily forcing
- Scenarios: base, CO₂+, LAI+, CO₂ & LAI+

Launiainen et al. 2015. Ecol. Mod; 2016 Global Change Biol.
3


Geophysical Research Abstracts
Vol. 20, EGU2018-18561, 2018
EGU General Assembly 2018
© Author(s) 2018. CC Attribution 4.0 license.



Does atmospheric CO₂ explain increased carbon sink at a boreal coniferous forest flux site?

Tiiia Grönholm (1), Samuli Launiainen (2), Gabriel Katul (3), Pasi Kolari (1), Toprak Aslan (1), Ivan Mamarella (1), Timo Vesala (1), and Pertti Hari (1)
(1) University of Helsinki, Institute for Atmosphere and Earth System Research, Finland (tiiia.gronholm@helsinki.fi), (2) Natural Resources Institute Finland, Finland, (3) Nicholas School of the Environment, Duke University, Durham, NC, USA

Climate impacts of bryophytes: from functional traits to global models



- Pinpoint the most important functional traits of the bryophyte layer?
 - Hydrological properties
 - Optical and thermal properties
 - Carbon usage
- Start with APES + develop description further bryophyte layer
 - Energy and water balance
 - Photosynthesis, respiration, growth
- Put mosses into a climate model
 - Orchidee?
 - Regional climate impacts



Antti-Jussi Kieloaho

Krycklan model workshop, 24.1.18 Uppsala


© Natural Resources Institute Finland

Response of drained peatland forest to management alternatives?




Kersti Haahti


Lettosuo Fluxnet site (FMI)



Measurements show water table level increase after treatments



Can modeling pinpoint reasons? What is a sufficient model description?



EC, chambers , WT depth, biomass

Coupled APES, 1-D Richards + Hooghoudt drainage equation

Validate for 5-year control, perturb system to replicate experiment

- Pinpoint reasons why changes took place
- Inter/extrapolate to other conditions



Kersti Haahti

Why water table, ET and CO₂ balance change?


Approach:

Couple APES, 1-D Richards
+ Hooghoud drainage
equation





Validate for 5-year control
data (WT, ET from EC)

Alter stand structure to
replicate experiment

- Pinpoint reasons why changes took place
- ... and perform model-based inter/extrapolation for other conditions



Can we upscale from point to landscape?



Working point
scale model

"simplifications where
possible"

Open GIS-data
meteorologic data
streamflow data


Distributed ET,
snow & soil water
predictions,
streamflow

SpatHy: (semi)distributed catchment model

INPUT:

- Multi-NFI: conifer vs. deciduous LAI
- Soil maps, peatland mask
- DEM
 - Catchment mask
 - Topographic wetness index TWI
 - For TOPMODEL – water budget
- Forcing by daily weather data
 - Precip, Rglob, Tair, RH

Catchment → (n x m) grid of buckets



GRID-OUTPUTS:


- ET components
- Root zone moisture
- Snow water equivalent
- Drainage to ground water
- Saturation deficit (→ WT)
- saturated gridcells, returnflow

CATCHMENT LEVEL

- Streamflow

Launiainen et al. 2018 GMD, in prep.

+ TOPMODEL for streamflow generation & saturated area



Upscaling of ET (P-M –based)






Figure 9. Annual total ET and its components: transpiration Tr, interception evaporation E and soil/ forest floor evaporation Ef as a fraction of annual P. Top row shows maximum annual LAI and the relative share of deciduous LAI / total LAI



Krycklan model workshop, 24.1.18 Uppsala

© Natural Resources Institute Finland


Can we predict partitioning of annual water budgets without calibration?



- Points → from measurements
- Red line + region = model + parameter uncertainty
- Model uncertainties: vegetation type, plant water use efficiency, interception efficiency, snow, peatland evaporation, open fjells, ...
- Measurement uncertainties:
 - Boundaries
 - Discharge
 - Precipitation from 10x10km FMI grid data



Example results

Saturation deficit during snowmelt





Blue ==
'wet'

During dry period



Saturated gridcells

24.1.201



NutSpatHy


- A grid-based distributed nutrient model for predicting nitrogen (N) and phosphorous (P) leaching in forested catchments
- Model key features:
 - Grid-based nutrient balance ($\Delta\text{storage} = \text{release} - \text{uptake}$)
 - Nutrient concentration prediction ($\text{conc} = \text{storage}/\text{water_volume}$)
 - Export loading simulation based on SpatHy runoff prediction
 - Spatiotemporal modelling of nutrient dynamics



Mingfu Guan





Ari Lauren





- N and P concentration**

Nutrient concentration is reasonably predicted considering the complex chemical process and a series of uncertainties of parameters and input data.



- Daily and monthly N and P export loading**

Good performance in the prediction of export loading verifies the controlling role of hydrology on nutrient transport processes.





Daily N and P export loading: NSC = 0.76, correlation coef.= 0.81


Dynamic forest trafficability prediction by fusion of open data, hydrologic forecasts and harvester-measured data




Aura Salmivaara



Spatial hydrology model simulations
at 16*16m spatial and daily temporal resolution
using weather observations and forecasts



Open spatial data




Forest machine measured data for improving prediction models


Rut depth
prediction
 $f(t, x, y)$

Salmivaara, A., Launiainen, S., Ala-Illomäki, J., Kulju, S., Laurén, A., Sirén, M., Tuominen, S., Finér, L., Uusitalo, J., Nevalainen, P., Pahikkala, T., and Heikkonen, J. 2017. *Dynamic forest trafficability prediction by fusion of open data, hydrologic forecasts and harvester-measured data*. Poster.

Dynamic forest trafficability prediction by fusion of open data, hydrologic forecasts and harvester-measured data



Predicted rut depth
in "dry" conditions (19.11.2016)
[cm]



Predicted rut depth
in "moist" conditions (18.5.2016)
[cm]



In "moist" conditions the area with ruts >10 cm covers 45% of the total area, while in "dry" conditions this proportion is 26%.

Summary

- We try to use mechanistic models to:
 - 1) Explain observations
 - 2) Offer ways to simplify problems
 - 3) Stimulate thinking
 - 4) Benefit practitioners?
 - 5) Find unpleasant truths (from data ...and models)
- We:
 - 1) Do ecohydrology – link water, carbon and energy, in future also biogeochemistry
 - 2) Aim to go spatial by merging GIS and mechanistic models
 - 3) Need to strengthen landscape approach (all interesting happens between local and global scale... which is often ignored)
- **4) Want to work with you and Krycklan data!**
- **Tandem Forest values call (DL end of March)**

Thank you!

Krycklan model workshop, 24.1.18 Uppsala



