

Question 7

Solve the following questions from the Discrete Math zyBook:

Exercise 6.1.5, sections b-d

A 5-card hand is dealt from a perfectly shuffled deck of playing cards. What is the probability of each of the following events?

$$\text{b. } \frac{C(13,1) * C(4,3) * C(12,2) * C(4,1) * C(4,1)}{C(52,5)} = \frac{54912}{2,598,960} = \frac{88}{4165}$$

$$\text{c. } \frac{C(4,1) * C(13,5)}{C(52,5)} = \frac{4 * 1287}{C(52,5)} = \frac{5148}{2598960} = \frac{33}{16660}$$

$$\text{d. } \frac{C(13,1) * C(4,2) * C(12,3) * C(4,1) * C(4,1) * C(4,1)}{C(52,5)} = \frac{1098240}{2598960} = \frac{352}{833}$$

Exercise 6.2.4, sections a-d

A 5-card hand is dealt from a perfectly shuffled deck of playing cards. What is the probability of each of the following events?

$$\text{a. } 1 - \frac{C(39,5)}{C(52,5)} = \frac{575757}{2598960} = \frac{2109}{9520}$$

$$\text{b. } 1 - \frac{C(13,5) * 4^5}{C(52,5)}$$

$$\frac{C(13,5) * 4^5}{C(52,5)} = \frac{1317888}{2598960}$$

$$1 - \frac{1317888}{2598960} = \frac{2112}{4165}$$

$$\begin{aligned} \text{c. } & \frac{C(13,1) * C(39,4)}{C(52,5)} + \frac{C(13,1) * C(39,4)}{C(52,5)} - \frac{C(13,1) * C(13,1) * C(26,3)}{C(52,5)} \\ &= \frac{65351}{99960} \end{aligned}$$

$$\text{d. } 1 - \frac{C(26,5)}{C(52,5)}$$

$$\frac{C(26,5)}{C(52,5)} = \frac{65780}{2598960} = \frac{253}{9996}$$

$$1 - \frac{253}{9996} = \frac{9743}{9996}$$

Question 8

Solve the following questions from the Discrete Math zyBook:

Exercise 6.3.2, sections a-e

$$\text{a. } p(A) = \frac{6!}{7!} = \frac{1}{7}$$

$$p(B) = \frac{7!}{2!} = \frac{1}{2}$$

$$p(C) = \frac{5!}{7!} = \frac{1}{42}$$

$$\text{b. } p(A|C) = \frac{|A \cap C|}{|C|} = \frac{2! * 3!}{5!} = \frac{1}{10}$$

$$\text{c. } p(B|C) = \frac{|B \cap C|}{|C|} = \frac{\frac{5!}{2}}{5!} = \frac{1}{2}$$

$$\text{d. } p(A|B) = \frac{|A \cap B|}{|B|} = \frac{3 * 5!}{\frac{7!}{2!}} = \frac{1}{7}$$

e. A and C are not independent, because $P(A|C) \neq P(A)$

B and C are independent, because $P(B|C) = P(B) = \frac{1}{2}$

A and B are independent, because $P(A|B) = P(A) = \frac{1}{7}$

Exercise 6.3.6, sections b, c

$$\text{b. First 5 heads} = \frac{1}{3} * \frac{1}{3} * \frac{1}{3} * \frac{1}{3} * \frac{1}{3} = \left(\frac{1}{3}\right)^5$$

$$\text{Last 5 Tails} = \frac{2}{3} * \frac{2}{3} * \frac{2}{3} * \frac{2}{3} * \frac{2}{3} = \left(\frac{2}{3}\right)^5$$

$$\left(\frac{2}{3}\right)^5 * \left(\frac{1}{3}\right)^5 = \frac{32}{59049}$$

$$\text{c. } \frac{1}{3} * \left(\frac{2}{3}\right)^9 = \frac{512}{59049}$$

Exercise 6.4.2, section a

a. A = *The dice chosen is fair*

\bar{A} = *The biased dice is chosen*

$$P(A) = P(\bar{A}) = \frac{1}{2}$$

B = *the dice came up 4, 3, 6, 6, 5, 5*

$$P(B|\bar{A}) = \left(\frac{3}{20}\right)^4 * \left(\frac{1}{4}\right)^2$$

$$P(B|A) = \left(\frac{1}{6}\right)^6$$

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\bar{A})P(\bar{A})}$$

$$= \frac{\left(\frac{1}{6}\right)^6 \frac{1}{2}}{\left(\frac{1}{6}\right)^6 * \frac{1}{2} * \left(\frac{3}{20}\right)^4 \left(\frac{1}{4}\right)^2 \left(\frac{1}{2}\right)} = \frac{40000}{99049} = 0.404$$

Question 9:

Solve the following questions from the Discrete Math zyBook:

Exercise 6.5.2, sections a, b

a. The range of \mathcal{A} is $\{0,1,2,3,4\}$

b.

$$(0, \frac{C(48,5)}{C(52,5)}), (1, \frac{C(4,1)*C(48,4)}{C(52,5)}), (2, \frac{C(4,2)*C(48,3)}{C(52,5)}), (3, \frac{C(4,3)*C(48,2)}{C(52,5)}), \\ (4, \frac{C(4,4)*C(48,1)}{C(52,5)})$$

Exercise 6.6.1, section a

a.

0, 1, or 2 potential choices

$C(10,2)$ potential outcomes

$$0 = C(3,2)$$

$$1 = (7,1)$$

$$2 = (7,2)$$

$$(0 * \frac{3}{45}) + (1 * \frac{21}{45}) + (2 * \frac{21}{45})$$

$$E[G] = 1.4$$

Exercise 6.6.4, sections a, b

a. Range of $X = \{1, 4, 9, 16, 25, 36\}$

$$E[X] = (1 * \frac{1}{6}) + (4 * \frac{1}{6}) + (9 * \frac{1}{6}) + (16 * \frac{1}{6}) + (25 * \frac{1}{6}) + (36 * \frac{1}{6})$$

$$E[X] = 15.166$$

b. Range of $Y = \{0, 1, 4, 9\}$

$$E[Y] = (0 * \frac{1}{8}) + (1 * \frac{1}{8}) + (4 * \frac{1}{8}) + (9 * \frac{1}{8})$$

$$E[Y] = 3$$

Exercise 6.7.4, section a

$$\text{a. } P(\textit{child gets coat}) = \frac{1}{10}$$

$$\frac{1}{10} * 10 = 1$$

Question 10:

Solve the following questions from the Discrete Math zyBook:

Exercise 6.8.1, sections a-d

The probability that a circuit board produced by a particular manufacturer has a defect is 1%. You can assume that errors are independent, so the event that one circuit board has a defect is independent of whether a different circuit board has a defect.

a. $b(2; 100, \frac{1}{100}) = C(100, 2) * (\frac{1}{100})^2 (\frac{99}{100})^{98} = 0.18$

b. Find the probability with 0 defects and 1 defect and subtract it from 1

$$1 - (C(100, 0) * (\frac{1}{100})^0 (\frac{99}{100})^{100} + C(100, 1) * (\frac{1}{100})^1 (\frac{99}{100})^{99})$$

c. $100 * \frac{1}{100} = 1$

d.

Probability that 1 batch out of 50 (2 out of 100) has a defect:

Find the probability that none have defects:

$$b(0; 50, \frac{1}{100}) = C(50, 0) (\frac{99}{100})^{50}$$

Subtract it from 1:

$$1 - b(0; 50, \frac{1}{100}) = C(50, 0) (\frac{99}{100})^{50}$$

Expected numbers of circuit boards with defects out of 100:

$$2 * (50 * \frac{1}{100}) = 1$$

The probability that at least 2 have defects differ if they are made separately versus made in batches, but the expected number of circuit boards with defects are the same.

Exercise 6.8.3, section b

b. You reach an incorrect conclusion if the number of heads in the coin flips are more than 3.

$$P(0; 10, 0.3) = C(10, 0)^0 (0.3)(0.7)^{10}$$

$$P(1; 10, 0.3) = C(10, 1)^1 (0.3)(0.7)^9$$

$$P(2; 10, 0.3) = C(10, 2)^2 (0.3)(0.7)^8$$

$$P(3; 10, 0.3) = C(10, 1)^3 (0.3)(0.7)^7$$

Add up the sum and subtract the sum from 1 = 0.35