

Statistical Foundations of Data Science

Combinatorics Part 1

University of the Witwatersrand

2022

Review Question

- Find the coefficient of x^5 in the expansion of $(2 + x)^7$

Lesson Plan

Statistical
Foundations
of Data
Science

- Review Question

Lesson Plan

Statistical
Foundations
of Data
Science

- Review Question
- Pigeonhole Principle

Lesson Plan

Statistical
Foundations
of Data
Science

- Review Question
- Pigeonhole Principle
- Basic Counting techniques

Lesson Plan

Statistical
Foundations
of Data
Science

- Review Question
- Pigeonhole Principle
- Basic Counting techniques
- Inclusion Exclusion

Review Question

- Find the coefficient of x^5 in the expansion of $(2 + x)^7$

Review Question

- Find the coefficient of x^5 in the expansion of $(2 + x)^7$
- Use the binomial theorem to see
$$(2 + x)^7 = \binom{7}{0}2^7x^0 + \binom{7}{1}2^6x^1 + \binom{7}{2}2^5x^2 + \binom{7}{3}2^4x^3 + \binom{7}{4}2^3x^4 + \binom{7}{5}2^2x^5 + \binom{7}{6}2^1x^6 + \binom{7}{7}2^0x^7$$

Review Question

- Find the coefficient of x^5 in the expansion of $(2 + x)^7$
- Use the binomial theorem to see
$$(2 + x)^7 = \binom{7}{0}2^7x^0 + \binom{7}{1}2^6x^1 + \binom{7}{2}2^5x^2 + \binom{7}{3}2^4x^3 + \binom{7}{4}2^3x^4 + \binom{7}{5}2^2x^5 + \binom{7}{6}2^1x^6 + \binom{7}{7}2^0x^7$$
- $\binom{7}{5}2^2 = 21 \times 4 = 84$

Hairy Question

- There are two people in Randburg with the same number of hairs on their heads!

Hairy Question

- There are two people in Randburg with the same number of hairs on their heads!
- I haven't counted the hairs on every head in Randburg and neither has anyone else

Hairy Question

- There are two people in Randburg with the same number of hairs on their heads!
- I haven't counted the hairs on every head in Randburg and neither has anyone else
- Still we know that essentially no one has over 200000 hairs on their head and the population of Randburg is a bit over 300000 people.

Other examples of the pigeonhole principle

- You play eight games of soccer. Two of them must happen on the same day of the week.

Other examples of the pigeonhole principle

- You play eight games of soccer. Two of them must happen on the same day of the week.
- There are 370 students in a school. Show that two of them have birthdays on the same day.

Other examples of the pigeonhole principle

- You play eight games of soccer. Two of them must happen on the same day of the week.
- There are 370 students in a school. Show that two of them have birthdays on the same day.
- There are 740 students in a school. Show that three of them have birthdays on the same day.

Other examples of the pigeonhole principle

- 25 points are placed in a 6×16 rectangle. Show that there are two between which the distance is not greater than $2\sqrt{2}$

Other examples of the pigeonhole principle

- 25 points are placed in a 6×16 rectangle. Show that there are two between which the distance is not greater than $2\sqrt{2}$
- In the Wits maths sciences there are 2022 undergrad students. Some of them are FB friends, others are not. Friendship is symmetric (if I'm friends with you you're friends with me). Show that there are two students who're friends with the same number of other students.

Basic Counting

- Addition Rule and Product Rule

Basic Counting

- Addition Rule and Product Rule
- Addition example. There are 30 CSAM faculty members and 150 students. No one is both faculty and a student. How many people are affiliated to the school? Answer 180 by addition.

Basic Counting

- Addition Rule and Product Rule
- Addition example. There are 30 CSAM faculty members and 150 students. No one is both faculty and a student. How many people are affiliated to the school? Answer 180 by addition.
- Product example. There are 30 CSAM faculty members and 150 students. No one is both faculty and a student. How many ways can we have a faculty member meet with a student? Answer 4500 by multiplication.

Basic Counting

- How many strings of length 20 are there consisting of the letters A , C , G and T (as in DNA)?

Basic Counting

- How many strings of length 20 are there consisting of the letters A , C , G and T (as in DNA)?
- 4^{20}

Basic Counting

- How many strings of length 20 are there consisting of the letters A , C , G and T (as in DNA)?
- 4^{20}
- How many strings of length 5 if we allow all 26 letters?

Basic Counting

- How many strings of length 20 are there consisting of the letters A , C , G and T (as in DNA)?
- 4^{20}
- How many strings of length 5 if we allow all 26 letters?
- 26^5

Basic Counting

- How many strings of length 20 are there consisting of the letters A , C , G and T (as in DNA)?
- 4^{20}
- How many strings of length 5 if we allow all 26 letters?
- 26^5
- What if we don't allow a letter to be repeated?

Basic Counting

- How many strings of length 20 are there consisting of the letters A , C , G and T (as in DNA)?
- 4^{20}
- How many strings of length 5 if we allow all 26 letters?
- 26^5
- What if we don't allow a letter to be repeated?
- $26 \times 25 \times 24 \times 23 \times 22$

Permutations

- How many ways can we order the 52 cards in a deck of cards?

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc
- $52! = 52 \times 51 \times 50 \times 49 \times \dots \times 1$

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc
- $52! = 52 \times 51 \times 50 \times 49 \times \dots \times 1$
- How many ways to line up 20 children. $20!$.

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc
- $52! = 52 \times 51 \times 50 \times 49 \times \dots \times 1$
- How many ways to line up 20 children. $20!$.
- How many ways to write the numbers 1 through 5 in some order? $5! = 120$. e.g 1, 2, 3, 4, 5 or 4, 3, 2, 1, 5 or 2, 4, 3, 5, 1.

Permutations

- How many ways can we order the 52 cards in a deck of cards?

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc
- $52! = 52 \times 51 \times 50 \times 49 \times \dots \times 1$

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc
- $52! = 52 \times 51 \times 50 \times 49 \times \dots \times 1$
- How many ways to line up 20 children. $20!$.

Permutations

- How many ways can we order the 52 cards in a deck of cards?
- 52 for the first card, 51 for the second etc
- $52! = 52 \times 51 \times 50 \times 49 \times \dots \times 1$
- How many ways to line up 20 children. $20!$.
- How many ways to write the numbers 1 through 5 in some order? $5! = 120$. e.g 1, 2, 3, 4, 5 or 4, 3, 2, 1, 5 or 2, 4, 3, 5, 1.

Permute function

- We call nPk the number of ways to select an ordered top k from n .

Permute function

- We call nPk the number of ways to select an ordered top k from n .
- $nPk = \frac{n!}{(n-k)!}$

Permute function

- We call nPk the number of ways to select an ordered top k from n .
- $nPk = \frac{n!}{(n-k)!}$
- Ten children are in a class. How many ways can the gold, silver and bronze medals be awarded in a race?

Permute function


- We call nPk the number of ways to select an ordered top k from n .
- $nPk = \frac{n!}{(n-k)!}$
- Ten children are in a class. How many ways can the gold, silver and bronze medals be awarded in a race?
- $10 \times 9 \times 8 = \frac{10!}{7!} = 10P3$.

Inclusion Exclusion

- $|A \cup B| = |A| + |B| - |A \cap B|$
- $|A_1 \cup A_2| = |A_1| + |A_2| - |A_1 \cap A_2|$

Inclusion Exclusion


Statistical
Foundations
of Data
Science



Inclusion Exclusion

Statistical
Foundations
of Data
Science


$$A_1 + A_2$$



Inclusion Exclusion

Statistical
Foundations
of Data
Science

$$A_1 + A_2 - A_1 \cap A_2$$



Inclusion Exclusion

- What about three sets?

Inclusion Exclusion

- What about three sets?
-

$$\begin{aligned}|A_1 \cup A_2 \cup A_3| &= |(A_1 \cup A_2) \cup A_3| \\&= |(A_1 \cup A_2)| + |A_3| \\&\quad - |(A_1 \cup A_2) \cap A_3| \\&= |(A_1 \cup A_2)| + |A_3| \\&\quad - |(A_1 \cap A_3) \cup (A_2 \cap A_3)|\end{aligned}$$

Inclusion Exclusion



$$\begin{aligned} |A_1 \cup A_2 \cup A_3| &= |(A_1 \cup A_2)| + |A_3| \\ &\quad - |(A_1 \cap A_3) \cup (A_2 \cap A_3)| \\ &= |A_1| + |A_2| - |A_1 \cap A_2| + |A_3| \\ &\quad - |(A_1 \cap A_3) \cup (A_2 \cap A_3)| \\ &\quad + |A_1 \cap A_3| + |A_2 \cap A_3| \\ &\quad - |A_1 \cap A_3 \cap A_2 \cap A_3| \end{aligned}$$


Inclusion Exclusion



$$\begin{aligned}|A_1 \cup A_2 \cup A_3| &= |A_1| \cup |A_2| + |A_3| \\ &\quad - |A_1 \cap A_2| - |A_1 \cap A_3| \\ &\quad - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|\end{aligned}$$


Inclusion Exclusion

Statistical
Foundations
of Data
Science




Inclusion Exclusion

Statistical
Foundations
of Data
Science




Inclusion Exclusion

Statistical
Foundations
of Data
Science




Inclusion Exclusion

Statistical
Foundations
of Data
Science



Inclusion Exclusion


Statistical
Foundations
of Data
Science



Inclusion Exclusion

Statistical
Foundations
of Data
Science


$$A_1 + A_2 + A_3 - A_1 \cap A_2 - A_1 \cap A_3$$



Inclusion Exclusion


Statistical
Foundations
of Data
Science

$$A_1 + A_2 + A_3 - A_1 \cap A_2 - A_1 \cap A_3 - A_2 \cap A_3 + A_1 \cap A_2 \cap A_3$$



Inclusion Exclusion


Statistical
Foundations
of Data
Science



Inclusion Exclusion


Statistical
Foundations
of Data
Science

$$A_1 + A_2 + A_3 - A_1 \cap A_2 - A_1 \cap A_3$$



Inclusion Exclusion

$$A_1 + A_2 + A_3 - A_1 \cap A_2 - A_1 \cap A_3 - A_2 \cap A_3 + A_1 \cap A_2 \cap A_3$$



Inclusion Exclusion

- How many of the whole numbers from 1 to 100 are divisible by 2, 3 or 5.

Inclusion Exclusion

- How many of the whole numbers from 1 to 100 are divisible by 2, 3 or 5.
- Make A_1 , A_2 and A_3 the whole numbers that are divisible by 2, 3 and 5 respectively.

Inclusion Exclusion

- How many of the whole numbers from 1 to 100 are divisible by 2, 3 or 5.
- Make A_1 , A_2 and A_3 the whole numbers that are divisible by 2, 3 and 5 respectively.
- $500 + 333 + 200 - 166 - 100 - 66 + 33 = 734$

Inclusion Exclusion

Statistical
Foundations
of Data
Science

- What about more sets?

Inclusion Exclusion

- What about more sets?

- General formula is

$$|\cup_{i=1}^n A_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{t \in k \text{ element sets}} |\cap_t A_t|$$