

Statistical Foundations of Data Science

Continuous Distributions

University of the Witwatersrand

2025

Review Question

- A five card poker hand is drawn. What's the probability that all five cards are the same suit.

Lesson Plan

- Review Question

Lesson Plan

- Review Question
- Continuous distributions

Lesson Plan

- Review Question
- Continuous distributions
- Exponential distribution

Lesson Plan

- Review Question
- Continuous distributions
- Exponential distribution
- Preview of the normal and t-distributions

Review Question

- A five card poker hand is drawn. What's the probability that all five cards are the same suit.

Review Question

- A five card poker hand is drawn. What's the probability that all five cards are the same suit.
- $\frac{12}{51} \times \frac{11}{50} \times \frac{10}{49} \times \frac{9}{48} = 0.00198079231$

Continuous distributions

- What's the probability a baby is born weighing exactly 3.21kg?

Continuous distributions

- What's the probability a baby is born weighing exactly 3.21kg?
- "Exactly" makes things hard because of course that'll effectively never happen.

Continuous distributions

- What's the probability a baby is born weighing exactly 3.21kg?
- "Exactly" makes things hard because of course that'll effectively never happen.
- Lots of variables like this.

Continuous distributions

- What's the probability a baby is born weighing exactly 3.21kg?
- "Exactly" makes things hard because of course that'll effectively never happen.
- Lots of variables like this.
- Instead we can sensibly ask. "What's the probability the Junior's weight is between 3.2 and 3.3 kg".

Continuous distributions

- What's the probability a baby is born weighing exactly 3.21kg?
- "Exactly" makes things hard because of course that'll effectively never happen.
- Lots of variables like this.
- Instead we can sensibly ask. "What's the probability the Junior's weight is between 3.2 and 3.3 kg".
- More generally for a continuous random variable we'll have an integrable function $f(x)$ and $\mathbb{P}(X \in (a, b)) = \int_a^b f(x)dx$

Continuous distributions

- What's the probability a baby is born weighing exactly 3.21kg?
- "Exactly" makes things hard because of course that'll effectively never happen.
- Lots of variables like this.
- Instead we can sensibly ask. "What's the probability the Junior's weight is between 3.2 and 3.3 kg".
- More generally for a continuous random variable we'll have an integrable function $f(x)$ and $\mathbb{P}(X \in (a, b)) = \int_a^b f(x)dx$
- $f(x)$ is called a "density function"

Continuous distributions - density properties

- $f(x) \geq 0$ (analog of $p(x) \geq 0$)

Continuous distributions - density properties

- $f(x) \geq 0$ (analog of $p(x) \geq 0$)
- $\int_{-\infty}^{\infty} f(x)dx = 1$ (analog of $\sum_x p(x) = 1$)

Continuous distributions - density properties

- $f(x) \geq 0$ (analog of $p(x) \geq 0$)
- $\int_{-\infty}^{\infty} f(x)dx = 1$ (analog of $\sum_x p(x) = 1$)
- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf(x)dx$

Continuous distributions - density properties

- $f(x) \geq 0$ (analog of $p(x) \geq 0$)
- $\int_{-\infty}^{\infty} f(x)dx = 1$ (analog of $\sum_x p(x) = 1$)
- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf(x)dx$
- Variance formulas hold.

Continuous distributions - density properties

- $f(x) \geq 0$ (analog of $p(x) \geq 0$)
- $\int_{-\infty}^{\infty} f(x)dx = 1$ (analog of $\sum_x p(x) = 1$)
- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf(x)dx$
- Variance formulas hold.
- We're also interested in $F(c) = \mathbb{P}(X \leq c) = \int_{-\infty}^c f(x)dx$

Example

- Find c such that

$$f(x) = \begin{cases} cx & x \in (0, 2) \\ 0 & \text{elsewhere} \end{cases}$$

is a distribution. Find its mean and variance.

Example

- Find c such that

$$f(x) = \begin{cases} cx & x \in (0, 2) \\ 0 & \text{elsewhere} \end{cases}$$

is a distribution. Find its mean and variance.

- $\int_0^2 cx dx = 1$

Example

- Find c such that

$$f(x) = \begin{cases} cx & x \in (0, 2) \\ 0 & \text{elsewhere} \end{cases}$$

is a distribution. Find its mean and variance.

- $\int_0^2 cx dx = 1$
- $c \frac{x^2}{2} \Big|_0^2 = 1$

Example

- Find c such that

$$f(x) = \begin{cases} cx & x \in (0, 2) \\ 0 & \text{elsewhere} \end{cases}$$

is a distribution. Find its mean and variance.

- $\int_0^2 cx dx = 1$
- $c \frac{x^2}{2} \Big|_0^2 = 1$
- $c = \frac{1}{2}$

Example

- $\mathbb{E}[X] = \int_0^2 xf(x)dx = \int_0^2 \frac{x^2}{2}dx = \frac{x^3}{6}\bigg|_0^2 = \frac{4}{3}$

Example

- $\mathbb{E}[X] = \int_0^2 xf(x)dx = \int_0^2 \frac{x^2}{2}dx = \frac{x^3}{6}\bigg|_0^2 = \frac{4}{3}$
- $\mathbb{E}[X^2] = \int_0^2 x^2f(x)dx = \int_0^2 \frac{x^3}{2}dx = \frac{x^4}{8}\bigg|_0^2 = 2$

Example

- $\mathbb{E}[X] = \int_0^2 xf(x)dx = \int_0^2 \frac{x^2}{2}dx = \frac{x^3}{6}\bigg|_0^2 = \frac{4}{3}$
- $\mathbb{E}[X^2] = \int_0^2 x^2f(x)dx = \int_0^2 \frac{x^3}{2}dx = \frac{x^4}{8}\bigg|_0^2 = 2$
- $V(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = 2 - \frac{4^2}{3} = \frac{2}{9}$

(Continuous) Uniform distribution



$$f(x) = \begin{cases} \frac{1}{b-a} & x \in (a, b) \\ 0 & \text{elsewhere} \end{cases}$$

(Continuous) Uniform distribution



$$f(x) = \begin{cases} \frac{1}{b-a} & x \in (a, b) \\ 0 & \text{elsewhere} \end{cases}$$

- $\mathbb{E}[X] = \int_a^b x \frac{1}{b-a} dx = \frac{x^2}{2(b-a)} \Big|_a^b = \frac{b^2 - a^2}{2(b-a)} = \frac{a+b}{2}$

(Continuous) Uniform distribution



$$f(x) = \begin{cases} \frac{1}{b-a} & x \in (a, b) \\ 0 & \text{elsewhere} \end{cases}$$

- $\mathbb{E}[X] = \int_a^b x \frac{1}{b-a} dx = \frac{x^2}{2(b-a)} \Big|_a^b = \frac{b^2-a^2}{2(b-a)} = \frac{a+b}{2}$

- $\mathbb{E}[X^2] = \int_a^b x^2 \frac{1}{b-a} dx = \frac{x^3}{3(b-a)} \Big|_a^b = \frac{b^3-a^3}{3(b-a)} = \frac{a^2+ab+b^2}{3}$

(Continuous) Uniform distribution



$$f(x) = \begin{cases} \frac{1}{b-a} & x \in (a, b) \\ 0 & \text{elsewhere} \end{cases}$$

- $\mathbb{E}[X] = \int_a^b x \frac{1}{b-a} dx = \frac{x^2}{2(b-a)} \Big|_a^b = \frac{b^2-a^2}{2(b-a)} = \frac{a+b}{2}$
- $\mathbb{E}[X^2] = \int_a^b x^2 \frac{1}{b-a} dx = \frac{x^3}{3(b-a)} \Big|_a^b = \frac{b^3-a^3}{3(b-a)} = \frac{a^2+ab+b^2}{3}$
- $V(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \frac{a^2+ab+b^2}{3} - \left(\frac{a+b}{2}\right)^2$

(Continuous) Uniform distribution



$$f(x) = \begin{cases} \frac{1}{b-a} & x \in (a, b) \\ 0 & \text{elsewhere} \end{cases}$$

- $\mathbb{E}[X] = \int_a^b x \frac{1}{b-a} dx = \frac{x^2}{2(b-a)} \Big|_a^b = \frac{b^2-a^2}{2(b-a)} = \frac{a+b}{2}$
- $\mathbb{E}[X^2] = \int_a^b x^2 \frac{1}{b-a} dx = \frac{x^3}{3(b-a)} \Big|_a^b = \frac{b^3-a^3}{3(b-a)} = \frac{a^2+ab+b^2}{3}$
- $V(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \frac{a^2+ab+b^2}{3} - \left(\frac{a+b}{2}\right)^2$
- $V(X) = \frac{(b-a)^2}{12}$

(Continuous) Uniform distribution



$$F(x) = \begin{cases} 0 & x \in x < a \\ \frac{x-a}{b-a} & x \in (a, b) \\ 1 & x > b \end{cases}$$

(Continuous) Uniform distribution



$$F(x) = \begin{cases} 0 & x \in x < a \\ \frac{x-a}{b-a} & x \in (a, b) \\ 1 & x > b \end{cases}$$

- $\int_a^c \frac{1}{b-a} dx = \frac{x-a}{b-a}$

Exponential distribution

- Continuous analogue of the geometric

Exponential distribution

- Continuous analogue of the geometric



$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

Exponential distribution

- Continuous analogue of the geometric



$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$



$$F(x) = \begin{cases} 1 - e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

Exponential distribution

- Continuous analogue of the geometric



$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$



$$F(x) = \begin{cases} 1 - e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

- $\mathbb{E}[X] = \int_0^{\infty} x \lambda e^{-\lambda x} dx$

Exponential distribution

- Continuous analogue of the geometric



$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$



$$F(x) = \begin{cases} 1 - e^{-\lambda x} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

- $\mathbb{E}[X] = \int_0^\infty x \lambda e^{-\lambda x} dx$
- Let $y = \lambda x$ so $\frac{dy}{dx} = \lambda$ and $\mathbb{E}[X] = \int_0^\infty y e^{-y} \frac{dy}{\lambda} = \frac{1}{\lambda}$

Exponential distribution

- $\mathbb{E}[X^2] = \int_0^\infty x^2 \lambda e^{-\lambda x} dx = \int_0^\infty \frac{1}{\lambda} (\lambda x)^2 e^{-\lambda x} d$

Exponential distribution

- $\mathbb{E}[X^2] = \int_0^\infty x^2 \lambda e^{-\lambda x} dx = \int_0^\infty \frac{1}{\lambda} (\lambda x)^2 e^{-\lambda x} d$
- Let $y = \lambda x$ so $\frac{dy}{dx} = \lambda$ and $\mathbb{E}[X^2] = \int_0^\infty \frac{1}{\lambda} y^2 e^{-y} \frac{dy}{\lambda} = \frac{2}{\lambda^2}$

Exponential distribution

- $\mathbb{E}[X^2] = \int_0^\infty x^2 \lambda e^{-\lambda x} dx = \int_0^\infty \frac{1}{\lambda} (\lambda x)^2 e^{-\lambda x} d$
- Let $y = \lambda x$ so $\frac{dy}{dx} = \lambda$ and $\mathbb{E}[X^2] = \int_0^\infty \frac{1}{\lambda} y^2 e^{-y} \frac{dy}{\lambda} = \frac{2}{\lambda^2}$
- $V(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \frac{1}{\lambda^2}$

Gamma distribution

- Sum of Geometric distributions.

Gamma distribution

- Sum of Geometric distributions.
- Can be thought of as waiting for the α^{th} event.

Gamma distribution

- Sum of Geometric distributions.
- Can be thought of as waiting for the α^{th} event.
-

$$f(x) = \begin{cases} \frac{\lambda^\alpha x^{\alpha-1} e^{-\lambda x}}{\Gamma(\alpha)} & \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

Gamma distribution

- Sum of Geometric distributions.
- Can be thought of as waiting for the α^{th} event.
-

$$f(x) = \begin{cases} \frac{\lambda^\alpha x^{\alpha-1} e^{-\lambda x}}{\Gamma(\alpha)} & \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

- Proof is via convolution.

Gamma distribution

- Sum of Geometric distributions.
- Can be thought of as waiting for the α^{th} event.
-

$$f(x) = \begin{cases} \frac{\lambda^\alpha x^{\alpha-1} e^{-\lambda x}}{\Gamma(\alpha)} & \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

- Proof is via convolution.
- $\mathbb{E}[X] = \frac{\alpha}{\lambda}$ by linearity of expectation

Gamma distribution

- Sum of Geometric distributions.
- Can be thought of as waiting for the α^{th} event.
-

$$f(x) = \begin{cases} \frac{\lambda^\alpha x^{\alpha-1} e^{-\lambda x}}{\Gamma(\alpha)} & \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

- Proof is via convolution.
- $\mathbb{E}[X] = \frac{\alpha}{\lambda}$ by linearity of expectation
- $V(X) = \frac{\alpha}{\lambda^2}$

Beta distribution



$$f(x) = \begin{cases} \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} & x \in (0, 1) \\ 0 & \textit{elsewhere} \end{cases}$$

Beta distribution



$$f(x) = \begin{cases} \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} & x \in (0, 1) \\ 0 & \textit{elsewhere} \end{cases}$$

- Generalizes the standard Uniform (parameters 0 and 1).

Beta distribution



$$f(x) = \begin{cases} \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} & x \in (0, 1) \\ 0 & \text{elsewhere} \end{cases}$$

- Generalizes the standard Uniform (parameters 0 and 1).
- Usually it's generate $\alpha + \beta - 1$ uniforms and choose the α smallest or β largest.

Beta distribution



$$f(x) = \begin{cases} \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)} & x \in (0, 1) \\ 0 & \text{elsewhere} \end{cases}$$

- Generalizes the standard Uniform (parameters 0 and 1).
- Usually it's generate $\alpha + \beta - 1$ uniforms and choose the α smallest or β largest.
- $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ is a scaling factor.

Beta distribution



$$f(x) = \begin{cases} \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} & x \in (0, 1) \\ 0 & \text{elsewhere} \end{cases}$$

- Generalizes the standard Uniform (parameters 0 and 1).
- Usually it's generate $\alpha + \beta - 1$ uniforms and choose the α smallest or β largest.
- $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ is a scaling factor.
- It's also the continuous analog of the choose function.

Beta distribution

- $$\mathbb{E}[X] = \int_0^1 x \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+1-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+1,\beta)}{B(\alpha,\beta)} = \frac{\alpha}{\alpha+\beta}$$

Beta distribution

- $\mathbb{E}[X] = \int_0^1 x \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+1-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+1,\beta)}{B(\alpha,\beta)} = \frac{\alpha}{\alpha+\beta}$
- $\mathbb{E}[X^2] = \int_0^1 x^2 \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+2-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+2,\beta)}{B(\alpha,\beta)} = \frac{\alpha(\alpha-1)}{(\alpha+\beta)(\alpha+\beta+1)}$

Beta distribution

- $\mathbb{E}[X] = \int_0^1 x \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+1-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+1,\beta)}{B(\alpha,\beta)} = \frac{\alpha}{\alpha+\beta}$
- $\mathbb{E}[X^2] = \int_0^1 x^2 \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+2-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+2,\beta)}{B(\alpha,\beta)} = \frac{\alpha(\alpha-1)}{(\alpha+\beta)(\alpha+\beta+1)}$
- $V(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$

Beta distribution

- $\mathbb{E}[X] = \int_0^1 x \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+1-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+1,\beta)}{B(\alpha,\beta)} = \frac{\alpha}{\alpha+\beta}$
- $\mathbb{E}[X^2] = \int_0^1 x^2 \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+2-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+2,\beta)}{B(\alpha,\beta)} = \frac{\alpha(\alpha-1)}{(\alpha+\beta)(\alpha+\beta+1)}$
- $V(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
- Lastly let's show that that is the right scaling factor!

Beta distribution

- $\mathbb{E}[X] = \int_0^1 x \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+1-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+1,\beta)}{B(\alpha,\beta)} = \frac{\alpha}{\alpha+\beta}$
- $\mathbb{E}[X^2] = \int_0^1 x^2 \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \int_0^1 \frac{x^{\alpha+2-1}(1-x)^{\beta-1}}{B(\alpha,\beta)} dx = \frac{B(\alpha+2,\beta)}{B(\alpha,\beta)} = \frac{\alpha(\alpha-1)}{(\alpha+\beta)(\alpha+\beta+1)}$
- $V(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
- Lastly let's show that that is the right scaling factor!
-

$$B(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt, \quad \text{for } x, y > 0. \quad (1)$$

Beta distribution

- Our goal is to show that:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad (2)$$

Beta distribution

- Our goal is to show that:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad (2)$$

- For integer α and β this can be done by repeated integration by parts.

Beta distribution

- Our goal is to show that:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad (2)$$

- For integer α and β this can be done by repeated integration by parts.
- Consider the product of two Gamma functions:

$$\Gamma(x)\Gamma(y) = \int_0^\infty u^{x-1}e^{-u} du \int_0^\infty v^{y-1}e^{-v} dv. \quad (3)$$

Beta distribution

- Our goal is to show that:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad (2)$$

- For integer α and β this can be done by repeated integration by parts.
- Consider the product of two Gamma functions:

$$\Gamma(x)\Gamma(y) = \int_0^\infty u^{x-1}e^{-u} du \int_0^\infty v^{y-1}e^{-v} dv. \quad (3)$$

- Introducing the change of variables:

$$u = st, \quad v = s(1-t), \quad s = u + v, \quad (4)$$

Beta distribution

- Our goal is to show that:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad (2)$$

- For integer α and β this can be done by repeated integration by parts.
- Consider the product of two Gamma functions:

$$\Gamma(x)\Gamma(y) = \int_0^\infty u^{x-1}e^{-u} du \int_0^\infty v^{y-1}e^{-v} dv. \quad (3)$$

- Introducing the change of variables:

$$u = st, \quad v = s(1-t), \quad s = u + v, \quad (4)$$

- The Jacobian is given by:

$$J = \begin{vmatrix} \frac{\partial u}{\partial s} & \frac{\partial v}{\partial s} \\ \frac{\partial u}{\partial t} & \frac{\partial v}{\partial t} \end{vmatrix} = \begin{vmatrix} t & s \\ 1-t & -s \end{vmatrix} = s. \quad (5)$$

Beta distribution

- This gives us:

$$\Gamma(x)\Gamma(y) = \int_0^\infty s^{x+y-1} e^{-s} ds \int_0^1 t^{x-1} (1-t)^{y-1} dt. \quad (6)$$

Beta distribution

- This gives us:

$$\Gamma(x)\Gamma(y) = \int_0^\infty s^{x+y-1} e^{-s} ds \int_0^1 t^{x-1} (1-t)^{y-1} dt. \quad (6)$$

- Showing that

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}. \quad (7)$$

Normal distribution

- Sometimes we like to add up random variables. For example to find an "average".

Normal distribution

- Sometimes we like to add up random variables. For example to find an "average".
- This leads to the celebrated Central Limit Theorem. Which roughly says that if we add some i.i.d random variables together that the sample mean will be approximately distributed normally.

Normal distribution

- Sometimes we like to add up random variables. For example to find an "average".
- This leads to the celebrated Central Limit Theorem. Which roughly says that if we add some i.i.d random variables together that the sample mean will be approximately distributed normally.
- Caveats exist here. We need finite moments and how good the approximation is will depend on how many X_i we're adding.

Normal distribution

- Two parameter family (mean and variance).


Normal distribution

- Two parameter family (mean and variance).
- $f(x) = \frac{1}{\sqrt{2\sigma\pi}} e^{(\frac{x-\mu}{\sigma})^2}$


Normal distribution

- Two parameter family (mean and variance).
- $f(x) = \frac{1}{\sqrt{2\sigma\pi}} e^{(\frac{x-\mu}{\sigma})^2}$
- $F(X)$ doesn't have a nice closed form. We'll use a table to look up values.

Normal distribution



Normal distribution



- Approximates the standard-normal (mean 0, variance 1)

- Approximates the standard-normal (mean 0, variance 1)
- We'll talk about how/why next lecture.

- Approximates the standard-normal (mean 0, variance 1)
- We'll talk about how/why next lecture.
- $f(X)$ is really ugly, and in practice we'll use a table.

Student T-distribution

