

Patient Information

Patient Name (optional)

smith,joe

MRN (optional)

12345678

Height (cm)

180

Weight (kg)

75

Age (years)

55

Gender

Male



Creatinine (μmol/L) (optional)

80

Treatment Start Date

2026-07-29



Diagnosis (optional)

Primary diagnosis

Allergies (optional)

Drug allergies and reactions

Calculated Values

BSA (Mosteller)

1.94 m²

$\sqrt{[(\text{height} \times \text{weight}) / 3600]}$

CrCl (Cockcroft-Gault)

97.9 mL/min

$((140 - \text{age}) \times \text{weight} \times \text{gender}) / (0.814 \times \text{creatinine})$

 Ready to Generate

All validations passed. Ready to generate PPO and MAR.

Regimen Selection

Filter by Site Group

All Sites



Select Regimen

AC - Doxorubicin/Cyclophosphamide



AC - Doxorubicin/Cyclophosphamide

Doxorubicin/Cyclophosphamide

Cycle: 21 days

Treatment Days: 1

Cycle Number

1

Total Cycles

Total planned cycles

Dose Adjustments

Percent of Ideal Dose (%)

-

100

+

%

Individual Drug Adjustments

DOXOrubicin hcl

Percentage: -

100

+

%

cyclophosphamide

Percentage: -

100

+

%

Medication Selection

Premedications



☒ ondansetron hcl

Dose: 16 mg PO

Treatment Days: Day 1

Frequency: Once

Timing: Pre-chemotherapy

Instructions: Once before chemotherapy

☒ **aprepitant**

Dose: 125 mg PO

Treatment Days: Day 1

Frequency: Once

Timing: Pre-chemotherapy

Instructions: Once before chemotherapy

☒ **dexamethasone**

Dose: 12 mg PO

Treatment Days: Day 1

Frequency: Once

Timing: Pre-chemotherapy

Instructions: Once before chemotherapy

☒ **olanzapine**

Dose: 5 mg PO

Treatment Days: Day 1

Frequency: Once

Timing: Pre-chemotherapy

Instructions: Once before chemotherapy

Primary Drugs

☒ DOXOrubicin hcl **HAZARDOUS**

Dose: 60 mg/m² PIV

Treatment Days: Day 1

Concentration: 2 mg/mL

Frequency: Once

Instructions: Infuse IV slowly at a rate of 5 mL / minute; total dose in ____ syringe(s); ____ mg / ____ mL per syringe; syringe ____ of ____

Per Day Calculation: 60 mg/m² × 1.94 m² × 100% = 116.4 → **116 mg** per day

Verification: 116 mg ÷ 1.94 m² ÷ 100% = 59.8 mg/m² (prescribed: 60 mg/m²) ✓

☒ cyclophosphamide **HAZARDOUS**

Dose: 600 mg/m² IV

Treatment Days: Day 1

Concentration: 20 mg/mL

Frequency: Once

Duration: 60 minutes

Instructions: Give IV over 60 minute(s); once

Per Day Calculation: 600 mg/m² × 1.94 m² × 100% = 1164 → **1160 mg** per day

Verification: 1160 mg ÷ 1.94 m² ÷ 100% = 597.9 mg/m² (prescribed: 600 mg/m²) ✓

Take-Home Medications

☒ pegfilgrastim LAPELGA

Dose: 6 mg SC

Treatment Days: Day 1

Frequency: Once

Timing: Post-chemotherapy

Instructions: Starting 24 hours after chemotherapy, inject the contents of one syringe (0.6mL) subcutaneously once

☒ filgrastim **Multi-day: Days 3-10**

Dose: 300 mcg SC

Treatment Days: Days 3-10

Frequency: Daily

Timing: Post-chemotherapy

Instructions: Starting Day 3 of cycle, subcutaneously daily for 8 days (Days 3-10).

☒ dexamethasone **Multi-day: Days 2-4**

Dose: 8 mg PO

Treatment Days: Days 2-4

Frequency: Daily AM

Instructions: Daily in the morning for 3 days, starting day 2

☒ aprepitant **Multi-day: Days 2-3**

Dose: 80 mg PO

Treatment Days: Days 2-3

Frequency: Daily AM

Instructions: Daily in the morning for 2 days, starting day 2

☒ olanzapine **Multi-day: Days 2-4**

Dose: 5 mg PO

Treatment Days: Days 2-4

Frequency: Daily PM

Instructions: Starting day 2, take 5mg daily in the evening for 3 days and take 2.5mg, twice daily, as required for nausea or vomiting