

CSCI 3104 PS4b

Luna McBride

TOTAL POINTS

37 / 45

QUESTION 1

10 pts

1.1 2 / 5

✓ + 1 pts Maintained separate sets of explored and unexplored vertices

+ 2 pts Minimization and picking of the next vertex from the unexplored set and adding it to explored set

✓ + 1 pts Partially correct way of picking next vertex from the unexplored set and adding it to explored or explanation not clear. Please refer to the solution file.

+ 2 pts Updating the max-weight values of unexplored vertices after a new vertex is added to the explored set.

+ 1 pts Partially correct update of max-weight values. This is applicable if there is a minor mistake while maintaining the max weight values. Please refer to the solution file.

+ 0 pts Incorrect or not attempted

1.2 0 / 5

+ 1 pts Identified the loop invariant or understood the fact that we have solved the problem for the explored vertices.

+ 2 pts Identified the fact that when a new vertex is added to explored it is chosen such that there is no max-weight path to the selected node from a vertex that does not belong to explored set

+ 2 pts Proof by contradiction or any other strategy

+ 1 pts Proof by contradiction or any other strategy partially correct

+ 1 pts Partially correct Identified the method of picking the right vertex from the unexplored set.

✓ + 0 pts Incorrect or not attempted

QUESTION 2

11 pts

2.1 4 / 4

✓ + 4 pts Correct order and correct set of edges: AE

-> EF -> FB/EB -> FG -> GH -> GC -> GD

+ 3 pts Slightly incorrect set of edges/order

+ 2 pts Partially correct set of edges/partially correct order

+ 1 pts Incorrect set of edges or incorrect correct order

+ 0 pts Incorrect

2.2 7 / 7

✓ + 2 pts Correct set of edges: AE -> EF -> FB/EB -> FG -> GH -> GC -> GD or other alternate solution

✓ + 2 pts Correct order of edges

✓ + 3 pts Correct Partitions: [[A,B,C,D], [E,F,G,H]], [[A,B,C,D,E], {F,G,H}], [[A,E,F,G,H], {B,C,D}], [[A,B,E], {C,D,F,G,H}], [[A,B,E,F,G], {C,D,H}], [[A,B,E,F,G,H], {C,D}], [[A,B,C,E,F,G,H], {D}] or other alternate solution

+ 1.5 pts Mostly correct set of edges

+ 1.5 pts Mostly correct order

+ 2.5 pts Mostly correct partitions

+ 1 pts Partially correct set of edges

+ 1 pts Partially correct order

+ 1.5 pts Partially correct partitions

+ 0 pts Incorrect

QUESTION 3

3 10 / 10

✓ + 2 pts Got the answer right (case 1: If reducing from T then it should not change, case 2: otherwise it could change)

✓ + 8 pts case 2 : gave a great counterexample or a

reasonable explanation.

- + **4 pts** case 1: No logical errors in the proof.
- + **2 pts** case 1: gave a reasonable premise
- + **2 pts** case 1: the reason why T will still be a MST is clear.
- + **0 pts** no solution / wrong
- + **8 pts** Got the idea right but need a more rigorous solution next time. (Please see the answer and try to imitate that)

QUESTION 4

14 pts

4.1 7 / 7

- ✓ + **3 pts** Get conclusion correctly. (No)
- ✓ + **4 pts** Give counterexample correctly.
- + **2 pts** Know the difference between MLST and MST, but failed to give a counterexample.
- + **0 pts** Incorrect/Not attempted.

4.2 7 / 7

- ✓ + **2 pts** Identified the contradiction correctly
- ✓ + **5 pts** Correct proof
 - + **3 pts** Proof is not explained clearly
 - + **3 pts** Knows the difference between MST and MLST but failed to provide correct proof
- + **0 pts** Incorrect answer or not attempted
- + **4 pts** Partially correct proof

Instructions for submitting your solution:

- The solutions **should be typed** and we cannot accept hand-written solutions. Here's a short intro to Latex.
 - You should submit your work through **Gradescope** only.
 - If you don't have an account on it, sign up for one using your CU email. You should have gotten an email to sign up. If your name based CU email doesn't work, try the identikey@colorado.edu version.
 - Gradescope will only accept **.pdf** files (except for code files that should be submitted separately on Gradescope if a problem set has them) and **try to fit your work in the box provided**.
 - You cannot submit a pdf which has less pages than what we provided you as Gradescope won't allow it.
 - Verbal reasoning is typically insufficient for full credit. Instead, write a logical argument, in the style of a mathematical proof.
 - For every problem in this class, you must justify your answer: show how you arrived at it and why it is correct. If there are assumptions you need to make along the way, state those clearly.
 - You may work with other students. However, **all solutions must be written independently and in your own words**. Referencing solutions of any sort is strictly prohibited. You must explicitly cite any sources, as well as any collaborators.
-

1. (10 pts) For a directed graph with positive weights, we define the max-weight of a path from s to d as the maximum of the edge weights along the path. For example, if the path from A to D has edges and weights of $e_{AB} = 5$, $e_{BC} = 4$, and $e_{CD} = 1$, the length of the path is defined as $e_{AB} + e_{BC} + e_{CD}$, and the max-weight is 5.
 - (a) (5 pts) Give an algorithm to compute the smallest max-weight paths from a source vertex s to all other vertices. In this problem, you are changing the definition of length of the path from A to D to $\max(e_{AB}, e_{BC}, e_{CD})$ (Hint: Your algorithm should be a modification of Dijkstra's algorithm presented in Lecture.)

Solution.

ModDijkstra(G,S):

```
- max=0
- Initialize dist, prev for all vertices plus max edge weight
-- > dist(s)=0
-- > dist(v)=∞
-- > Q=min priority queue
-- > Q.add(v)
-- while Q!empty:
--- / u=Q.pop()
--- / for each vert in u.adjacent:
---- / -- > d=dist(v)+edge(u,v)
---- / -- > if d<dist(v):
---- / -- / -- > dist(v)=d
---- / -- / -- > prev(v)=u

-- for i in range 0, len(vertices):
--- / d=dist(v[i])
--- / if max < d:
---- / -- > max=d
- return G
```

(b) (5 pts) Prove the correctness of your algorithm.

Solution.

Loop Invariance for the added for loop: $\text{dist}(v[0]), \text{dist}(v[1]) \dots \text{dist}(v[i-1]) \leq \text{max}$ for the already completed version of Dijkstra's algorithm.

Base case: $\text{dist}(v[0])$ is greater than 0, which max is then set equal to. max is then equal to the only vertex checked, which fits into the \geq title.

Maintainence: For every iteration of the loop, max is set to the biggest value it comes across. Any value less than the current max can then be ignored, as it is not the max . Thus, as the largest value is contained in max , which means every value $\text{dist}(v[0-(i-1)])$ is necessarily less than or equal to max .

Termination: On the last iteration, the max either keeps the last value if it is the largest or ignores it because it is less than the current max . In either case, the largest value overall is now contained within the max variable. This means that every single distance value for every vertex is either less than or equal to max .


Since the shortest path has already been obtained before this invariance relation comes into play and it does obtain the maximum as shown in the invariance, then the max value is the maximum value needed within the shortest path.

Name: Luna McBride
ID: 107607144

CSCI 3104, Algorithms
Problem Set 4b (45 points)

Profs. Hoenigman & Agrawal
Fall 2019, CU-Boulder

2. (11 pts) Based on the following graph :



- (a) (4 pts) In what order would Prim's algorithm add edges to the MST if we start at vertex A ?

Solution.

Prim: Order: A, E, F, B, G, H, C, D

Start: A Root = A, Cost = 0

AB: Cost = 6, B.Par = A

AE: Cost = 1, E.Par = A

Q.pop() \rightarrow v = E A, E & Q

EB: Cost = 2, B.Par = E

EF: Cost = 1, F.Par = E

Q.pop() \rightarrow v = F A, E, F & Q

FB: Cost = 2

FC: Cost = 5, C.Par = F

FG: Cost = 3, G.Par = F

Q.pop() \rightarrow v = B A, E, F, B & Q

BC: Cost = 5

Q.pop() \rightarrow v = G A, E, F, B, G & Q

GC: Cost = 4, C.Par = G

GH: Cost = 3, H.Par = G

GD: Cost = 5, D.Par = G

Q.pop() \rightarrow v = H A, B, E, F, G, H & Q \leftarrow alphabetical

HD: Cost = 7

Q.pop() \rightarrow v = C A, B, C, E, F, G, H & Q

CD: Cost = 6

Q.pop() \rightarrow v = D A, B, C, D, E, F, G, H & Q

All other values not in above

- (b) (7 pts) In what order Kruskal's would add the edges to the MST? For each edge added by Kruskal's sequentially, give a cut that justifies its addition.

Solution.

From left to right (following value assention):

For A: $p_1=A$, $p_2=B,C,D,E,F,G,H$. Edges in cut: $AB=6$, $AE=1$. $6>1$, so AE is chosen.

$p_1=E$, $p_2=A,B,C,D,F,G,H$. AE is already chosen, so Edges in cut: $EB=2$, $EF=1$. $2>1$, so EF is chosen.

Next: $p_1=A,B,C,D$, $p_2=E,F,G,H$. Disregarding the already chosen AE , the smallest value in the cut is 2, both going to B. It could choose either (but not both), so I will go left to right and say $EB=2$ is chosen.

Next: $p_1=A,B,E,F$, $p_2=C,D,G,H$. The lowest in the cut is 3, so next chosen is $FG=3$.

Next: $p_1=A,B,C,E,F,G$, $p_2=D,H$ (as there is a 3 in there). The value of 3 exists in this set, which is less than 5 and 6. Thus, $GH=3$ will be chosen

Next: $p_1=A,B,C,D$, $p_2=E,F,G,H$. Previously looked at lines AE, EB, BF are ignored via cycles and already chosen values. The value CG is less than the other values in the set, so $CG=4$ is chosen.


Next: $p_1=D$, $p_2=A,B,C,E,F,G,H$ (as it is the last node unconnected). The values here are 6,7,5. Since 5 is the lowest, $GD=5$ will be chosen.

All other edges will create loops, so the final edge list is $[AE, EF, BE, FG, GH, CG, DG]$ at a weight of 19 (same ones listed above in the same order, but I put the edges in alphabetical order)

3. (10 pts) Let T be a MST of a given graph G . Will T still be the MST if we reduce the weight of exactly one of the edges in G by a constant c ? Prove your answer.

Solution.

Well, let us say this assumption is true for any edge. Take a look at this image:



This shows a basic graph and MST. For AB and BC, since these are already the required minimum, any lowering of them will not change the shape. Next, we have AC.

$$AC = 16 - c, AB = 3, BC = 2$$


For the MST to stay an MST, for all $c < \text{edgeweight}$ (it must have a positive edge weight at all times), $AC \geq AB$ or BC , or $16 - c \geq 3$ or $16 - c \geq 2$, since otherwise this spanning tree is not the minimum by definition since there is a smaller edge.

Let us take AB as an example, since it is the largest between AB and BC

$$16 - c \geq 3 \rightarrow 16 \geq 3 + c \rightarrow 13 \geq c \text{ for AB to remain a minimum}$$

However, 13 does not cover all possible c values. Since $c < 16$, this leaves all values $13 < c < 16$ that will create a smaller edge weight, which would turn that into an edge needed in the MST, negating the previous MST, putting it as just a spanning tree.

Therefore, by contradiction, T will not necessarily be a MST if we reduce one of the graph's edges by a constant c




4. (14 pts) One of the uses of MSTs is finding a set of edges that span a network for minimum cost. Network problems could also have another type of objective: designing a spanning tree for which the most expensive edge is minimized. Specifically, let $G = (V, E)$ be a connected graph with n vertices, m edges, and positive edge costs that you may assume are all distinct. Let $T = (V, E)$ be a spanning tree of G ; we define the **limiting edge** of T to be the edge of T with the greatest cost. A spanning tree T of G is a minimum-limiting spanning tree if there is no spanning tree T' of G with a cheaper limiting edge.

- (a) (7 pts) Is every minimum-limiting tree of G an MST of G ? Prove or give a counterexample.


Solution.

Assuming that all MLT are MST, Say we have this graph:



With this tree, edges DE and AB must necessarily be a part of all spanning trees, as otherwise, not all vertices are connected. That means the limiting edge is going to be DE, or 9, for every spanning tree, and thus every single spanning tree is also a limiting tree.

Taking this, we can look at this spanning tree:



This is a minimum limiting tree, since all trees in this graph must necessarily be a minimum limiting tree. However, since the smallest edge has been removed, it cannot be a minimum spanning tree.

Therefore, by contradiction, all MLT are not necessarily MST.

- (b) (7 pts) Prove that every MST of G is a minimum-limiting tree of G . [Hint: Let T be an MST of G , and let T' be a minimum-limiting tree of G . If T is not a minimum-limiting tree, can we replace the heaviest edge of T ? Think about how to use T' here.]

Solution. Say, within our tree T , we have our largest edge E . Let us say this is a larger edge than the minimum limiting edge from T' , E' , such that $E \geq E'$.

Since E is larger than E' , we should try and replace value E for an E to make $E=E'$. By the nature of a MST, however, any edge replacement in the graph for this edge will be a higher value than this edge, as otherwise this spanning tree will not be a minimum spanning tree. With no lower edges to come by and E being the highest in our minimum tree, E fits the definition of a limiting edge. Thus, E cannot be higher than E' , as otherwise it will not fit the definition of a MST. E cannot be lower than E' , as that would go against the definition of a limiting edge and make E' not a limiting edge.


Therefore, E must equal E' and the MST must also be an MLT for all limiting edges E .

Ungraded questions - These questions are for your practice. We won't grade them or provide a typed solution but we are open to discuss these in our OHs and you should take feed backs on your approach during the OHs. These questions are part of the syllabus.

1. Suppose you are given the minimum spanning tree T of a given graph G (with n vertices and m edges) and a new edge $e = (u, v)$ of weight w that will be added to G . Give an efficient algorithm to find the MST of the graph $G \cup e$, and prove its correctness. Your algorithm should run in $O(n)$ time.

Solution.

2. Based on the following graph :



- (a) Run Kruskal's and find the MST. You can break the ties however you want. Draw the MST that you found and also find it's total weight.

Solution.


- (b) Run Prim's starting from vertex A and find the MST. You can break the ties however you want. Draw the MST that you found and also find its total weight. Is the total weight same as what you get from the above?

Solution.

3. Consider the following unweighted graph, and assign the edge weights (using positive integer weights only), such that the following conditions are true regarding minimum spanning trees (MST) and single-source shortest path (SSSP) trees:

- The MST is distinct from any of the seven SSSP trees.
- The order in which Prim's algorithm adds the safe edges is different from the order in which Kruskal's algorithm adds them.

Justify your solution by (i) giving the edges weights, (ii) showing the corresponding MST and all the SSSP trees, and (iii) giving the order in which edges are added by each of the three algorithms.



Solution.