

# DateLife Workflows

Luna L. Sanchez Reyes

2019-03-01

## Taxon Fringilidae

### I. Query data

There are 475 species in the Open Tree of Life Taxonomy for the taxon Fringilidae. Information on time of divergence is available for 286 of these species across 13 published and peer-reviewed chronograms from the following studies:

1. Hooper, Daniel M., Trevor D. Price. 2017. Chromosomal inversion differences correlate with range overlap in passerine birds. *Nature Ecology & Evolution* 1 (10): 1526-1534
2. Hedges, S. Blair, Julie Marin, Michael Suleski, Madeline Paymer, Sudhir Kumar. 2015. Tree of life reveals clock-like speciation and diversification. *Molecular Biology and Evolution* 32 (4): 835-845
3. Claramunt, Santiago, Joel Cracraft. 2015. A new time tree reveals Earth history's imprint on the evolution of modern birds. *Science Advances* 1 (11): e1501005-e1501005
4. Gibb, Gillian C., Ryan England, Gerrit Hartig, P.A. (Trish) McLenachan, Briar L. Taylor Smith, Bennet J. McComish, Alan Cooper, David Penny. 2015. New Zealand passerines help clarify the diversification of major songbird lineages during the Oligocene. *Genome Biology and Evolution* 7 (11): 2983-2995.
5. Barker, F. Keith, Kevin J. Burns, John Klicka, Scott M. Lanyon, Irby J. Lovette. 2015. New insights into New World biogeography: An integrated view from the phylogeny of blackbirds, cardinals, sparrows, tanagers, warblers, and allies. *The Auk* 132 (2): 333-348.
6. Jetz, W., G. H. Thomas, J. B. Joy, K. Hartmann, A. O. Mooers. 2012. The global diversity of birds in space and time. *Nature* 491 (7424): 444-448
7. Barker, F. K., K. J. Burns, J. Klicka, S. M. Lanyon, I. J. Lovette. 2013. Going to extremes: contrasting rates of diversification in a recent radiation of New World passerine birds. *Systematic Biology* 62 (2): 298-320.
8. Burns, Kevin J., Allison J. Shultz, Pascal O. Title, Nicholas A. Mason, F. Keith Barker, John Klicka, Scott M. Lanyon, Irby J. Lovette. 2014. Phylogenetics and diversification of tanagers (Passeriformes: Thraupidae), the largest radiation of Neotropical songbirds. *Molecular Phylogenetics and Evolution* 75: 41-77.

**9. Price, Trevor D., Daniel M. Hooper, Caitlyn D. Buchanan, Ulf S. Johansson, D. Thomas Tietze, Per Alström, Urban Olsson, Mousumi Ghosh-Harihar, Farah Ishtiaq, Sandeep K. Gupta, Jochen Martens, Bettina Harr, Pratap Singh, Dhananjai Mohan. 2014. Niche filling slows the diversification of Himalayan songbirds. Nature 509: 222-225.**

All source chronograms are fully ultrametric. The proportion of Fringilidae queried species found across source chronograms is as follows:

|   | Trees                           | Tips | Resolved |
|---|---------------------------------|------|----------|
| 1 | Open Tree of Life Subtree       | 474  | 100%     |
| 2 | Dated Open Tree of Life Subtree | 475  | 79%      |
| 3 | Median Summary Chronogram       | 286  | 100%     |
| 4 | SDM Summary Chronogram          | 286  | 100%     |




Figure 1: Fringilidae Species Dated Open Tree of Life Induced Subtree. This chronogram was obtained with `get_dated_otol_induced_subtree()` function.




Figure 2: Fringilidae lineage through time (LTT) plots from source chronograms, summary median chronogram and dated Open Tree of Life chronogram.

## II. Summarize results.

### II.A. Diagnosing clustering issues.

We identified some issues with chronograms coming from SDM and Median summary matrices. First, clustering algorithms implemented to go from a summary distance matrix to a tree return trees that are too old (generally with UPGMA algorithms) or non-ultrametric (generally with Neighbour Joining algorithms). In most studied cases, UPGMA returns fully ultrametric trees but with very old ages (we had to multiply the matrix by 0.25 to get ages approximate to source chronograms ages, however this is a number chosen at random, it was just the number that worked well. NJ returned reasonable ages, but trees are way non ultrametric, as you can see in Fig. 3 and Fig. 4.

This taxon's SDM matrix has some negative values in the following taxa: *Carduelis uropygialis*, *Spinus crassirostris*. This taxon's Median matrix has NO negative values.

### II.B. Age distributions form Median and SDM summary trees.

Comparison of summary chronograms reconstructed with min and max ages.




Figure 3: Fringilidae lineage through time (LTT) plots from source chronograms and Median summary matrix converted to phylo with different methods (NJ and UPGMA). Clustering algorithms used often are returning non-ultrametric trees or with maximum ages that are just off (too old or too young). So we developed an alternative algorithm in **datelife** to go from a summary matrix to a fully ultrametric tree.




Figure 4: Fringilidae lineage through time (LTT) plots from source chronograms and SDM summary matrix converted to phylo with different methods (NJ and UPGMA). As you can note, dashed lines and solid lines from trees coming out from both types of clustering algorithms implemented are mostly overlapping. This means that removing negative values does not change results from clustering algorithms much. Clustering algorithms used often are returning non-ultrametric trees or with maximum ages that are just off (too old or too young). So we developed an alternative algorithm in **datelife** to go from a summary matrix to a fully ultrametric tree.




Figure 5: Fringilidae lineage through time (LTT) plots from source chronograms and Median summary matrix converted to phylo with `datelife` algorithm.




Figure 6: Fringilidae lineage through time (LTT) plots from source chronograms and SDM summary matrix converted to phylo with `datelife` algorithm.

### III. Create new data

As an example, we're gonna date the Open Tree Synthetic tree (mainly because the taxonomic tree is usually less well resolved.)

Now, let's say you like the Open Tree of Life Taxonomy and you want to stick to that tree. Dates from available studies were tested over the Open Tree of Life Synthetic tree of Fringilidae and a tree was constructed, but all branch lengths are NA. We also tried each source chronogram independently, with the Dated OToL and with each other, as a form of cross validation in Table 2. This is not working perfectly yet, but we are developping new ways to use all calibrations efficiently.

Table 2: Was it successful to use each source chronogram independently as calibration (CalibN) against the Dated Open Tree of Life (dOToL) and each other (ChronoN)?

|         | dOToL | Chr1 | Chr2 | Chr3 | Chr4 | Chr5 | Chr6 | Chr7 | Chr8 | Chr9 | Chr10 | Chr11 | Chr12 | Chr13 |
|---------|-------|------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|
| Calib1  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib2  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib3  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib4  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib5  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib6  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib7  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib8  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib9  | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib10 | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib11 | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib12 | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |
| Calib13 | yes   | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes  | yes   | yes   | yes   | yes   |

### III. Simulate data

An alternative to generate a dated tree from a set of taxa is to take the available information and simulate into it the missing data. We will take the median and sdm summary chronograms to date the Synthetic tree of Life:

```
#> Error in paste0("\n![" , figcap_lttplot_sdm, "](plots/" , taxon, "_LTTplot_sdm.pdf)\n"): object 'fig'
#> Error in cat(lttplot): object 'lttplot' not found
```



## Appendix

The following species were completely absent from the chronogram data base: *Acanthis cabaret*, *Acanthis rostrata*, *Akialoa ellisiana*, *Akialoa lanaiensis*, *Akialoa obscura*, *Buarremon apertus*, *Bucanetes crassirostris*, *Calcarius coloratus*, *Cardinalis carneus*, *Cardinalis peninsulae*, *Carduelis ankoberensis*, *Carduelis elegans*, *Carduelis ultima*, *Carpodacus beicki*, *Carpodacus davidianus*, *Carpodacus deserticolor*, *Carpodacus dubius*, *Carpodacus formosanus*, *Carpodacus henrici*, *Carpodacus longirostris*, *Carpodacus lucifer*, *Carpodacus portenkoi*, *Carpodacus rhodopeplus*, *Carpodacus roseatus*, *Carpodacus rubicundus*, *Carpodacus stoliczkae*, *Carpodacus verreauxii*, *Carpodacus waltoni*, *Caryothraustes brasiliensis*, *Caryothraustes scapularis*, *Chaunoproctus ferreorostris*, *Chloridops kona*, *Chloris heinrichi*, *Chloris turkestanica*, *Chrysocorythus mindanensis*, *Ciridops anna*, *Coccothraustes japonicus*, *Coccothraustes migratorius*, *Corytus rhenana*, *Crithagra albifrons*, *Crithagra ankoberensis*, *Crithagra buehneri*, *Crithagra canicapilla*, *Crithagra capistrata*, *Crithagra concolor*, *Crithagra deserti*, *Crithagra donaldsoni*, *Crithagra flavigula*, *Crithagra frontalis*, *Crithagra granti*, *Crithagra hewitti*, *Crithagra hildegardeae*, *Crithagra kikuyensis*, *Crithagra koliensis*, *Crithagra leucoptera*, *Crithagra marshalli*, *Crithagra menachensis*, *Crithagra montanorum*, *Crithagra mozambica*, *Crithagra reichenowi*, *Crithagra rothschildi*, *Crithagra rufobrunnea*, *Crithagra symonsi*, *Crithagra thomensis*, *Crithagra tristriata*, *Crithagra xantholaema*, *Crithagra xanthopygia*, *Cyanerpes holti*, *Cyanerpes isthmicus*, *Cyanerpes microrhynchus*, *Cyanocompsa argentina*, *Cyanocompsa rothschildii*, *Drepanis coccinea*, *Drepanis funerea*, *Drepanis pacifica*, *Dysmorodrepanis munroi*, *Emberiza buturlini*, *Emberiza ciodes*, *Emberiza ciopsis*, *Emberiza continentalis*, *Emberiza elegantula*, *Emberiza erythrogenys*, *Emberiza flemingorum*, *Emberiza fronto*, *Emberiza kuatunensis*, *Emberiza lydiae*, *Emberiza meridionalis*, *Emberiza militaris*, *Emberiza musica*, *Emberiza neobscura*, *Emberiza nivenorum*, *Emberiza omissa*, *Emberiza omoensis*, *Emberiza orientalis*, *Emberiza ornata*, *Emberiza pyrrhulinus*, *Emberiza rufibarba*, *Emberiza ruficularis*, *Emberiza sahari*, *Emberiza semenowi*, *Emberiza sloggetti*, *Emberiza sordida*, *Emberiza vincenti*, *Emberiza zaidamensis*, *Embernagra gossei*, *Eophona magnirostris*, *Eophona sowerbyi*, *Erythrospiza phaenicoptera*, *Euphonia aurantiicollis*, *Euphonia carnegiei*, *Euphonia flavifrons*, *Euphonia gnatho*, *Euphonia nitida*, *Euphonia olivacea*, *Euphonia praetermissa*, *Euphonia purpurascens*, *Euphonia rufivertex*, *Euphonia serrirostris*, *Euphonia tavarae*, *Fringilla albicollis*, *Fringilla bella*, *Fringilla brissonii*, *Fringilla nortoniensis*, *Fringilla palmae*, *Fringilla polatzeki*, *Fringilla syriaca*, *Fringillaria goslingi*, *Fringillaria poliopleura*, *Haemorhous californicus*, *Haemorhous griscomi*, *Hemignathus affinis*, *Hemignathus hanapepe*, *Hemispingus castaneicollis*, *Hemispingus macrophrys*, *Hemispingus ochraceus*, *Hemispingus urubambae*, *Hesperiphona abeillei*, *Hesperiphona cobanensis*, *Hesperiphona montana*, *Himatione fraithii*, *Leucosticte brunneonucha*, *Leucosticte wallowa*, *Leucosticte walteri*, *Linaria harterti*, *Linaria johannis*, *Linaria rufostrigata*, *Linaria yemenensis*, *Linurgus kilimensis*, *Loxia cardinalis*, *Loxia cyanea*, *Loxia dominica*, *Loxia mesamericana*, *Loxops ochraceus*, *Loxops wolstenholmei*, *Melopyrrha taylori*, *Mycerobas melanozanthos*, *Passerina lazula*, *Passerina pallidior*, *Passerina purpurascens*, *Peucaea cohaerens*, *Peucaea ibarrorum*, *Peucaea vulcanica*, *Peucedramus micrus*, *Pheucticus aurantiacus*, *Pinicola eschatosia*, *Plectrophenax townsendi*, *Psittirostra psittacea*, *Pyrrhula cineracea*, *Pyrrhula owstoni*, *Pyrrhula rosacea*, *Pyrrhula steerei*, *Pyrrhula uchidai*, *Rhodacanthis flaviceps*, *Rhodacanthis palmeri*, *Rhodopechys alienus*, *Rhodopechys sanguineus*, *Rhynchostruthus louisae*, *Rhynchostruthus percivali*, *Rhynchostruthus socotranus*, *Serinus huillensis*, *Spinus atriceps*, *Spinus colombiana*, *Spinus dominicensis*, *Spinus longirostris*, *Spinus nigricauda*, *Spinus oleacea*, *Spinus perplexa*, *Spinus stejnegeri*, *Viridonia sagittirostris*