

Data Wrangling: The Portal Data Set

Luna L Sanchez Reyes

2023-03-02

1. Intro to the Portal data set

Homework: create an intro describing the location of the experiment and the different experimental treatments. Paper here <https://esajournals.onlinelibrary.wiley.com/doi/full/10.1890/15-2115.1>

2. Load the data

There are three different data sets, so to load them we need to create three data frames

```
surveys <- read.csv(file = "../data-raw/surveys.csv")
head(surveys)
```

```
##   record_id month day year plot_id species_id sex hindfoot_length weight
## 1         1     7  16 1977      2         NL   M             32      NA
## 2         2     7  16 1977      3         NL   M             33      NA
## 3         3     7  16 1977      2         DM   F             37      NA
## 4         4     7  16 1977      7         DM   M             36      NA
## 5         5     7  16 1977      3         DM   M             35      NA
## 6         6     7  16 1977      1         PF   M             14      NA
```

```
species <- read.csv(file = "../data-raw/species.csv")
head(species)
```

```
##   species_id      genus      species      taxa
## 1         AB  Amphisipiza  bilineata  Bird
## 2         AH Ammospermophilus  harrisi Rodent
## 3         AS  Ammodramus  savannarum  Bird
## 4         BA   Baiomys  taylori Rodent
## 5         CB Campylorhynchus  brunneicapillus  Bird
## 6         CM  Calamospiza  melanocorys  Bird
```

```
plots <- read.csv(file = "../data-raw/plots.csv")
head(plots)
```

```
##   plot_id      plot_type
## 1         1 Spectab enclosure
## 2         2      Control
## 3         3 Long-term Krat Enclosure
## 4         4      Control
## 5         5      Rodent Enclosure
## 6         6 Short-term Krat Enclosure
```

3. The dplyr package for data wrangling

Subset columns from a data frame with the function `select()`

```
head(surveys)
```

```
##   record_id month day year plot_id species_id sex hindfoot_length weight
## 1         1     7  16 1977      2         NL   M          32         NA
## 2         2     7  16 1977      3         NL   M          33         NA
## 3         3     7  16 1977      2         DM   F          37         NA
## 4         4     7  16 1977      7         DM   M          36         NA
## 5         5     7  16 1977      3         DM   M          35         NA
## 6         6     7  16 1977      1         PF   M          14         NA
```

```
surveys_subset <- select(surveys, month, day, year)
```

Create new variables from existing variables or transform existing variables with `mutate()`

The `hindfoot_length` variable is measured in mm. I want a new variable that stores hindfoot length in cm.

```
mutate(head(surveys), hindfoot_length_cm = hindfoot_length/10)
```

```
##   record_id month day year plot_id species_id sex hindfoot_length weight
## 1         1     7  16 1977      2         NL   M          32         NA
## 2         2     7  16 1977      3         NL   M          33         NA
## 3         3     7  16 1977      2         DM   F          37         NA
## 4         4     7  16 1977      7         DM   M          36         NA
## 5         5     7  16 1977      3         DM   M          35         NA
## 6         6     7  16 1977      1         PF   M          14         NA
##   hindfoot_length_cm
## 1                 3.2
## 2                 3.3
## 3                 3.7
## 4                 3.6
## 5                 3.5
## 6                 1.4
```

```
head(surveys)
```

```
##   record_id month day year plot_id species_id sex hindfoot_length weight
## 1         1     7  16 1977      2         NL   M          32         NA
## 2         2     7  16 1977      3         NL   M          33         NA
## 3         3     7  16 1977      2         DM   F          37         NA
## 4         4     7  16 1977      7         DM   M          36         NA
## 5         5     7  16 1977      3         DM   M          35         NA
## 6         6     7  16 1977      1         PF   M          14         NA
```

```
surveys_mutated <- mutate(surveys, hindfoot_length_cm = hindfoot_length/10)
head(surveys_mutated)
```

```
##   record_id month day year plot_id species_id sex hindfoot_length weight
## 1         1     7  16 1977      2         NL   M          32         NA
## 2         2     7  16 1977      3         NL   M          33         NA
## 3         3     7  16 1977      2         DM   F          37         NA
## 4         4     7  16 1977      7         DM   M          36         NA
## 5         5     7  16 1977      3         DM   M          35         NA
```

```
## 6      6      7 16 1977      1      PF  M      14      NA
## hindfoot_length_cm
## 1      3.2
## 2      3.3
## 3      3.7
## 4      3.6
## 5      3.5
## 6      1.4
```

Sorting or ordering data with the function `arrange()`

If we want to order the data frame values based on the weight variable:

```
surveys_arranged <- arrange(surveys, weight)
head(surveys_arranged)
```

```
## record_id month day year plot_id species_id sex hindfoot_length weight
## 1      218     9 13 1977      1      PF  M      13      4
## 2     4052     4  5 1981      3      PF  F      15      4
## 3     4290     4  6 1981      4      PF      NA      4
## 4     5346     2 22 1982     21      PF  F      14      4
## 5     7084    11 22 1982      3      PF  F      16      4
## 6     8736    12  8 1983     19      RM  M      17      4
```

Order values in descendant order with the function `desc()`

```
surveys_arranged <- arrange(surveys, desc(weight))
head(surveys_arranged)
```

```
## record_id month day year plot_id species_id sex hindfoot_length weight
## 1     33049    11 17 2001     12      NL  M      33     280
## 2     12871     5 28 1987      2      NL  M      32     278
## 3     15459     1 11 1989      9      NL  M      36     275
## 4      2133    10 25 1979      2      NL  F      33     274
## 5     12729     4 26 1987      2      NL  M      32     270
## 6     13114     7 26 1987      2      NL  M      NA     269
```

Filter values with the function `filter()`

Filter the data frame to keep rows with weight values that are equal to 4:

```
filter(surveys, weight == 4)
```

```
## record_id month day year plot_id species_id sex hindfoot_length weight
## 1      218     9 13 1977      1      PF  M      13      4
## 2     4052     4  5 1981      3      PF  F      15      4
## 3     4290     4  6 1981      4      PF      NA      4
## 4     5346     2 22 1982     21      PF  F      14      4
## 5     7084    11 22 1982      3      PF  F      16      4
## 6     8736    12  8 1983     19      RM  M      17      4
## 7     9790     1 19 1985     16      RM  F      16      4
## 8     9794     1 19 1985     24      RM  M      16      4
## 9     9799     1 19 1985     19      RM  M      16      4
## 10    9823     1 19 1985     23      RM  M      16      4
```

```
## 11      9853      1 19 1985      17      RM  M      16      4
## 12      9909      1 20 1985      15      RM  F      15      4
## 13      9937      2 16 1985      21      RM  M      16      4
## 14     10119      3 17 1985      10      RM  M      16      4
## 15     10439      5 24 1985       7      RM  M      16      4
## 16     28126      6 28 1998      15      PF  M      NA      4
## 17     29906     10 10 1999       4      PP  M      21      4
```

```
surveys_filtered <- filter(surveys, weight != 4)
head(surveys_filtered)
```

```
##  record_id month day year plot_id species_id sex hindfoot_length weight
## 1         63     8 19 1977      3         DM  M          35      40
## 2         64     8 19 1977      7         DM  M          37      48
## 3         65     8 19 1977      4         DM  F          34      29
## 4         66     8 19 1977      4         DM  F          35      46
## 5         67     8 19 1977      7         DM  M          35      36
## 6         68     8 19 1977      8         DO  F          32      52
```

```
surveys_filtered <- filter(surveys, weight > 200)
head(surveys_filtered)
```

```
##  record_id month day year plot_id species_id sex hindfoot_length weight
## 1         588     2 18 1978      2         NL  M          NA     218
## 2         646     2 20 1978     18         NL  M          32     228
## 3         655     3 11 1978      3         NL  M          32     232
## 4         825     4 10 1978     18         NL  M          NA     225
## 5         845     5  6 1978      2         NL  M          32     204
## 6         848     5  6 1978     22         NL  M          32     212
```

Filter with more complex conditions

I want values that have weight larger than 200 AND also are females

```
surveys_filtered <- filter(surveys, weight > 200, sex == "F")
head(surveys_filtered)
```

```
##  record_id month day year plot_id species_id sex hindfoot_length weight
## 1         875     5 17 1978      5         NL  F          33     212
## 2        1731     4 28 1979     12         NL  F          32     239
## 3        2081    10 24 1979     12         NL  F          32     211
## 4        2133    10 25 1979      2         NL  F          33     274
## 5        2247    11 18 1979     12         NL  F          33     217
## 6        2305     1 15 1980     12         NL  F          32     214
```

```
surveys_filtered <- filter(surveys, weight > 200 & sex == "F")
head(surveys_filtered)
```

```
##  record_id month day year plot_id species_id sex hindfoot_length weight
## 1         875     5 17 1978      5         NL  F          33     212
## 2        1731     4 28 1979     12         NL  F          32     239
## 3        2081    10 24 1979     12         NL  F          32     211
## 4        2133    10 25 1979      2         NL  F          33     274
## 5        2247    11 18 1979     12         NL  F          33     217
## 6        2305     1 15 1980     12         NL  F          32     214
```

Now, I want values that have weight larger than 200 OR are also females:

```
surveys_filtered <- filter(surveys, weight > 200 | sex == "F")
head(surveys_filtered)
```

Filtering NA values

NA is a special value in R. We can't use logical statements with it, we have to use the `is.na()` function:

```
surveys_filtered <- filter(surveys, !is.na(weight))
head(surveys_filtered)
```

| ## | record_id | month | day | year | plot_id | species_id | sex | hindfoot_length | weight |
|------|-----------|-------|-----|------|---------|------------|-----|-----------------|--------|
| ## 1 | 63 | 8 | 19 | 1977 | 3 | DM | M | 35 | 40 |
| ## 2 | 64 | 8 | 19 | 1977 | 7 | DM | M | 37 | 48 |
| ## 3 | 65 | 8 | 19 | 1977 | 4 | DM | F | 34 | 29 |
| ## 4 | 66 | 8 | 19 | 1977 | 4 | DM | F | 35 | 46 |
| ## 5 | 67 | 8 | 19 | 1977 | 7 | DM | M | 35 | 36 |
| ## 6 | 68 | 8 | 19 | 1977 | 8 | DO | F | 32 | 52 |