

# Emerging Directions in Deep Learning

Manifolds - key concepts and examples

Mihai-Sorin Stupariu, 2024-2025

## Introduction

# Motivation - why do we need Geometric Machine Learning?

Face detection. 3D shapes and lack of flatness.




Fig. 10. An example of the system in action. Top row: input polygonal mesh, projected range image and eyes and noses candidates. Second row: a candidate face triangle and the associated range image that was classified as non-face. Third and fourth rows: the two face triangles correctly classified as faces.

[Colombo et al., 2006]

# Motivation - why do we need Geometric Machine Learning?

Facial expressions. Again curvatures, this time in a ML pipeline!




Fig. 2. Various examples of DMCMs after normalization, applied to 3D expressional faces. From top to bottom, expressions are Anger (AN), Disgust (DI), Fear (FE), Happiness (HA), Sadness (SA) and Surprise (SU). From left to the right, images are the original range image, then the DMCMs following sets of radii  $S_1, S_2, S_3, S_4, S_5, S_6$  and  $S_7$  according to section III-B.

[Lemaire et al., 2013]

# Motivation - why do we need Geometric Machine Learning?

Facial expressions. Use convolutions or 3D irregular structures.




Fig. 11. The geometry signatures on the 3D face. Each row shows different expressions of the same subject.

[Jin et al., 2018]

# Motivation - why do we need Geometric Machine Learning?

Facial expressions. Use a manifold embedded in a high dimensional feature space.





Fig. 6. Facial expression recognition result with manifold visualization.

[Chang et al., 2006]

# Motivation - why do we need Geometric Machine Learning?


Molecular Biology. DG-GL. Again curvatures and other feature embedding.



# Motivation - why do we need Geometric Machine Learning?

Pedestrian detection. The manifold of interest is known - space of matrices.

TUZEL ET AL.: PEDESTRIAN DETECTION VIA CLASSIFICATION ON RIEMANNIAN MANIFOLDS



[Tuzel et al., 2008]

# What is a manifold?


- A (**smooth**) **manifold of dimension  $n$**  (Romanian=*varietate diferențiabilă n-dimensională*): topological space  $M$  that ‘locally’ (around each point) is homeomorphic to  $\mathbb{R}^n$  (via maps called **charts**) and such that the overlapping charts induce smooth functions between open sets of  $\mathbb{R}^n$ .

# What is a manifold?


- A (**smooth**) **manifold of dimension  $n$**  (Romanian=*varietate diferențiabilă n-dimensională*): topological space  $M$  that ‘locally’ (around each point) is homeomorphic to  $\mathbb{R}^n$  (via maps called **charts**) and such that the overlapping charts induce smooth functions between open sets of  $\mathbb{R}^n$ .
- A collection on charts covering the manifold is an **atlas**.

# What is a manifold?

- A (**smooth**) **manifold of dimension  $n$**  (Romanian=*varietate diferențiabilă n-dimensională*): topological space  $M$  that ‘locally’ (around each point) is homeomorphic to  $\mathbb{R}^n$  (via maps called **charts**) and such that the overlapping charts induce smooth functions between open sets of  $\mathbb{R}^n$ .
- A collection on charts covering the manifold is an **atlas**.




# Examples (I) Smooth curves



Circle



Ellipse

Parabola



Helix

# Counterexamples



## Examples (II) Smooth surfaces



Sphere



Cylinder



Hyperboloid


Is this a smooth surface?




# How many clusters?



# How many clusters?



# ... or other practical problems



## Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)

# Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)
- The  $n$ -dimensional sphere  $S^n$  (subspace of  $\mathbb{R}^{n+1}$ )

## Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)
- The  $n$ -dimensional sphere  $S^n$  (subspace of  $\mathbb{R}^{n+1}$ )
- Rotation spaces  $\text{SO}(2) = S^1$ ,  $\text{SO}(3)$

## Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)
- The  $n$ -dimensional sphere  $S^n$  (subspace of  $\mathbb{R}^{n+1}$ )
- Rotation spaces  $\text{SO}(2) = S^1$ ,  $\text{SO}(3)$
- Some manifolds arise as hypersurfaces (vanishing locus of a smooth function, e.g.  $xy - 1 = 0$ ), however, not any vanishing locus of a smooth function is a smooth manifold (e.g.  $xy = 0$ )

## Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)
- The  $n$ -dimensional sphere  $S^n$  (subspace of  $\mathbb{R}^{n+1}$ )
- Rotation spaces  $\text{SO}(2) = S^1$ ,  $\text{SO}(3)$
- Some manifolds arise as hypersurfaces (vanishing locus of a smooth function, e.g.  $xy - 1 = 0$ ), however, not any vanishing locus of a smooth function is a smooth manifold (e.g.  $xy = 0$ )
- Parametric representations: there exist  $f_1, \dots, f_d$  smooth (suitable) such that each sample  $\mathbf{x}$  can be represented as  
$$\mathbf{x} = (f_1(u_1, \dots, u_p), \dots, f_d(u_1, \dots, u_p)),$$
 for suitably chosen parameters  $u_1, \dots, u_p$

## Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)
- The  $n$ -dimensional sphere  $S^n$  (subspace of  $\mathbb{R}^{n+1}$ )
- Rotation spaces  $\text{SO}(2) = S^1$ ,  $\text{SO}(3)$
- Some manifolds arise as hypersurfaces (vanishing locus of a smooth function, e.g.  $xy - 1 = 0$ ), however, not any vanishing locus of a smooth function is a smooth manifold (e.g.  $xy = 0$ )
- Parametric representations: there exist  $f_1, \dots, f_d$  smooth (suitable) such that each sample  $\mathbf{x}$  can be represented as  
$$\mathbf{x} = (f_1(u_1, \dots, u_p), \dots, f_d(u_1, \dots, u_p)),$$
 for suitably chosen parameters  $u_1, \dots, u_p$
- The embedding of a manifold in an Euclidean space  $\mathbb{R}^d$  is important. The sphere  $S^2$ , the cylinder, the hyperboloid can be embedded in  $\mathbb{R}^3$ . Examples of surfaces that cannot be embedded in  $\mathbb{R}^3$ : **the Möbius strip, the Klein bottle.**

## Examples of manifolds

- The space  $\mathbb{R}^n$  (Euclidean space)
- The  $n$ -dimensional sphere  $S^n$  (subspace of  $\mathbb{R}^{n+1}$ )
- Rotation spaces  $\text{SO}(2) = S^1$ ,  $\text{SO}(3)$
- Some manifolds arise as hypersurfaces (vanishing locus of a smooth function, e.g.  $xy - 1 = 0$ ), however, not any vanishing locus of a smooth function is a smooth manifold (e.g.  $xy = 0$ )
- Parametric representations: there exist  $f_1, \dots, f_d$  smooth (suitable) such that each sample  $\mathbf{x}$  can be represented as  
$$\mathbf{x} = (f_1(u_1, \dots, u_p), \dots, f_d(u_1, \dots, u_p)),$$
 for suitably chosen parameters  $u_1, \dots, u_p$
- The embedding of a manifold in an Euclidean space  $\mathbb{R}^d$  is important. The sphere  $S^2$ , the cylinder, the hyperboloid can be embedded in  $\mathbb{R}^3$ . Examples of surfaces that cannot be embedded in  $\mathbb{R}^3$ : **the Möbius strip, the Klein bottle.**
- ... etc.

# Distance computations, vectors, inner product

- **Q:** What are the keys of the distance computations?

# Distance computations, vectors, inner product

- **Q:** What are the keys of the distance computations?

- In  $\mathbb{R}^d$  one has

$$d(A, B)^2 = \| \overrightarrow{AB} \| = \langle \overrightarrow{AB}, \overrightarrow{AB} \rangle = \overrightarrow{AB} \cdot \overrightarrow{AB}$$

# Distance computations, vectors, inner product

- **Q:** What are the keys of the distance computations?

- In  $\mathbb{R}^d$  one has

$$d(A, B)^2 = \| \overrightarrow{AB} \| = \langle \overrightarrow{AB}, \overrightarrow{AB} \rangle = \overrightarrow{AB} \cdot \overrightarrow{AB}$$

- Recall that for  $x, y \in \mathbb{R}^d$  one has  $\langle x, y \rangle = \sum_{i=1}^d x_i y_i$ . More general, one can consider **inner product**

$$\langle x, y \rangle = \sum_{i,j=1}^d g_{ij} x_i y_j,$$

given by a matrix  $(g_{ij})$  that satisfies several conditions (symmetric, positive definite).

# Distance computations, vectors, inner product

- **Q:** What are the keys of the distance computations?

- In  $\mathbb{R}^d$  one has

$$d(A, B)^2 = \| \overrightarrow{AB} \| = \langle \overrightarrow{AB}, \overrightarrow{AB} \rangle = \overrightarrow{AB} \cdot \overrightarrow{AB}$$

- Recall that for  $x, y \in \mathbb{R}^d$  one has  $\langle x, y \rangle = \sum_{i=1}^d x_i y_i$ . More general, one can consider **inner product**

$$\langle x, y \rangle = \sum_{i,j=1}^d g_{ij} x_i y_j,$$

given by a matrix  $(g_{ij})$  that satisfies several conditions (symmetric, positive definite).

- In general, an inner product allows you to compute **distances** and **angles**.

# Distance computations, vectors, inner product

- **Q:** What are the keys of the distance computations?

- In  $\mathbb{R}^d$  one has

$$d(A, B)^2 = \| \overrightarrow{AB} \| = \langle \overrightarrow{AB}, \overrightarrow{AB} \rangle = \overrightarrow{AB} \cdot \overrightarrow{AB}$$

- Recall that for  $x, y \in \mathbb{R}^d$  one has  $\langle x, y \rangle = \sum_{i=1}^d x_i y_i$ . More general, one can consider **inner product**

$$\langle x, y \rangle = \sum_{i,j=1}^d g_{ij} x_i y_j,$$

given by a matrix  $(g_{ij})$  that satisfies several conditions (symmetric, positive definite).

- In general, an inner product allows you to compute **distances** and **angles**.
- **A:** The vector space structure and the inner product.

# Distance computations, vectors, inner product

- **Q:** What are the keys of the distance computations?

- In  $\mathbb{R}^d$  one has

$$d(A, B)^2 = \| \overrightarrow{AB} \| = \langle \overrightarrow{AB}, \overrightarrow{AB} \rangle = \overrightarrow{AB} \cdot \overrightarrow{AB}$$

- Recall that for  $x, y \in \mathbb{R}^d$  one has  $\langle x, y \rangle = \sum_{i=1}^d x_i y_i$ . More general, one can consider **inner product**

$$\langle x, y \rangle = \sum_{i,j=1}^d g_{ij} x_i y_j,$$

given by a matrix  $(g_{ij})$  that satisfies several conditions (symmetric, positive definite).


- In general, an inner product allows you to compute **distances** and **angles**.
- **A:** The vector space structure and the inner product.
- If a manifold is embedded in  $\mathbb{R}^d$ , although one may consider vectors with origin at a given point, the end point of the vector might not be situated on the manifold.

## Tangent vectors, tangent space

- Let  $M$  be a  $n$ -dimensional manifold. For each point  $x \in M$ , one can define a vector space of dimension  $n$ , called the tangent space at  $M$  in  $x$  (usually denoted by  $T_x M$ ).

# Tangent vectors, tangent space

- Let  $M$  be a  $n$ -dimensional manifold. For each point  $x \in M$ , one can define a vector space of dimension  $n$ , called the tangent space at  $M$  in  $x$  (usually denoted by  $T_x M$ ).
- Examples: (i)  $\mathbb{R}^d$  (in this case for each point the tangent space is  $\mathbb{R}^d$  itself), (ii) the circle, (iii) the sphere



A point  $P$  on the sphere and the tangent plane to the sphere at  $P$ .

## Distance measurement on manifolds: Riemannian metrics


- To give a Riemannian metric  $g$  on a manifold means to indicate an inner product in each tangent space (to each point), such that certain compatibility conditions are verified. A Riemannian manifold  $(M, g)$  is a manifold endowed with a Riemannian metric

## Distance measurement on manifolds: Riemannian metrics

- ▶ To give a Riemannian metric  $g$  on a manifold means to indicate an inner product in each tangent space (to each point), such that certain compatibility conditions are verified. A Riemannian manifold  $(M, g)$  is a manifold endowed with a Riemannian metric
- ▶ The same space can be endowed with different metrics!


# Distance measurement on manifolds: Riemannian metrics

- ▶ To give a Riemannian metric  $g$  on a manifold means to indicate an inner product in each tangent space (to each point), such that certain compatibility conditions are verified. A Riemannian manifold  $(M, g)$  is a manifold endowed with a Riemannian metric
- ▶ The same space can be endowed with different metrics!
- ▶ “Hierarchy”: topological space / smooth manifold / Riemannian manifold




# Distance measurement on manifolds: Riemannian metrics

- ▶ To give a Riemannian metric  $g$  on a manifold means to indicate an inner product in each tangent space (to each point), such that certain compatibility conditions are verified. A Riemannian manifold  $(M, g)$  is a manifold endowed with a Riemannian metric
- ▶ The same space can be endowed with different metrics!
- ▶ “Hierarchy”: topological space / smooth manifold / Riemannian manifold




- ▶ Metrics on a circle (sphere): the chordal distance and the arc (geodesic) distance



# Distance measurement on manifolds: geodesics, exponential and logarithm

- Given a Riemannian manifold  $(M, g)$ , take  $x \in M$  arbitrary and  $v \in T_x M$ . There exists a unique **geodesic**  $\gamma_v$  such that  $\gamma_v(0) = x$  and  $\gamma'_v(0) = v$ . There exists a map, called **exponential map** (it is defined locally!),  $\exp_x : T_x M \rightarrow M$ , such that  $\exp_x(v) = \gamma_v(1)$ . Its inverse is called **logarithm**. A geodesic is, locally, a path of shortest length.




The exponential and the logarithm map are locally defined and they are inverse to each other.

# Examples

- Geodesics of the Euclidean space  $\mathbb{R}^d$  are line segments


## Examples

- Geodesics of the Euclidean space  $\mathbb{R}^d$  are line segments
- Geodesics of the sphere  $S^2$  are the great circles. Are the geodesics unique?



## Examples

- Geodesics of the Euclidean space  $\mathbb{R}^d$  are line segments
- Geodesics of the sphere  $S^2$  are the great circles. Are the geodesics unique?



- Given two points on a cylinder: how can one construct a geodesic?



# The groups $SO(3)$ , $E(3)$ and movements

Structure from motion. (Relative) rotations can be modeled by using the group  $SO(3)$  (rotations in the 3-dimensional space), which also has a (natural) structure of 3-dimensional manifold. Could you describe the group  $SO(2)$  (plane rotations)?



(a)



(b)



(c)

Source: [Özyeşil et al., 2017, cf. Snavely et al., 2006], see also Hartley

# SPD-matrices

SPD (Symmetric positive definite) matrices

$$SPD(n) = \{S \in \mathcal{M}_n(\mathbb{R}) | S^T = S, \forall x \neq 0, x^T S x > 0\}.$$




Fig. 13. The brain regions used in the evaluation of the default mode network. a) The brain network includes the superior frontal gyrus (SFG), medial orbitofrontal cortex (MOF), parahippocampus (PHC), precuneus (PrC) and inferior parietal lobe (IPL) according to the Desikan-Killiany atlas [Desikan et al., 2006]. The brain regions in the left hemisphere are presented. The first letters "l" and "r" in each region's name indicate left and right hemisphere. b) The geometric average of a group of 734 functional connectivity matrices (FC) is presented. All square pixels in the connectivity matrix indicate edges among the brain regions. Dots in (b) indicate edges that show 5% differences between Euclidean average and geometric average of group connectivity matrices. c) Functional connectivity matrices of 30 exemplary subjects are displayed.

Source: [You and Park, 2021]

# Reference

[Miolane et al., 2020] and the references therein. This work is the reference paper for the `geomstats` package.