

Quantum Field Theory II

派蒙の素论派学习笔记

Author Be Like:



2026 年 6 月 26 日

University of Hamburg
Department of Physics

目录

1	Path integral method	5
1.	Path integral in quantum mechanics	5
1.1	Functional method	5
1.2	Generalization to quantum system with coordinates $\{q_i\}$ and conjugate momenta $\{p_i\}$	7
2.	Path integral in the Scalar field	8
2.1	Correlation Function	9
2.2	Feynman Rules	11
2.3	Generating Functional	13
2.4	Feynman Rules for the Free Theory	14
2.5	Interaction theory: ϕ^4 theory as an example	15
2.6	Other Type of Generating functional	17
3.	Quantization of EM Field in Path Integral Formalism	18
3.1	Faddeev-Popov Quantization	18
3.2	Gauge Fixing and the Photon Propagator	20
3.3	Standard Gauge Choices	21
4.	Path Integral for Fermions	21
4.1	Grassmann Numbers and Algebra	21
4.2	Berezin Integration	22
4.3	Gaussian Integrals and Determinants	22
4.4	Transformation of Variables	22
4.5	Comparison: Bosonic vs. Fermionic Case	23
5.	Path Integral Formulation for Dirac Spinors	23
5.1	Fermion Propagator	23
5.2	Generating Functional for Fermion Fields	23
5.3	Derivation of the Propagator	24
6.	Symmetries in Path Integral Formalism	24
6.1	Schwinger-Dyson Equations	24
6.2	Conservation Laws and Noether's Theorem	26
6.3	Ward-Takahashi Identity in QED	28
2	Renormalization	29
1.	Renormalization in ϕ^3 -theory	29
1.1	One-loop self-energy	29
1.2	Wick rotation	29

1.3	Degree of divergence	36
1.4	Arbitrariness in a renormalization graph	38
1.5	Dimensional regularization	39
1.6	Massless theories	45
1.7	Differentiation with respect to external momenta	49
1.8	Composite operators	51
1.9	Renormalization of composite operators	54
2.	Renormalization in ϕ^4 -theory	58
2.1	Analysis of dimensions in view of renormalization	59
2.2	Renormalized perturbation theory	60
3.	Wilson's approach to renormalization	63
4.	Callan-Symanzik Equation	67
4.1	Alternative choice for renormalization in ϕ^4 -theory	67
4.2	β -function in massless ϕ^4 -theory	69
4.3	Analogous analysis for massless QED	70
5.	Evolution of coupling	72
5.1	Solution of Callan-Symanzik equation	72
5.2	Renormalization group equation	74
5.3	Higgs mass and fate of universe	75
6.	Renormalization for local operators and evolution of mass parameter	77
6.1	Renormalized Green's functions with local operator $\mathcal{O}$ in ϕ^4 -theory . . .	78
6.2	Renormalization for local operators in massless ϕ^4 -theory	79
6.3	Evolution of mass parameter	80
3	Non-Abelian gauge theories	83
1.	Recap of gauge theory and brief look at Lie group theory	83
1.1	Recap: Abelian gauge theory	83
1.2	SU(N) gauge theory	84
1.3	Recap for essential Lie group theory	85
1.4	Representations of Lie algebra	88
1.5	Inner product of generators	90
1.6	Representation of Lorentz group	92
1.7	Integration over gauge group	93
2.	Quantization of non-Abelian gauge theory	94
2.1	Preparation	94
2.2	Quantization by path integral	95
2.3	Feynman rules for non-Abelian gauge theory	99
3.	BRST Symmetry	100
3.1	Problem of negative norm	100
3.2	Infinitesimal BRST transformation	101
3.3	Factorization of Hilbert space via BRST operator	105
4.	Renormalization of non-Abelian gauge theory	107
4.1	Why the Faddeev–Popov theory is renormalizable	108

4.2	Counter terms of Faddeev–Popov Lagrangian	118
4.3	One-loop structure of non-Abelian gauge theory	120
4.4	Callan–Symanzik equation in non-Abelian gauge theory	131
5.	Quantum Chromodynamics: QCD	134
5.1	What QCD is	135
5.2	Why color is needed	135
5.3	No free quarks and confinement	138
5.4	Renormalization constants and the running coupling	139
5.5	Solving the one-loop running explicitly	140
5.6	The basic idea of a conformal window	141
5.7	Classical running in dimensional regularization	142
5.8	Heavy-quark thresholds and matching of the running coupling	142
5.9	Cross-section predictions and infrared safety	144
4	Standard Model	147
1.	Spontaneous Symmetry Breaking	147
1.1	A real scalar with a discrete symmetry	147
1.2	A complex scalar and global $U(1)$ breaking	148
1.3	Expansion around the chosen vacuum	150
1.4	N complex scalar fields	151
1.5	Goldstone theorem	152
1.6	Proof by the mass matrix	153
2.	The Higgs Mechanism	154
2.1	Difference between Higgs mechanism and Goldstone’s theorem	154
2.2	The Abelian Higgs model	155
2.3	The same physics in Cartesian fields	157
2.4	What the propagator is telling us	158
2.5	$SU(2)$ gauge symmetry breaking	160

Path integral method

1. Path integral in quantum mechanics

1.1 Functional method

Functional S maps a function to numbers, i.e.

$$S : x(t) \rightarrow S[x(t)]. \quad (1.1)$$

As we all know, the Hamilton operator in 1-dim is $H = \frac{p^2}{2m} + V(x)$, now let us consider the amplitude for motion of a particle from x_a to x_b . For this, we introduce the evolution operator $U(x_a, x_b; t)$ in Heisenburg picture, so we have

$$U(x_a, x_b; T) = \langle x_b | e^{-iHT/\hbar} | x_a \rangle, \quad (1.2)$$

where the Hamiltonian **is already quantized**, but in the following calculation, the Hamiltonian that in the path integral **is classical** due to the path integral itself is the process of quantization. **So we don't need to quantize anything inside of the integration.** So the path integral is directly defined from this operator. The basic idea is to cut the space and time into small pieces, and add them together, i.e.,

$$U(x_a, x_b; T) = \int \mathcal{D}x(t) e^{iS[x(t)]/\hbar}, \quad (1.3)$$

where $S[x(t)]$ is the action that depends on the paths. In QM, we can expand it as

$$S[x(t)] = \int_0^T dt L = \int_0^T dt \left[\frac{1}{2} m v^2 - V(x) \right]. \quad (1.4)$$

And the integration measure $\mathcal{D}x(t)$ accounts for all possible paths from x_a to x_b , i.e.,


$$\sum_{\substack{\text{all paths from} \\ x_a \text{ to } x_b}} \longrightarrow \int \mathcal{D}x(t). \quad (1.5)$$

Now we need to prove that the definition we gave that can derive the same quantization result. Firstly, we discretize the time interval from 0 to T in steps $\Delta t = \varepsilon$:

$$S = \int_0^T dt \left[\frac{1}{2} m \dot{x}^2 - V(x) \right] = \sum_k \left\{ \frac{m}{2} \frac{(x_{k+1} - x_k)^2}{\varepsilon} - \varepsilon V \left(\frac{x_{k+1} + x_k}{2} \right) \right\}. \quad (1.6)$$

The path integral build up from N discrete time intervals

$$\mathcal{D}x(t) \equiv \frac{1}{c(\varepsilon)} \int \frac{dx_1}{c(\varepsilon)} \frac{dx_2}{c(\varepsilon)} \cdots \frac{dx_N}{c(\varepsilon)} = \frac{1}{c(\varepsilon)} \prod_{k=1}^N \int_{-\infty}^{+\infty} \frac{dx_k}{c(\varepsilon)}, \quad (1.7)$$



where $\frac{1}{c(\varepsilon)}$ is an undetermined normalization factor. Now we consider the evolution operator $U(x_a, x_b; T)$ evolve from $T - \varepsilon$ to T in the interval $[x_a, x_b]$, which is the last step in (1.7):

$$\begin{aligned} U(x_a, x_b; T) &= \langle x_b | e^{-iH[(T-\varepsilon)+\varepsilon]/\hbar} | x_a \rangle = \int dx' \langle x_b | e^{-iH\varepsilon/\hbar} | x' \rangle \langle x' | e^{-iH(T-\varepsilon)/\hbar} | x_a \rangle \\ &= \int_{-\infty}^{+\infty} \frac{dx'}{c(\varepsilon)} \exp \left[\frac{i}{\hbar} \frac{m(x_b - x')^2}{2\varepsilon} - \frac{i}{\hbar} \varepsilon V \left(\frac{x_b + x'}{2} \right) \right] U(x_a, x'; T - \varepsilon) \end{aligned} \quad (1.8)$$

Notice that we can expand $U(x_a, x'; T - \varepsilon)$ around x_b

$$U(x_a, x'; T - \varepsilon) = \left(1 + (x' - x_b) \frac{\partial}{\partial x_b} + \frac{1}{2} (x' - x_b)^2 \frac{\partial^2}{\partial x_b^2} + \dots \right) U(x_a, x_b; T - \varepsilon) \quad (1.9)$$

and for $\varepsilon \rightarrow 0$, the potential term can also expand as

$$\exp \left\{ \frac{i}{\hbar} \varepsilon V \left(\frac{x_b - x_{x'}}{2} \right) \right\} \stackrel{x' \rightarrow x_b}{=} e^{\frac{i}{\hbar} \varepsilon V(x_b)} = 1 - \frac{i}{\hbar} \varepsilon V(x_b) + \dots \quad (1.10)$$

Thus, (1.9) can be rewritten as

$$\begin{aligned} U(x_a, x_b; T) &= \int_{-\infty}^{+\infty} \frac{dx'}{c(\varepsilon)} \exp \left\{ \frac{i}{\hbar} \frac{m(x_b - x')^2}{\varepsilon} \right\} \left[1 - \frac{i\varepsilon}{\hbar} V(x_b) + \dots \right] \\ &\quad \times \left[1 + (x' - x_b) \frac{\partial}{\partial x_b} + \frac{1}{2} (x' - x_b)^2 \frac{\partial^2}{\partial x_b^2} + \dots \right] U(x_a, x_b; T - \varepsilon) \end{aligned} \quad (1.11)$$

To do this integration the Gauss integration formulas are very useful in the next:

$$\int_{-\infty}^{+\infty} d\xi e^{-b\xi^2} = \sqrt{\frac{\pi}{b}}; \quad (1.12)$$

$$\int_{-\infty}^{+\infty} d\xi \xi e^{-b\xi^2} = 0; \quad (1.13)$$

$$\int_{-\infty}^{+\infty} d\xi \xi^2 e^{-b\xi^2} = \frac{1}{2b} \sqrt{\frac{\pi}{b}}. \quad (1.14)$$

Thus,

$$U(x_a, x_b; T) = \frac{1}{c(\varepsilon)} \sqrt{\frac{\pi}{b}} \left[1 - \frac{i\varepsilon}{\hbar} V(x_b) + \frac{1}{4b} \frac{\partial^2}{\partial x_b^2} + \mathcal{O}(\varepsilon^2) \right] U(x_a, x_b; T - \varepsilon), \quad (1.15)$$

where $b = \frac{-im}{2\varepsilon\hbar}$. When $\varepsilon \rightarrow 0$,

$$\begin{aligned} U(x_a, x_b; T) &= \overbrace{\frac{1}{c(\varepsilon)} \sqrt{\frac{\pi}{b}}}^{=1} U(x_a, x_b; T - \varepsilon) \\ \Rightarrow c(\varepsilon) &= \sqrt{\frac{\pi}{b}}. \end{aligned} \quad (1.16)$$

So we have the Schrödinger equation for the evolution operator

$$i\hbar \frac{U(x_a, x_b; T) - U(x_a, x_b; T - \varepsilon)}{\varepsilon} = \left(V(x_b) - \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_b^2} \right) U(x_a, x_b; T - \varepsilon). \quad (1.17)$$

So we have derived the familiar differential equation with the canonical quantization did. Now we also need to consider the boundary condition. For $T \rightarrow 0$, canonical quantization requires $U(x_a, x_b; 0) = \langle x_b | \mathbb{1} | x_a \rangle \delta(x_a - x_b)$. For path integral, we firstly take a piece of time slice, since $T \rightarrow 0$ there is no difference for labeling the space interval, so we can just take $\delta x = x_b - x_a$:

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{c(\varepsilon)} \exp \left[\frac{im}{\hbar} \frac{(x_b - x_a)^2}{\varepsilon} \right] = \delta(x_b - x_a) \quad (1.18)$$

Thus,

$$\lim_{\varepsilon \rightarrow 0} \int \mathcal{D}x(t) e^{iS[x(t)]/\hbar} = \lim_{\varepsilon \rightarrow 0} \frac{1}{c(\varepsilon)} \exp \left[\frac{im}{\hbar} \frac{(x_b - x_a)^2}{\varepsilon} \right] \rightarrow \delta(x_a - x_b) \quad (1.19)$$

- Path integral formulas quantize system through summation/integration over all possible paths; weighted by the phase $e^{-iS[x(t)]/\hbar}$.
- In the limit $\hbar \rightarrow 0$ (classical physics) path integral is dominated by classical path with $\delta S = 0$.

1.2 Generalization to quantum system with coordinates $\{q_i\}$ and conjugate momenta $\{p_i\}$

Now we consider the general case for a quantum system with Hamiltonian $H(q, p)$, and evolution operator $U(q_a, q_b; T) = \langle q_b | e^{-iHT} | q_a \rangle$. Split up $f(q, p) = e^{-iHT}$ in N time intervals $\Delta t = \varepsilon$

$$e^{-iHT} = \prod_{k=1}^N e^{-iH\varepsilon} = e^{-iH\varepsilon} \mathbb{1} e^{-iH\varepsilon} \mathbb{1} \dots \mathbb{1} e^{-iH\varepsilon}, \quad (1.20)$$

with $\mathbb{1} = \left(\prod_i \int dq_k^i \right) |q_k\rangle \langle q_k|$. At k -th position,

$$\langle q_{k+1} | f(q) | q_k \rangle \xrightarrow{\varepsilon=0} \langle q_{k+1} | 1 - i\varepsilon H + \mathcal{O}(\varepsilon^2) | q_k \rangle \quad (1.21)$$

For a function that its variable is only q , we have

$$\langle q_{k+1} | f(q) | q_k \rangle = f(q_k) \prod_i \delta(q_k^i - q_{k+1}^i) = f\left(\frac{q_k + q_{k+1}}{2}\right) \left(\prod_i \int \frac{dp_i}{2\pi} \right) \exp \left\{ i \sum_i p_k^i (q_{k+1}^i - q_k^i) \right\}, \quad (1.22)$$

and for a function that its variable is only p , we have

$$\langle q_{k+1} | f(p) | q_k \rangle = \left(\prod_i \int \frac{dp_k^i}{2\pi} \right) f(p_k) \exp \left\{ i \sum_i p_k^i (q_{k+1}^i - q_k^i) \right\}. \quad (1.23)$$

In general, we normally face the more complicated case that using the Weyl ordering is convenient to keep the properties we have in the above, i.e.,

$$\langle q_{k+1} | \frac{1}{4} (q^2 p^2 + 2qp^2 q + p^2 q^2) | q_k \rangle = \left(\frac{q_{k+1} + q_k}{2} \right)^2 \langle q_k + 1 | p^2 | q_k \rangle, \quad (1.24)$$

where the Weyl order is defined as

$$W(p, q) = \frac{1}{2} \{x, p\}. \quad (1.25)$$

Assume $H(p, q)$ is Weyl ordered, i.e.,

$$\langle q_{k+1} | H(p, q) | q_k \rangle = \left(\prod_i \int \frac{dp_k^i}{2\pi} \right) H \left(\frac{q_{k+1} + q_k}{2}, p_k \right) \exp \left\{ i \sum_i p_k^i (q_{k+1}^i - q_k^i) \right\}. \quad (1.26)$$

Thus

$$\langle q_{k+1} | e^{-iH\varepsilon} | q_k \rangle = \left(\prod_i \int \frac{dp_k^i}{2\pi} \right) \exp \left\{ -i\varepsilon H \left(\frac{q_{k+1} + q_k}{2}, p_k \right) \right\} \exp \left\{ i \sum_i p_k^i (q_{k+1}^i - q_k^i) \right\}. \quad (1.27)$$

Take $q_0 = q_a$ and $q_N = q_b$, the integration are

$$U(q_0, q_N; T) = \left(\prod_{i,k} \int dq_k^i \int \frac{dp_k^i}{2\pi} \right) \exp \left\{ i \sum_k \left[\sum_i p_k^i (q_{k+1}^i - q_k^i) - \varepsilon H \left(\frac{q_{k+1} + q_k}{2}, p_k \right) \right] \right\}. \quad (1.28)$$

In path integral notation

$$U(q_0, q_N; T) = \left(\prod_{i,k} \int \mathcal{D}q^i(t) \int \mathcal{D}p^i(t) \right) \exp \left\{ i \int_0^T dt \left[\sum_i p^i \dot{q}^i - H(p, q) \right] \right\}. \quad (1.29)$$

2. Path integral in the Scalar field

For a scalar field $\phi(\vec{x}, t)$, we can simply change the variables $\vec{q}_i(t) \mapsto \phi(\vec{x}, t)$, the evolution operator is then became:

$$U(\phi_a, \phi_b; T) = \langle \phi_b(\vec{x}) | e^{-i\hat{H}T} | \phi_a(\vec{x}) \rangle. \quad (1.30)$$

The action S and Hamiltonian H for the scalar field are defined as:

$$S = \int d^4x \left(\frac{1}{2} (\partial \cdot \phi)^2 - V(\phi) \right), \quad (1.31)$$

$$H = \int d^3x \left(\frac{1}{2} \pi^2 + \frac{1}{2} (\nabla \phi)^2 + V(\phi) \right). \quad (1.32)$$

The path integral form can then be simply seen from (1.29) as

$$\langle \phi_b(\vec{x}) | e^{-i\hat{H}T} | \phi_a(\vec{x}) \rangle = \int_{\phi(0, \vec{x})=\phi_a}^{\phi(T, \vec{x})=\phi_b} \mathcal{D}\phi(x) \int \mathcal{D}\pi(x) \exp \left\{ i \int_0^T dt \int d^3x \left(\pi \dot{\phi} - \frac{1}{2} \pi^2 - \frac{1}{2} (\nabla \phi)^2 - V(\phi) \right) \right\} \quad (1.33)$$

$$= \int \mathcal{D}\phi(x) \int \mathcal{D}(\pi(x) - \dot{\phi}) \exp \left\{ i \int_0^T dt \int d^3x \left[-\frac{1}{2} (-2\pi \dot{\phi} + \pi^2 + \dot{\phi}^2 - \dot{\phi}^2) - \frac{1}{2} (\nabla \phi)^2 - V(\phi) \right] \right\} \quad (1.34)$$

To cancel the $\int \mathcal{D}\pi(x)$, we change the variable $\mathcal{D}\pi(x) \rightarrow \mathcal{D}(\pi(x) - \partial_0\phi)$, i.e.,

$$\langle \phi_b(\vec{x}) | e^{-i\hat{H}T} | \phi_a(\vec{x}) \rangle = \int \mathcal{D}\phi(x) \int \mathcal{D}(\pi(x) - \dot{\phi}) \exp \left\{ i \int_0^T dt \int d^3x \left[-\frac{1}{2} (\pi - \dot{\phi})^2 + \frac{1}{2} (\partial\phi)^2 - V(\phi) \right] \right\} \quad (1.35)$$

$$= \int \mathcal{D}\phi(x) \int \mathcal{D}\tilde{\pi}(x) \exp \left\{ i \int_0^T dt \int d^3x \left[-\frac{1}{2} \tilde{\pi}(x)^2 + \frac{1}{2} (\partial\phi)^2 - V(\phi) \right] \right\}. \quad (1.36)$$

Thus

$$\langle \phi_b(\vec{x}) | e^{-i\hat{H}T} | \phi_a(\vec{x}) \rangle = \mathcal{C} \int \mathcal{D}\phi(x) \exp \left\{ i \int_0^T dt \int d^3x \left[\frac{1}{2} (\partial\phi)^2 - V(\phi) \right] \right\}, \quad (1.37)$$

where

$$\mathcal{C} = \int \mathcal{D}\tilde{\pi}(x) \exp \left\{ -i \int_0^T dt \int d^3x \frac{1}{2} \tilde{\pi}(x)^2 \right\} \quad (1.38)$$

So we can directly see it is

$$\langle \phi_b(\vec{x}) | e^{-i\hat{H}T} | \phi_a(\vec{x}) \rangle = \mathcal{C} \int_{\phi(0,\vec{x})=\phi_a}^{\phi(T,\vec{x})=\phi_b} \mathcal{D}\phi(x) \exp \left\{ i \int d^4x \mathcal{L} \right\} \quad (1.39)$$

For some reason that we will consider later, the normalization constant $\mathcal{C}$ is ignored in the next calculation.

2.1 Correlation Function

Now we consider the two-point correlation function. For that, we firstly consider the structure

$$\int_{\phi(-T,\vec{x})=\phi_a}^{\phi(T,\vec{x})=\phi_b} \mathcal{D}\phi(x) \phi(x_1) \phi(x_2) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\}, \quad (1.40)$$

where we assume $x_2^0 > x_1^0$. Now the boundary condition changes from $-T$ to T . So we can separate the integration in three area $-T \rightarrow x_1^0$, $x_1^0 \rightarrow x_2^0$ and $x_2^0 \rightarrow T$ such that we can fix the field at x_1^0 and x_2^0 as $\phi_1(\vec{x})$ and $\phi_2(\vec{x})$ respectively. This operation means no matter how the field evolve from $-T$ to T , we can always separate the integration with 2 steps: First, fixing the field at these two time point and integrate the three area. Second, we integrate all possible situation of these two field.

$$\int \mathcal{D}\phi(x) = \int \mathcal{D}\phi_1(\vec{x}) \int \mathcal{D}\phi_2(\vec{x}) \int_{\substack{\phi(x_1^0,\vec{x})=\phi_1(\vec{x}) \\ \phi(x_2^0,\vec{x})=\phi_2(\vec{x})}} \mathcal{D}\phi(x). \quad (1.41)$$

So we have

$$\begin{aligned} & \int_{\phi(-T,\vec{x})=\phi_a}^{\phi(T,\vec{x})=\phi_b} \mathcal{D}\phi(x) \phi(x_1) \phi(x_2) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\} \\ &= \int \mathcal{D}\phi_1(\vec{x}) \int \mathcal{D}\phi_2(\vec{x}) \int \mathcal{D}\phi(x) \phi(x_1) \phi(x_2) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\}. \end{aligned} \quad (1.42)$$

Because we have fixed the time point x_1^0 and x_2^0 , so $\phi(x_1)$ and $\phi(x_2)$ can be directly picked from the integration $\int \mathcal{D}\phi$ and becomes $\phi_1(\vec{x}_1)$ and $\phi_2(\vec{x}_2)$, i.e.,

$$\begin{aligned} & \int_{\phi(-T, \vec{x})=\phi_a}^{\phi(T, \vec{x})=\phi_b} \mathcal{D}\phi(x) \phi(x_1) \phi(x_2) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\} \\ &= \int \mathcal{D}\phi_1(\vec{x}) \int \mathcal{D}\phi_2(\vec{x}) \phi_1(\vec{x}_1) \phi_2(\vec{x}_2) \left[\int \mathcal{D}\phi(x) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\} \right] \end{aligned} \quad (1.43)$$

$$= \int \mathcal{D}\phi_1(\vec{x}) \int \mathcal{D}\phi_2(\vec{x}) \phi_1(\vec{x}_1) \phi_2(\vec{x}_2) \left[\langle \phi_b | e^{-i\hat{H}(2T)} | \phi_a \rangle \right] \quad (1.44)$$

$$= \int \mathcal{D}\phi_1(\vec{x}) \int \mathcal{D}\phi_2(\vec{x}) \phi_1(\vec{x}_1) \phi_2(\vec{x}_2) \langle \phi_b | e^{-i\hat{H}(T-x_2^0)} | \phi_2 \rangle \langle \phi_2 | e^{-i\hat{H}(x_2^0-x_1^0)} | \phi_1 \rangle \langle \phi_1 | e^{-i\hat{H}(x_1^0+T)} | \phi_a \rangle. \quad (1.45)$$

Using $\hat{\phi}_S(\vec{x})|\phi_i\rangle = \phi_i(\vec{x})|\phi_i\rangle$ and $\int \mathcal{D}\phi_i |\phi_i\rangle \langle \phi_i| = \mathbb{1}$:

$$\int_{\phi(-T, \vec{x})=\phi_a}^{\phi(T, \vec{x})=\phi_b} \mathcal{D}\phi(x) \phi(x_1) \phi(x_2) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\} = \langle \phi_b | e^{-i\hat{H}(T-x_2^0)} \hat{\phi}_S(\vec{x}_2) e^{-i\hat{H}(x_2^0-x_1^0)} \hat{\phi}_S(\vec{x}_1) e^{-i\hat{H}(x_1^0+T)} | \phi_a \rangle. \quad (1.46)$$

Defining the Heisenberg picture operator $\hat{\phi}_H(x) = e^{i\hat{H}x^0} \hat{\phi}_S(\vec{x}) e^{-i\hat{H}x^0}$:

$$\int_{\phi(-T, \vec{x})=\phi_a}^{\phi(T, \vec{x})=\phi_b} \mathcal{D}\phi(x) \phi(x_1) \phi(x_2) \exp \left\{ i \int_{-T}^T d^4x \mathcal{L} \right\} = \langle \phi_b | e^{-i\hat{H}T} \hat{\phi}_H(x_2) \hat{\phi}_H(x_1) e^{-i\hat{H}T} | \phi_a \rangle. \quad (1.47)$$

For a general time ordering $x_2^0 > x_1^0$, this becomes the time-ordered product:

$$\langle \phi_b | e^{-i\hat{H}T} T \{ \hat{\phi}_H(x_1) \hat{\phi}_H(x_2) \} e^{-i\hat{H}T} | \phi_a \rangle. \quad (1.48)$$

Using the ground state projection

$$\lim_{T \rightarrow \infty(1-i\epsilon)} e^{-i\hat{H}T} | \phi_a \rangle = \lim_{T \rightarrow \infty(1-i\epsilon)} \sum_n e^{-iE_n T} | n \rangle \langle n | \phi_a \rangle = \langle \Omega | \phi_a \rangle e^{-iE_0 \infty(1-i\epsilon)} | \Omega \rangle, \quad (1.49)$$

and similarly,

$$\lim_{T \rightarrow \infty(1-i\epsilon)} \langle \phi_b | e^{-i\hat{H}T} = \lim_{T \rightarrow \infty(1-i\epsilon)} \sum_n \langle \phi_b | n \rangle \langle n | e^{-iE_n T} = \langle \phi_b | \Omega \rangle \langle \Omega | e^{-iE_0 \infty(1-i\epsilon)}. \quad (1.50)$$

So we have

$$\langle \phi_b | e^{-i\hat{H}T} T \{ \hat{\phi}_H(x_1) \hat{\phi}_H(x_2) \} e^{-i\hat{H}T} | \phi_a \rangle = \langle \phi_b | \phi_a \rangle \langle \Omega | T \{ \hat{\phi}_H(x_1) \hat{\phi}_H(x_2) \} | \Omega \rangle. \quad (1.51)$$

Thus, we can finally reach the correlation function:

$$\langle \Omega | T \{ \hat{\phi}_H(x_1) \hat{\phi}_H(x_2) \} | \Omega \rangle = \frac{\langle \phi_b | e^{-i\hat{H}T} T \{ \hat{\phi}_H(x_1) \hat{\phi}_H(x_2) \} e^{-i\hat{H}T} | \phi_a \rangle}{\langle \phi_b | \phi_a \rangle}. \quad (1.52)$$

So in the end, we have

$$\langle \Omega | T \{ \hat{\phi}_H(x_1) \hat{\phi}_H(x_2) \} | \Omega \rangle = \lim_{T \rightarrow \infty(1-i\epsilon)} \frac{\int \mathcal{D}\phi \phi(x_1) \phi(x_2) e^{i \int_{-T}^T d^4x \mathcal{L}}}{\int \mathcal{D}\phi e^{i \int_{-T}^T d^4x \mathcal{L}}}. \quad (1.53)$$

2.2 Feynman Rules

Now we want to get the Feynman rules from (1.53), so we need to calculate the numerator and the denominator respectively. For that, we firstly consider the free action of the scalar field:

$$S_0 = \int d^4x \mathcal{L}_0 = \int d^4x \left(\frac{1}{2} (\partial_\mu \phi)^2 - \frac{1}{2} m^2 \phi^2 \right). \quad (1.54)$$

The general Gaussian integration formulas could be useful in the next:

$$\int d^n x e^{-x_i A_{ij} x_j} = \text{const.} \cdot (\det A)^{-1/2}; \quad (1.55)$$

$$\int \mathcal{D}\phi e^{\frac{i}{2} \int d^4x \phi (\partial^2 - m^2) \phi} = \text{const.} \cdot (\det(\partial^2 - m^2))^{-1/2}. \quad (1.56)$$

The idea is firstly discretizing the path integral and then take the limit to get it back to the continuous form. So we work in the lattice $\{x_i\}$, where we can define the discrete field to replace the continuous variable:

$$\mathcal{D}\phi = \prod_i d\phi(x_i). \quad (1.57)$$

And the field can be expressed by the Fourier series:

$$\phi(x_i) = \frac{1}{V} \sum_n e^{-ik_n \cdot x_i} \phi(k_n); \quad k_n^\mu = \frac{2\pi n^\mu}{L}, \quad (1.58)$$

where n^μ is integers and $|k^\mu| < \pi/\varepsilon$, ε is the distance between two close lattice points. $V = L^4$ is the volume of the space. Because of $\phi(x_n)$ is real, so

$$\begin{aligned} \phi^*(x_i) &= \frac{1}{V} \sum_n e^{+ik_n^* \cdot x_i} \phi^*(k_n) \\ &= \frac{1}{V} \sum_n e^{-i(-k_n) \cdot x_i} \phi(-k_n), \end{aligned} \quad (1.59)$$

which means $\phi^*(k_n) = \phi(-k_n)$. So the integral measurement becomes

$$\mathcal{D}\phi(x) = \prod_i d\phi(x_i) = |\det(M)| \prod_n d\phi(k_n) = \prod_{k_n^0 > 0} d\text{Re } \phi(k_n) d\text{Im } \phi(k_n), \quad (1.60)$$

where M is the Fourier transformation matrix, so the Jacobian determinant is 1. And the last equality is because $d\phi(k_n)$ is an area measurement in the complex plane, so this small piece of area measurement is $d\text{Re } \phi(k_n) d\text{Im } \phi(k_n)$. And the free action in momentum space can be rewritten as

$$S_0 = -\frac{1}{V} \sum_{k_n^0 > 0} \frac{1}{2} (m^2 - k_n^2) |\phi_n|^2 = -\frac{1}{V} \sum_{k_n^0 > 0} (m^2 - k_n^2) [(\text{Re } \phi_n)^2 + (\text{Im } \phi_n)^2]. \quad (1.61)$$

Therefore, we can together all these parts to express the denominator of (1.53), then using the Gaussian integration formula to calculate the integral:

$$\int \mathcal{D}\phi e^{iS_0} = \prod_{k_n^0 > 0} \left(\int d \operatorname{Re} \phi_n \exp \left[-\frac{i}{V} (m^2 - k_n^2) (\operatorname{Re} \phi_n)^2 \right] \right) \left(\int d \operatorname{Im} \phi_n \exp \left[-\frac{i}{V} (m^2 - k_n^2) (\operatorname{Im} \phi_n)^2 \right] \right) \quad (1.62)$$

$$= \prod_{k_n^0 > 0} \sqrt{\frac{-i\pi V}{m^2 - k_n^2}} \sqrt{\frac{-i\pi V}{m^2 - k_n^2}} \quad (1.63)$$

$$= \prod_{k_n^0 > 0} \sqrt{\frac{-i\pi V}{m^2 - k_n^2}} \sqrt{\frac{-i\pi V}{m^2 - (-k_n)^2}} \quad (1.64)$$

$$= \prod_{\text{all } k_n} \sqrt{\frac{-i\pi V}{m^2 - k_n^2}}, \quad (1.65)$$

where the last step is counting from the upper half plane to the whole complex k_n plane. And to avoid the divergence, we take $k^0 \rightarrow k^0(1 + i\epsilon)$ in the denominator. For the expression for the propagator, we still need to calculating $\phi(x_1)\phi(x_2)$ for the numerator of (1.53), so we have

$$\phi(x_1)\phi(x_2) = \frac{1}{V^2} \sum_m e^{-ik_m x_1} \phi_m \sum_l e^{-ik_l x_2} \phi_l. \quad (1.66)$$

So the numerator becomes

$$\begin{aligned} \int \mathcal{D}\phi \phi(x_1)\phi(x_2) e^{i \int_{-T}^T d^4x \mathcal{L}} &= \frac{1}{V^2} \sum_{m,l} e^{-i(k_m x_1 + k_l x_2)} \left(\prod_{n|k_n^0 > 0} \int d \operatorname{Re} \phi_n d \operatorname{Im} \phi_n \right) \\ &\quad \times (\operatorname{Re} \phi_m + i \operatorname{Im} \phi_m)(\operatorname{Re} \phi_l + i \operatorname{Im} \phi_l) \\ &\quad \times \exp \left\{ -\frac{i}{V} \sum_{n|k_n^0 > 0} (m^2 - k_n^2) [(\operatorname{Re} \phi_n)^2 + (\operatorname{Im} \phi_n)^2] \right\}. \end{aligned} \quad (1.67)$$

Notice that for $m \neq l$ and $m \neq -l$, the term

$$(\operatorname{Re} \phi_m + i \operatorname{Im} \phi_m)(\operatorname{Re} \phi_l + i \operatorname{Im} \phi_l) \exp \{ (\operatorname{Re} \phi_n)^2 + (\operatorname{Im} \phi_n)^2 \} \quad (1.68)$$

is **(odd function) $\times$ (even function) = (odd function)**, so the integral is 0. And for $m = -l$, because of $\phi(-k) = \phi^*(k)$, which means $\phi_l = \phi_{-m} = \phi_m^*$, the production term is then

$$(\phi_m) \cdot (\phi_l) = \phi_m \cdot \phi_m^* = |\phi_m|^2 = (\operatorname{Re} \phi_m)^2 + (\operatorname{Im} \phi_m)^2. \quad (1.69)$$

So the integration is $\sim \int x^2 e^{-ax^2}$, which is non-zero. Thus only the terms with $k_m = -k_l$ contribute to the integral. Finally, we apply the Gaussian integration formula to (1.67), then we have

$$\int \mathcal{D}\phi \phi(x_1)\phi(x_2) e^{i \int_{-T}^T d^4x \mathcal{L}} = \frac{1}{V^2} \sum_m e^{-ik_m(x_1 - x_2)} \left(\prod_{k_n^0 > 0} \frac{-i\pi V}{m^2 - k_n^2} \right) \frac{-iV}{m^2 - k_m^2 - i\epsilon} \quad (1.70)$$

$$= \frac{1}{V^2} \sum_m e^{-ik_m(x_1 - x_2)} \left(\prod_{\text{all } k_n} \sqrt{\frac{-i\pi V}{m^2 - k_n^2}} \right) \frac{-iV}{m^2 - k_m^2 - i\epsilon} \quad (1.71)$$

Finally, we got the Feynman propagator through dividing (1.71) by (1.65):

$$\langle 0|T\{\phi(x_1)\phi(x_2)\}|0\rangle = \frac{\int \mathcal{D}\phi \phi(x_1)\phi(x_2)e^{i\int_{-T}^T d^4x \mathcal{L}}}{\int \mathcal{D}\phi e^{i\int_{-T}^T d^4x \mathcal{L}}} \quad (1.72)$$

$$= \frac{\frac{1}{V^2} \sum_m e^{-ik_m(x_1-x_2)} \left(\prod_{\text{all } k_n} \sqrt{\frac{-i\pi V}{m^2-k_n^2}} \right) \frac{-iV}{m^2-k_m^2-i\epsilon}}{\prod_{\text{all } k_n} \sqrt{\frac{-i\pi V}{m^2-k_n^2}}} \quad (1.73)$$

$$= \sum_m \frac{1}{V} \frac{i}{k_m^2 - m^2 - i\epsilon} e^{-ik_m(x_1-x_2)} \quad (1.74)$$

$$= \int \frac{d^4k}{(2\pi)^4} \frac{i}{k^2 - m^2 + i\epsilon} e^{-ik(x_1-x_2)} = D_F(x_1 - x_2) \quad (1.75)$$

2.3 Generating Functional

Now, we consider a more convenient and general method to calculate the Feynman rules. But it might be not very direct to understanding. First, we introduce $J(x)$ as a source in the outside to define the generating functional. For this, it's convenient to review some properties of the functional derivative:

$$\frac{\delta}{\delta J(x)} J(y) = \delta^{(4)}(x-y) \implies \frac{\delta}{\delta J(x)} \int d^4y J(y) \phi(y) = \phi(x). \quad (1.76)$$

And some basic examples of the functional derivative:

1. Exponential function:

$$\frac{\delta}{\delta J(x)} \exp \left\{ i \int d^4y J(y) \phi(y) \right\} = i\phi(x) \exp \left\{ i \int d^4y J(y) \phi(y) \right\}. \quad (1.77)$$

2. Source terms containing derivatives (obtained through integration by parts):

$$\frac{\delta}{\delta J(x)} \int d^4y \partial_\mu J(y) V^\mu(y) = -\frac{\delta}{\delta J(x)} \int d^4y J(y) \partial_\mu V^\mu(y) = -\partial_\mu V^\mu(x). \quad (1.78)$$

3. Functional composite differentiation (chain rule):

$$\frac{\delta F[\phi]}{\delta J(x)} = \int d^4z \frac{\delta F[\phi]}{\delta \phi(z)} \frac{\delta \phi(z)}{\delta J(x)} \quad \left(\text{Generalization of } \frac{\partial x_i(z_k)}{\partial y_j} = \sum_k \frac{\partial x_i}{\partial z_k} \frac{\partial z_k}{\partial y_j} \right). \quad (1.79)$$

Define the **Generating Functional** $Z[J]$ as:

$$Z[J] = \int \mathcal{D}\phi \exp \left\{ i \int d^4x (\mathcal{L} + J(x)\phi(x)) \right\}. \quad (1.80)$$

Using $Z[J]$, we can conveniently calculate the n -point correlation function:

$$\langle \Omega | T\{\phi(x_1)\phi(x_2)\} | \Omega \rangle = \frac{1}{Z[0]} \left(-i \frac{\delta}{\delta J(x_1)} \right) \left(-i \frac{\delta}{\delta J(x_2)} \right) Z[J] \Big|_{J=0} \quad (1.81)$$

2.4 Feynman Rules for the Free Theory

For a free scalar field, the Lagrangian is

$$\mathcal{L}_0 = \frac{1}{2} [(\partial_\mu \phi)(\partial^\mu \phi) + (-m^2 + i\epsilon)\phi^2] \quad (1.82)$$

$$= \frac{1}{2} [\partial_\mu(\phi \partial^\mu \phi) - (\phi \partial^2 \phi) + (-m^2 + i\epsilon)\phi^2] \quad (1.83)$$

$$= \frac{1}{2} \phi(-\partial^2 - m^2 + i\epsilon)\phi. \quad (1.84)$$

Define the operator $K_x = -\partial^2 - m^2 + i\epsilon$, and the action with source can be written as:

$$S_J = \int d^4x (\mathcal{L}_0 + J\phi) = \int d^4x \left(\frac{1}{2} \phi K_x \phi + J\phi \right) \quad (1.85)$$

$$= \int d^4x d^4y \left(\frac{1}{2} \phi(x) K(x, y) \phi(y) \right) + \int d^4x J(x) \phi(x), \quad (1.86)$$

where $K(x, y) = \delta^{(4)}(x - y) K_y$. Now the generating functional is:

$$Z_0[J] = \int \mathcal{D}\phi \exp \left\{ i \int d^4x \left(\frac{1}{2} \phi K \phi + J\phi \right) \right\}. \quad (1.87)$$

Now we change the variable $\phi \rightarrow \phi' = \phi + K^{-1}J$, so the measurement keeps unchanged $\mathcal{D}\phi' = \mathcal{D}\phi$:

$$Z_0[J] = \int \mathcal{D}\phi' \left| \det \frac{\delta\phi'}{\delta\phi} \right| \exp \left\{ i \int d^4x \left[\frac{1}{2} (\phi' - K^{-1}J) K (\phi' - K^{-1}J) + J(\phi' - K^{-1}J) \right] \right\} \quad (1.88)$$

$$= \left(\int \mathcal{D}\phi' \exp \left\{ \frac{i}{2} \int d^4x \phi' K \phi' \right\} \right) \exp \left\{ -\frac{i}{2} \int d^4x d^4y J(x) K^{-1}(x, y) J(y) \right\} \quad (1.89)$$

$$= Z_0[J = 0] \exp \left\{ -\frac{i}{2} \int d^4x d^4y J(x) K^{-1}(x, y) J(y) \right\}, \quad (1.90)$$

where $\left| \det \frac{\delta\phi'}{\delta\phi} \right| = 1$. And notice for

$$J(y) = K(x, y) (K^{-1}(x, y) J(y)) \quad (1.91)$$

$$= (-\partial_x^2 - m^2 + i\epsilon) K^{-1}(x, y) \delta^{(4)}(x - y) J(y) \quad (1.92)$$

$$= (-\partial_x^2 - m^2 + i\epsilon) K^{-1}(x, y) J(x). \quad (1.93)$$

Thus, K^{-1} satisfies the Green's function (except an $-i$ factor in the right hand side):

$$(-\partial_x^2 - m^2 + i\epsilon) K^{-1}(x, y) = \delta^{(4)}(x - y). \quad (1.94)$$

Take a sharp look! So we can directly realize that the Feynman propagator $-iD_F(x - y)$ also satisfies this equation, so we realize that $K^{-1}(x - y)$ is nothing but $-iD_F(x - y)$. So the **generating functional in free theory** can be then expressed as

$$Z_0[J] = Z_0[0] \exp \left\{ -\frac{1}{2} \int d^4x d^4y J(x) D_F(x - y) J(y) \right\}. \quad (1.95)$$

So the correlation function can be obtained from (1.81):

$$\langle 0 | T \{ \phi(x_1) \phi(x_2) \} | 0 \rangle = \frac{1}{Z_0[0]} \left(-i \frac{\delta}{\delta J(x_1)} \right) \left(-i \frac{\delta}{\delta J(x_2)} \right) Z_0[J] \Big|_{J=0} \quad (1.96)$$

$$= \left(-i \frac{\delta}{\delta J(x_1)} \right) \left(-i \frac{\delta}{\delta J(x_2)} \right) \exp \left\{ -\frac{1}{2} \int d^4x d^4y J(x) D_F(x-y) J(y) \right\} \Big|_{J=0} \quad (1.97)$$

$$= \left(-\frac{\delta}{\delta J(x_1)} \right) \left[-\frac{1}{2} \left(\int d^4y D_F(x_2-y) J(y) + \int d^4x J(x) D_F(x-x_2) \right) \right] \times \exp \left\{ -\frac{1}{2} \int d^4x d^4y J(x) D_F(x-y) J(y) \right\} \Big|_{J=0} \quad (1.98)$$

$$= \frac{\delta}{\delta J(x_1)} \left[\int d^4y D_F(y-x_2) J(y) \right] \frac{Z_0[J]}{Z_0[0]} \Big|_{J=0} \quad (1.99)$$

$$= D_F(x_1-x_2) \frac{Z_0[J]}{Z_0[0]} \Big|_{J=0} + \left[\int d^4y D_F(y-x_2) J(y) \right] \left[\int d^4y D_F(y-x_1) J(y) \right] \frac{Z_0[J]}{Z_0[0]} \Big|_{J=0} \quad (1.100)$$

$$= D_F(x_1-x_2). \quad (1.101)$$

Technically, the last step we do, which impose the boundary condition to cancel the second term is not very strict. In general, one can not impose the boundary condition before actually calculating the integration. More precisely, the reason why we have confidence to do so is based on the property that we can let $f \in L^2(\mathbb{R}^{1,3})$ be fixed, and define a functional

$$F_f : L^2(\mathbb{R}^{1,3}) \rightarrow \mathbb{C}, \quad F_f[J] := \int_{\mathbb{R}^{1,3}} f(x) J(x) dx. \quad (1.102)$$

Then F_f is a continuous linear functional on $L^2(\mathbb{R}^{1,3})$. Indeed, by the Cauchy–Schwarz inequality,

$$|F_f[J]| \leq \|f\|_{L^2} \|J\|_{L^2}, \quad (1.103)$$

so F_f is bounded and hence continuous. In particular, for any sequence (or net) $J_n \rightarrow 0$ in $L^2(\mathbb{R}^{1,3})$, one has

$$\lim_{n \rightarrow \infty} F_f[J_n] = F_f[0] = 0. \quad (1.104)$$

Therefore, the notation $F_f[J] \big|_{J=0}$ is to be understood as the evaluation of the functional F_f at the zero element of $L^2(\mathbb{R}^{1,3})$, i.e.

$$F_f[J] \big|_{J=0} := F_f(0). \quad (1.105)$$

Moreover, by continuity, this evaluation coincides with the limit along any sequence $J_n \rightarrow 0$ in the L^2 topology:

$$F_f(0) = \lim_{J \rightarrow 0} F_f[J], \quad (1.106)$$

where the limit is taken with respect to the L^2 norm.

2.5 Interaction theory: ϕ^4 theory as an example

For theories involving interactions, such as $\mathcal{L} = \mathcal{L}_0 - \frac{\lambda}{4!} \phi^4$ and $\mathcal{L}_0 = \frac{1}{2}(\partial_\mu \phi)^2 - \frac{1}{2}m^2 \phi^2$, the generating functional can be written as:

$$Z[J] = \int \mathcal{D}\phi \exp \left\{ i \int d^4x \left(-\frac{\lambda}{4!} \phi^4 + \mathcal{L}_0 + J(x) \phi(x) \right) \right\}. \quad (1.107)$$

Notice that

$$\begin{aligned}\frac{\delta}{\delta J(x)} Z_0[J] &= \frac{\delta}{\delta J(x)} \int \mathcal{D}\phi \exp \left\{ -i \int d^4x \mathcal{L}_0 + J(x)\phi(x) \right\} \\ &= \phi(x) \int \mathcal{D}\phi \exp \left\{ -i \int d^4x \mathcal{L}_0 + J(x)\phi(x) \right\} \\ &= \phi(x) Z_0[J].\end{aligned}\tag{1.108}$$

So we have

$$\left(-i \frac{\delta}{\delta J(x)} \right)^4 Z_0[J] = \phi^4 Z_0[J],\tag{1.109}$$

which means

$$Z[J] = \int \mathcal{D}\phi \left\{ 1 + \left(-i \int d^4x \frac{\lambda}{4!} \phi^4 \right) + \left(-i \int d^4x \frac{\lambda}{4!} \phi^4 \right)^2 + \dots \right\} \exp \left\{ -i \int d^4x [\mathcal{L}_0 + J(x)\phi(x)] \right\}\tag{1.110}$$

$$\begin{aligned}&= \int \mathcal{D}\phi \left\{ 1 + \left[-i \int d^4x \frac{\lambda}{4!} \left(-i \frac{\delta}{\delta J(x)} \right)^4 \right] + \left[-i \int d^4x \frac{\lambda}{4!} \left(-i \frac{\delta}{\delta J(x)} \right)^4 \right]^2 + \dots \right\} \\ &\quad \times \exp \left\{ -i \int d^4x [\mathcal{L}_0 + J(x)\phi(x)] \right\}\end{aligned}\tag{1.111}$$

$$= \exp \left\{ -i \int d^4x \frac{\lambda}{4!} \left(-i \frac{\delta}{\delta J(x)} \right)^4 \right\} \int \mathcal{D}\phi \exp \left\{ -i \int d^4x [\mathcal{L}_0 + J(x)\phi(x)] \right\}\tag{1.112}$$

$$= \exp \left\{ -i \int d^4x \frac{\lambda}{4!} \left(-i \frac{\delta}{\delta J(x)} \right)^4 \right\} Z_0[J],\tag{1.113}$$

where we expanded the exponential term of interaction part as the perturbed series, and using the derivative relation (1.109). So we link the interaction theory generating functional with the free theory generating functional:

$$Z[J] = \left(1 - \frac{i\lambda}{4!} \int d^4x \left(-i \frac{\delta}{\delta J(x)} \right)^4 + \dots \right) Z_0[J].\tag{1.114}$$

So the correlation function can be expanded by the free propagator as

$$\langle \Omega | T \{ \phi(x_1) \phi(x_2) \} | \Omega \rangle = \frac{1}{Z[0]} \left(-i \frac{\delta}{\delta J(x_1)} \right) \left(-i \frac{\delta}{\delta J(x_2)} \right) Z[J] \Big|_{J=0}\tag{1.115}$$

$$= \frac{1}{Z[0]} \left(-i \frac{\delta}{\delta J(x_1)} \right) \left(-i \frac{\delta}{\delta J(x_2)} \right) \left(1 - \frac{i\lambda}{4!} \int d^4x \left(-i \frac{\delta}{\delta J(x)} \right)^4 + \dots \right) Z_0[J] \Big|_{J=0}\tag{1.116}$$

$$= \frac{1}{Z[0]} \left[(-i)^2 \frac{\delta^2 Z_0[J]}{\delta J(x_1) \delta J(x_2)} + \left(-\frac{i\lambda}{4!} \right) \int d^4x (-i)^6 \frac{\delta^6 Z_0[J]}{\delta J(x)^4 \delta J(x_1) \delta J(x_2)} + \dots \right] \Big|_{J=0}\tag{1.117}$$

$$= \frac{1}{Z[0]} \langle 0 | T \{ \phi(x_1) \phi(x_2) \} | 0 \rangle + \frac{1}{Z[0]} \left(-i \frac{\lambda}{4!} \right) \int d^4x \langle 0 | T \{ \phi^4(x) \phi(x_1) \phi(x_2) \} | 0 \rangle + \dots\tag{1.118}$$

$$= \frac{1}{Z[0]} \langle 0 | T \left\{ \phi(x_1) \phi(x_2) \exp \left\{ i \int d^4x \left(-\frac{\lambda}{4!} \phi^4 \right) \right\} \right\} | 0 \rangle.\tag{1.119}$$

Comparing with the result we got in the canonical quantization

$$\langle \Omega | T \{ \phi(x_1) \phi(x_2) \} | \Omega \rangle = \frac{\langle 0 | T \{ \phi(x_1) \phi(x_2) \exp \{ i \int d^4x \left(-\frac{\lambda}{4!} \phi^4 \right) \} \} | 0 \rangle}{\langle 0 | T \{ \exp \{ i \int d^4x \left(-\frac{\lambda}{4!} \phi^4 \right) \} \} | 0 \rangle}, \quad (1.120)$$

we find in the interaction case, $Z[0]$ has different meaning comparing with $Z_0[0]$ in the free case, i.e.,

$$Z[0] = T \exp \left\{ i \int d^4x \left(-\frac{\lambda}{4!} \phi^4 \right) \right\}, \quad (1.121)$$

which included all the **Vacuum Bubbles**.

2.6 Other Type of Generating functional

We define the **Energy Functional** for the connected graphs $E[J]$ as

$$E[J] = i \ln Z[J]. \quad (1.122)$$

So we have

$$\frac{\delta E[J]}{\delta J(x)} = \frac{i}{Z[0]} \frac{\delta Z[J]}{\delta J(x)} = -\frac{\langle \Omega | \phi(x) | \Omega \rangle}{\langle \Omega | \Omega \rangle} = -\phi_{\text{classical}}(x), \quad (1.123)$$

where the second step is because $\delta Z[J] = \langle \Omega | i\phi(x) | \Omega \rangle$. And in physics, it is defined as the **Classical Field**. And the definition of $E[J]$ can be easily seen that it's very similar with the free energy in classical statistical mechanics with the partition function $Z[J]$. So using the Legendre transformation, we can define the **Effective Action**, which looks like the Gibbs free energy $\Gamma[\phi_{cl}]$ as

$$\Gamma[\phi_{cl}] = -E[J] - \int d^4x J(x) \phi_{cl}(x). \quad (1.124)$$

Base on this, we can see there is a symmetric relation:

$$\frac{\delta \Gamma[\phi_{cl}]}{\delta \phi_{cl}(x)} = -J(x); \quad \frac{\delta E[J]}{\delta J(x)} = -\phi_{cl}(x). \quad (1.125)$$

Taking another functional derivative with respect to $J(y)$ on both sides yields:

$$\Rightarrow \frac{\delta}{\delta J(y)} \frac{\delta}{\delta \Phi_{cl}(x)} \Gamma[\Phi_{cl}] = -\frac{\delta J(x)}{\delta J(y)} = -\delta(x - y). \quad (1.126)$$

Using the chain rule for functional differentiation $\frac{\delta}{\delta J(y)} = \int d^4z \frac{\delta \Phi_{cl}(z)}{\delta J(y)} \frac{\delta}{\delta \Phi_{cl}(z)}$, we obtain the following identity:

$$-\delta(x - y) = \int d^4z \frac{\delta \Phi_{cl}(z)}{\delta J(y)} \frac{\delta^2 \Gamma[\Phi_{cl}]}{\delta \Phi_{cl}(z) \delta \Phi_{cl}(x)}. \quad (1.127)$$

We can rewrite the first term as $\frac{\delta \Phi_{cl}(z)}{\delta J(y)} = \frac{\delta^2 W[J]}{\delta J(y) \delta J(z)}$, where $W[J]$ is the generating functional of connected Green's functions (denoted as $E[J]$ in the notes):

$$-\delta(x - y) = \int d^4z \underbrace{\frac{\delta^2 W[J]}{\delta J(y) \delta J(z)}}_{-iG(y-z)} \underbrace{\frac{\delta^2 \Gamma[\Phi_{cl}]}{\delta \Phi_{cl}(z) \delta \Phi_{cl}(x)}}_{iG^{-1}(z-x)}. \quad (1.128)$$

Here, the first part corresponds to the connected two-point function $-iG(y-z)$, and the second part corresponds to the inverse propagator $iG^{-1}(z-x)$. Thus, the equation simplifies to:

$$\delta(x-y) = \int d^4z [-iG(y-z)] [iG^{-1}(z-x)]. \quad (1.129)$$

The n -th derivative of $E[J]$ gives the connected correlation function

$$\left. \frac{\delta^n E[J]}{\delta J(x_1) \dots \delta J(x_n)} \right|_{J=0} = (i)^{n+1} \langle \Omega | T\{\phi(x_1) \dots \phi(x_n)\} | \Omega \rangle_{\text{connect}}. \quad (1.130)$$

And the n -th derivative of $\Gamma[\phi_{cl}]$ gives the correlation function for 1PI graphs

$$\frac{\delta^n \Gamma[\phi_{cl}]}{\delta \phi_{cl}(x_1) \dots \delta \phi_{cl}(x_n)} = -i \langle \Omega | T\{\phi(x_1) \dots \phi(x_n)\} | \Omega \rangle_{1PI}. \quad (1.131)$$

3. Quantization of EM Field in Path Integral Formalism

The classical action for the electromagnetic field is given by:

$$S[A] = -\frac{1}{4} \int d^4x F_{\mu\nu} F^{\mu\nu}, \quad (1.132)$$

where $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$. Integrating by parts, we can rewrite the action as:

$$S[A] = \frac{1}{2} \int d^4x A_\mu D^{\mu\nu} A_\nu, \quad \text{with} \quad D^{\mu\nu} = g^{\mu\nu} \square - \partial^\mu \partial^\nu. \quad (1.133)$$

In the path integral formulation, the vacuum-to-vacuum transition amplitude is given by:

$$\int \mathcal{D}A e^{iS[A]}. \quad (1.134)$$

But the problem the measure $\mathcal{D}A = \mathcal{D}A^0 \mathcal{D}A^1 \mathcal{D}A^2 \mathcal{D}A^3$ is **not** well defined. The integral is ill-defined because of the redundancy due to infinite gauge copies. Under a local gauge transformation:

$$A_\mu \rightarrow A_\mu^\alpha = A_\mu + \frac{1}{e} \partial_\mu \alpha(x), \quad (1.135)$$

where $\alpha(x)$ is an arbitrary scalar function. For a pure gauge choice where $A_\mu = \frac{1}{e} \partial_\mu \alpha$, we find that the field strength $F_{\mu\nu} = 0$, leading to:

$$S[A] = 0 \quad \Rightarrow \quad e^{iS[A]} = 1. \quad (1.136)$$

This implies that we have an unbounded path integral because the integrand does not decay along the gauge orbits.

3.1 Faddeev-Popov Quantization

Solution: We need to divide out the gauge degrees of freedom in the measure $\mathcal{D}A$. This procedure is known as the **Faddeev-Popov quantization**.

Let $G(A)$ be a function that fixes the gauge (for example, $G(A) = \partial_\mu A^\mu = 0$ for the Lorentz gauge). We want to define $\int \mathcal{D}A e^{iS[A]}$ with a constraint such that the path integral only sums over field configurations fulfilling $G(A) = 0$.

To motivate this, recall the standard Gauss integration with matrix M (rank N):

$$(\det M)^{-1/2} = (2\pi)^{-N/2} \int_{-\infty}^{+\infty} dx_1 \dots dx_N e^{-\frac{1}{2}x^T M x}. \quad (1.137)$$

Generalizing this to a functional determinant (Path integral):

$$(\det M)^{-1/2} = \int \mathcal{D}\phi e^{-\frac{1}{2} \int d^4x \phi(x) M(x) \phi(x)}, \quad (1.138)$$

where M is some differential operator, e.g., $M(x) = \square + m_\phi^2$.

Next, consider the Dirac δ -distribution on an N -dimensional space for an arbitrary function $\vec{g}(x_1, \dots, x_N)$:

$$\int_{-\infty}^{+\infty} dx_1 \dots dx_N \delta^{(N)}[\vec{g}(x_1, \dots, x_N)] \det \left[\frac{\partial g_i}{\partial x_j} \right] = 1 \quad (1.139)$$

In 1-dimension, this reduces to the property:

$$\int dx \delta(ax) a = \int dx \frac{1}{a} \delta(x) a = 1. \quad (1.140)$$

Extending this to a functional Dirac distribution for gauge fields, we have the Faddeev-Popov identity:

$$\int \mathcal{D}\alpha(x) \underbrace{\delta[G(A^\alpha)] \det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right]}_{\text{Faddeev-Popov det.}} = 1. \quad (1.141)$$

Quantization of electrodynamics proceeds by inserting this identity “1” into the path integral:

$$I = \int \mathcal{D}A e^{iS[A]} = \int \mathcal{D}A (1) e^{iS[A]} = \int \mathcal{D}A \left(\int \mathcal{D}\alpha \delta[G(A^\alpha)] \det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right] \right) e^{iS[A]}. \quad (1.142)$$

If $G(A^\alpha)$ is the Lorentz gauge condition, we have:

$$G(A^\alpha) = \partial_\mu A^\mu + \frac{1}{e} \square \alpha. \quad (1.143)$$

Then the functional derivative becomes:

$$\det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right] = \det \left(\frac{\square}{e} \right). \quad (1.144)$$

Notice that in quantum electrodynamics, this determinant is invariant under gauge transformations and independent of A_μ .

Due to gauge invariance, we have $\mathcal{D}A = \mathcal{D}A^\alpha$ and $S[A] = S[A^\alpha]$. Therefore, the path integral can be rewritten as:

$$\begin{aligned} I &= \int \mathcal{D}\alpha \det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right] \int \mathcal{D}A \delta[G(A^\alpha)] e^{iS[A]} \\ &= \int \mathcal{D}\alpha \left(\det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right] \int \mathcal{D}A^\alpha \delta[G(A^\alpha)] e^{iS[A^\alpha]} \right) \\ &= \int \mathcal{D}\alpha \tilde{I}. \end{aligned} \quad (1.145)$$

By dropping the infinite volume factor of the gauge group $\int \mathcal{D}\alpha$, we define the physical path integral for QED as:

$$\tilde{I} = \det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right] \int \mathcal{D}A^\alpha \delta[G(A^\alpha)] e^{iS[A]}. \quad (1.146)$$

3.2 Gauge Fixing and the Photon Propagator

Our next task is to absorb the Dirac constraint $\delta[G(A^\alpha)]$ into a modified action. Let us generalize the gauge fixing condition by setting:

$$G(A^\alpha) = \partial^\mu A_\mu(x) - \omega(x), \quad (1.147)$$

where $\omega(x)$ is an arbitrary scalar function. We also set the determinant remains independent of ω as the form:

$$\det \left[\frac{\delta G(A^\alpha)}{\delta \alpha} \right] = \det \left(\frac{\square}{e} \right). \quad (1.148)$$

So the functional integration $\tilde{I}$ becomes

$$\int \mathcal{D}A e^{-iS[A]} = \det \left(\frac{\square}{e} \right) \int \mathcal{D}A^\alpha \delta[\partial^\mu A_\mu - \omega(x)] e^{iS[A]}. \quad (1.149)$$

Since this formula is satisfied for any $\omega(x)$, so we can integrating over all possible functions $\omega(x)$ with a Gaussian weight centered at $\omega = 0$:

$$I' = N(\xi) \int \mathcal{D}\omega e^{-i \int d^4x \frac{\omega(x)^2}{2\xi}} \tilde{I}, \quad (1.150)$$

where $N(\xi)$ is a normalization factor. Then we substitute $\tilde{I}$ and explicitly show the delta function:

$$\begin{aligned} I' &= N(\xi) \det \left(\frac{\square}{e} \right) \int \mathcal{D}A e^{iS[A]} \int \mathcal{D}\omega e^{-i \int d^4x \frac{\omega^2}{2\xi}} \underbrace{\delta[\partial^\mu A_\mu - \omega]}_{\text{linear covariant gauge}} \\ &= N(\xi) \det \left(\frac{\square}{e} \right) \int \mathcal{D}A e^{iS[A]} e^{-\frac{i}{2\xi} \int d^4x (\partial^\mu A_\mu)^2} \\ &= N(\xi) \det \left(\frac{\square}{e} \right) \int \mathcal{D}A e^{iS'[A]}. \end{aligned} \quad (1.151)$$

The modified action $S'[A]$ becomes:

$$S'[A] = \frac{1}{2} \int d^4x \left[A_\mu D^{\mu\nu} A_\nu - \frac{1}{\xi} (\partial^\mu A_\mu)^2 \right] = \frac{1}{2} \int d^4x A_\mu \tilde{D}^{\mu\nu} A_\nu, \quad (1.152)$$

which implies the new differential operator:

$$\tilde{D}^{\mu\nu} = g^{\mu\nu} \square - \left(1 - \frac{1}{\xi} \right) \partial^\mu \partial^\nu. \quad (1.153)$$

From this modified action, we can derive the generating functional as

$$Z[J_\mu] = \int \mathcal{D}A \exp \left\{ i \int d^4x \left(\frac{1}{2} A_\mu \tilde{D}^{\mu\nu} A_\nu + J^\mu(x) A_\mu(x) \right) \right\}. \quad (1.154)$$

Then transform the field as $A_\mu \rightarrow A'_\mu = A_\mu + \tilde{D}_\mu^{-1\nu} J_\nu$, we have

$$Z[J_\mu] = \int \mathcal{D}A \exp \left\{ i \int d^4x \left(\frac{1}{2} (A'_\mu - \tilde{D}_\mu^{-1\lambda} J_\lambda) \tilde{D}^{\mu\nu} (A'_\nu - \tilde{D}_\nu^{-1\lambda} J_\lambda) + J^\mu(x) (A'_\mu - \tilde{D}_\mu^{-1\lambda} J_\lambda) \right) \right\} \quad (1.155)$$

$$= \left(\int \mathcal{D}A' \exp \left\{ \frac{i}{2} A'_\mu \tilde{D}^{\mu\nu} A'_\nu \right\} \right) \exp \left\{ -\frac{i}{2} \int d^4x d^4y (J_\mu(x) \tilde{D}^{-1\mu\nu} J_\nu(y)) \right\} \quad (1.156)$$

$$= Z[0] \exp \left\{ -\frac{i}{2} \int d^4x d^4y (J_\mu(x) \tilde{D}^{-1\mu\nu} J_\nu(y)) \right\} \quad (1.157)$$

$$= Z[0] \exp \left\{ -\frac{i}{2} \int d^4x d^4y (J_\mu(x) \tilde{G}^{\mu\nu} J_\nu(y)) \right\}, \quad (1.158)$$

where we have ignored the determinant due to $\det\left(\frac{\delta A'}{\delta A}\right) = 1$. And the photon propagator is the inverse of $\tilde{D}^{\mu\nu}$ in momentum space:

$$G^{\mu\nu}(x-y) = -i \int \frac{d^4k}{(2\pi)^4} e^{-ik(x-y)} \frac{g^{\mu\nu} - (1-\xi) \frac{k^\mu k^\nu}{k^2}}{k^2 + i\epsilon}. \quad (1.159)$$

The effective action then can be constructed as same as (1.124)

$$\Gamma[A_{cl}^\mu] = -iZ[J_\mu] - \int d^4x J_\mu(x) A_{cl}^\mu. \quad (1.160)$$

$S'[A]$ with $\tilde{D}^{\mu\nu}$ now has no problems with the longitudinal degrees of freedom for photons.

3.3 Standard Gauge Choices

Depending on the choice of the parameter ξ and conditions, we have different gauges in the linear covariant gauge family:

- $\xi = 0$: **Landau gauge**, corresponding to the condition $\partial^\mu A_\mu = 0$.
- $\xi = 1$: **Feynman gauge**.
- $i = 1, 2, 3$, $\partial^i A_i = 0$: **Coulomb gauge**.
- $n^\mu A_\mu = 0$: **Axial gauge**.

Finally, the path integral in QED for calculating an n -point correlator is written as:

$$\langle \Omega | T \hat{O}(A) | \Omega \rangle = \lim_{T \rightarrow \infty(1-i\epsilon)} \frac{\int \mathcal{D}A \hat{O}(A) \exp \left\{ i \int_{-T}^T d^4x \left[\mathcal{L}(x) - \frac{1}{2\xi} (\partial^\mu A_\mu)^2 \right] \right\}}{\int \mathcal{D}A \exp \left\{ i \int_{-T}^T d^4x \left[\mathcal{L}(x) - \frac{1}{2\xi} (\partial^\mu A_\mu)^2 \right] \right\}}. \quad (1.161)$$

4. Path Integral for Fermions

To define the path integral for fermions, we consider the transition amplitude:

$$Z = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp \left\{ i \int d^4x \mathcal{L}(\psi, \bar{\psi}) \right\}, \quad \mathcal{L} = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi. \quad (1.162)$$

Our task is to make sense of this integral, which requires treating fermion fields as Grassmann-valued quantities.

4.1 Grassmann Numbers and Algebra

Grassmann numbers θ satisfy the anti-commutation relation:

$$\{\theta_i, \theta_j\} = 0 \quad \Rightarrow \quad \theta^2 = 0. \quad (1.163)$$

A differential operator $\frac{d}{d\theta}$ is defined such that it also anti-commutes:

$$\left\{ \frac{d}{d\theta}, \theta \right\} = \frac{d}{d\theta} \theta + \theta \frac{d}{d\theta} 1 = 1. \quad (1.164)$$

Any Grassmann-valued function $f(\theta)$ can be defined through a power series. Due to $\theta^2 = 0$, the series terminates at the linear term:

$$f(\theta) = a + \beta\theta, \quad (1.165)$$

where a is a regular number (real/complex) and β is another Grassmann-valued quantity. Taking the derivative:

$$\frac{d}{d\theta} f(\theta) = \frac{d}{d\theta} (a + \beta\theta) = 0 + \frac{d}{d\theta} (\beta\theta) = -\beta \frac{d\theta}{d\theta} = -\beta. \quad (1.166)$$

Note the sign change because $\frac{d}{d\theta}$ must pass β . Other properties include:

$$\frac{d^2}{d\theta^2} f(\theta) = 0, \quad \frac{d}{d\theta} a = \left\{ \frac{d}{d\theta}, \beta \right\} = 0. \quad (1.167)$$

4.2 Berezin Integration

Integration over Grassmann variables (Berezin integration) is defined to be identical to differentiation:

$$\int d\theta \, 1 = 0, \quad \int d\theta \, \theta = 1. \quad (1.168)$$

For multiple Grassmann variables θ_i with internal symmetry ($i = 1, \dots, N$):

$$\left\{ \frac{d}{d\theta_i}, \theta_j \right\} = \delta_{ij}, \quad \left\{ \frac{d}{d\theta_i}, \frac{d}{d\theta_j} \right\} = 0, \quad \int d\theta_i = 0, \quad \int d\theta_i \, \theta_j = \delta_{ij}. \quad (1.169)$$

Example: Consider a two-dimensional integral:

$$\int d\theta_1 d\theta_2 \, \theta_1 \theta_2 = \int d\theta_1 \left(\int d\theta_2 \, \theta_1 \theta_2 \right) = \int d\theta_1 \, (-\theta_1) = -1. \quad (1.170)$$

4.3 Gaussian Integrals and Determinants

For a complex Grassmann pair $\eta, \bar{\eta}$, we represent the determinant of a matrix M_{ij} :

$$\int \left[\prod_{j=1}^N d\eta_j d\bar{\eta}_j \right] \exp \left(\sum_{k,l} \bar{\eta}_k M_{kl} \eta_l \right) = \det[M]. \quad (1.171)$$

Auxiliary Calculation (1D Example):

$$\int d\bar{\eta} d\eta \, e^{-\bar{\eta}b\eta} = \int d\bar{\eta} d\eta \, (1 - \bar{\eta}b\eta) = \int d\bar{\eta} d\eta \, (1 + \eta\bar{\eta}b) = \int d\bar{\eta} \, (\bar{\eta}b) = b. \quad (1.172)$$

Similarly, for the insertion of variables:

$$\int d\bar{\eta} d\eta \, \eta\bar{\eta} e^{-\bar{\eta}b\eta} = \int d\bar{\eta} d\eta \, \eta\bar{\eta} (1 - \bar{\eta}b\eta) = \int d\bar{\eta} d\eta \, \eta\bar{\eta} = 1 = \frac{1}{b} \cdot b. \quad (1.173)$$

4.4 Transformation of Variables

Under a linear transformation $\theta_i = U_{ij}\theta'_j$, the product of Grassmann variables transforms with the determinant:

$$\prod_{k=1}^N \theta'_k = \frac{1}{N!} \varepsilon^{i_1 \dots i_N} U_{i_1 j_1} \dots U_{i_N j_N} \theta_{j_1} \dots \theta_{j_N} = \det[U] \prod_{k=1}^N \theta_k. \quad (1.174)$$

This is inverse to the bosonic case where the Jacobian appears in the denominator.

4.5 Comparison: Bosonic vs. Fermionic Case

For a Hermitian matrix M and source w, ξ :

- **Bosonic Case:**

$$F(M, \omega) = \int d^N x \exp \left(-\frac{1}{2} x^T M x + \omega^T x \right) = \pi^{N/2} (\det M)^{-1/2} \exp \left(\frac{1}{2} \omega^T M^{-1} \omega \right). \quad (1.175)$$

- **Fermionic Case:**

$$F(M, \xi, \bar{\xi}) = \int \prod d\eta_i d\bar{\eta}_i \exp (\bar{\eta} M \eta + \bar{\xi} \eta + \bar{\eta} \xi) = \det[M] \exp (-\bar{\xi} M^{-1} \xi). \quad (1.176)$$

5. Path Integral Formulation for Dirac Spinors

A Dirac spinor $\psi_a(x)$ consists of Grassmann-valued quantities:

$$\psi(x) = \sum_{a=1}^4 \psi_a(x), \quad \text{where } \psi_a \text{ are Grassmann variables.} \quad (1.177)$$

The Dirac Lagrangian is $\mathcal{L}_{\text{Dirac}} = \bar{\psi}_a(x) (i\gamma_{ab}^\mu \partial_\mu - m\delta_{ab}) \psi_b(x)$. The vacuum amplitude is:

$$\int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left[i \int d^4x \mathcal{L}(\bar{\psi}, \psi) \right] = \text{const} \cdot \det(i\not{\partial} - m). \quad (1.178)$$

5.1 Fermion Propagator

In the path integral formulation, the propagator is defined as the time-ordered product:

$$\langle 0 | T \{ \psi(x_1) \bar{\psi}(x_2) \} | 0 \rangle = \frac{\int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp [i \int d^4x \bar{\psi}(i\not{\partial} - m)\psi] \psi(x_1) \bar{\psi}(x_2)}{\int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp [i \int d^4x \bar{\psi}(i\not{\partial} - m)\psi]}. \quad (1.179)$$

The calculation yields:

$$\langle 0 | T \{ \psi(x_1) \bar{\psi}(x_2) \} | 0 \rangle = \frac{1}{\text{const} \cdot \det(i\not{\partial} - m)} \cdot \frac{\text{const} \cdot \det(i\not{\partial} - m)}{-i(i\not{\partial} - m)_{ab}} = \frac{i}{(i\not{\partial} - m)_{ab}} = S_{F,ab}(x_1 - x_2). \quad (1.180)$$

5.2 Generating Functional for Fermion Fields

Analogous to the scalar case, we introduce Grassmann-valued sources $\eta(x), \bar{\eta}(x)$:

$$Z[\eta, \bar{\eta}] = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left\{ i \int d^4x [\mathcal{L}(\bar{\psi}, \psi) + \bar{\psi}(x)\eta(x) + \bar{\eta}(x)\psi(x)] \right\}. \quad (1.181)$$

The functional derivative with respect to these sources follows:

$$\frac{\delta \eta(x)}{\delta \eta(y)} = \delta(x - y). \quad (1.182)$$

By completing the square, we can express $Z[\eta, \bar{\eta}]$ in terms of the propagator:

$$Z[\eta, \bar{\eta}] = Z[\eta = 0, \bar{\eta} = 0] \exp \left[- \int d^4x d^4y \bar{\eta}(x) S_F(x - y) \eta(y) \right]. \quad (1.183)$$

5.3 Derivation of the Propagator

The propagator can finally be extracted via functional derivatives:

$$\langle 0 | T \{ \psi(x_1) \bar{\psi}(x_2) \} | 0 \rangle = \frac{1}{Z[0,0]} \left(-i \frac{\delta}{\delta \bar{\eta}(x_1)} \right) \left(-i \frac{\delta}{\delta \eta(x_2)} \right) Z[\eta, \bar{\eta}] \Big|_{\eta, \bar{\eta}=0} = S_F(x_1 - x_2). \quad (1.184)$$

6. Symmetries in Path Integral Formalism

The path integral approach provides a direct link between symmetries and physical observables through the following hierarchy: **Variational calculus** → **Schwinger-Dyson equations** → **Noether's theorem** → **Ward identities in QED**.

6.1 Schwinger-Dyson Equations

For simplicity, we still consider the scalar field theory as an example. By doing the partial integration, the action can be rewritten as:

$$S[\phi] = \int d^4x \mathcal{L}[\phi] = \int d^4x \left\{ \frac{1}{2} (\partial_\mu \phi)^2 - \frac{1}{2} m^2 \phi^2 \right\} = -\frac{1}{2} \int d^4x \phi (\square + m^2) \phi. \quad (1.185)$$

The n -point correlator is given by:

$$\langle \Omega | T \{ \phi(x_1) \dots \phi(x_n) \} | \Omega \rangle = \frac{\int \mathcal{D}\phi \phi_1 \dots \phi_n e^{iS[\phi]}}{\int \mathcal{D}\phi e^{iS[\phi]}}. \quad (1.186)$$

We derive classical equations of motion from a variation of fields: $\phi(x) \rightarrow \phi'(x) = \phi(x) + \varepsilon(x)$, i.e., $\delta\phi(x) = \varepsilon(x)$. The integration measurement is invariant due to ε is infinitesimal: $\mathcal{D}\phi' = \mathcal{D}\phi$. And the action changes as: $S[\phi'] = S[\phi + \varepsilon] = S[\phi] + \delta S$. So we have

$$\int \mathcal{D}\phi \phi_1 \dots \phi_n e^{iS[\phi]} = \int \mathcal{D}\phi' \phi'_1 \dots \phi'_n e^{iS[\phi + \varepsilon]} \quad (1.187)$$

$$= \int \mathcal{D}\phi e^{i(S[\phi] + \delta S)} (\phi_1 + \varepsilon_1) \dots (\phi_n + \varepsilon_n), \quad (1.188)$$

where we have denoted that $\phi_i \equiv \phi(x_i)$ and $\varepsilon_i \equiv \varepsilon(x_i)$. Then we expand ε to the first order, the exponential is $e^{i(S + \delta S)} \simeq e^{iS} (1 + i\delta S)$. And for δS , we have

$$\delta S = \int d^4y \left(\frac{\partial \mathcal{L}}{\partial \phi} \delta\phi + \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} \partial_\mu \delta\phi \right) \quad (1.189)$$

$$= \int d^4y [-m^2 \phi \varepsilon + \partial^\mu \phi \partial_\mu \varepsilon] \quad (1.190)$$

$$= - \int d^4y [\varepsilon(y) (\square_y + m^2) \phi(y)]. \quad (1.191)$$

Thus, put these result into (1.188), we have:

$$\begin{aligned} & \int \mathcal{D}\phi e^{i(S[\phi] + \delta S)} (\phi_1 + \varepsilon_1) \dots (\phi_n + \varepsilon_n) \\ &= \int \mathcal{D}\phi e^{iS[\phi]} \left[1 - i \int d^4y \varepsilon(y) (\square_y + m^2) \phi(y) \right] [(\phi_1 \dots \phi_n) + (\varepsilon_1 \phi_2 \dots \phi_n) + \dots + (\phi_1 \dots \varepsilon_n)] \end{aligned} \quad (1.192)$$

$$\begin{aligned}
&= \int \mathcal{D}\phi e^{iS[\phi]} [(\phi_1 \dots \phi_n) + (\varepsilon_1 \phi_2 \dots \phi_n) + \dots + (\phi_1 \dots \varepsilon_n)] \\
&\quad - \int \mathcal{D}\phi e^{iS[\phi]} \int d^4y i [\varepsilon(y)(\square_y + m^2)\phi(y)] (\phi_1 \dots \phi_n) + \mathcal{O}(\varepsilon^2).
\end{aligned} \tag{1.193}$$

Subtracting the first term $\int \mathcal{D}\phi \phi_1 \dots \phi_n e^{iS[\phi]}$ with (1.187), we have

$$0 = -i \int \mathcal{D}\phi e^{iS[\phi]} \left\{ \int d^4y \varepsilon(y) [(\square_y + m^2)\phi(y)] \phi_1 \dots \phi_n + i \sum_{i=1}^n \phi_1 \dots \varepsilon_i \dots \phi_n \right\}, \tag{1.194}$$

where ε_i will replace the corresponding ϕ_j when $i = j$ in the summation. And we can notice that $\varepsilon(x_i) = \int d^4y \varepsilon(y) \delta^{(4)}(x_i - y)$, so we have

$$\int d^4y \varepsilon(y) \left\{ -i \int \mathcal{D}\phi e^{iS[\phi]} \left[[(\square_y + m^2)\phi(y)] \phi_1 \dots \phi_n + i \sum_{i=1}^n \phi_1 \dots \delta^{(4)}(y - x_i) \dots \phi_n \right] \right\} = 0. \tag{1.195}$$

For arbitrary ε holds this relation, the only possible case is

$$\int \mathcal{D}\phi e^{iS[\phi]} \left[[(\square_y + m^2)\phi(y)] \phi_1 \dots \phi_n + i \sum_{i=1}^n \phi_1 \dots \delta^{(4)}(y - x_i) \dots \phi_n \right] = 0. \tag{1.196}$$

Example: Consider two-point correlator, i.e., $n = 1$:

$$(\square_y + m^2) \int \mathcal{D}\phi e^{iS[\phi]} \phi(y) \phi(x_1) = -i \delta^{(4)}(y - x_1) \int \mathcal{D}\phi e^{iS[\phi]} = -i \delta^{(4)}(y - x_1) Z[J = 0] \tag{1.197}$$

Putting it back in the expression for the correlator:

$$(\square_y + m^2) \underbrace{\langle \Omega | T \{ \phi(y) \phi(x_1) \} | \Omega \rangle}_{G_F(y-x_1)} = -i \delta^{(4)}(y - x_1). \tag{1.198}$$

Thus,

$$G_F(y - x_1) = -i \delta^{(4)}(y - x_1) (\square_y + m^2)^{-1}. \tag{1.199}$$

Similarly, we would like to generalize this form into n-point correlator. From (1.196) we have

$$\int \mathcal{D}\phi e^{iS[\phi]} (\square_y + m^2) \phi(y), (\phi_1 \dots \phi_n) = -i \int \mathcal{D}\phi e^{iS[\phi]} \left(\sum_{i=1}^n \phi_1 \dots \delta^{(4)}(y - x_i) \dots \phi_n \right). \tag{1.200}$$

Notice $-(\square_y + m^2)\phi(y)$ in the right-hand side is nothing but the variation of the action:

$$\frac{\delta S[\phi]}{\delta \phi(y)} = \int d^4x \left(\frac{\delta \mathcal{L}}{\delta \phi} \delta(x - y) + \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi} \partial_\mu \delta(x - y) \right) = \frac{\delta \mathcal{L}}{\delta \phi(y)} - \partial_\mu \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi(y)}, \tag{1.201}$$

the result of this variation always the equation of motion (or the Euler-Lagrange equation, if you want). So Without loss of generality, we can replace $-(\square_y + m^2)\phi(y)$ by $\frac{\delta S[\phi]}{\delta \phi(y)}$ in (1.200), i.e.,

$$\frac{\int \mathcal{D}\phi e^{iS[\phi]} \left(\frac{\delta S[\phi]}{\delta \phi(y)} \phi_1 \dots \phi_n \right)}{\int \mathcal{D}\phi e^{iS[\phi]}} = \frac{\int \mathcal{D}\phi e^{iS[\phi]} \left(i \sum_{i=1}^n \phi_1 \dots \delta^{(4)}(y - x_i) \dots \phi_n \right)}{\int \mathcal{D}\phi e^{iS[\phi]}}. \tag{1.202}$$

We then get the **Schwinger-Dyson equation** for n -point functions:

$$\left\langle \Omega \left| \frac{\delta S[\phi]}{\delta \phi(y)} T\{\phi_1 \dots \phi_n\} \right| \Omega \right\rangle = i \sum_{i=1}^n \langle \Omega | T\{\phi_1 \dots \delta^{(4)}(y - x_i) \dots \phi_n\} | \Omega \rangle. \quad (1.203)$$

We can simply denote it as:

$$\left\langle \frac{\delta S[\phi]}{\delta \phi(y)} T\{\phi_1 \dots \phi_n\} \right\rangle = i \sum_{i=1}^n \langle T\{\phi_1 \dots \delta^{(4)}(y - x_i) \dots \phi_n\} \rangle. \quad (1.204)$$

6.2 Conservation Laws and Noether's Theorem

Consider a complex scalar field $\mathcal{L} = \partial_\mu \phi \partial^\mu \phi^* - m^2 \phi \phi^*$. Under the global $U(1)$ transformation $\phi \rightarrow \phi' = e^{i\alpha} \phi$, the variation is:

$$\delta \phi = i\alpha \phi; \quad \text{for } \alpha \in \mathbb{R}, \quad (1.205)$$

with the unchanged Lagrangian. For a local transformation $\alpha \rightarrow \alpha(x)$, we have $\delta \phi(x) = i\alpha(x)\phi(x)$, so the variation for the Lagrangian is

$$\begin{aligned} \delta \mathcal{L} &= \mathcal{L}[\phi'] - \mathcal{L}[\phi] \\ &= \partial_\mu \phi \partial^\mu \phi^* + i\partial_\mu \alpha(x) (\phi \partial^\mu \phi^* - \phi^* \partial^\mu \phi) + \mathcal{O}(\alpha^2) - m^2 \phi \phi^* - \partial_\mu \phi \partial^\mu \phi^* + m^2 \phi \phi^* \\ &= \partial_\mu \alpha(x) j^\mu(x), \end{aligned} \quad (1.206)$$

where $j^\mu(x) = i(\phi \partial^\mu \phi^* - \phi^* \partial^\mu \phi)$. An n -point correlator is invariant under this transformation

$$\int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} (\phi_1 \dots \phi_{n/2}) (\phi_1^* \dots \phi_{n/2}^*) = \int \mathcal{D}\phi' \mathcal{D}\phi'^* e^{iS'[\phi', \phi'^*]} (\phi'_1 \dots \phi'_{n/2}) (\phi'^*_1 \dots \phi'^*_{n/2}), \quad (1.207)$$

Notice that $e^{iS'[\phi', \phi'^*]} = e^{i(S+\delta S)} \simeq e^{iS}(1 + i\delta S)$. Expand the right-hand side to order $\alpha(x)$, and subtracting with the left-hand side, we have

$$0 = \int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} (1 + i\delta S) ((1 + i\alpha)\phi_1 \dots (1 + i\alpha)\phi_n^*) \quad (1.208)$$

$$\begin{aligned} &= \int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} \left\{ i\delta S(\phi_1 \dots \phi_n^*) \right. \\ &\quad \left. + [i\alpha\phi_1] \dots [\phi_n^*] + \dots + [\phi_1 \dots (\pm i\alpha\phi_i^{(*)}) \dots \phi_n^*] + \dots + [\phi_1 \dots (-i\alpha\phi_n^*)] \right\} \end{aligned} \quad (1.209)$$

$$= \int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} \left\{ i \int d^4y \partial_\mu \alpha(y) j^\mu(y) (\phi_1 \dots \phi_n^*) + \sum_{i=1}^n \phi_1 \dots (\pm i\alpha\phi_i^{(*)}) \dots \phi_n^* \right\}, \quad (1.210)$$

where $\pm i\phi^{(*)}_i$ denote that for ϕ_i^* taking $-i$ and for ϕ_i taking $+i$. After partial integration to factor out $\alpha(y)$:

$$0 = \int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} \left\{ \int d^4y \alpha(y) \partial_\mu j^\mu(y) \phi_1 \dots \phi_n + i \sum_{i=1}^n \phi_1 \dots (\pm i) \left[\int d^4y \delta(y - x_i) \alpha(y) \right] \phi_i^{(*)} \dots \phi_n \right\} \quad (1.211)$$

$$= \int d^4y \alpha(y) \int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} \left\{ -i\partial_\mu j^\mu(y) \phi_1 \dots \phi_n + i \sum_{i=1}^n \phi_1 \dots (\pm i) \delta(y - x_i) \phi_i^{(*)} \dots \phi_n \right\}. \quad (1.212)$$

Similarly, for any infinitesimally α holding this relation, so the only possible way is

$$\int \mathcal{D}\phi \mathcal{D}\phi^* e^{iS[\phi, \phi^*]} \left\{ -i\partial_\mu j^\mu(y) \phi_1 \dots \phi_n + i \sum_{i=1}^n \phi_1 \dots (\pm i) \delta(y - x_i) \phi_i^{(*)} \dots \phi_n \right\} = 0. \quad (1.213)$$

We can also use the correlator notation:

$$\langle \partial_\mu j^\mu(y) \phi(x_1) \dots \phi(x_n) \rangle = -i \sum_{i=1}^n \langle \phi(x_1) \dots (\pm i) \delta(y - x_i) \phi_i^{(*)} \dots \phi(x_n) \rangle. \quad (1.214)$$

Now we generalize this relation to a set of real fields ϕ_a . For a local transformation $\phi_a(x) \rightarrow \phi'_a(x) = \phi_a(x) + \varepsilon \Delta \phi_a(x)$, if the global transformation gives $\mathcal{L} \rightarrow \mathcal{L} + \varepsilon \partial_\mu J^\mu$, then for a local $\varepsilon(x)$ we have

$$\partial_\mu \phi'_a = \partial_\mu (\phi_a + \varepsilon \Delta \phi_a) = \partial_\mu \phi_a + \varepsilon \partial_\mu (\Delta \phi_a) + (\partial_\mu \varepsilon) \Delta \phi_a. \quad (1.215)$$

So when we expand the Lagrangian, we have:

$$\mathcal{L}(\phi', \partial \phi') = \mathcal{L}(\phi, \partial \phi) + \sum_a \left(\frac{\partial \mathcal{L}}{\partial \phi_a} \delta \phi_a + \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} \delta (\partial_\mu \phi_a) \right), \quad (1.216)$$

then plug the variation $\delta \phi_a = \varepsilon \Delta \phi_a$ into it and using the Euler-Lagrange equation:

$$\delta \mathcal{L} = \sum_a \left[\frac{\partial \mathcal{L}}{\partial \phi_a} (\varepsilon \Delta \phi_a) + \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} (\varepsilon \partial_\mu (\Delta \phi_a) + (\partial_\mu \varepsilon) \Delta \phi_a) \right] \quad (1.217)$$

$$= \sum_a \left[\partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} (\varepsilon \Delta \phi_a) + \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} (\varepsilon \partial_\mu (\Delta \phi_a) + (\partial_\mu \varepsilon) \Delta \phi_a) \right] \quad (1.218)$$

$$= \sum_a \left[\varepsilon \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} \Delta \phi_a \right) + (\partial_\mu \varepsilon) \Delta \phi_a \right]. \quad (1.219)$$

Then we get the variation for the Lagrangian:

$$\mathcal{L}(\phi_a) \rightarrow \mathcal{L}(\phi'_a) = \mathcal{L}(\phi_a) + \varepsilon(x) \partial_\mu J^\mu + \partial_\mu \varepsilon(x) \sum_a \Delta \phi_a \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} \quad (1.220)$$

$$= \mathcal{L}(\phi_a) + \varepsilon(x) \partial_\mu J^\mu - \varepsilon(x) \partial_\mu \left(\sum_a \Delta \phi_a \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} \right) \quad (1.221)$$

$$= \mathcal{L}(\phi_a) - \varepsilon(x) \partial_\mu \left(\sum_a \Delta \phi_a \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} - J^\mu \right), \quad (1.222)$$

which appears an extra term due to the infinitesimal transformation rely on the coordinate. Therefore, the Noether current j^μ can be defined from the variation for the action

$$S[\phi'_a] = \int d^4y [\mathcal{L}[\phi_a] - \varepsilon(y) \partial_\mu j^\mu(y)] = S[\phi_a] - \int d^4y \varepsilon(y) \partial_\mu j^\mu(y). \quad (1.223)$$

Thus, the Noether current j^μ is

$$j^\mu = \sum_a \Delta \phi_a \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_a)} - J^\mu. \quad (1.224)$$

Put this current back to (1.214), we get the conserved symmetry Schwinger-Dyson equation as:

$$\langle \partial_\mu j^\mu(y) \phi_{a_1}(x_1) \dots \phi_{a_n}(x_n) \rangle = -i \sum_{i=1}^n \langle \phi_{a_1}(x_1) \dots \delta(y - x_i) \Delta \phi_{a_i} \dots \phi_{a_n}(x_n) \rangle \quad (1.225)$$

6.3 Ward-Takahashi Identity in QED

Consider the QED Lagrangian with fermions:

$$\mathcal{L}[\psi, \bar{\psi}, A_\mu] = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi + A_\mu j^\mu, \quad (1.226)$$

where $j^\mu = e\bar{\psi}\gamma^\mu\psi$. The global $U(1)$ transformation gives

$$\psi \rightarrow e^{i\alpha}\psi \implies \delta\psi = i\alpha\psi. \quad (1.227)$$

So the local transformation gives

$$\mathcal{L}[\psi', \bar{\psi}'] = \mathcal{L}[\psi, \bar{\psi}] - (\partial_\mu \alpha)j^\mu. \quad (1.228)$$

Applying Schwinger-Dyson equation with two fields, we have

$$\partial_\mu \langle 0|T\{j^\mu(y)\psi(x_1)\bar{\psi}(x_2)\}|0\rangle = \delta^{(4)}(y-x_1)\langle 0|T\{\psi(x_1)\bar{\psi}(x_2)\}|0\rangle - \delta^{(4)}(y-x_2)\langle 0|T\{\psi(x_1)\bar{\psi}(x_2)\}|0\rangle. \quad (1.229)$$

The minus sign shows that ψ follows **Fermi statistics**. Take a Fourier transform to momentum space:

$$k_\mu \langle 0|T\{j^\mu(k)\psi(q)\bar{\psi}(p)\}|0\rangle = e \left(\langle 0|T\{\psi(q-k)\bar{\psi}(p)\}|0\rangle - \langle 0|T\{\psi(q)\bar{\psi}(p+k)\}|0\rangle \right), \quad (1.230)$$

which is nothing but the **Ward-Takahashi identity** in QED that we are familiar with:

$$k_\mu \mathcal{M}^\mu(k; q, p) = e\mathcal{M}_0(p, q-k) - e\mathcal{M}_0(q, p+k). \quad (1.231)$$

It can also be expressed by the Feynman diagram:

$$k_\mu \left(\begin{array}{c} \text{Diagram 1: A shaded circle with an incoming wavy line from the left labeled } \mu \text{ and } k, \text{ and two outgoing fermion lines labeled } q \text{ and } p. \end{array} \right) = e \left(\begin{array}{c} \text{Diagram 2: A shaded circle with an incoming fermion line from the bottom labeled } p, \text{ and two outgoing lines labeled } q-k=p \text{ and } q. \end{array} \right) - \left(\begin{array}{c} \text{Diagram 3: A shaded circle with an incoming fermion line from the bottom labeled } p+k=q, \text{ and two outgoing lines labeled } q \text{ and } p. \end{array} \right). \quad (1.232)$$

Especially for the one loop case:

$$k_\mu \left(\begin{array}{c} \text{Diagram 1: A shaded circle with an incoming wavy line from the left labeled } \mu \text{ and } k, \text{ and two outgoing fermion lines labeled } q \text{ and } p. \end{array} \right) = e \left(\begin{array}{c} \text{Diagram 2: A shaded circle with an incoming fermion line from the bottom labeled } p, \text{ and two outgoing lines labeled } q-k=p \text{ and } q. \end{array} \right) - \left(\begin{array}{c} \text{Diagram 3: A shaded circle with an incoming fermion line from the bottom labeled } q=k+p, \text{ and two outgoing lines labeled } q \text{ and } p. \end{array} \right). \quad (1.233)$$

Renormalization

1. Renormalization in ϕ^3 -theory


1.1 One-loop self-energy

Simply speaking, renormalization is to solve the problem of divergent when we try to solve the loop graphs by subtracting the divergent part in the expression. In the following, we will use ϕ^3 -theory as an example to discuss the renormalization.

Firstly, the Lagrangian of ϕ^3 -theory is:

$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}m^2\phi^2 - \frac{g}{3!}\phi^3. \quad (2.1)$$

As the simplest case, the 1-loop correlation for the propagator is (notice that in this graph, there are **all inner lines**, i.e., propagators):



where the momentum k is **off-shelled**, which means k doesn't follow the relation $k^2 + E_k^2 = m^2$, so technically k could be very large. But we don't need to worry about it, because k only appears in the loop, which means that it must follow the 4-momentum conservation. Thus we need to sum over all possible k values, which finally will contribute a integral measurement when we calculate the loop part. Thus, generally, we can write the whole graph as

$$\frac{i}{p^2 - m^2 + i\epsilon} [-i\Sigma_1(p^2)] \frac{i}{p^2 - m^2 + i\epsilon}, \quad (2.2)$$

where $-i\Sigma_1(p^2)$ represent the loop part, i.e.,

$$-i\Sigma_1(p^2) = \frac{g^2}{2} \int \frac{d^4k}{(2\pi)^4} \frac{i}{k^2 - m^2 + i\epsilon} \frac{i}{(p+k)^2 - m^2 + i\epsilon}, \quad (2.3)$$

where we will note the integral measurement $\frac{d^4k}{(2\pi)^4} \rightarrow d^4k$ for simple. Next, we will focus on this integral as an example to talk about how to solve this type of integral.

1.2 Wick rotation

The integral in (2.3) over the loop energy can be also written as

$$I(p, \vec{k}) = \int_{-\infty}^{+\infty} dk^0 \frac{i}{(k^0)^2 - E_k^2 + i\epsilon} \frac{i}{(p^0 + k^0)^2 - E_{p+k}^2 + i\epsilon}, \quad (2.4)$$

where

$$E_k = \sqrt{\vec{k}^2 + m^2}, \quad E_{p+k} = \sqrt{(\vec{p} + \vec{k})^2 + m^2}. \quad (2.5)$$

The variable k^0 is now regarded as a complex variable. The four poles are

$$k^0 = +E_k - i\epsilon, \quad k^0 = -E_k + i\epsilon, \quad (2.6)$$

$$k^0 = -p^0 + E_{p+k} - i\epsilon, \quad k^0 = -p^0 - E_{p+k} + i\epsilon. \quad (2.7)$$

The small $i\epsilon$ is important: it tells us which poles sit slightly above or slightly below the real axis. Wick rotation means changing the integration path in the complex k^0 plane. Originally the integral is along the real axis,

$$C_M : \quad k^0 \in (-\infty, +\infty).$$

In Euclidean variables we use

$$k^0 = i\omega, \quad dk^0 = i d\omega,$$

so the new path is the imaginary axis,

$$C_E : \quad k^0 = i\omega, \quad \omega \in (-\infty, +\infty).$$

In the figures below, the **blue path** is the real-axis Minkowski contour C_M , the **green path** is the imaginary-axis Euclidean contour C_E , and the crosses are poles.

There is one orientation point which is very easy to mix up. The Euclidean contour itself is oriented upward, because

$$\omega : -\infty \rightarrow +\infty \implies k^0 = i\omega : -i\infty \rightarrow +i\infty.$$

But when we prove the Wick rotation by closing the contour, the closed contour is

$$C_{\text{closed}} = C_M + A_+ - C_E + A_-,$$

where A_+ runs from $+R$ to $+iR$ in the first quadrant, while A_- runs from $-iR$ back to $-R$ in the third quadrant. Thus the imaginary-axis part of the closed contour is the downward path $-C_E$. After moving it to the other side of the equation, the final Euclidean integral is again along C_E , upward. The orange dashed arrows in the figures show these auxiliary closing arcs, not the direction of the final integral along C_E .

The phrase “change the contour” means the following simple thing: we slide the integration line from one path to another in the complex k^0 plane. If the sliding line does not hit any pole, the integral is unchanged, apart from the Jacobian $dk^0 = i d\omega$. If the line hits a pole, then we must bend the path around that pole, and the difference is a residue term. Symbolically,

$$\int_{C_{\text{old}}} f(k^0) dk^0 - \int_{C_{\text{new}}} f(k^0) dk^0 = 2\pi i \sum_{\text{poles swept by the contour}} \text{Res } f, \quad (2.8)$$

up to the sign determined by the orientation of the contour.

Case 1: p^0 is imaginary.

Take, for example, $p^0 = -iP$ with $P > 0$. Then $-p^0 = +iP$, so the two poles from the second propagator are shifted upward:

$$k^0 = iP + E_{p+k} - i\epsilon, \quad k^0 = iP - E_{p+k} + i\epsilon.$$

This is the Euclidean starting point. The current integration path is C_E , not C_M .




Fig. 1: p^0 imaginary. The first-quadrant pole is harmless, because the contour currently being used is only the green line C_E .

The point which is easy to miss is this: the pole in the first quadrant does not automatically give a residue. A residue appears only when the contour is actually moved through that pole, or when the pole moves through the contour. In this first figure we are not rotating C_E into C_M while keeping p^0 imaginary. We are simply defining the Euclidean integral on the green line.

Case 2: p^0 is real and $|p^0| < m$.

Now keep the green contour C_E fixed, and analytically continue p^0 from imaginary value back to a real value. The poles move because their positions contain p^0 . If $|p^0| < m$, then

$$E_{p+k} \geq m > |p^0|,$$

so

$$-p^0 + E_{p+k} > 0, \quad -p^0 - E_{p+k} < 0.$$

Thus the second pair of poles ends up in the same pattern as the first pair: one pole is slightly below the real axis on the right, and one pole is slightly above the real axis on the left.




Fig. 2: p^0 real and $|p^0| < m$. No pole lies in the region swept when the blue contour is rotated into the green contour.

Therefore, for $|p^0| < m$, the real-axis integral and the imaginary-axis integral are related by an ordinary Wick rotation. On the green contour,

$$k^0 = i\omega, \quad k^2 = (k^0)^2 - \vec{k}^2 = -(\omega^2 + \vec{k}^2) < 0,$$

and similarly $(p+k)^2 < 0$ in the large Euclidean region. This is why power counting becomes Euclidean.

What does “a pole crosses a contour” mean?

A pole is not just a fixed dot: its position depends on the external energy p^0 . For example, if $p^0 = -P$ with $P > 0$, then one pole is

$$k^0 = -p^0 - E_{p+k} + i\epsilon = P - E_{p+k} + i\epsilon. \quad (2.9)$$

As P increases, this pole moves horizontally. If $P < E_{p+k}$, it is on the left side of the imaginary axis. If $P = E_{p+k}$, it sits on the imaginary-axis contour. If $P > E_{p+k}$, it has moved to the right side of the imaginary axis. This is what we mean by saying that the pole has crossed C_E .

In formula language, crossing the imaginary axis means

$$\text{Re } k_{\text{pole}}^0 \text{ changes sign.}$$

Crossing the real axis would similarly mean

$$\text{Im } k_{\text{pole}}^0 \text{ changes sign.}$$

But only crossing the contour that we are actually using produces a residue.

Case 3: p^0 is real and $|p^0| > m$.

Take the negative-energy example $p^0 = -P$ with $P > m$, which is the case drawn in the book. The pole

$$k^0 = P - E_{p+k} + i\epsilon$$

crosses the green contour whenever

$$P > E_{p+k} \iff (p^0)^2 > m^2 + (\vec{p} + \vec{k})^2.$$

When this happens, the contour cannot pass straight through the pole. It must detour around it, and this detour gives an extra residue, usually called the pole term.




Fig. 3: $p^0 = -P$ and $P > m$. A pole crosses the green contour, so the Wick-rotated answer contains an extra residue term.

However, this pole term can occur only when

$$(p^0)^2 > m^2 + (\vec{p} + \vec{k})^2.$$

For very large $|\vec{k}|$, the right-hand side is very large, so the inequality cannot hold. Therefore the pole term comes only from a finite region of $\vec{k}$, not from the UV region $|\vec{k}| \rightarrow \infty$. The UV divergence is still captured by the Euclidean large-momentum integral along C_E .

So the divergence in Eq.(2.3) is called the UV divergence, which can be seen by firstly taking the polar coordinate: $d^4k_E = k_E^3 dk_E d\Omega$, thus in large k

$$\Sigma_1 \sim \int d^d k \frac{1}{k^4} \sim \int dk d\Omega \frac{1}{k} \sim \int \frac{dk}{k} \sim \ln \Lambda \rightarrow \infty. \quad (2.10)$$

As we said above, renormalization is a way to cancel the divergent. So first of all, we need to recognize which part is finite and which part is divergent. Thus, we introduce a finite momentum μ to avoid new divergent in $k = 0$, so we have

$$\begin{aligned} \Sigma_1 = \frac{ig^2}{32\pi^4} \left[\int d^4k \left\{ \frac{1}{(k^2 - m^2 + i\epsilon)((p+k)^2 - m^2 + i\epsilon)} - \frac{1}{(k^2 - \mu^2 + i\epsilon)^2} \right\} \right. \\ \left. + \int d^4k \frac{1}{(k^2 - \mu^2 + i\epsilon)^2} \right]. \end{aligned} \quad (2.11)$$

In a large k , the expression inside the first integral behaves as

$$\frac{1}{k^4} \left(1 + O\left(\frac{1}{k^2}\right) \right) - \frac{1}{k^4} = O\left(\frac{1}{k^6}\right). \quad (2.12)$$

Thus $\int d^4k/k^6$ is UV finite, and the logarithmic UV divergence has been isolated in the second integral. This second integral is independent of the external momentum p , which is why the corresponding counterterm is local.

To understand the divergent integral, we need to work in a space-time lattice with spacing a , the free propagator is then $S_F(k; \mu, a)$, so we have

$$\Sigma_1 = \frac{ig^2}{32\pi^4} \left[\int d^4k S_F(k; m, a) S_F(p+k; m, a) - S_F^2(k; \mu, a) + \int d^4k S_F^2(k; m, a) \right] \quad (2.13)$$

$$\equiv \Sigma_{1,\text{fin}}(p, m, \mu, a) + \Sigma_{1,\text{div}}(m, \mu, a). \quad (2.14)$$

The $\Sigma_{1,\text{div}}(m, \mu, a)$ in the lattice diverges as

$$\Sigma_{1,\text{div}}(m, \mu, a) = \frac{ig^2}{32\pi^4} \int d^4k S_F^2(k; \mu, a) \quad (2.15)$$

$$= -\frac{ig^2}{32\pi^4} \int_{\omega^2 + \vec{k}^2 < \frac{1}{a^2}} d\omega d^3\vec{k} \frac{1}{(\omega^2 + \vec{k}^2 + \mu^2)^2} \quad (2.16)$$

$$= -\frac{ig^2}{16\pi^2} \int_0^{1/a} dk \frac{k^3}{(k^2 + \mu^2)^2} = -\frac{g^2}{16\pi^2} \ln \frac{1}{a}, \quad \text{as } a \rightarrow 0. \quad (2.17)$$

This result just as same as our analysis in Eq.(2.10).

Thus we can interpret the divergence in a more general way. We consider all the loop corrections for the propagator. Let $i\Sigma$ be the sum of 1-PI, the propagator is then

$$G_2(p^2) = \frac{p}{\longrightarrow} + \frac{p}{\longrightarrow} \textcircled{1\text{PI}} \frac{p}{\longrightarrow} + \frac{p}{\longrightarrow} \textcircled{1\text{PI}} \textcircled{1\text{PI}} \frac{p}{\longrightarrow} + \dots \quad (2.18)$$

$$= \frac{i}{p^2 - m^2} + \frac{i}{p^2 - m^2} (-i\Sigma) \frac{i}{p^2 - m^2} + \frac{i}{p^2 - m^2} (-i\Sigma) \frac{i}{p^2 - m^2} (-i\Sigma) \frac{i}{p^2 - m^2} + \dots \quad (2.19)$$

$$= \frac{i}{p^2 - m^2} + \frac{i}{p^2 - m^2} \frac{\Sigma}{p^2 - m^2} + \frac{i}{p^2 - m^2} \left(\frac{\Sigma}{p^2 - m^2} \right)^2 + \dots \quad (2.20)$$

$$= \frac{i}{p^2 - m^2} \frac{p^2 - m^2}{p^2 - m^2 - \Sigma} \quad (2.21)$$

$$= \frac{i}{p^2 - m^2 - \Sigma} \equiv \frac{i}{p^2 - m_{\text{ph}}^2}, \quad (2.22)$$

where we have used $\sum_{i=0}^{\infty} a_i = \frac{1}{1-a}$. And $m_{\text{ph}}^2 = m^2 + \Sigma(p^2 = m_{\text{ph}}^2, m^2)$ in the pole define as $p^2 = m_{\text{ph}}^2$. The physical mass m_{ph} is the mass that can be directly observed. So the goal is changing Σ_1 to Σ_{1R} , which all the mass in it should be physical mass, i.e.,

$$\Sigma_1 = \frac{ig^2}{32\pi^4} \int d^4k \frac{1}{(k^2 - m^2 + i\epsilon)((p+k)^2 - m^2 + i\epsilon)} \quad (2.23)$$

$$\Rightarrow \Sigma_1 = \frac{ig^2}{32\pi^4} \int d^4k \frac{1}{(k^2 - m_{\text{ph}}^2 + i\epsilon)((p+k)^2 - m_{\text{ph}}^2 + i\epsilon)}. \quad (2.24)$$

For that, we suppose the mass term in Lagrangian satisfies

$$\frac{1}{2}m_0^2\phi^2 = \frac{1}{2}m_{\text{ph}}^2\phi^2 + \frac{1}{2}\delta m^2\phi^2 \quad (2.25)$$

The last term is called the **counterterm**, which will contribute a graph of

$$\text{---} \otimes \text{---} = -i\delta m^2.$$

Thus, the renormalized self-energy to $O(g^2)$ is defined as

$$-i\Sigma_{1R} = \frac{p}{\longrightarrow} \textcircled{\text{loop}} \frac{p}{\longrightarrow} + \frac{p}{\longrightarrow} \otimes \frac{p}{\longrightarrow} = -i\Sigma_1 - i\delta m^2 \quad (2.26)$$

$$\Rightarrow \Sigma_{1R} = \Sigma_1 + \delta m^2 = (\Sigma_{1,\text{div}} + \delta m^2) + \Sigma_{1,\text{fin}} \quad (2.27)$$

Since the counterterm δm^2 can be chosen freely, we can then always choose $\delta m^2 = -\Sigma_{1,\text{div}}$, such that the UV-divergent cancels.

Therefore, to calculate Σ_{1R} , we have many ways, one way is firstly differentiate with respect to p^2 , and use the fact that only Σ_1 contains p^μ , so we have

$$\frac{\partial \Sigma_{1R}}{\partial p^2} = \frac{\partial p^\mu}{\partial p^2} \frac{\partial}{\partial p^\mu} \Sigma_{1R} = \frac{p^\mu}{2p^2} \frac{\partial}{\partial p^\mu} \Sigma_1 \quad (2.28)$$

$$= \frac{ig^2}{32\pi^4} \frac{p^\mu}{2p^2} \int d^4k \frac{\partial}{\partial p^\mu} \left[\frac{1}{(k^2 - m^2 + i\epsilon)((p+k)^2 - m^2 + i\epsilon)} - \frac{1}{(k^2 - \mu^2 + i\epsilon)^2} \right] \quad (2.29)$$

$$= \frac{ig^2}{64\pi^4} \frac{p^\mu}{2p^2} \int d^4k \frac{-2(p+k)_\mu}{(k^2 - m^2 + i\epsilon)((p+k)^2 - m^2 + i\epsilon)^2} \quad (2.30)$$

$$= \frac{-ig^2}{32\pi^4 p^2} \int d^4k \frac{p \cdot (p+k)}{(k^2 - m^2 + i\epsilon)((p+k)^2 - m^2 + i\epsilon)^2}. \quad (2.31)$$

To calculate this integral, we need to introduce a Feynman parameter x and use the fact:

$$\frac{1}{AB^n} = \int_0^1 dx \frac{nx^{n-1}}{(xB + (1-x)A)^{n+1}}. \quad (2.32)$$

Thus,

$$\frac{\partial \Sigma_{1R}}{\partial p^2} = \frac{-ig^2}{32\pi^4 p^2} \int_0^1 dx \int d^4k \frac{2p \cdot (p+k)x}{(-m_{\text{ph}}^2 + p^2x + 2p \cdot kx + k^2 + i\epsilon)^3} \quad (2.33)$$

$$= \frac{ig^2}{16\pi^4 p^2} \int_0^1 dx \int d^4k \frac{p \cdot (p+k)x}{(m_{\text{ph}}^2 - p^2x - 2p \cdot kx - k^2 - i\epsilon)^3} \quad (2.34)$$

$$= \frac{ig^2}{16\pi^4 p^2} \int_0^1 dx \int d^4k \frac{p \cdot (p+k+px-px)x}{(m_{\text{ph}}^2 - p^2x - (k+px)^2 + p^2x^2 - i\epsilon)^3}, \quad (2.35)$$

let $k' = k + px$

$$\frac{\partial \Sigma_{1R}}{\partial p^2} = \frac{ig^2}{16\pi^4 p^2} \int_0^1 dx \int d^4k' \frac{p^2(1-x)x + p \cdot k'x}{(m_{\text{ph}}^2 - p^2x(1-x) - k'^2 - i\epsilon)^3} \quad (2.36)$$

$$= \frac{ig^2}{16\pi^4 p^2} \int_0^1 dx \int d^4k' \frac{p^2x(1-x)}{(m_{\text{ph}}^2 - p^2x(1-x) - k'^2)^3} \quad (2.37)$$

$$= ig^2 \int_0^1 dx x(1-x) \int d^4k' \frac{1}{(k'^2 - \Delta)^3}. \quad (2.38)$$

Use the integral fact

$$\int \frac{d^d l}{(2\pi)^d} \frac{1}{(l^2 - \Delta)^m} = \frac{i(-1)^m}{(4\pi)^{d/2}} \frac{\Gamma(m - d/2)}{\Gamma(m)} \Delta^{d/2-m}, \quad (2.39)$$

with $\Delta = m_{\text{ph}}^2 - p^2x(1-x)$. We then have

$$\frac{\partial \Sigma_{1R}}{\partial p^2} = -\frac{g^2}{32\pi^2} \int_0^1 dx \frac{x(1-x)}{m_{\text{ph}}^2 - p^2x(1-x)} \quad (2.40)$$

$$= \frac{g^2}{32\pi^2} \frac{\partial}{\partial p^2} \int_0^1 dx \ln[m_{\text{ph}}^2 - p^2x(1-x)]. \quad (2.41)$$

Thus, we can integrate p^2 in both side, then we read


$$\Sigma_{1R} = \frac{g^2}{32\pi^2} \int_0^1 dx \ln[m_{\text{ph}}^2 - p^2x(1-x)] + C. \quad (2.42)$$

The boundary condition is at $p^2 = m_{\text{ph}}^2$, $\Sigma_{1R} = 0$, so we can determine the integration constant, therefore, we obtain the final result as:


$$\Sigma_{1R} = \frac{g^2}{32\pi^2} \int_0^1 dx \ln \left[\frac{m_{\text{ph}}^2 - p^2 x(1-x)}{m_{\text{ph}}^2(1-x+x^2)} \right]. \quad (2.43)$$

For higher order graphs, the same idea is used inside larger graphs. If a one-loop bubble appears as a subgraph, we renormalize that small part first:

1. First draw the graph before the subgraph is subtracted. The one-loop bubble may sit vertically on another line, or several bubbles may sit on the same propagator line:



2. Then replace one or more divergent bubbles by their counterterm. The cross means “use the local counterterm instead of the loop”:



3. After adding the original graph and these counterterm graphs, the divergent bubble is replaced by its finite renormalized value $-\text{i}\Sigma_{1R}$.

In words, we do not wait until the whole large graph is finished. We remove the divergence as soon as a divergent subgraph appears. This is why counterterm graphs have to be included together with ordinary loop graphs.

1.3 Degree of divergence

$$\Sigma_1(p^2, m^2, d) = \frac{\text{i}g^2}{2(2\pi)^d} \int d^d k \frac{1}{(k^2 - m^2 + \text{i}\epsilon)((p+k)^2 - m^2 + \text{i}\epsilon)} \text{ has } d-4 \text{ dim for } k \quad (2.44)$$

we call it as degree of divergence of the graph.

$$\text{If } d < 4 \implies \text{graph is convergent.} \quad (2.45)$$

$$d \geq 4 \implies \text{graph is divergent} \implies \text{needs to be renormalized} \quad (2.46)$$

Differentiating once

$$\frac{\partial}{\partial p^\mu} \Sigma_1 = \frac{-\text{i}g^2}{(2\pi)^d} \int d^d k \frac{(p+k)_\mu}{(k^2 - m^2 + \text{i}\epsilon)((p+k)^2 - m^2 + \text{i}\epsilon)^2} \quad (2.47)$$

degree of divergence is $d-5$. if $d=5$, it diverges logarithmically.

Symmetrizing by $k \rightarrow -k$ will kill the divergence. Differentiating again:

$$\frac{\partial^2}{\partial p^\mu \partial p^\nu} \Sigma_1 = \frac{\text{i}g^2}{(2\pi)^d} \int d^d k \frac{2(p+k)_\mu(p+k)_\nu - g_{\mu\nu}((p+k)^2 - m^2)}{(k^2 - m^2)^2((p+k)^2 - m^2)^3} \quad (2.48)$$

The degree of divergence is $d-6$. Integrating twice will get Σ_{1R} with 2 constants in the form: $A + B_\mu p^\mu$. Since Σ must be Lorentz invariant, thus $B_\mu = 0$. Thus only the mass term is the counterterm.

If $d = 6$, differentiating Σ_1 3 times will get finite results with degree of divergence $d - 7$. Integrating gives the constant $A + Bp^2 + C_\mu p^\mu$, where $C_\mu = 0$ because of Lorentz invariance condition. Thus, the counterterm includes the mass and another proportion to p^2 , i.e.

$$\delta m^2 - \delta Z p^2 \quad (2.49)$$

generated from the Lagrangian

$$-\frac{1}{2}\delta m^2 \phi^2 + \frac{1}{2}\delta Z (\partial \cdot \phi)^2 \subset \mathcal{L} \quad (2.50)$$

where δZ is an example of wave function renormalization. We can understand it from the bare propagator:

$$G_{2(0)} = \langle 0 | T \{ \phi_0(p) \phi_0(0) \} | 0 \rangle \quad (2.51)$$

$$= Z \langle 0 | T \{ \tilde{\phi}(p) \phi(0) \} | 0 \rangle = Z G_2 \quad (\text{renormalized propagator}) \quad (2.52)$$

For p^2 not at the pole ($p^2 = m_{ph}^2$), so the propagator cannot be written as

$$G_2(p^2 = m_{ph}^2) = \frac{i}{p^2 - m_{ph}^2} \quad (2.53)$$

but only be

$$G_2(p^2) = \frac{i}{p^2 - m_0^2 - \Sigma_1(p^2)} = \frac{i}{p^2 - m_0^2 - \Sigma_1(m_{ph}^2) + \Sigma_1(m_{ph}^2) - \Sigma_1(p^2)} \quad (2.54)$$

$$\text{where } \Sigma_1(m_{ph}^2) = m_{ph}^2 - m_0^2, \quad \Sigma_{1R}(p^2) = \Sigma_1(p^2) - \Sigma_1(m_{ph}^2) \quad (2.55)$$

$$= \frac{i}{p^2 - m_{ph}^2 - \Sigma_{1R} + i\epsilon} \quad (2.56)$$

where $\Sigma_{1R}(p^2) = \Sigma_1 + \text{counterterm} = \Sigma_{1(0)} + \delta m^2 + \delta Z p^2$

$$\implies G_{2(0)} = \frac{iZ}{p^2 - m_{ph}^2 - \Sigma_{1(0)} + \delta m^2 + \delta Z p^2} = \frac{iZ}{p^2 - m_{ph}^2 - \Sigma_{1R} + i\epsilon} \quad (2.57)$$

where $\Sigma_{1R}(m_{ph}^2) = 0$ (**On-shell renormalization condition**).

The residue of $G_{2(0)}$ is $R_{(0)}$ around $p^2 = m_{ph}^2$

$$\implies G_{2(0)} = \frac{iR_{(0)}}{p^2 - m_{ph}^2} + \text{finite} \quad \text{as } p^2 \rightarrow m_{ph}^2 \quad (2.58)$$

In bare theory: $G_{2(0)} = \frac{iR_{(0)}}{p^2 - m_{ph}^2} + \text{finite}$, residue = $0 \leq R_{(0)} < 1$. In interaction theory: $G(p) = \frac{i}{p^2 - m_{ph}^2}$, residue = 1. For renormalized propagator $G_2 = \frac{i}{p^2 - m_{ph}^2 - \Sigma_{1R}}$. There are two on-shell requirements. The first one fixes the pole position,

$$\Sigma_{1R}(m_{ph}^2) = 0, \quad (2.59)$$

and the second one fixes the residue of the pole to be one. To see the second condition explicitly, expand Σ_{1R} around $p^2 = m_{ph}^2$:

$$\Sigma_{1R}(p^2) = \Sigma_{1R}(m_{ph}^2) + (p^2 - m_{ph}^2)\Sigma'_{1R}(m_{ph}^2) + O((p^2 - m_{ph}^2)^2) \quad (2.60)$$

$$= (p^2 - m_{ph}^2)\Sigma'_{1R}(m_{ph}^2) + O((p^2 - m_{ph}^2)^2). \quad (2.61)$$

Hence, near the pole,

$$G_2(p^2) \simeq \frac{i}{(p^2 - m_{ph}^2)[1 - \Sigma'_{1R}(m_{ph}^2)]}. \quad (2.62)$$

If we choose the renormalized field so that the pole residue is one, then the second on-shell condition is

$$\Sigma'_{1R}(m_{ph}^2) = 0. \quad (2.63)$$

Using $G_{2(0)} = ZG_2$, we have

$$G_{2(0)}(p^2 \rightarrow m_{ph}^2) = ZG_2(p^2 \rightarrow m_{ph}^2) = \frac{iZ}{p^2 - m_{ph}^2} \quad (2.64)$$

So if we require the residue = 1, then

$$Z = R_{(0)} \quad (2.65)$$

Thus we can require Z defined by $\phi = Z^{-1/2}\phi_0$ satisfies

$$Z = Z \times \text{finite constant} \quad (2.66)$$

For bare propagator, $\Sigma_{1(0)}(p^2) \simeq \Sigma_{1(0)}(m_{ph}^2) + (p^2 - m_{ph}^2)\frac{\partial}{\partial p^2}\Sigma_{1(0)}(m_{ph}^2)$

$$G_{2(0)} = \frac{i}{p^2 - m_0^2 - \Sigma_{1(0)}} \simeq \frac{i}{(p^2 - m_{ph}^2)(1 - \Sigma'_{1(0)})} \quad (2.67)$$

$$Z = \frac{1}{1 - \Sigma'_{1(0)}} = R_{(0)} \quad (2.68)$$

For bare ϕ^3 : $\mathcal{L} = \frac{1}{2}(\partial\phi_0)^2 - \frac{1}{2}m_0^2\phi_0^2 - \frac{g_0}{3!}\phi_0^3$ def: field strength renormalization: $\phi = Z^{-1/2}\phi_0$

$$\implies \mathcal{L} = \frac{1}{2}Z(\partial\phi)^2 - \frac{1}{2}Zm_0^2\phi^2 - \frac{g_0}{3!}Z^{3/2}\phi^3 \quad (2.69)$$

$$= \frac{1}{2}(1 - \delta Z)(\partial\phi)^2 - \frac{1}{2}(m_{ph}^2 + \delta m^2)\phi^2 - \frac{1}{3!}(g_{ph} + \delta g)\phi^3 \quad (2.70)$$

where we have defined:
$$\begin{cases} Z = 1 - \delta Z \\ Zm_0^2 = m_{ph}^2 + \delta m^2 \\ Z^{3/2}g_0 = g_{ph} + \delta g \end{cases}$$

$$\implies \mathcal{L} = \frac{1}{2}(\partial\phi)^2 - \frac{1}{2}m_{ph}^2\phi^2 - \frac{g_{ph}}{3!}\phi^3 - \frac{1}{2}\delta Z(\partial\phi)^2 - \frac{1}{2}\delta m^2\phi^2 - \frac{1}{3!}\delta g\phi^3 \quad (2.71)$$

1.4 Arbitrariness in a renormalization graph

Consider 1-loop graph in $d = 4$, we choose $\delta Z = 0$ so the counterterm is δm^2 . The important point is that the divergent part of the counterterm is fixed by the requirement of finiteness, while its finite part is a convention. Different conventions are called different renormalization schemes. They do not change the bare theory; they only change which finite parameter we decide to call the renormalized mass.

$$\Sigma_{1R}(p^2) = \Sigma_{1,fin}(p^2) + (\delta m^2 + \Sigma_{1,div}) \quad (2.72)$$

In the mass-shell scheme we require that the pole of the full propagator is at $p^2 = m_{ph}^2$. This is the condition

$$\Sigma_{1R}(m_{ph}^2) = 0. \quad (2.73)$$

Substituting $p^2 = m_{ph}^2$ into the previous line gives

$$\implies \delta m^2 = -\Sigma_{1,div} - \Sigma_{1,fin}(p^2 = m_{ph}^2) \quad (2.74)$$

$$\implies \Sigma_{1R}(p^2) = \Sigma_{1,fin}(p^2) - \Sigma_{1,fin}(p^2 = m_{ph}^2) \quad (2.75)$$

$$\implies G_2 = \frac{i}{p^2 - m_{ph}^2 - \Sigma_{1R}} = \frac{i}{p^2 - m_{ph}^2 - \Sigma_{1,fin}(p^2) + \Sigma_{1,fin}(m_{ph}^2) + O(g^4)} \quad (2.76)$$

$$m_0^2 = m_{ph}^2 + \delta m^2 = m_{ph}^2 - \Sigma_{1,div} - \Sigma_{1,fin}(p^2 = m_{ph}^2) + O(g^4)$$

One can also compute with different finite part to δm^2 . Generally, the counterterm only requires to cancel the divergent. So it can be written as

$$\delta m^2 = -\Sigma_{1,div} + \text{finite} \quad (2.77)$$

Thus, we can choose

$$\delta m^2 = -\Sigma_{1,div} + g^2 C \quad (2.78)$$

where C is a const. The self-energy is now

$$\Sigma_{1R}^{(C)}(p^2, m^2) = \Sigma_{1,fin}(p^2) + g^2 C \quad (2.79)$$

Here m is the mass parameter defined by this new scheme. It is not automatically equal to the physical pole mass. The pole mass is obtained only after solving

$$p^2 - m^2 - \Sigma_{1R}^{(C)}(p^2, m^2) = 0 \quad (2.80)$$

order by order in g .

$$\implies G_2 = \frac{i}{p^2 - m^2 - \Sigma_{1R}^{(C)}(p^2, m^2)} = \frac{i}{p^2 - m^2 - \Sigma_{1,fin}(p^2) - g^2 C + O(g^4)} \quad (2.81)$$

$$m_0^2 = m^2 - \Sigma_{1,div} - g^2 C + O(g^4) \quad (2.82)$$

Thus, we have at least **two ways** to renormalize the theory that gives the same result if m_0^2 are the same. So obviously we have

$$m^2 = m_{ph}^2 + g^2 C - \Sigma_{1,fin}(p^2 = m_{ph}^2) + O(g^4) \quad (2.83)$$

This equation is the finite redefinition between two schemes. The left hand side is the scheme-dependent mass parameter, while m_{ph} is defined by the pole and is observable.

1.5 Dimensional regularization

As another way to deal with the divergent, Dimensional regularization is more convenient. Take the dimension as a regulator d which is a continuous variable. In the end, the cut-off will be removed by taking the limit $d \rightarrow 4$. We also use 1-loop in ϕ^3 as an example. The integral is first evaluated where it converges, and the answer is then analytically continued as a function

of d . Therefore the divergence no longer appears as a large cut-off Λ ; it appears as a pole of a Gamma function at the physical value of d .

$$\Sigma_1(p^2, m^2, d) = \frac{ig^2}{2} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[k^2 - m^2 + i\epsilon][(p+k)^2 - m^2 + i\epsilon]} \quad (2.84)$$

The Gaussian integral is

$$\int d^n x e^{-x^2} = (\sqrt{\pi})^n = \pi^{n/2} \quad (2.85)$$

Thus, after Wick rotation, $k^0 = i\omega$, we have

$$\int d^d k e^{k^2} = i \int d\omega \int d^{d-1} \vec{k} e^{-(\omega^2 + \vec{k}^2)} \quad (2.86)$$

$$= i \left(\int d\omega e^{-\omega^2} \right) \left(\int d^{d-1} \vec{k} e^{-\vec{k}^2} \right) \quad (2.87)$$

$$= i\sqrt{\pi} \cdot \pi^{\frac{d-1}{2}} = i\pi^{d/2} \quad (2.88)$$

Notice that

$$\frac{i}{m^2 - k^2 - i\epsilon} = \int_0^\infty da e^{-ia(m^2 - k^2 - i\epsilon)} \quad (2.89)$$

Thus,

$$\Sigma_1 = \frac{ig^2}{2} \int \frac{d^d k}{(2\pi)^d} \left(\int_0^\infty da e^{-ia(m^2 - k^2 - i\epsilon)} \right) \left(\int_0^\infty db e^{-ib(m^2 - (p+k)^2 - i\epsilon)} \right) \quad (2.90)$$

$$= \frac{ig^2}{2(2\pi)^d} \int d^d k \int_0^\infty da \int_0^\infty db e^{-i[(a+b)m^2 + (a+b)k^2 + bp^2 + 2bp \cdot k]} \quad (2.91)$$

$$= \frac{ig^2}{2(2\pi)^d} \int d^d k \int_0^\infty da \int_0^\infty db e^{-i[(a+b)m^2 + (a+b)(k + \frac{b}{a+b}p)^2 + bp^2(1 - \frac{b}{a+b})]} \quad (2.92)$$

$$\text{take } k' = k + \frac{b}{a+b}p, \quad z = a+b, \quad x = \frac{a}{z} \implies a = xz, \quad b = (1-x)z$$

$$\implies dadb = J dx dz, \quad J = \begin{vmatrix} \frac{\partial a}{\partial x} & \frac{\partial a}{\partial z} \\ \frac{\partial b}{\partial x} & \frac{\partial b}{\partial z} \end{vmatrix} = \begin{vmatrix} z & x \\ -z & 1-x \end{vmatrix} = z \quad (2.93)$$

$$\implies dadb = z dx dz, \quad z \in (0, \infty), \quad x = \frac{a}{a+b} \in (0, 1) \quad (2.94)$$

$$\implies \Sigma_1 = \frac{ig^2}{2(2\pi)^d} \int d^d k' \int_0^1 dx \int_0^\infty dz \cdot z \cdot e^{-izm^2 + izk'^2 + izp^2 x(1-x)} \quad (2.95)$$

$$\text{take } k'' = z^{1/2} k'$$

$$= \frac{ig^2}{2(2\pi)^d} \int_0^1 dx \int_0^\infty dz \cdot z^{1-d/2} e^{-iz(m^2 - p^2 x(1-x))} \left(\int d^d k'' e^{ik''^2} \right) \quad (2.96)$$

$$= \frac{g^2}{2(4\pi)^{d/2}} \int_0^1 dx \int_0^\infty d(\alpha z) (\alpha z)^{(2-d/2)-1} e^{-\alpha z} \quad [\Gamma(t) = \int_0^\infty dz \frac{z^{t-1}}{e^z}] \quad (2.97)$$

$$= -\frac{g^2}{2(4\pi)^{d/2}} \Gamma(2 - \frac{d}{2}) \int_0^1 dx [m^2 - p^2 x(1-x)]^{d/2-2} \quad (2.98)$$

We can use Feynman parameter to get the same result:

$$\Sigma_1 = \frac{ig^2}{2} \int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 - m^2 + i\epsilon)((p+k)^2 - m^2 + i\epsilon)} \quad (2.99)$$

$$= \frac{ig^2}{2(2\pi)^d} \int d^d k \int_0^1 dx \frac{1}{[k^2 - m^2 + xp^2 + 2xp \cdot k]^2} \quad (2.100)$$

$$= \frac{ig^2}{2(2\pi)^d} \int d^d k \int_0^1 dx \frac{1}{[(k+xp)^2 + p^2 x(1-x) - m^2]^2} \quad (2.101)$$

take $k' = xp + k$, and do Wick rotation, i.e. $k^0 = ik_E^0 \implies k'^2 = -k_E^2$

$$= \frac{-ig^2 \cdot i}{2(2\pi)^d} \int_0^1 dx \int d^d k_E \frac{1}{[-k_E^2 - m^2 + p^2 x(1-x)]^2} \quad (2.102)$$

$$= \frac{-g^2}{2(2\pi)^d} \int_0^1 dx \int d^d k_E \frac{1}{(k_E^2 + \Delta)^2}, \quad \Delta = m^2 - p^2 x(1-x) \quad (2.103)$$

$$= \frac{-g^2}{2(4\pi)^{d/2}} \Gamma(2 - \frac{d}{2}) \int_0^1 dx \frac{1}{(m^2 - p^2 x(1-x))^{2-d/2}} \quad (2.104)$$

Now, the divergence only in the Γ -function's poles, at $d = 4, 6, 8, \dots$. The advantage of dimensional regularization is:

1. Calculation procedure is unchanged.
2. Preserve Poincaré invariance and gauge symmetries.
3. It can also be useful for IR-divergence (infra-red), e.g. $\frac{ig^2}{8\pi^2} \int d^4 k \frac{1}{(k^2 + i\epsilon)((p+k)^2 + i\epsilon)}$

Minimal subtraction

Now we can compute the renormalized self-energy from Σ_1 above, i.e.

$$\Sigma_{1R}^{(ph)} = \Sigma_1(p^2, m_{ph}^2, g, d) - \Sigma_1(m_{ph}^2, m_{ph}^2, g, d) \quad (2.105)$$

$$= -\frac{g^2}{2(4\pi)^{d/2}} \Gamma(2 - \frac{d}{2}) \int_0^1 dx \{ [m_{ph}^2 - p^2 x(1-x)]^{d/2-2} - [m_{ph}^2(1-x+x^2)]^{d/2-2} \} \quad (2.106)$$

To expand it, set $d = 4 - \epsilon$. The two elementary expansions are

$$\Gamma(z) = \frac{1}{z} - \gamma_E + O(z), \quad z \rightarrow 0, \quad (2.107)$$

$$A^\delta = e^{\delta \ln A} = 1 + \delta \ln A + O(\delta^2), \quad \delta \rightarrow 0. \quad (2.108)$$

Thus

$$\Gamma\left(2 - \frac{d}{2}\right) = \Gamma\left(\frac{\epsilon}{2}\right) = \frac{2}{\epsilon} - \gamma_E + O(\epsilon), \quad (2.109)$$

$$[m_{ph}^2 - p^2 x(1-x)]^{\frac{d}{2}-2} = 1 - \frac{\epsilon}{2} \ln[m_{ph}^2 - p^2 x(1-x)] + O(\epsilon^2). \quad (2.110)$$

Similarly,

$$[m_{ph}^2(1-x+x^2)]^{\frac{d}{2}-2} = 1 - \frac{\epsilon}{2} \ln[m_{ph}^2(1-x+x^2)] + O(\epsilon^2). \quad (2.111)$$

Subtracting the two terms in the curly bracket removes the order-one part before the pole multiplies it. This is why the mass-shell subtraction gives a finite answer:

$$\Sigma_{1R}^{(ph)} = \frac{g^2}{32\pi^2} \left(\frac{2}{\epsilon} - \gamma_E \right) \frac{\epsilon}{2} \int_0^1 dx \ln \frac{m_{ph}^2 - p^2 x(1-x)}{m_{ph}^2(1-x+x^2)} + O(\epsilon) \quad (2.112)$$

$$= \frac{g^2}{32\pi^2} \int_0^1 dx \ln \frac{m_{ph}^2 - p^2 x(1-x)}{m_{ph}^2(1-x+x^2)} + O(\epsilon) \quad (2.113)$$

This is the mass-shell scheme result: $\Sigma_{1R}^{(ph)}(m_{ph}^2) = 0$, the pole residue is normalized to one, and the mass parameter is the physical mass m_{ph} . which is as same as the result we have done before.

Notice that $\delta m^2 = \Sigma_{1R} - \Sigma_1(p^2) = -\Sigma_1(m_{ph}^2)$

$$= \frac{g^2}{2(4\pi)^{d/2}} \Gamma(2 - \frac{d}{2}) \int_0^1 dx [m_{ph}^2(1-x+x^2)]^{\frac{d}{2}-2} \quad (2.114)$$

when $d = 4$, $\int_0^1 dx [\dots]^0 = 1$, $\Gamma(2 - \frac{d}{2}) \simeq \frac{1}{2-d/2} - \gamma_E \implies \delta m^2 = \frac{g^2/32\pi^2}{2-d/2} \implies$

Minimal Subtraction

Thus $\Sigma_{1R} = \Sigma_1(p^2) + \delta m^2$ ($\Sigma_{1R}(p^2 = m^2) \neq 0$, Counterterm only cancels divergence)

$$\Sigma_{1R}^{(MS)} = -\frac{g^2}{2(4\pi)^{d/2}} \Gamma\left(2 - \frac{d}{2}\right) \int_0^1 dx [m^2 - p^2 x(1-x)]^{\frac{d}{2}-2} + \frac{g^2}{32\pi^2} \frac{1}{2-d/2} \quad (2.115)$$

$$= -\frac{g^2}{32\pi^2} \left(1 + \frac{\epsilon}{2} \ln 4\pi\right) \left(\frac{2}{\epsilon} - \gamma_E\right) \int_0^1 dx \left[1 - \frac{\epsilon}{2} \ln \Delta(x)\right] + \frac{g^2}{32\pi^2} \frac{2}{\epsilon} + O(\epsilon) \quad (2.116)$$

$$= \frac{g^2}{32\pi^2} \int_0^1 dx [\ln \Delta(x) + \gamma_E - \ln 4\pi] + O(\epsilon), \quad \Delta(x) = m^2 - p^2 x(1-x). \quad (2.117)$$

However, there are mass dimension in logarithm which is no-physics. The reason is we need to change the dimension of g .

$$\frac{g}{3!} \phi^3 \subset \mathcal{L}, \quad [\mathcal{L}] = [m]^d, \quad [\phi] = [m]^{\frac{d-2}{2}}$$

$$\implies [g] = [m]^{\frac{6-d}{2}}, \quad \text{when } d = 4, [g] = [m] \quad (2.118)$$

Thus, we can rewrite g as

$$g \rightarrow \mu^{\frac{4-d}{2}} g = \mu^{\epsilon/2} g \quad (2.119)$$

where the g shows on the right hand side has dimension $[m]^1$. This can also check on $\delta m^2 \propto g^2$, in which g is rescaled. So we have

$$\Sigma_{1R} = \Sigma_1(p^2) + \delta m^2 \quad (2.120)$$

$$= -\frac{g^2 \mu^\epsilon}{2(4\pi)^{d/2}} \Gamma\left(2 - \frac{d}{2}\right) \int_0^1 dx [\Delta(x)]^{\frac{d}{2}-2} + \frac{g^2}{32\pi^2} \frac{2}{\epsilon} \quad (2.121)$$

$$= -\frac{g^2}{32\pi^2} \left(\frac{2}{\epsilon} - \gamma_E + \ln 4\pi + 2 \ln \mu\right) \int_0^1 dx \left[1 - \frac{\epsilon}{2} \ln \Delta(x)\right] + \frac{g^2}{32\pi^2} \frac{2}{\epsilon} + O(\epsilon) \quad (2.122)$$

$$\implies \Sigma_{1R}^{(MS)} = \frac{g^2}{32\pi^2} \int_0^1 dx \left[\ln \left(\frac{m^2 - p^2 x(1-x)}{4\pi \mu^2} \right) + \gamma_E \right] \quad (2.123)$$

This renormalization prescription is called **Minimal Subtraction (MS)**, where counterterms are pure poles at the physical value of d .

$d = 6$ case

Now we apply it to $d = 6$. Since the superficial degree of divergence is

$$\omega(\Sigma_1) = d - 4 = 2, \quad (2.124)$$

we need to differentiate three times with respect to the external momentum in order to make the integral convergent. When we integrate back, the ambiguity is a polynomial of degree two in p . Lorentz invariance removes the linear term, so the counterterm has the form

$$\delta m^2 - p^2 \delta Z \quad (2.125)$$

The coupling is also continued away from six dimensions by

$$g_0 = \mu^{3-d/2}(g + \delta g), \quad \text{poles at } d = 6. \quad (2.126)$$

And for $\Gamma(z - 1)$, we have

$$\Gamma(z + 1) = z\Gamma(z) \quad (2.127)$$

$$\implies \Gamma(z - 1) = \frac{1}{z - 1}\Gamma(z) = \frac{1}{z - 1}\left(\frac{1}{z} - \gamma_E + O(z)\right) \quad (2.128)$$

$$= -(1 + z + \cdots + z^n)\left(\frac{1}{z} - \gamma_E + O(z)\right) \quad (2.129)$$

$$= \left(\gamma_E - \frac{1}{z} + O(z)\right) + (-1 + O(z)) + \cdots = -\frac{1}{z} + \gamma_E - 1 + O(z) \quad (2.130)$$

Thus, for $d = 6 - \epsilon$, we have

$$\Sigma_1 = -\frac{g^2}{2(4\pi)^{d/2}}\Gamma(2 - \frac{d}{2}) \int_0^1 dx [m^2 - p^2 x(1 - x)]^{\frac{d}{2}-2} \quad (2.131)$$

$$= -\frac{g^2}{128\pi^3}\mu^{6-d}(4\pi)^{\epsilon/2}\Gamma(\frac{\epsilon}{2} - 1) \int_0^1 dx (m^2 - p^2 x(1 - x))^{1-\epsilon/2} \quad (2.132)$$

$$= -\frac{g^2}{128\pi^3}\left(-\frac{2}{\epsilon} + \gamma_E - 1 - \ln(4\pi\mu^2)\right) \int_0^1 dx \Delta(x) \quad (2.133)$$

$$- \frac{g^2}{128\pi^3} \int_0^1 dx \Delta(x) \ln \Delta(x) + O(\epsilon), \quad \Delta(x) = m^2 - p^2 x(1 - x). \quad (2.134)$$

$$\Sigma_1 = -\frac{g^2}{128\pi^3}\left\{-\frac{2}{\epsilon}\left(m^2 - \frac{p^2}{6}\right) + (\gamma_E - 1)\left(m^2 - \frac{p^2}{6}\right) + \int_0^1 dx \Delta(x) \ln \frac{\Delta(x)}{4\pi\mu^2}\right\}. \quad (2.135)$$

Thus the counterterm will cancel the divergence. So we have

$$\Sigma_{1R} = \Sigma_1 + \delta m^2 - \delta Z p^2 \quad (2.136)$$

where


$$\delta m^2 = \frac{g^2}{64\pi^3} \frac{m^2}{d - 6} + O(g^4) \quad (2.137)$$

$$\delta Z = \frac{g^2}{6 \cdot 64\pi^3} \frac{1}{d - 6} + O(g^4) \quad (2.138)$$

Now we can consider δg . As a counterterm, it relates with the term in $\mathcal{L}$

$$\frac{g}{3!}\phi^3 \subset \mathcal{L} \quad (2.139)$$

The term δg cancels the UV divergence of the three-point vertex. So the ordinary tree vertex is corrected by the one-loop triangle, and this loop divergence is then canceled by a local three-leg counterterm:



The counterterm has no loop momentum. It is local, so in momentum space it is just a constant for this logarithmic divergence. To extract only this local UV pole, we may set the external momenta to zero after checking that the superficial divergence is logarithmic. Momentum-dependent terms are less divergent and only change the finite part. Thus, the loop is

$$\Sigma_1^{(3)} = (-ig\mu^{3-d/2})^3 \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 - m^2} \frac{i}{(k+p_1)^2 - m^2} \frac{i}{(k-p_2)^2 - m^2} \quad (2.140)$$

$$\xrightarrow{p \rightarrow 0} (-ig\mu^{3-d/2})^3 \int \frac{d^d k}{(2\pi)^d} \frac{-i}{(k^2 - m^2)^3} = (g\mu^{3-d/2})^3 \int \frac{d^d k_E}{(2\pi)^d} \frac{-i}{(k_E^2 + m^2)^3} \quad (2.141)$$

$$= (g\mu^{3-d/2})^3 \frac{-i}{(4\pi)^{d/2}} \frac{\Gamma(3-d/2)}{2} m^{d-6}, \quad \text{take } d = 6 - \epsilon \quad (2.142)$$

$$\simeq g^3 \mu^{3\epsilon/2} \frac{-i}{(4\pi)^3} \left(1 + \frac{\epsilon}{2} \ln 4\pi\right) \frac{1}{2} \left(\frac{2}{\epsilon} - \gamma_E\right) (1 - \epsilon \ln m) \quad (2.143)$$

$$\simeq \frac{-ig^3}{2(4\pi)^3} \mu^{3\epsilon/2} \left(\frac{2}{\epsilon} - \gamma_E + \ln 4\pi - \ln m^2\right) \quad (2.144)$$

$$\simeq \frac{-ig^3}{2(4\pi)^3} \left(1 + \frac{3}{2}\epsilon \ln \mu\right) \left(\frac{2}{\epsilon} + \text{finite}\right) \quad (2.145)$$

$$= \frac{-ig^3}{2(4\pi)^3} \left(\frac{2}{\epsilon} + 3 \ln \mu + \text{finite}\right) \quad (2.146)$$

To cancel the divergence, we introduce δg as

$$\frac{g}{3!}\phi^3 \rightarrow \frac{g}{3!}\phi^3 + \frac{\delta g}{3!}\phi^3 \quad (2.147)$$

where the last term should cancel the divergent $\frac{2}{\epsilon}$ and keep the order of unit mass to 1. So we have:

$$-i\delta g = \frac{ig^3}{2(4\pi)^3} \left(\frac{2}{\epsilon} + \ln \mu\right) = \frac{ig^3}{2(4\pi)^3} \frac{2}{\epsilon} \left(1 + \frac{\epsilon}{2} \ln \mu\right) \quad (2.148)$$

In this case, $\Sigma_1^{(3)} - i\delta g = \frac{ig^3}{2(4\pi)^3} (\ln \mu + \text{finite})$

$$\Rightarrow \text{the counterterm is } \delta g = \frac{-g^3}{64\pi^3} \frac{1}{\epsilon} \mu^{3-d/2} = \frac{g^3}{64\pi^3} \frac{1}{d-6} \mu^{3-d/2} \quad (2.149)$$

1.6 Massless theories

In Mass-shell scheme

$$\delta m_{ph}^2 = \frac{g^2}{2(4\pi)^{d/2}} \Gamma(2 - \frac{d}{2}) \int_0^1 dx [m_{ph}^2(1 - x + x^2)]^{\frac{d}{2}-2} \quad (2.150)$$

$$\xrightarrow{m_{ph} \rightarrow 0, d \rightarrow 4} \frac{g^2}{32\pi^2} (1 + \frac{\epsilon}{2} \ln 4\pi) \left(\frac{2}{\epsilon} - \gamma_E \right) (1 - \epsilon \ln m_{ph}) \quad (2.151)$$

$$= \frac{g^2}{32\pi^2} \left(\frac{2}{\epsilon} - \gamma_E + \ln 4\pi - \ln m_{ph}^2 \right) \quad (2.152)$$

$$= \frac{g^2}{32\pi^2} \left(\frac{1}{2 - d/2} - \ln(m_{ph}^2) + \text{finite} \right) \quad (2.153)$$

which fails when $m_{ph} \rightarrow 0$, i.e. IR div. We can see from MS subtraction from $d = 4 - \epsilon$ in above, i.e.

$$\Sigma_{1R}^{(MS)} = \frac{g^2}{32\pi^2} \left(\ln \frac{-p^2}{4\pi\mu^2} + \gamma_E - 2 \right) \quad (2.154)$$

So, here we know that the reason that the mass-shell scheme failed is because of the force is long range, so the propagator pole is not a simple pole, physically means that the particle will be never free from the other's influence.

Renormalization

The Lagrangian can be written as

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_b + \mathcal{L}_{ct} \quad (2.155)$$

where $\mathcal{L}_0$ is the free Lagrangian

$$\mathcal{L}_0 = \frac{1}{2}(\partial\phi)^2 - \frac{1}{2}m^2\phi^2 \quad (2.156)$$

$\mathcal{L}_b$ is the basic interaction

$$\mathcal{L}_b = -\frac{g}{3!}\phi^3 \quad (2.157)$$

$\mathcal{L}_{ct}$ is the counterterm Lagrangian

$$\mathcal{L}_{ct} = \delta Z \frac{1}{2}(\partial\phi)^2 - \delta m^2 \frac{1}{2}\phi^2 - \frac{\delta g}{3!}\phi^3 - \delta h\phi \quad (2.158)$$

where δh is to cancel the tadpole graphs. For order g^4 , we must draw not only the genuine two-loop graphs. We must also draw the graphs where a lower-order divergent subgraph has already been replaced by its counterterm. The relevant classes are:

1. Two one-loop bubbles on the same propagator line:

$$\text{---}\bigcirc\text{---}\bigcirc\text{---} + \text{---}\bigcirc\times\text{---} + \times\bigcirc\text{---} + \text{---}\times\times\text{---} + \dots$$

2. A two-loop self-energy graph with a one-loop subdivergence:

3. A one-loop graph with a vertex correction:

Each cross stands for a local counterterm. The point of the list is simple: every divergent subgraph needs its own subtraction. We have ignored the tadpoles, because they are local one-point divergences and can be canceled by $\delta h\phi$:

The form of **divergent $\times$ logarithm** is the **non-local subdivergence**. The form of **polynomial of p** is the **local divergence**.

Two loop self-energy in ϕ^3

Now we consider the graph (2) and (3) in $d = 6$.

graph (2)

Graph (a) is the genuine two-loop graph. Graph (b) subtracts the one-loop subdivergence inside (a). Graph (c) is the remaining local wave-function counterterm. We need all three graphs before the final answer is finite.

$$\begin{aligned} \Sigma_{2a} = & (ig\mu^{3-d/2})^4 \frac{1}{2} \int \frac{d^d k}{(2\pi)^d} \frac{d^d l}{(2\pi)^d} \frac{i}{k^2 - m^2 + i\epsilon} \frac{i}{l^2 - m^2 + i\epsilon} \\ & \times \frac{i}{(k-l)^2 - m^2 + i\epsilon} \frac{i}{(k+p)^2 - m^2 + i\epsilon}. \end{aligned} \quad (2.159)$$

For convenience, we consider the massless case, i.e.

$$\Sigma_{2a} = \frac{1}{2} g^4 \mu^{12-2d} \int \frac{d^d k}{(2\pi)^d} \frac{d^d l}{(2\pi)^d} \frac{1}{(k^2 + i\epsilon)(l^2 + i\epsilon)((k-l)^2 + i\epsilon)((k+p)^2 + i\epsilon)} \quad (2.160)$$

After Wick rotation, we have

$$\Sigma_{2a} = -\frac{1}{2}g^4\mu^{12-2d} \int \frac{d^d k_E}{(2\pi)^d} \frac{d^d l_E}{(2\pi)^d} \frac{1}{[l_E^2(k_E - l_E)^2][k_E^2(k_E + p_E)^2]} \quad (2.161)$$

$$= -\frac{1}{2}g^4\mu^{12-2d} \int \frac{d^d k_E}{(2\pi)^d} \frac{1}{k_E^2(k_E + p_E)^2} \left(\int \frac{d^d l_E}{(2\pi)^d} \frac{1}{l_E^2(k_E - l_E)^2} \right) \quad (2.162)$$

$$= -\frac{1}{2}g^4\mu^{12-2d} \int \frac{d^d k_E}{(2\pi)^d} \frac{1}{k_E^2(k_E + p_E)^2} i\pi^{d/2} \frac{\Gamma(2-d/2)}{(2\pi)^d} \int_0^1 dx (-k_E^2 x(1-x))^{d/2-2} \quad (2.163)$$

$$= -\frac{1}{2}g^4\mu^{12-2d} \int \frac{d^d k_E}{(2\pi)^d} \frac{1}{k_E^2(k_E + p_E)^2} i\pi^{d/2} \Gamma(2-d/2) \frac{\Gamma(\frac{d}{2}-1)^2}{\Gamma(d-2)} (-k_E^2)^{d/2-2} \quad (2.164)$$

$$= \frac{ig^4}{2^{13}\pi^9} (16\pi^3\mu^4)^{3-d/2} \frac{\Gamma(2-d/2)\Gamma(\frac{d}{2}-1)^2}{\Gamma(d-2)} \int d^d k_E \frac{1}{(-k_E^2)^{3-d/2}(k_E + p_E)^2} \quad (2.165)$$

For the integral, one can introduce a Feynman parameter

$$\frac{1}{A^{4-d/2}B} = \frac{\Gamma(5-d/2)}{\Gamma(4-d/2)} \int_0^1 \frac{x^{3-d/2}}{[Ax + B(1-x)]^{5-d/2}} \quad (2.166)$$

Thus, the result is

$$\Sigma_{2a} = \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{2} \left(\frac{-p^2}{4\pi\mu^2} \right)^{d-6} \frac{\Gamma(2-d/2)\Gamma(5-d)\Gamma(\frac{d}{2}-1)^3\Gamma(d-4)}{\Gamma(d-2)\Gamma(4-d/2)\Gamma(3d/2-5)} \quad (2.167)$$

$$\equiv \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{2} \left(\frac{-p^2}{4\pi\mu^2} \right)^{d-6} \Gamma(2-d/2)\Gamma(5-d)K(d) \quad (2.168)$$

where the **overall divergence** is contained in $\Gamma(5-d)$. And the **subdivergence** is contained in $\Gamma(2-d/2)$. One can also notice that there is IR divergence since we are talking about the massless fields when $p \rightarrow 0$. It can go away if we consider the massive theory. But that will be no explicit formula for Σ_{2a} .

Now, we expand Σ_{2a} in $\epsilon = 6-d$. To keep track of the non-local part, write $L_p = \ln(-p^2/(4\pi\mu^2))$. The factors singular at $d=6$ are the two gamma functions:

$$\Sigma_{2a} = \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{2} \left(\frac{-p^2}{4\pi\mu^2} \right)^{d-6} \Gamma(2-d/2)\Gamma(5-d)K(d) \quad (2.169)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{36} \left(\frac{1}{\epsilon} - \frac{\epsilon}{2}L_p - \frac{\epsilon^2}{4}L_p^2 \right) \left(\frac{2}{\epsilon} + \gamma_E - 1 \right) \left(-\frac{1}{\epsilon} + \gamma_E - 1 \right) K_d \quad (2.170)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{36} \left[\frac{1}{\epsilon^2} - \frac{1}{\epsilon} \left(L_p + \frac{3}{2}(\gamma_E - 1) \right) + \frac{1}{2}L_p^2 + \frac{3}{2}(\gamma_E - 1)L_p + \frac{1}{2}(\gamma_E - 1)^2 \right] \quad (2.171)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{36} \left[\frac{1}{(d-6)^2} + \frac{L_p + \text{Const.}}{d-6} + \frac{1}{2}L_p^2 + \text{Const.} L_p + \text{Const.} + O(d-6) \right]. \quad (2.172)$$

The term proportional to $L_p/(d-6)$ is important: it is a pole multiplying a non-polynomial function of the external momentum. A local counterterm cannot cancel such a term directly. Therefore it must be canceled by the lower-order counterterm graph (b), which subtracts the

divergent one-loop subgraph before the remaining overall divergence is removed. Notice for $m = 0$, the counterterm only contains $\delta_2 Z = \frac{g^2}{6 \cdot 64\pi^3} \frac{1}{d-6} + O(g^4)$.

$$\Sigma_{2b} = (-ig\mu^{3-d/2})^2 \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 - m^2} \frac{i}{k^2 - m^2} \frac{i}{(k+p)^2 - m^2} \delta_1 Z \cdot k^2 \quad (2.173)$$

$$= ig^2 \mu^{6-d} \delta_1 Z \int \frac{d^d k_E}{(2\pi)^d} \frac{k_E^2}{(k_E^2)^2 (k_E + p)^2} \quad (2.174)$$

$$= ig^2 \mu^{6-d} \delta_1 Z \frac{i\pi^{d/2}}{(2\pi)^d} \Gamma(2-d/2) \frac{\Gamma(\frac{d}{2}-1)^2}{\Gamma(d-2)} (p^2)^{d/2-2} \quad (2.175)$$

$$= \left(\frac{g^2}{64\pi^3} \right) \left(\frac{p^2}{6} \right) \frac{\Gamma(2-d/2) \Gamma^2(\frac{d}{2}-1)}{(d-6) \Gamma(d-2)} \left(\frac{-p^2}{4\pi\mu^2} \right)^{d/2-3} \quad (2.176)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \left(\frac{p^2}{36} \right) \left(-\frac{2}{\epsilon} + \gamma_E - 1 \right) \frac{1}{\epsilon} \left(1 - \frac{\epsilon}{2} L_p + \frac{\epsilon^2}{8} L_p^2 \right) \quad (2.177)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \left(\frac{p^2}{36} \right) \left[-\frac{2}{\epsilon^2} + \frac{\gamma_E - 1 + L_p}{\epsilon} - \frac{1}{2} (\gamma_E - 1) L_p - \frac{1}{4} L_p^2 \right] \quad (2.178)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{36} \left[-\frac{2}{(d-6)^2} + \frac{-L_p + \text{const.}}{d-6} - \frac{1}{4} L_p^2 + \text{const.} L_p + O(d-6) \right]. \quad (2.179)$$

Thus, we have

$$\begin{aligned} & \Sigma_{2a} + \Sigma_{2b} \\ &= \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{36} \left[\frac{-1}{(d-6)^2} + \frac{1}{d-6} \text{Const.} + \frac{1}{4} \ln^2 \frac{-p^2}{4\pi\mu^2} + \text{Const.} \ln \frac{-p^2}{4\pi\mu^2} + \text{const.} + O(d-6) \right] \end{aligned} \quad (2.180)$$

Now, we can see the non-local divergence has been canceled

$$\frac{1}{d-6} \left(\ln \frac{-p^2}{4\pi\mu^2} + \text{Const.} \right) \rightarrow \frac{1}{d-6} \text{const.} \quad (2.181)$$

The non-local subdivergence is, from its name, non-locally, which means the divergent is not because of the superposition of two space-time point on one-propagator, like the usual divergent. It has to be cancelled, otherwise we have to add extra terms onto the Lagrangian in this form

$$\phi(x) \ln(\square) \phi(x) \quad (2.182)$$

for the subdivergence $\frac{1}{\epsilon} \ln(p^2)$, which is unacceptable. Since it has been cancelled, the overall divergent part left can be easily cancelled by (c), the renormalization of wave function on the counterterm is then

$$\delta_2 Z = \left(\frac{g^2}{64\pi^3} \right)^2 \frac{1}{36} \left(-\frac{1}{d-6} + \frac{\text{Const.}}{d-6} \right) \quad (2.183)$$

Thus, the renormalized result is

$$\Sigma_{2R}^{(MS)} = \Sigma_{2a} + \Sigma_{2b} + \Sigma_{2c} = \Sigma_{2a} + \Sigma_{2b} + (-p^2 \delta_2 Z) \quad (2.184)$$

$$= \left(\frac{g^2}{64\pi^3} \right)^2 \frac{p^2}{36} \left(\frac{1}{4} \ln^2 \frac{-p^2}{4\pi\mu^2} + \text{Const.} \ln \frac{-p^2}{4\pi\mu^2} + \text{const.} \right) \quad (2.185)$$

1.7 Differentiation with respect to external momenta

Now, we want to show that the counterterm of 1-PI graph is a polynomial in its external momenta with degree equal to the degree of divergence.

First, the degree of divergence for Σ_{2a} is $d - 4$. To make it be negative in $d = 6$, we differentiate Σ_{2a} three times.

$$\Sigma_{2a}(p) \sim \int d^d k_E d^d l_E \frac{1}{[l_E^2(k_E - l_E)^2][k_E^2(k_E + p_E)^2]} \quad (2.186)$$

$$\Rightarrow \frac{\partial^3 \Sigma_{2a}}{\partial p^\mu \partial p^\nu \partial p^\rho} \sim \int d^d k \underbrace{\left(\int d^d l \frac{1}{l^2(k-l)^2} \right)}_{\text{subgraph}} \frac{\partial^3}{\partial p^3} \left(\frac{1}{k^2(k+p)^2} \right) \quad (2.187)$$


Notice that

$$\frac{\partial}{\partial p^\mu} \frac{1}{p^2 - m^2} = -\frac{1}{(p^2 - m^2)^2} \frac{\partial}{\partial p^\mu} (p^2 - m^2) = -\frac{2p_\mu}{(p^2 - m^2)^2}. \quad (2.188)$$


Thus differentiating a propagator produces one more propagator power and a numerator proportional to the momentum through that line. In graph language we denote this by adding a marked insertion on the differentiated line:

$$\frac{\partial}{\partial p^\mu} (\text{---}) \longrightarrow \text{---} \quad (2.189)$$

The three red dots below mean that the external-momentum dependent propagator has been differentiated three times. They are not new interaction vertices; they only remind us where the powers of momentum in the numerator come from. Hence $\frac{\partial^3 \Sigma_{2a}}{\partial p^\mu \partial p^\nu \partial p^\rho}$ should then be



and $\frac{\partial^3 \Sigma_{2b}}{\partial p^\mu \partial p^\nu \partial p^\rho}$ should be



Thus, $(a''') + (b''')$ is finite. So the overall counterterm is quadratic in P , i.e.

$$A(d)p^2 + B^\mu(d)p_\mu + C(d) \quad (2.190)$$

the Lorentz invariance requires $B^\mu = 0$

$$\Rightarrow A(d)p^2 + C(d) \quad (2.191)$$

Now, we prove it. Consider $l \gg k \rightarrow \infty$ ($l \sim k \rightarrow \infty$ is overall divergence cancelled by differentiation; k fixed, $l \rightarrow \infty$ cancelled by (b'''))

$$(a''') \sim \int^\infty dk \frac{1}{k^4} \int_{l \gg k} dl \cdot l \sim l^2 \rightarrow \text{quadratic diverges.} \quad (2.192)$$

The remaining question is the region $l \gg k$. In this region the loop momentum l is much larger than the momentum k entering the subgraph, so the inner loop can be expanded in powers of k/l . Let

$$I(k) = \int d^6 l \frac{1}{l^2(l+k)^2}. \quad (2.193)$$

Here we keep $d = 6$, because this is the case where the two-loop graph has the subdivergence we want to subtract. The useful expansion is

$$(l+k)^2 = l^2 + 2l \cdot k + k^2 = l^2 \left(1 + \frac{2l \cdot k + k^2}{l^2} \right), \quad (2.194)$$

$$\frac{1}{(l+k)^2} = \frac{1}{l^2} \left[1 - \frac{2l \cdot k + k^2}{l^2} + \left(\frac{2l \cdot k + k^2}{l^2} \right)^2 + \dots \right]. \quad (2.195)$$

This is the same as the Taylor expansion

$$I(k) = I(0) + k^\mu \frac{\partial I}{\partial k^\mu} \Big|_{k=0} + \frac{1}{2} k^\mu k^\nu \frac{\partial^2 I}{\partial k^\mu \partial k^\nu} \Big|_{k=0} + \dots. \quad (2.196)$$

Now look term by term. In six dimensions $d^6 l = \Omega_5 l^5 dl$, and angular symmetry gives $\langle (l \cdot k)^2 \rangle = l^2 k^2 / 6$. Therefore

$$k^0 : \quad \int d^6 l \frac{1}{l^4} \sim \Omega_5 \int^\Lambda dl l \sim \Lambda^2, \quad \text{quadratic divergence,} \quad (2.197)$$

$$k^1 : \quad - \int d^6 l \frac{2l \cdot k}{l^6} = -2k_\mu \int d^6 l \frac{l^\mu}{l^6} = 0, \quad \text{odd integrand,} \quad (2.198)$$

$$\begin{aligned} k^2 : \quad & \int d^6 l \left(-\frac{k^2}{l^6} + \frac{4(l \cdot k)^2}{l^8} \right) \sim \Omega_5 \int^\Lambda dl l^5 \left(-\frac{k^2}{l^6} + \frac{4}{l^8} \frac{l^2 k^2}{6} \right) \\ & = -\frac{\Omega_5 k^2}{3} \int^\Lambda \frac{dl}{l} \sim -\frac{\Omega_5 k^2}{3} \ln \Lambda, \quad \text{logarithmic divergence.} \end{aligned} \quad (2.199)$$

So the counterterm in graph (b) removes exactly these local divergent pieces. The k^0 term is a constant, and the k^2 term is quadratic in the external momentum. After this subtraction, only the dependence on the matching scale k remains:

$$k^0 : \quad \int_1^\Lambda dl l - \text{subtraction} = O(k^2), \quad (2.200)$$

$$k^2 : \quad \int_k^\Lambda \frac{dl}{l} - \ln \Lambda = -\ln k = O(\ln k). \quad (2.201)$$

Terms of order k^3 and higher are already finite in the inner loop. They do not add new ultraviolet divergences. Thus the worst remaining behavior in the outer integral is

$$\int dk \frac{1}{k^4} (k^2 \ln k) = \int dk \frac{\ln k}{k^2} \quad (2.202)$$

which is finite.

1.8 Composite operators


Goal: Renormalize products of operators, e.g., $\phi(x)\phi(y)$. When the separation $x-y$ becomes small, the product can be represented by local operators at one point. Equivalently, after Fourier transform, this is an expansion in the limit where the momentum flowing through the pair is much larger than the other external momenta:

$$\phi(x)\phi(y) \sim C_1(x-y)\not{1} + C_{\phi^2}(x-y)[\phi^2(y)] + C_\phi(x-y)[\phi(y)] + \dots \quad (2.203)$$

where $[A(x)]$ denotes the renormalized operator, while $A(x)$ without brackets is the unrenormalized composite operator. The functions C_i are c-numbers and are called Wilson coefficients.

operator product expansion

Consider the graphs for four-point function in ϕ^3 -theory. In the labels of the figures, we write $P = p_1 + p_2$.



where $q^\mu \rightarrow \infty$, and p_1, p_2 fixed. The words “hard” and “soft” only tell us which part carries the large momentum. In (a) and (b), all three propagators on the top line carry momenta of order q . In (c), only the two upper propagators are hard; the lower line with momentum $P = p_1 + p_2$ is soft and will become a local operator. The expansion used below is simply

$$\frac{1}{(q+p)^2 - m^2} = \frac{1}{q^2} \left[1 - \frac{2q \cdot p + p^2 - m^2}{q^2} + O\left(\frac{1}{q^2}\right) \right]. \quad (2.204)$$

For the soft denominators it is useful to write

$$P = p_1 + p_2, \quad D_1 = p_1^2 - m^2, \quad D_2 = p_2^2 - m^2, \quad D_P = P^2 - m^2.$$

Only the propagators carrying q are expanded; D_1, D_2, D_P are kept fixed.

$$(a) = (-ig)^2 \frac{i}{q^2 - m^2} \frac{i}{D_1} \frac{i}{(q+p_1)^2 - m^2} \frac{i}{(q+P)^2 - m^2} \frac{i}{D_2} \quad (2.205)$$

$$\xrightarrow{q \rightarrow \infty} ig^2 \left(\frac{i}{D_1} \frac{i}{D_2} \right) \frac{1}{(q^2)^3} + O\left(\frac{1}{q^7}\right), \quad (2.206)$$

$$(b) \xrightarrow{q \rightarrow \infty} ig^2 \left(\frac{i}{D_1} \frac{i}{D_2} \right) \frac{1}{(q^2)^3} + O\left(\frac{1}{q^7}\right), \quad (2.207)$$

$$\Rightarrow (a) + (b) \simeq 2ig^2 \left(\frac{i}{D_1} \frac{i}{D_2} \right) \frac{1}{(q^2)^3}, \quad (2.208)$$

$$(c) = (-ig)^2 \frac{i}{q^2 - m^2} \frac{i}{(q+P)^2 - m^2} \frac{i}{D_P} \frac{i}{D_1} \frac{i}{D_2}. \quad (2.209)$$

$$(c) = \frac{-ig^2}{(q^2)^2} \left[1 - \frac{2q \cdot P}{q^2} + \frac{2m^2 - P^2}{q^2} + \frac{4(q \cdot P)^2}{(q^2)^2} + \dots \right] \times \frac{1}{(P^2 - m^2)(p_1^2 - m^2)(p_2^2 - m^2)}, \quad P = p_1 + p_2. \quad (2.210)$$

Now, we will show that it can be expanded like an operator-product. For (a) + (b),

$$\begin{aligned}
 (a) + (b) &\simeq \underbrace{\frac{2ig^2}{(q^2)^3}}_{\text{hard coefficient}} \underbrace{\left(\frac{i}{p_1^2 - m^2} \frac{i}{p_2^2 - m^2} \right)}_{\text{soft matrix element}} \\
 &= C_{\phi^2}(q) \langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \frac{1}{2} \phi^2(0) \} | 0 \rangle. \tag{2.211}
 \end{aligned}$$

$\Rightarrow$ For this four-point correlation function, after the operator $\frac{1}{2}\phi^2(0)$ is fixed at the origin, only two spacetime points are left free: x and y (or, in momentum space, p_1 and p_2). Hence

$$\langle 0 | T \{ \phi(x) \phi(y) (\frac{1}{2} \phi(0) \phi(0)) \} | 0 \rangle \tag{2.212}$$

$$= \langle 0 | T \{ \phi(x) \phi(y) (\frac{1}{2} \phi(0) \phi(0)) \} | 0 \rangle_c + \langle 0 | T \{ \phi(x) \phi(0) \} | 0 \rangle \langle 0 | T \{ \phi(y) \phi(0) \} | 0 \rangle \tag{2.213}$$

$$= \frac{1}{2} D(x) D(y) + \frac{1}{2} D(y) D(x) = D(x) D(y) \tag{2.214}$$

In momentum space:

$$\int d^4x d^4y e^{-ip_1 \cdot x} e^{-ip_2 \cdot y} D(x) D(y) = \frac{i}{p_1^2 - m^2} \frac{i}{p_2^2 - m^2} \tag{2.215}$$


$$\Rightarrow \int d^4x d^4y e^{-ip_1 \cdot x} e^{-ip_2 \cdot y} \langle 0 | T \{ \phi(x) \phi(y) \frac{1}{2} \phi^2(0) \} | 0 \rangle \tag{2.216}$$

$$= \langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \frac{1}{2} \phi^2(0) \} | 0 \rangle \tag{2.217}$$

Here $\phi(0)$ is not Fourier transformed, because it is the local operator inserted at the origin. To see why the factor $\frac{1}{2}\phi^2(0)$ gives this rule, use the functional integral definition:

$$\langle 0 | T \{ \phi(x) \phi(y) \frac{1}{2} \phi^2(0) \} | 0 \rangle = N \int [dA] A(x) A(y) \frac{1}{2} A^2(0) e^{iS} \tag{2.218}$$

Here ϕ denotes the quantum operator, while A is the classical field inside the path integral. At lowest order, the insertion of $\frac{1}{2}\phi^2(0)$ therefore gives one local black-dot vertex with two scalar lines:



This insertion $\frac{1}{2}\phi^2(0)$ is our first example of a composite operator. Therefore

$$(a) + (b) \sim \frac{2ig^2}{(q^2)^3} \langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \frac{1}{2} \phi^2(0) \} | 0 \rangle \tag{2.219}$$

$$= \frac{2ig^2}{(q^2)^3} \left(\text{V-vertex diagram} \right) \tag{2.220}$$

For (c), firstly, notice

$$\begin{aligned}
 \langle 0 | T \{ \phi(x) \phi(y) \phi(z) \} | 0 \rangle_{\text{interaction vacuum}} &= \langle 0 | T \{ \phi(x) \phi(y) \phi(z) \int d^4w \frac{-ig}{3!} \phi^3(w) \} | 0 \rangle_{\text{free vacuum}} \\
 &\tag{2.221}
 \end{aligned}$$

$$= -ig \int d^4w D(x-w) D(y-w) D(z-w) \tag{2.222}$$

Use

$$D(u - v) = \int \frac{d^4 k}{(2\pi)^4} e^{ik \cdot (u-v)} \frac{i}{k^2 - m^2}.$$

In momentum space, with $P = p_1 + p_2$,

$$\langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \phi(x) \} | 0 \rangle = \int d^4 y d^4 z e^{-ip_1 \cdot y - ip_2 \cdot z} \langle 0 | T \{ \phi(x) \phi(y) \phi(z) \} | 0 \rangle \quad (2.223)$$

$$= -ig \left(\frac{i}{p_1^2 - m^2} \frac{i}{p_2^2 - m^2} \right) \int d^4 w e^{-iP \cdot w} D(x - w) \quad (2.224)$$

$$= -ig \left(\frac{i}{p_1^2 - m^2} \frac{i}{p_2^2 - m^2} \right) \int \frac{d^4 k}{(2\pi)^4} \frac{i e^{ik \cdot x}}{k^2 - m^2} \int d^4 w e^{-i(P+k) \cdot w} \quad (2.225)$$

$$= -ig \left(\frac{i}{p_1^2 - m^2} \frac{i}{p_2^2 - m^2} \right) \frac{i}{P^2 - m^2} e^{-iP \cdot x}. \quad (2.226)$$

Thus, we can take $x = 0$ such that the $e^{-i(p_1+p_2) \cdot x} = 1$. And also, if we only consider the propagator, the $-ig$ belongs to the vertex, so it's not considered here. When this matrix element is written as a local operator at $x = 0$, the momentum P is represented by derivatives acting on the local field:

$$P_\mu \rightarrow i\partial_\mu, \quad P^2 \rightarrow -\square.$$

$$(c) \sim \frac{ig^2}{(q^2)^2} \left[1 - \frac{2q \cdot P}{q^2} + \frac{2m^2 - P^2}{q^2} + \frac{4(q \cdot P)^2}{(q^2)^2} + \dots \right] \left[\frac{-i}{(P^2 - m^2)(p_1^2 - m^2)(p_2^2 - m^2)} \right] \quad (2.227)$$

$$= \frac{ig^2}{(q^2)^2} \left[1 - \frac{2iq^\mu \partial_\mu}{q^2} + \frac{2m^2 + \square}{q^2} - \frac{4q^\mu q^\nu \partial_\mu \partial_\nu}{(q^2)^2} + \dots \right] \langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \phi(x) \} | 0 \rangle \Big|_{x=0} \quad (2.228)$$

$$= \frac{ig^2}{(q^2)^2} \sum \text{coefficients} \times \text{derivatives} \left(\text{diagram} \right) \quad (2.229)$$

Thus, we have

$$\begin{aligned} & \langle 0 | T \{ \tilde{\phi}(q) \phi(0) \tilde{\phi}(p_1) \tilde{\phi}(p_2) \} | 0 \rangle \\ & \sim \sum_i C_i(q) \langle 0 | T \{ O_i(0) \tilde{\phi}(p_1) \tilde{\phi}(p_2) \} | 0 \rangle. \end{aligned} \quad (2.230)$$

The sum is over a set of local operators O_i . Each of these is either the elementary field ϕ , one of its derivatives, or a composite operator, such as ϕ^2 . e.g.

$$i = 1 \quad O_1 = \frac{1}{2} \phi^2 \quad ((a), (b)) \quad (2.231)$$

$$i = 2 \quad O_2 = \phi \quad ((c) \text{ leading order}) \quad (2.232)$$

$$i = 3 \quad O_3 = \partial_\mu \phi \text{ or } \square \phi \quad ((c) \text{ higher order}) \quad (2.233)$$

The functions $C_i(q)$ are the **Wilson coefficients**. They contain the short-distance, high-momentum part of the graph. The matrix elements of O_i contain the long-distance part.


1.9 Renormalization of composite operators

Renormalization of ϕ^2

Consider ϕ^3 theory in 6-dim.

$$\langle 0 | T \{ \phi(x) \phi(y) \frac{1}{2} \phi^2(z) \} | 0 \rangle \quad (2.234)$$

The connected graphs up to order g^2 are shown below. In these pictures, the black dot is the insertion of the composite operator $\frac{1}{2}\phi^2$. A crossed mark, when it appears later, means an ordinary counterterm from the Lagrangian.



To work on momentum space, we have

$$\begin{aligned} G &= \langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \frac{1}{2} \phi^2(z) \} | 0 \rangle \\ &= \int d^d x d^d y e^{i(p_1 \cdot x + p_2 \cdot y)} \langle 0 | T \{ \phi(x) \phi(y) \frac{1}{2} \phi^2(z) \} | 0 \rangle. \end{aligned} \quad (2.235)$$

The lowest order graph, i.e., (a) is

$$\begin{aligned} S_i &= \frac{i}{p_i^2 - m^2 + i\epsilon}, \quad S_{12} = S_1 S_2. \\ G_a &= S_{12}. \end{aligned} \quad (2.236)$$

For the order g^2 graphs, the large- k behavior is

- (b) $\sim \int d^6 k \frac{1}{k^2} \frac{1}{k^2} \frac{1}{k^2} \sim \int dk \frac{1}{k} \implies$ logarithmically divergent.
- (c), (d), (e) $\sim \int d^6 k \frac{1}{k^4} \sim \int dk k \implies$ quadratically divergent.

Graphs (d) and (e) are ordinary self-energy corrections on the external lines. Their divergences are cancelled by the usual wave-function and mass counterterms. They do not give a new counterterm for $[\phi^2]$.

Graphs (b) and (c) are different. Their short-distance divergence sits at the composite-operator insertion itself, so the bare operator ϕ^2 is not enough. We must add local counterterms to define the finite operator $[\phi^2]$.

$$G_b = S_{12} (-ig)^2 \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 - m^2 + i\epsilon} \frac{i}{(p_1 - k)^2 - m^2 + i\epsilon} \frac{i}{(p_2 + k)^2 - m^2 + i\epsilon}. \quad (2.237)$$

After the continuation $g \rightarrow g\mu^{(6-d)/2}$, the loop part is

$$\begin{aligned} G_b &= S_{12} \left(\frac{ig^2 \mu^{6-d}}{(2\pi)^d} \right) \\ &\times \int d^d k \frac{1}{(k^2 - m^2)[(p_1 - k)^2 - m^2][(p_2 + k)^2 - m^2]}. \end{aligned} \quad (2.238)$$

$$\begin{aligned}
& \frac{1}{(k^2 - m^2)[(p_1 - k)^2 - m^2][(p_2 + k)^2 - m^2]} \\
&= 2 \int_0^1 dx \int_0^{1-x} dy \left[(1-x-y)(k^2 - m^2) + x((p_1 - k)^2 - m^2) + y((p_2 + k)^2 - m^2) \right]^{-3}
\end{aligned} \tag{2.239}$$

$$= 2 \int_0^1 dx \int_0^{1-x} dy \left[k^2 - m^2 - 2(xp_1 - yp_2) \cdot k + xp_1^2 + yp_2^2 \right]^{-3}. \tag{2.240}$$

Completing the square with $k' = k - xp_1 + yp_2$ and $P = p_1 + p_2$ gives

$$\begin{aligned}
& 2 \int_0^1 dx \int_0^{1-x} dy \left[(k - xp_1 + yp_2)^2 - (xp_1 - yp_2)^2 + xp_1^2 + yp_2^2 - m^2 \right]^{-3} \\
&= 2 \int_0^1 dx \int_0^{1-x} dy \left[k'^2 - \Delta_b(x, y) \right]^{-3},
\end{aligned} \tag{2.241}$$

$$\Delta_b(x, y) = m^2 - p_1^2 x(1-x-y) - p_2^2 y(1-x-y) - P^2 xy. \tag{2.242}$$

After Wick rotation, the momentum integral is

$$\begin{aligned}
& \int_0^1 dx \int_0^{1-x} dy \int d^d k_E \frac{2}{(k_E^2 + \Delta_b)^3} \\
&= \int_0^1 dx \int_0^{1-x} dy \frac{(2\pi)^d}{(4\pi)^{d/2}} \Gamma\left(3 - \frac{d}{2}\right) \Delta_b^{d/2-3}.
\end{aligned} \tag{2.243}$$

Therefore

$$\begin{aligned}
G_b &= S_{12} \frac{g^2}{64\pi^3} \Gamma\left(3 - \frac{d}{2}\right) \\
&\quad \times \int_0^1 dx \int_0^{1-x} dy \left(\frac{\Delta_b(x, y)}{4\pi\mu^2} \right)^{d/2-3}.
\end{aligned} \tag{2.244}$$

And for (c), use the same P and set

$$D_{12P} = (p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2).$$

Then

$$G_c = (-ig\mu^{\frac{6-d}{2}})^2 \frac{-i}{D_{12P}} \frac{1}{2!} \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 - m^2} \frac{i}{(k+P)^2 - m^2} \tag{2.245}$$

$$\begin{aligned}
&= \frac{-ig^2\mu^{6-d}}{D_{12P}} \int_0^1 dx \int \frac{d^d k}{(2\pi)^d} \left[(k+xP)^2 - \Delta_c(x) \right]^{-2}, \\
&\quad \Delta_c(x) = m^2 - P^2 x(1-x)
\end{aligned} \tag{2.246}$$

$$= \frac{\frac{1}{2}g^2\mu^{6-d}}{D_{12P}} \int_0^1 dx \int \frac{d^d k_E}{(2\pi)^d} \frac{1}{(k_E^2 + \Delta_c(x))^2} \tag{2.247}$$

$$= \frac{\frac{1}{2}g^2(\mu^2)^{3-d/2}}{D_{12P}} \frac{1}{(4\pi)^{d/2}} \Gamma\left(2 - \frac{d}{2}\right) \int_0^1 dx \Delta_c(x)^{d/2-2} \tag{2.248}$$

$$= \frac{g^2}{D_{12P}} \frac{1}{128\pi^3} \Gamma\left(2 - \frac{d}{2}\right) \int_0^1 dx \frac{\Delta_c(x)^{d/2-2}}{(4\pi\mu^2)^{d/2-3}} \tag{2.249}$$

$$\begin{aligned}
\Rightarrow G_c &= \frac{-g\mu^{3-d/2}}{D_{12P}} \frac{-g\mu^{d/2-3}}{128\pi^3} \Gamma\left(2 - \frac{d}{2}\right) \\
&\quad \times \int_0^1 dx \frac{\Delta_c(x)^{d/2-2}}{(4\pi\mu^2)^{d/2-3}}
\end{aligned} \tag{2.250}$$

G_b and G_c both diverge, so the bare composite operator ϕ^2 does not define finite Green functions by itself. For the operator-product expansion we need a local operator with the same quantum numbers and a finite insertion. In the $\overline{\text{MS}}$ scheme, the required counterterms are obtained by replacing each divergent short-distance loop, i.e.,



The counterterm is: **for (b):**

$$G_b = S_{12} \frac{g^2}{64\pi^3} \Gamma\left(3 - \frac{d}{2}\right) \int_0^1 dx \int_0^{1-x} dy \left(\frac{\Delta_b(x, y)}{4\pi\mu^2} \right)^{d/2-3} \quad (2.251)$$

$$= S_{12} \frac{g^2}{64\pi^3} \int_0^1 dx \int_0^{1-x} dy \left[\frac{2}{\epsilon} - \gamma_E - \ln \frac{\Delta_b(x, y)}{4\pi\mu^2} \right] + O(\epsilon) \quad (2.252)$$

$$= \frac{1}{(p_1^2 - m^2)(p_2^2 - m^2)} \frac{g^2}{64\pi^3(d-6)} + \text{finite}. \quad (2.253)$$

$$\Rightarrow \text{Counterterm} = S_{12} \frac{g^2}{64\pi^3(d-6)}. \quad (2.254)$$

for (c):

$$G_c = \frac{-g\mu^{3-d/2}}{(p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2)} \frac{-g\mu^{d/2-3}}{128\pi^3} \Gamma\left(2 - \frac{d}{2}\right) \int_0^1 dx \frac{\Delta_c(x)^{d/2-2}}{(4\pi\mu^2)^{d/2-3}}, \quad (2.255)$$

$$\Delta_c(x) = m^2 - P^2 x(1-x), \quad P = p_1 + p_2,$$

$$G_c = \frac{-g\mu^{3-d/2}}{(p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2)} \frac{-g\mu^{d/2-3}}{128\pi^3} \left(-\frac{2}{\epsilon}\right) \int_0^1 dx \Delta_c(x) + \text{finite} \quad (2.256)$$

$$= \frac{-g\mu^{3-d/2}}{(p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2)} \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} \left(m^2 - \frac{P^2}{6}\right) + \text{finite}. \quad (2.257)$$

$$\Rightarrow \text{Counterterm} = \frac{-g\mu^{3-d/2}}{(p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2)} \left\{ \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} \left(m^2 - \frac{P^2}{6}\right) \right\}. \quad (2.258)$$

Thus, the renormalized value at $d = 6$ is

$$R(G_b) = \frac{i^2}{(p_1^2 - m^2)(p_2^2 - m^2)} \frac{g^2}{64\pi^3} \left[-\frac{\gamma_E}{2} - \int_0^1 dx \int_0^{1-x} dy \ln \frac{\Delta_b(x, y)}{4\pi\mu^2} \right], \quad (2.259)$$

$$R(G_c) = \frac{-g}{(p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2)} \frac{-g}{128\pi^3} \left[(\gamma_E - 1) \left(m^2 - \frac{P^2}{6}\right) + \int_0^1 dx \Delta_c(x) \ln \frac{\Delta_c(x)}{4\pi\mu^2} \right]. \quad (2.260)$$

Now, in last section, we found that

$$\begin{aligned} (a) + (b) &= 2ig^2 \left(\frac{i}{p_1^2 - m^2} \frac{i}{p_2^2 - m^2} \right) \frac{1}{(q^2)^3} \\ &= \frac{2ig^2}{(q^2)^3} \langle 0 | T \{ \tilde{\phi}(p_1) \tilde{\phi}(p_2) \frac{1}{2} \phi^2(0) \} | 0 \rangle = \frac{2ig^2}{(q^2)^3} \left(\text{diagram} \right). \end{aligned} \quad (2.261)$$

comparing with the counterterm, we find

$$\begin{aligned} & -\frac{1}{(p_1^2 - m^2)(p_2^2 - m^2)} \frac{g^2}{64\pi^3(d-6)} \\ & = \frac{g^2}{64\pi^3(d-6)} \langle 0 | T\{\tilde{\phi}(p_1)\tilde{\phi}(p_2)\frac{1}{2}\phi^2(0)\} | 0 \rangle. \end{aligned} \quad (2.262)$$

And also

$$(c) = \frac{ig^2}{(q^2)^2} \left[1 - \frac{2q \cdot P}{q^2} + \frac{2m^2 - P^2}{q^2} + \frac{4(q \cdot P)^2}{(q^2)^2} + \dots \right] \frac{-i}{D_{12P}} \quad (2.263)$$

$$= \frac{ig^2}{(q^2)^2} \left[1 - \frac{2iq^\mu \partial_\mu}{q^2} + \frac{2m^2 + \square}{q^2} - \frac{4q^\mu q^\nu \partial_\mu \partial_\nu}{(q^2)^2} + \dots \right] \langle 0 | T\{\tilde{\phi}(p_1)\tilde{\phi}(p_2)\phi(x)\} | 0 \rangle \Big|_{x=0} \quad (2.264)$$

$$= \frac{ig^2}{(q^2)^2} \sum \text{coefficients} \times \text{derivatives} \left(\text{diagram} \right). \quad (2.265)$$

comparing with the counterterm, we find

$$\begin{aligned} & \frac{-g\mu^{3-d/2}}{(p_1^2 - m^2)(p_2^2 - m^2)(P^2 - m^2)} \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} \left(m^2 - \frac{P^2}{6} \right) \\ & = \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} \left(m^2 + \frac{1}{6}\square \right) \langle 0 | T\{\tilde{\phi}(p_1)\tilde{\phi}(p_2)\phi(0)\} | 0 \rangle. \end{aligned} \quad (2.266)$$

Thus, we have ($R(G_d)$ and $R(G_e)$ have no contribution due to the divergents are cancelled by the usual counterterms)

$$\begin{aligned} & G_a + R(G_b) + R(G_c) + R(G_d) + R(G_e) \\ & = \left(1 + \frac{g^2}{64\pi^3(d-6)} \right) \langle 0 | T\{\tilde{\phi}(p_1)\tilde{\phi}(p_2)\frac{1}{2}\phi^2(0)\} | 0 \rangle \\ & \quad + \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} \langle 0 | T\{\tilde{\phi}(p_1)\tilde{\phi}(p_2)\left(m^2 + \frac{1}{6}\square\right)\phi(0)\} | 0 \rangle + O(g^4). \end{aligned} \quad (2.267)$$

So what we are computing is a Green's function of the operator

$$\frac{1}{2}[\phi^2] \equiv \left(1 + \frac{g^2}{64\pi^3(d-6)} \right) \frac{1}{2}\phi^2 + \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} \left(m^2 + \frac{1}{6}\square \right) \phi + O(g^4) \quad (2.268)$$

$$= \left(1 + \frac{g^2}{64\pi^3(d-6)} \right) \frac{1}{2}\phi^2 + \frac{g\mu^{d/2-3}}{64\pi^3(d-6)} m^2 \phi + \frac{g\mu^{d/2-3}}{6 \cdot 64\pi^3(d-6)} \square \phi. \quad (2.269)$$

Here we use $[\phi^2]$ to denote a **renormalized operator**. We will see later that this result can be generalized to all orders, which can be written as

$$\frac{1}{2}[\phi^2] = Z_a \frac{1}{2}\phi^2 + \mu^{d/2-3} Z_b m^2 \phi + \mu^{d/2-3} Z_c \square \phi \quad (2.270)$$

where Z 's only depend on g and d . The reason that only these operators appear is the same power-counting logic used before: counterterms must be local, and an operator can mix only with operators whose dimension is not larger than its own. For ϕ^2 in six-dimensional ϕ^3 theory, the allowed lower-or-equal operators are ϕ^2 , $m^2\phi$, and $\square\phi$. We can also write the mixing as

$$\begin{pmatrix} \frac{1}{2}[\phi^2] \\ \phi \\ \square\phi \end{pmatrix} = \begin{pmatrix} Z_a & \mu^{d/2-3} Z_b m^2 & \mu^{d/2-3} Z_c \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{2}\phi^2 \\ \phi \\ \square\phi \end{pmatrix} \quad (2.271)$$

2. Renormalization in ϕ^4 -theory

Firstly, we briefly recall some basic idea about renormalization. For counting the divergent of a theory, one may use the superficial degree of divergence:

$$D = \text{power of } k \text{ in numerator} - \text{power of } k \text{ in denominator.} \quad (2.272)$$

Suppose the Loop integrals in $d = 4$ space-time dimension have superficial degree of divergence D , which can be easily seen from the directly calculating:

$$\int^{\Lambda} d^4k \frac{k^{a-4}}{k^b} = \int^{\Lambda} k^3 dk d\Omega \frac{k^{a-b}}{k^4} \sim \int^{\Lambda} dk \frac{k^{a-b}}{k} = \begin{cases} \Lambda^D & \text{for } D = a - b \neq 0 \\ \ln \Lambda & \text{for } D = 0 \end{cases}, \quad (2.273)$$

where Λ is the cut-off. But we would like to use some more general form to think about it. As an example, considering the ϕ^n -theory

$$\mathcal{L} = \frac{1}{2}(\partial_{\mu}\phi)(\partial^{\mu}\phi) - \frac{1}{2}m^2\phi^2 - \frac{\lambda}{n!}\phi^n, \quad (2.274)$$

in which the propagators have the form of $(k^2 - m^2)^{-1}$, vertices are $\sim \lambda$ that connect n lines. So we can define So in this case, we can

$$1. \quad L = P - V + 1$$

- L : loops (loop integral with d^4k)
- P : internal propagators (propagator with $(k^2)^{-1}$)
- V : vertices (vertices with energy-momentum conservation $\delta(\sum p_i)$)
- $+1$: global conservation of energy-momentum

$$2. \quad nV = 2P + N$$

- n : number of lines at each vertex
- P : internal propagators
- N : external lines (or external propagator)

So for each loop, the power of k in numerator is 4, which is contributed by an integral measurement $\sim d^4k$, and the power of k in denominator is contributed by the propagator. Therefore, the superficial degree of divergence is

$$D = dL - 2P; \quad d = 4. \quad (2.275)$$

Thus, base on the relation we introduced above, we have

$$\begin{aligned} D &= 4(P - V + 1) - 2P \\ &= 2P - 4V + 4 \\ &= (n - 4)V - N + 4. \end{aligned} \quad (2.276)$$

Thus, we find D is independent of P , but only depends on N for the case $n = 4$. But for a more general n in $\frac{\lambda}{n!}\phi^n$, we have

- $n = 2$: $\implies$ free theory (included in the mass term);

- $n = 3$: $D = 4 - N - V \implies$ **super renormalizable**
- $n = 4$: $D = -N + 4 \implies$ **renormalizable**
- $n > 4$: $D = (n - 4)V - N + 4 \implies$ **non-renormalizable**

For a theory that the signature in front of V is “ $-$ ”, we call it as **super renormalizable theory** because the divergence only appear in some leading order terms, with respect the order getting higher, the diagram will finally be convergent, in which $D < 0$. *So in the super renormalizable theory only contains finite number of divergent degree*, which means only need finite counterterms. Similarly, for a theory that calls **renormalizable**, it is because it has no relation with V , which means *no matter how many orders we are counting, the divergent degrees are the same!* Although in this case, it contains infinite divergent diagrams, we can still by re-define the “physical” parameter to include the divergence. In the end, it is quite obvious to understand why we call the last situation as **non-renormalizable**. Because the pre-signature of V is positive, which means with the order getting higher, the divergent will be keep increasing, *so we need infinite parameter to cancel the divergence*, which in most of cases are technically impossible.

Another case is QED, we have two type of propagators, but they all contribute $\sim k^{-2}$

- N_γ : external photon lines;
- N_e : external electron lines;
- P_e : electron propagators;
- P_γ : photon propagators;
- $V = 2P_\gamma + N_\gamma = \frac{1}{2}(2P_e + N_e)$: number of vertices;
- $L = P_e + P_\gamma - V + 1$: number of the loops.

$$\begin{aligned}
 D &= 4L - P_e - 2P_\gamma \\
 &= 4(P_e + P_\gamma - V + 1) - P_e - 2P_\gamma \\
 &= 3P_e + 2P_\gamma - 4V + 4 \\
 &= 4 - N_\gamma - \frac{3}{2}N_e.
 \end{aligned} \tag{2.277}$$

So as we discussed above, since D is non-related with V , so QED is a renormalizable theory.

2.1 Analysis of dimensions in view of renormalization

Now we consider another way to analyze the renormalizability. We use the dimension of mass as a standard. So

$$[\text{Length}] = [\text{Time}] = [\text{Energy}]^{-1} = [\text{Mass}]^{-1}. \tag{2.278}$$

We require the dimension for the action $[S] = 0$, thus

$$[d^4x] = -4, \quad [\mathcal{L}] = 4, \quad [\partial_\mu] = 1, \quad [\phi] = 1. \tag{2.279}$$

The last one is because $[\partial_\mu \phi \partial^\mu \phi] = 4$. The coupling constant $[\lambda]$ in $\frac{\lambda}{n!} \phi^n$ then has the dimension

$$[\lambda] = 4 - n. \tag{2.280}$$

Take a sharp look! This number, $4 - n$, is nothing but the prefactor of superficial degree of divergence in ϕ^n theory, i.e.,

$$D = 4 - N - \underbrace{(4 - n)}_{[\lambda]} V. \quad (2.281)$$

Thus, base on the discussion we did above, we find $[\lambda] < 0$ corresponds to a non-renormalizable theory. As an example, we consider the ϕ^4 -theory. So $n = 4$ brings the superficial degree of divergence is $D = 4 - N \geq 0$ for degree up to $N = 4$, which is a renormalizable theory.

$D = 4(\text{vacuum}) \quad D = 2 \text{ (self energy)} \quad D = 0$

We can also do the same thing for QED. For convenience, we rewrite the QED Lagrangian:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}(i\not{\partial} - m)\psi + eA_\mu\bar{\psi}\gamma^\mu\psi. \quad (2.282)$$

Similarly, since the action S is dimensionless, so

$$[\psi] = \frac{3}{2}, \quad [A_\mu] = 1. \quad (2.283)$$

Thus again, we can then see the coupling constant e is dimensionless, i.e., $[e] = 0$, which just corresponds to a renormalizable theory.

2.2 Renormalized perturbation theory

Now we still use ϕ^4 -theory as an example to show the renormalization procedure. The bare Lagrangian is

$$\mathcal{L}_{\text{bare}} = \frac{1}{2}(\partial_\mu\phi_0)(\partial^\mu\phi_0) - \frac{1}{2}m_0^2\phi_0^2 + \frac{1}{4!}\lambda_0\phi_0^4 \quad (2.284)$$

where the corner mark “0” represent the bare parameters, i.e., λ_0, m_0 , and ϕ_0 . This Lagrangian is nothing special, it just the original Lagrangian that we used in perturbation theory. Now we just give it a different name to separate with the renormalized Lagrangian that we will discuss later. The name “bare” represent this quantity is not actually the physical quantity that we observe because the bare parameters are actually infinity. These parameters appears only for cancel the infinity in the loop integral. The rest of this canceling is then the physical quantity, which is the renormalized quantity. This explanation shows the core idea of why do we need renormalization, i.e., the physical part of graphs needs to be observable, so we have to find a well-defined quantity that suitable for any order. But in this case, we may have to accept that many “constants” we thought unchanged will start to running with the higher energy scale being considered. And this will be described by the renormalization group equation.

The relation of the renormalized quantities with the bare quantities are

$$\phi_0 = \sqrt{Z}\phi_r, \quad Zm_0^2 = m^2 + \delta m^2, \quad Z^2\lambda_0 = \lambda + \delta\lambda, \quad (2.285)$$

where $Z = 1 + \delta Z$. And $\delta Z, \delta m^2$ and $\delta\lambda$ are called the **counterterms**. So the bare Lagrangian can be written as

$$\begin{aligned} \mathcal{L}_{\text{bare}} &= \mathcal{L}_{\text{ren}} + \mathcal{L}_{\text{count}} \\ &= \underbrace{\frac{1}{2}(\partial_\mu\phi_r)(\partial^\mu\phi_r) - \frac{1}{2}m^2\phi_r^2 + \frac{\lambda}{4!}\phi_r^4}_{\mathcal{L}_{\text{ren}}} + \underbrace{\frac{1}{2}\delta Z(\partial_\mu\phi_r)(\partial^\mu\phi_r) - \frac{1}{2}\delta m^2\phi_r^2 + \frac{1}{4!}\delta\lambda\phi_r^4}_{\mathcal{L}_{\text{counter term}}}. \end{aligned} \quad (2.286)$$

The counterterms also have corresponding Feynman rules

$$\longrightarrow \otimes \longrightarrow = i(p^2 \delta Z - \delta m^2); \quad \bigotimes = -i\delta\lambda. \quad (2.287)$$

And the Feynman rules of renormalized term is

$$\longrightarrow_p = \frac{i}{p^2 - m^2}; \quad \bigotimes = -mi\lambda. \quad (2.288)$$

The renormalization condition is usually called the **scheme**. The on-shell scheme for ϕ_r, m in two-point function is

$$\longrightarrow_p \bigotimes_p = \frac{i}{p^2 - m_0^2 - \Sigma(p^2)} \quad (2.289)$$

$$= \frac{i}{p^2 - m_{\text{phy}}^2} + \underbrace{A + B(p^2 - m_{\text{phy}}^2) + C(p^2 - m_{\text{phy}}^2)^2 + \dots}_{\text{Regular Terms}}, \quad (2.290)$$

where the $\Sigma(p^2)$ is the self-energy correction. The main point of this scheme is requiring the residue at the pole $p^2 = m_{\text{phys}}^2$ to be normalized to 1. The mission of the counterterms is to absorb UV divergences and ensure the self-energy $\Sigma(p^2)$ satisfies the specific renormalization conditions, rather than cancelling the regular terms.

Now for the four-point functions, we impose a similar renormalization condition on the scattering amplitude:

$$\left(\bigotimes \right)_{\text{Amp}} = i\lambda, \quad (2.291)$$

at the threshold $s = 4m^2$ and $t = u = 0$, which corresponds to the case of zero momentum transfer. The Mandelstam variables are defined as usual: $s = (p_1 + p_2)^2$, $t = (p_1 - p_3)^2$, and $u = (p_1 - p_4)^2$. This condition, together with the two-point function conditions mentioned earlier, defines the on-shell renormalization scheme. To implement this in practice, we follow these steps:

1. Calculate the divergent diagrams: Evaluate all Feynman diagrams contributing to the n -point functions at a given order in perturbation theory. A suitable regularization scheme (e.g., dimensional regularization) must be chosen to handle the UV divergences.
2. Determine the counterterms: Extract the specific values of the counterterms δm^2 , δZ , and $\delta\lambda$ by requiring that the renormalized Green's functions satisfy the chosen renormalization conditions (such as the one in (2.291)).

To conclude this section, we will briefly do the renormalization process for the 1-loop for ϕ^4 -theory. Firstly, we expand the two point function until λ^2 :

$$\bigotimes = \text{---} + \left(\text{---} \bigcirc \text{---} + \text{---} \otimes \text{---} \right) + O(\lambda^2), \quad (2.292)$$

3. Wilson's approach to renormalization

The main idea of Wilson's renormalization is introducing a cut-off Λ and compute how the theory changes at different length scales. For that, we need to working with the generating functional with cut-off and successive integration over large momenta/small length scales:

$$Z = \int \mathcal{D}\phi_\Lambda e^{-S[\phi]}, \quad (2.303)$$

where we used Wick rotation changing to Euclidean space-time:

$$x^0 \rightarrow ix^4, \quad d^4x \rightarrow id^4x_E. \quad (2.304)$$

Thus, the measurement is

$$\mathcal{D}\phi_\Lambda = \prod_{|k| < \Lambda} d\phi(k). \quad (2.305)$$

In this case, the modes of ϕ are well defined in Euclidean space-time such that we can make sure that when $k \rightarrow \infty$ all the components of momentum also go to infinity. This step is to avoid in Minkowski space-time $k^2 = (k^0)^2 - (k^i)^2$. So when $k^2 \rightarrow \infty$, it could be $k^0 \rightarrow \infty$ and $k^i \rightarrow 0$, which doesn't suitable for our analysis.

Now, the action and Lagrangian become


$$S[\phi] = \int d^4x \mathcal{L}, \quad (2.306)$$

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \frac{1}{2} m^2 \phi^2 + \frac{\lambda}{4!} \phi^4. \quad (2.307)$$

Suppose the mass $m \ll \Lambda$, and the coupling constant $\lambda \ll 1$, which means the last two terms in the Lagrangian can be treated as perturbation. For going to the perturbation theory, it's useful to split up fields $\phi \rightarrow \phi + \hat{\phi}$ as

$$\hat{\phi} = \begin{cases} \phi(k) & \text{for } b\Lambda \leq |k| \leq \Lambda \\ 0 & \text{for } |k| \leq b\Lambda \end{cases}; \quad \phi = \begin{cases} 0 & \text{for } b\Lambda \leq |k| \leq \Lambda \\ \phi(k) & \text{for } |k| \leq b\Lambda \end{cases}, \quad (2.308)$$

where $(b\Lambda, \Lambda)$ is a very small interval controlled by parameter $b \leq 1$



Then we can also split up the action as $S[\phi] \rightarrow S[\phi] + \hat{S}[\phi, \hat{\phi}]$, so we have

$$S[\phi] = \int d^4x \left[\frac{1}{2} \partial_\mu (\phi + \hat{\phi}) \partial^\mu (\phi + \hat{\phi}) + \frac{1}{2} m^2 (\phi + \hat{\phi})^2 + \frac{\lambda}{4!} (\phi + \hat{\phi})^4 \right] \quad (2.309)$$

$$= \int d^4x \left[\frac{1}{2} \left(\partial_\mu \phi \partial^\mu \phi + 2 \partial_\mu \phi \partial^\mu \hat{\phi} + \partial_\mu \hat{\phi} \partial^\mu \hat{\phi} \right) + \frac{1}{2} m^2 (\phi^2 + \hat{\phi}^2 + 2\phi\hat{\phi}) \right] \quad (2.310)$$

$$+ \frac{\lambda}{4!} \left(\phi^4 + \phi^2 \hat{\phi}^2 + 2\phi^3 \hat{\phi} + \hat{\phi}^2 \phi^2 + \hat{\phi}^4 + 2\phi \hat{\phi}^3 + 2\phi^3 \hat{\phi} + 2\phi \hat{\phi}^3 + 4\phi^2 \hat{\phi}^2 \right) \quad (2.311)$$

$$= \int d^4x \left[\underbrace{\frac{1}{2} (\partial_\mu \phi)^2 + \frac{1}{2} m^2 \phi^2 + \frac{\lambda}{4!} \phi^4}_{\mathcal{L}[\phi]} + \underbrace{\frac{1}{2} (\partial_\mu \hat{\phi})^2 + \frac{1}{2} m^2 \hat{\phi}^2 + \frac{\lambda}{4!} (6\phi^2 \hat{\phi}^2 + 4\phi^3 \hat{\phi} + 4\phi \hat{\phi}^3 + \hat{\phi}^4)}_{\hat{\mathcal{L}}[\phi, \hat{\phi}]} \right], \quad (2.312)$$

where we have dropped terms containing the products $\phi\hat{\phi}$ because Fourier components are orthogonal:

$$\int \frac{d^4k}{(2\pi)^4} \frac{d^4k'}{(2\pi)^4} \phi(k) \hat{\phi}(k') e^{i(k+k') \cdot x} \sim \delta^{(4)}(k+k') = 0 \quad (2.313)$$

Thus, we have

$$\hat{S}[\phi, \hat{\phi}] = \int d^4x (\hat{\mathcal{L}}_0[\hat{\phi}] + \hat{\mathcal{L}}_{\text{int}}[\phi, \hat{\phi}]), \quad (2.314)$$

and

$$\hat{\mathcal{L}}_0[\hat{\phi}] = \frac{1}{2} \partial_\mu \hat{\phi} \partial^\mu \hat{\phi}, \quad (2.315)$$

$$\hat{\mathcal{L}}_{\text{int}}[\phi, \hat{\phi}] = \frac{1}{2} m^2 \hat{\phi}^2 + \lambda \left(\frac{1}{3!} \phi \hat{\phi}^3 + \frac{1}{4} \phi^2 \hat{\phi}^2 + \frac{1}{3!} \phi^3 \hat{\phi} + \frac{1}{4!} \phi^4 \right). \quad (2.316)$$

Thus, the first part in $\hat{S}$ can be rewritten by the Fourier transformation as

$$\int d^4x \hat{\mathcal{L}}_0[\hat{\phi}] = \int \frac{d^4k d^4p}{(2\pi)^8} (ik_\mu \hat{\phi}(k)) (ip^\mu \hat{\phi}(p)) \int d^4x e^{i(k+p) \cdot x} \quad (2.317)$$

$$= \int \frac{d^4k d^4p}{(2\pi)^8} (ik_\mu) (ip^\mu) \hat{\phi}(k) \hat{\phi}(p) (2\pi)^4 \delta^{(4)}(k+p) \quad (2.318)$$

$$= \frac{1}{2} \int_{b\Lambda < |k| < \Lambda} \frac{d^4k}{(2\pi)^4} \hat{\phi}(-k) k^2 \hat{\phi}(k) \quad (2.319)$$

$$= \frac{1}{2} \int_{b\Lambda < |k| < \Lambda} \frac{d^4k}{(2\pi)^4} \hat{\phi}^*(k) k^2 \hat{\phi}(k). \quad (2.320)$$

This Gaussian integral form remind us to write a propagator for $\hat{\phi}$ from Wick contraction of $\hat{\phi}$ fields:

$$\overline{\hat{\phi}(k) \hat{\phi}(p)} = \langle \hat{\phi}(k) \hat{\phi}(p) \rangle = \frac{\int \mathcal{D}\hat{\phi} \hat{\phi}(k) \hat{\phi}(p) e^{-S_0[\hat{\phi}]}}{\int \mathcal{D}\hat{\phi} e^{-S_0[\hat{\phi}]}} = \frac{1}{k^2} (2\pi)^4 \delta^{(4)}(k+p) \Theta(k), \quad (2.321)$$

where

$$\Theta(k) = \begin{cases} 1 & \text{for } b\Lambda < |k| < \Lambda \\ 0 & \text{else} \end{cases}. \quad (2.322)$$

Therefore, we can separate the generating functional measurement with the **high energy modes** $\hat{\phi}$ and **low energy modes** $\phi_{b\Lambda}$ as $\mathcal{D}\phi_\Lambda = \mathcal{D}\phi_{b\Lambda} \mathcal{D}\hat{\phi}_\Lambda$. And we can integrate the high energy modes

$$Z = \int \mathcal{D}\phi_\Lambda e^{-S[\phi]} = \int \mathcal{D}\phi_{b\Lambda} \mathcal{D}\hat{\phi}_\Lambda e^{-S[\phi] - \hat{S}[\phi, \hat{\phi}]} = \int \mathcal{D}\phi_{b\Lambda} e^{-S[\phi]} \left(\int \mathcal{D}\hat{\phi}_\Lambda e^{-\hat{S}[\phi, \hat{\phi}]} \right) \quad (2.323)$$

$$= \int \mathcal{D}\phi_{b\Lambda} e^{-S_{\text{eff}}[\phi]}, \quad (2.324)$$

where left a **effective action** defined as

$$S_{\text{eff}}[\phi] = S[\phi] + \delta S[\phi], \quad (2.325)$$

in which $\delta S[\phi]$ can be treated as perturbation, thus it can be expanded as

$$e^{-\delta S[\phi]} = \int \mathcal{D}\hat{\phi}_\Lambda e^{-\hat{S}[\phi, \hat{\phi}]} = \int \mathcal{D}\hat{\phi}_\Lambda e^{-\int d^4x \hat{\mathcal{L}}[\phi, \hat{\phi}]} \quad (2.326)$$

$$\simeq \int \mathcal{D}\hat{\phi}_\Lambda \int d^4x \left[1 - \left(\hat{\mathcal{L}}_0 + \frac{1}{2}m^2\hat{\phi}^2 + \frac{\lambda}{3!}\phi\hat{\phi}^3 + \frac{\lambda}{4}\phi^2\hat{\phi}^2 + \frac{\lambda}{3!}\phi^3\hat{\phi} + \frac{\lambda}{4!}\hat{\phi}^4 \right) + \mathcal{O}(\lambda^2) \right] \quad (2.327)$$

$$= \int \mathcal{D}\hat{\phi}_\Lambda \int d^4x \left[1 - \underbrace{\left(\hat{\mathcal{L}}_0 + \frac{1}{2}m^2\hat{\phi}^2 + \frac{\lambda}{4!}\hat{\phi}^4 \right)}_{\substack{\text{Doesn't contain } \phi, \\ \text{which has no contribution for running}}} - \frac{\lambda}{4}\phi^2\hat{\phi}^2 + \mathcal{O}(\lambda^2) \right], \quad (2.328)$$

where in the third step we have dropped all the odd terms which don't contribute to the integral. The introducing of $\delta S[\phi]$ will change the parameters in the original action $S[\phi]$, which will finally leading the parameters start running. To show this, we now need to work in the perturbation scenario. Since we have assumed that $m \ll \Lambda$ and $\lambda \ll 1$, so we can expand $e^{-S_{\text{eff}}[\phi]}$ with Dyson series in terms of λ . For that, we can generally write the effective Lagrangian $\mathcal{L}_{\text{eff}}$ for effective action $S_{\text{eff}}[\phi]$ as:

$$\mathcal{L}_{\text{eff}}[\phi] = \frac{1}{2}(1 + \Delta Z)\partial_\mu\phi\partial^\mu\phi + \frac{1}{2}(m^2 + \Delta m^2)\phi^2 + \frac{1}{4!}(\lambda + \Delta\lambda)\phi^4 + (\text{higher dimensional operators}). \quad (2.329)$$

Now the mission is to determine ΔZ , Δm^2 and $\Delta\lambda$. We firstly consider the 1-loop order $\mathcal{O}(\lambda)$ correction for the propagator, which corresponds to the term $\frac{1}{4}\lambda\phi^2\hat{\phi}^2$, the graph is :

$$I_1 = \phi \text{ --- } \text{ (loop) } \text{ --- } \phi, \quad \mathcal{O}(\lambda) \quad (2.330)$$

where the double line represent the contraction of $\hat{\phi}$, which doesn't contribute the real physical observable, thus it needs to be integrated. Thus, we have

$$I_1 = -\frac{\lambda}{4} \int d^4x \phi^2 \hat{\phi} \hat{\phi} \quad (2.331)$$

$$= -\frac{\lambda}{4} \int d^4x \int \frac{d^4k_1}{(2\pi)^4} \cdots \frac{d^4k_4}{(2\pi)^4} e^{i(k_1+\cdots+k_4)\cdot x} \phi(k_1)\phi(k_2) \frac{1}{k_3^2} (2\pi)^4 \delta^{(4)}(k_3+k_4) \Theta(k_3), \quad (2.332)$$

integrate for x, k_4 :

$$I_1 = -\frac{\lambda}{4} \int \frac{d^4k_1}{(2\pi)^4} \cdots \frac{d^4k_3}{(2\pi)^4} \phi(k_1)\phi(k_2) \frac{1}{k_3^2} (2\pi)^4 \delta^{(4)}(k_1+k_2) \Theta(k_3) \quad (2.333)$$

$$= -\frac{1}{2} \mu \int \frac{d^4k_1}{(2\pi)^4} \phi(k_1)\phi(-k_1), \quad (2.334)$$

where

$$\mu = \frac{\lambda}{2} \int_{b\Lambda < |k| < \Lambda} \frac{d^4k}{(2\pi)^4} \frac{\Theta(k)}{k^2} \quad (2.335)$$

$$= \frac{\lambda}{2} \frac{1}{(2\pi)^4} \int d\Omega \int_{b\Lambda}^{\Lambda} k dk \quad (2.336)$$

$$= \frac{\lambda}{32\pi^2} (1 - b^2) \Lambda^2. \quad (2.337)$$

$$\exp\left\{-\int d^4x \frac{1}{2}\mu\phi^2\right\}. \quad (2.338)$$
$$\left(\begin{array}{c} \text{Diagram 1: A horizontal line with } \phi \text{ at both ends and a teardrop loop above it.} \\ \text{Diagram 2: Two intersecting lines forming an X, with a double-lined oval in the center.} \end{array} \right)^2,$$
$$I_2 = -\frac{1}{4!}\xi \int d^4x \phi^4(x), \quad (2.339)$$
$$\xi = -4! \cdot \underbrace{2}_{\substack{\text{2 possible} \\ \text{contractions} \\ \text{of } \phi^2(\phi\phi)^2}} \cdot \underbrace{\frac{1}{2!}}_{\substack{\text{symmetry factor} \\ \text{of diagram}}} \cdot \left(\frac{\lambda}{4}\right)^2 \int_{b\Lambda < |k| < \Lambda} \frac{d^4 k}{(2\pi)^4} \left(\frac{1}{k^2}\right)^2 \Theta(k) = -\frac{3\lambda^2}{16\pi^2} \textcolor{red}{\ln \frac{1}{b}} \quad (2.340)$$
$$k' = \frac{k}{b} \quad , \quad x' = bx \quad (2.341)$$
$$S_{\text{eff}} = \int d^4x' \frac{1}{b^4} \left(\frac{1}{2} (1 + \Delta Z) b^2 \partial'_\mu \phi \partial'^\mu \phi + \frac{1}{2} (m^2 + \Delta m^2) \phi^2 + \frac{1}{4!} (\lambda + \Delta \lambda) \phi^4 + \underbrace{\dots + \frac{1}{n!} (\lambda_n + \Delta \lambda_n) \phi^n + \dots}_{\text{higher dim. operators}} \right) \quad (2.342)$$

$$\lambda'_n = b^{n-4} \Delta \lambda_n \quad (2.350)$$

Generally, in D dimensions, we have:

$$m'^2 = b^{-2} \Delta m^2 \quad (2.351)$$

$$\lambda' = b^{D-4} \Delta \lambda \quad (2.352)$$

$$\lambda'_n = b^{D-4+n-4} \Delta \lambda_n \quad (2.353)$$

for $b < 1$ on $D = 4$:

- Super-renormalizable m^2 grows as $b^{-2} \longleftrightarrow$ relevant operator ϕ^2 .
- Renormalizable λ scales as $b^0 \longleftrightarrow$ marginal operators $\lambda \phi^4$.
- Non-renormalizable λ_n vanishes as $b^{n-4} \longleftrightarrow$ irrelevant operators $\lambda_n \phi^n, n > 4$.

$$\lambda' = \lambda - \frac{3}{16\pi^2} \lambda^2 \ln \left(\frac{1}{b} \right) + O(\lambda^3) \quad (2.354)$$

4. Callan-Symanzik Equation

Goal: Quantify scale dependence of Green's functions $G^{(n)}(x_1 \dots x_n, M)$, where M sets the scale of external momenta, which leads to the Callan-Symanzik equation.

4.1 Alternative choice for renormalization in ϕ^4 -theory

In ϕ^4 -theory, the 1-PI graphs of propagator can be renormalized by taking the renormalization scheme. We simply recall this process again, firstly, taking

$$\text{---} \bigcirc \text{1PI} \text{---} = -i\Sigma(p^2), \quad (2.355)$$

for all 1-PI contributions to propagators. Thus, the full propagator goes like:

$$\begin{aligned} \text{---} \text{---} \bigcirc \text{---} \text{---} &= \text{---} \text{---} + \text{---} \text{---} \bigcirc \text{1PI} \text{---} \text{---} + \text{---} \text{---} \bigcirc \text{1PI} \text{---} \bigcirc \text{1PI} \text{---} \text{---} + \dots \\ &= \frac{i}{p^2 - m^2 - \Sigma(p^2)}. \end{aligned} \quad (2.356)$$

Here we took the on-shell scheme:

$$\Sigma(p^2) \big|_{p^2=m_{ph}^2} = 0, \quad \frac{d}{dp^2} \Sigma(p^2) \big|_{p^2=m_{ph}^2} = 0. \quad (2.357)$$

This scheme force the residue around the pole to be unity. Then we obtained the renormalized propagator:

$$\text{---} \text{---} \bigcirc \text{---} \text{---} = \frac{i}{p^2 - m_{ph}^2}. \quad (2.358)$$

But this is not physical in many cases, say QCD and SM due to many of problems. For example, since the quarks are confined so normally we can hardly define the asymptotic state to imagine a free quark. And the massless field of SM in this scheme will meet IR-divergence, which will also lead us to many unnecessary problems. So we need to choose other better schemes for

a more general model. One alternative is to define counterterms at arbitrary scale M , at the external space-like momenta $p^2 = -M^2 < 0$, we take

$$\begin{aligned} \left(\begin{array}{c} \xrightarrow{p} \text{1PI} \xrightarrow{p} \\ p^2 = -M^2 \end{array} \right) &= 0, \quad \frac{d}{dp^2} \left(\begin{array}{c} \xrightarrow{p} \text{1PI} \xrightarrow{p} \\ p^2 = -M^2 \end{array} \right) = 0, \\ \left(\begin{array}{c} p_1 \searrow \quad \nearrow p_4 \\ \quad \bullet \\ p_2 \nearrow \quad \searrow p_3 \\ s=t=u=-M^2 \end{array} \right) &= i\lambda. \end{aligned} \quad (2.359)$$

where $(p_1 + p_2)^2 = s$, $(p_1 + p_3)^2 = t$, $(p_1 + p_4)^2 = u$. The reason why we choose the space-like momentum is to avoid poles. If we choose a time-like momentum, it for sure corresponds to physical particle, however also having divergence. The first condition $\Sigma(-M^2) = 0$ requires the correction for mass Σ_{finite} is 0 at this non-fixing scale M , and all the divergence Σ_{div} will be subtracted by the counterterms. So basically it looks very similar with the on-shell condition, the only thing that we change is changing the fixed physical mass m^2 to the unphysical mass scale $-M^2$.

This arbitrary parameter M is called the **renormalization scaling**. These conditions defined the only certain vale at a fixed point, which canceled all the divergent. So we say **it defines the theory in scale M** .

For seeing this scheme is indeed valid, we firstly look at the Green's function with renormalized fields $\phi = \sqrt{Z}\phi_r$ and coupling λ , and taking the renormalized scale as M , we have

$$\begin{aligned} G_{\text{ren}}^{(n)}(x_1 \dots x_n, \lambda, M) &= \langle \Omega | T \{ \phi_r(x_1) \dots \phi_r(x_n) \} | \Omega \rangle \\ &= Z^{-n/2} \langle \Omega | T \{ \phi(x_1) \dots \phi(x_n) \} | \Omega \rangle \\ &= Z^{-n/2} \underbrace{G_{\text{bare}}^{(n)}(x_1 \dots x_n, \lambda, M)}_{\text{bare Green's fun. independent of } M}. \end{aligned} \quad (2.360)$$

Now we change variables as $M \rightarrow M + \delta M$. This changing of our scaling will also change the other parameters to satisfy the renormalization condition. Thus it leads to $\lambda \rightarrow \lambda + \delta\lambda$ and $\phi_{\text{ren}} \rightarrow \phi'_{\text{ren}} = (1 + \delta\eta)\phi_{\text{ren}}$, thus

$$(Z')^{-1/2}\phi_0 = Z^{-1/2}(1 + \delta\eta)\phi_0 \implies Z' = Z(1 + \delta\eta)^{-2} \simeq Z(1 - 2\delta\eta) \equiv Z(1 + \delta Z), \quad (2.361)$$

where $\delta\eta = \frac{-\delta Z}{2}$. The latter gives the way of Green's function changing:

$$G_{\text{ren}}^{(n)} \rightarrow Z^{-n/2}(1 + \delta Z)^{-n/2} G_{\text{bare}}^{(n)} \quad (2.362)$$

$$\simeq Z^{-n/2} \left(1 - \frac{n}{2} \delta Z \right) G_{\text{bare}}^{(n)} \quad (2.363)$$

$$= Z^{-n/2} (1 + n\delta\eta) G_{\text{bare}}^{(n)} \quad (2.364)$$

$$= (1 + n\delta\eta) G_{\text{ren}}^{(n)}, \quad (2.365)$$

We will ignore the corner index “ren” in $G_{\text{ren}}^{(n)}$, **namely the renormalized Green's function denotes as $G^{(n)}$** . Thus from $\delta G_{\text{bare}} = 0$, we have

$$0 = \delta(Z^{n/2} G^{(n)}) = \underbrace{(\delta Z^{n/2}) G^{(n)}}_{n\delta\eta Z^{n/2}} + Z^{n/2} \frac{\partial}{\partial M} G^{(n)} \delta M + Z^{n/2} \frac{\partial}{\partial \lambda} G^{(n)} \delta \lambda. \quad (2.366)$$

Introducing the new variables:

$$\beta = \frac{M}{\delta M} \delta \lambda, \quad \gamma = \frac{M}{\delta M} \delta \eta \quad (2.367)$$

Thus, dropping the arbitrary δM , we obtain the **Callan-Symanzik equation**:

$$\left(M \frac{\partial}{\partial M} + \beta \frac{\partial}{\partial \lambda} + n \gamma \right) G^{(n)}(x_1 \dots x_n, \lambda, M) = 0, \quad (2.368)$$

where β and γ are universal for a given theory, which is independent for particular Green's function. This also means that these parameters are independent of any UV-cut off (dimensionless).

$$\beta = \beta(\lambda), \quad \gamma = \gamma(\lambda). \quad (2.369)$$

We can also directly compute $dG^{(n)}$ as

$$dG^{(n)} = (1 + n\delta\eta)G^{(n)} - G^{(n)} = n\delta\eta G^{(n)} = n \frac{dM}{M} \left(M \frac{d\eta}{dM} \right) G^{(n)} \quad (2.370)$$

$$= \frac{\partial G^{(n)}}{\partial M} dM + \frac{\partial G^{(n)}}{\partial \lambda} d\lambda \quad (2.371)$$

$$= \frac{\partial G^{(n)}}{\partial M} dM + \left(M \frac{d\lambda}{dM} \right) \frac{dM}{M} \frac{\partial G^{(n)}}{\partial \lambda} \quad (2.372)$$

$$= \frac{dM}{M} \left[M \frac{\partial}{\partial M} + \left(M \frac{d\lambda}{dM} \right) \frac{\partial}{\partial \lambda} \right] G^{(n)}, \quad (2.373)$$

from this construction, we can also obtain the Callan-Symanzik equation.

4.2 β -function in massless ϕ^4 -theory

Now let's focus some actual cases to find the *beta*-function and the anomalous dimension $\gamma(\lambda)$. For simplicity, we consider the 1-loop correction for **massless ϕ^4 -theory**. The Callan-Symanzik equation for $G^{(2)}$ gives us $\gamma = 0 + O(\lambda^2)$, because in 1-loop order, there's no contribution to Z , but only to δm^2 . Only until 2-loops or higher orders which contribute Z and also of course δm^2 .

$$\text{---} \bigcirc \text{1PI} \text{---} = \text{---} + \text{---} \bigcirc \text{---} + \frac{1}{2} \left(\text{---} \bigcirc \text{---} \right)^2 + \text{---} \bigcirc \text{---} + \text{---} \bigcirc \text{---} + O(\lambda^3). \quad (2.374)$$

For $\beta(\lambda)$ at $O(\lambda)$, we need to consider the 4-point function correction:

$$G^{(4)}(p) = \text{---} \times \text{---} + \left(\text{---} \bigcirc \text{---} + \text{---} \bigcirc \text{---} + \text{---} \bigcirc \text{---} \right) + \text{---} \bigcirc \text{---} + O(\lambda^3) \quad (2.375)$$

$$= -i\lambda + (-i\lambda)^2(iV(t) + iV(s) + iV(u)) - i\delta\lambda + O(\lambda^3), \quad (2.376)$$

where the renormalization condition are taken in $t = s = u = -M^2$. So for each diagram vertex correction, for $m = 0$ we have:

$$V(p^2) = \frac{1}{2} \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2} \frac{i}{(p-k)^2}. \quad (2.377)$$

Counterterm $\delta\lambda$ determined from amputated Green's function

$$G_{\text{amp}}^{(4)}|_{s=t=u=-M^2} \rightarrow \delta\lambda = (-i\lambda)^2 3V(-M^2). \quad (2.378)$$

Take $D = 4 - 2\epsilon$:

$$V(-M^2) = \lim_{\epsilon \rightarrow 0} \frac{1}{2} \frac{-1}{(4\pi)^{d/2}} \int_0^1 dx \frac{\Gamma(\epsilon)}{(x(1-x)M^2)^\epsilon} \quad (2.379)$$

$$= -\frac{1}{32\pi^2} \left(\frac{1}{\epsilon} - \ln M^2 + O(1) \right). \quad (2.380)$$

Thus

$$\delta\lambda = \frac{3\lambda^2}{32\pi^2} \left(\frac{1}{\epsilon} - \ln M^2 + O(1) \right). \quad (2.381)$$

This will lead to

$$G^{(4)} = -i\lambda + 3i(i\lambda)^2 V(p^2) - i\delta\lambda + \mathcal{O}(\lambda^3) \quad (2.382)$$

$$= -i\lambda + \frac{3i\lambda^2}{32\pi^2} \left(\frac{1}{\epsilon} - \ln p^2 - \frac{1}{\epsilon} + \ln M^2 \right) + \mathcal{O}(\lambda^3) \quad (2.383)$$

$$= -i\lambda + \frac{3i\lambda^2}{32\pi^2} \ln \left(\frac{M^2}{p^2} \right) + \mathcal{O}(\lambda^3) \quad (2.384)$$

Thus, we have

$$M \frac{\partial}{\partial M} G^{(4)} = \frac{3i\lambda^2}{16\pi^2} + \mathcal{O}(\lambda^3). \quad (2.385)$$

Thus, bring it into the Callan-Symanzik equation, we have

$$-\beta(\lambda) \frac{\partial}{\partial \lambda} G^{(4)} = i \frac{3\lambda^2}{16\pi^2} + \mathcal{O}(\lambda^3) \quad (2.386)$$

where $\gamma = 0 + \mathcal{O}(\lambda^2)$. Since we only consider order $\mathcal{O}(\lambda^2)$, therefore we only need to expand $G^{(4)}$ into $\mathcal{O}(\lambda)$, i.e.,

$$\frac{\partial}{\partial \lambda} G^{(4)} = -i + \frac{3i\lambda}{16\pi^2} \ln \left(\frac{M^2}{p^2} \right) + \mathcal{O}(\lambda^2) \simeq -i + \mathcal{O}(\lambda), \quad (2.387)$$

thus the β -function at 1-loop is:

$$\beta(\lambda) = \frac{3}{16\pi^2} \lambda^2 + \mathcal{O}(\lambda^3). \quad (2.388)$$

4.3 Analogous analysis for massless QED

Now we consider the QED, which as a good example to calculate the anomalous dimension $\gamma(\lambda)$. In QED, we need to renormalize the vertex(Z_1) and the propagator for electron(Z_2) and photon(Z_3). These three parameters defined the electron charge renormalization:

$$e_0 = Z_1^{-1} Z_2 Z_3^{1/2} e. \quad (2.389)$$

The Green's function $G^{(n_e, n_A)}(x_1 \dots x_n, e, M)$ now contains two fields, so the Callan-Symanzik equation is

$$\left(M \frac{\partial}{\partial M} + \beta(e) \frac{\partial}{\partial e} + n_e \gamma_2 + n_A \gamma_3 \right) G^{(n_e, n_A)}(x_1 \dots x_n, e, M) = 0, \quad (2.390)$$

where n_e denote for the electron fields, n_A for photon fields. And γ_2, γ_3 are anomalous dimensions of electron and photon field respectively

$$\gamma_i = \frac{1}{2} M \frac{\partial}{\partial M} \delta_i, \quad i = 2, 3. \quad (2.391)$$

The γ_i comes from the field renormalization, which related with Z_2 and Z_3 . And $Z_i = 1 + \delta_i$, where δ_i is the counterterms of fields. So for $i = 2$, i.e., the electron field

$$\begin{aligned} \delta_2 &= \frac{d}{d\bar{p}} \Sigma(p) \Big|_{p^2 = -M^2} = \frac{d}{d\bar{p}} \left(\text{---} \text{---} \text{---} \right) \\ &= \frac{-e^2}{16\pi^2} \left(\frac{\Gamma(\epsilon)}{(M^2)^{\epsilon/2}} + O(1) \right) \simeq \frac{-e^2}{16\pi^2} \left(\frac{2}{\epsilon} - \gamma_E \right) (1 - \epsilon \ln M) \\ &= \frac{-e^2}{16\pi^2} \left(\frac{2}{\epsilon} - \gamma_E - 2 \ln M \right), \end{aligned} \quad (2.392)$$

which leads to

$$\gamma_2 = \frac{e^2}{16\pi^2} \equiv \frac{\alpha}{4\pi}. \quad (2.393)$$

Also $i = 3$ is for the photon field

$$\begin{aligned} \delta_3 &= \Pi(p^2) \Big|_{p^2 = -M^2} = \left(\text{---} \text{---} \text{---} \right) = \left(g^{\mu\nu} - \frac{p^\mu p^\nu}{p^2} \right) \Pi(p^2) \\ &= \frac{-e^2}{12\pi^2} \left(\frac{\Gamma(\epsilon)}{(M^2)^{\epsilon/2}} + O(1) \right) \simeq \frac{-e^2}{12\pi^2} \left(\frac{2}{\epsilon} - \gamma_E - 2 \ln M \right). \end{aligned} \quad (2.394)$$

Thus,

$$\gamma_3 = \frac{e^2}{12\pi^2} \equiv \frac{\alpha}{4\pi} \left(\frac{4}{3} \right). \quad (2.395)$$

For calculating the β -function, we first briefly review a very important solution in QED renormalization. Consider the Ward-Takahashi identity

$$(p - q)_\mu \Gamma^\mu(p, q) = S_F^{-1}(p) - S_F^{-1}(q), \quad (2.396)$$

when we take the limit that $p \rightarrow q$ and move it to the right, thus we have

$$\Gamma^\mu(p, p) = \frac{\partial S_F^{-1}(p)}{\partial p_\mu}. \quad (2.397)$$

Notice $S_F^{-1}(p) = \gamma^\mu p_\mu - m - \Sigma(p)$, thus we have

$$\Gamma^\mu(p, p) = \gamma^\mu - \frac{\partial \Sigma(p)}{\partial p_\mu}. \quad (2.398)$$

Beside the momentum p , the propagator can be renormalized as

$$S_F^{-1}(p) = Z_2^{-1}(\gamma \cdot p - m_{\text{ph}}) + \dots, \quad (2.399)$$

which means

$$\frac{\partial S_F^{-1}}{\partial p_\mu} \approx Z_2^{-1} \gamma^\mu. \quad (2.400)$$

And for the renormalized vertex, we have

$$\Gamma^\mu = Z_1^{-1} \Gamma_R^\mu. \quad (2.401)$$

In low energy approximation, the vertex just $\Gamma_R^\mu \rightarrow \gamma^\mu$. Thus, comparing (2.398) and (2.401), we have

$$Z_1 = Z_2. \quad (2.402)$$

Thus, $\delta_1 = \delta_2$. Then we immediately have

$$0 = \frac{\partial e_0}{\partial M} = \frac{\partial}{\partial M} \left[e \left(1 - \delta_1 + \delta_2 + \frac{1}{2} \delta_3 \right) \right] \quad (2.403)$$

$$= e \frac{\partial}{\partial M} \left(1 - \delta_1 + \delta_2 + \frac{1}{2} \delta_3 \right) + \left(1 - \delta_1 + \delta_2 + \frac{1}{2} \delta_3 \right) \frac{\partial e}{\partial M} \quad (2.404)$$

$$\simeq e \frac{\partial}{\partial M} \left(-\delta_1 + \delta_2 + \frac{1}{2} \delta_3 \right) + \frac{\partial e}{\partial M}. \quad (2.405)$$

Therefore, we get the final result for the β -function:

$$\begin{aligned} \beta(e) &= M \frac{\partial}{\partial M} e = -e M \frac{\partial}{\partial M} \left(\underbrace{-\delta_1 + \delta_2}_{\delta_1 = \delta_2} + \frac{1}{2} \delta_3 \right) \\ &= \frac{e^3}{12\pi^2} \equiv e \frac{\alpha}{4\pi} \left(\frac{4}{3} \right). \end{aligned} \quad (2.406)$$

5. Evolution of coupling

5.1 Solution of Callan-Symanzik equation

Now we would like to solve the solutions of Callan-Symanzik equation. We firstly start with the 2-point function $G^{(2)}(p)$, which generally can be written as:

$$G^{(2)}(p) = \frac{i}{p^2} f \left(\frac{-p^2}{M^2} \right), \quad (2.407)$$

where $f \left(\frac{-p^2}{M^2} \right)$ is a dimensionless function, which only depends on $\frac{-p^2}{M^2}$. In momentum space, without the amputation, which means we add the external lines to contribute the momentum p in the propagator. We have:

$$M \frac{\partial}{\partial M} G^{(2)}(p) = \frac{i}{p^2} \left(\frac{2p^2}{M^2} \right) f' \left(\frac{-p^2}{M^2} \right), \quad (2.408)$$

$$p \frac{\partial}{\partial p} G^{(2)}(p) = -2G^{(2)}(p) - \frac{i}{p^2} \left(\frac{2p^2}{M^2} \right) f' \left(\frac{-p^2}{M^2} \right). \quad (2.409)$$

Therefore, we have:

$$M \frac{\partial}{\partial M} G^{(2)}(p) = \left(-p \frac{\partial}{\partial p} - 2 \right) G^{(2)}(p). \quad (2.410)$$

Putting this results into the Callan-Symanzik equation for 2-point function, we then have:

$$\left(p \frac{\partial}{\partial p} - \beta \frac{\partial}{\partial \lambda} + 2 - 2\gamma(\lambda) \right) G^{(2)}(p^2, \lambda, M) = 0. \quad (2.411)$$

This is an equation for a current. We can firstly write the solution in free theory, i.e. $\lambda = 0$ which means $\beta = \gamma = 0$:

$$G^{(2)}(p) = \frac{i}{p^2} \bar{G}_0, \quad (2.412)$$

where $\bar{G}_0$ is a constant function in p^2 . But one can not happy if we only have the free theory. Therefore, we need to solve the renormalization group equation with a more general form. For that, we firstly introduce some new variables to make the PDE simpler:

$$t = \ln \left(\frac{p}{M} \right), \quad (2.413)$$

$$v(x) = -\beta(\lambda), \quad (2.414)$$

$$\rho(x) = 2\gamma(\lambda) - 2, \quad (2.415)$$

$$x = \lambda. \quad (2.416)$$

Now we call the new function as $D(t, x)$, which is nothing but the $G^{(2)}$. So (2.411) becomes

$$\left(\frac{\partial}{\partial t} + v(x) \frac{\partial}{\partial x} - \rho(x) \right) D(t, x) = 0. \quad (2.417)$$

The solution of $D(t, x)$ is somehow like a current with the current density

$$j(t, x) = \rho(t, x) v(t, x). \quad (2.418)$$

So the solution of a current can be expressed as

$$D(t, x) = D_0(\bar{x}(t, x)) \exp \left[\int_0^t dt' \rho(\bar{x}(t', x)) \right], \quad (2.419)$$

with the boundary condition and the initial condition:

$$\frac{\partial}{\partial t'} \bar{x}(t', x) = -v(\bar{x}), \quad \bar{x}(0, x) = x, \quad (2.420)$$

where $\bar{x}(t, x)$ is the position of current volum element at t and x . So the Callan-Symanzik equation has the solution:

$$G^{(2)}(p^2, \lambda, M) = \bar{G}_0(\bar{\lambda}(p, \lambda)) \exp \left[\int_{p'=M}^{p'=p} d \ln \left(\frac{p'}{M} \right) 2 [1 - \bar{\gamma}(\bar{\lambda}(p', \lambda))] \right]. \quad (2.421)$$

It is important to note that $\bar{G}_0(\bar{\lambda}(p, \lambda))$ is an unknown function, which does not directly corresponds to the propagator. To fix this function, we firstly put (2.407) into (2.411):

$$0 = \left(p \frac{\partial}{\partial p} - \beta \frac{\partial}{\partial \lambda} + 2 - 2\gamma(\lambda) \right) \left[\frac{i}{p^2} f \left(\frac{p^2}{M^2}, \lambda \right) \right] \quad (2.422)$$

$$= \frac{i}{p^2} \left(-2 + p \frac{\partial}{\partial p} - \beta \frac{\partial}{\partial \lambda} + 2 - 2\gamma(\lambda) \right) f \left(\frac{p^2}{M^2}, \lambda \right) \quad (2.423)$$

Thus, we have

$$\left(p \frac{\partial}{\partial p} - \beta \frac{\partial}{\partial \lambda} - 2\gamma(\lambda)\right) f\left(\frac{p^2}{M^2}, \lambda\right) = 0. \quad (2.424)$$

So the solution is similar with $G^{(2)}$:

$$f\left(\frac{p^2}{M^2}, \lambda\right) = \bar{G}_0(\bar{\lambda}(p, \lambda)) \exp\left[-2 \int_M^p d \ln\left(\frac{p'}{M}\right) \gamma(\bar{\lambda})\right]. \quad (2.425)$$

Therefore, we can take the another form of the solution by combine the solution of f and (2.407)

$$G^{(2)}(p^2, \lambda, M) = \frac{i}{p^2} \bar{G}_0(\bar{\lambda}(p, \lambda)) \exp\left[-2 \int_M^p d \ln\left(\frac{p'}{M}\right) \gamma(\bar{\lambda}(p', \lambda))\right]. \quad (2.426)$$

Now, we can take the boundary $p = M$, which all the integration part will vanish. So we have

$$\bar{G}_0(\lambda) = -iM^2 \cdot G^{(2)}(M^2, \lambda, M). \quad (2.427)$$

We can then go to the specific case and use the perturbative expansion of propagator to get each order of $\bar{G}_0(\lambda)$.

5.2 Renormalization group equation

Now the boundary condition and the initial condition in (2.420) are changed to the so-called **renormalization group equation**:

$$\frac{d}{d \ln(p/M)} \bar{\lambda}(p, \lambda) = \beta(\bar{\lambda}), \quad \bar{\lambda}(M, \lambda) = \lambda. \quad (2.428)$$

This equation shows that the flow of the running coupling constant $\bar{\lambda}(p)$ with the speed $\beta(\bar{\lambda})$. And we set the initial energy scale is M , and then set the value at this point is the original coupling constant λ . Now we expand $\bar{\lambda}$ to the first order and use (2.388):

$$\frac{d}{d \ln(p/M)} \bar{\lambda}(p, \lambda) = \beta(\bar{\lambda}) = \frac{3}{16\pi^2} \bar{\lambda}^2 + O(\bar{\lambda}^3). \quad (2.429)$$

So we actually are solving the ODE

$$\frac{d\bar{\lambda}}{dt} = A\bar{\lambda}^2 \quad (2.430)$$

It is easy to solve it by integrating

$$\int_{\bar{\lambda}(M)=\lambda}^{\bar{\lambda}(p)} \bar{\lambda}'^{-2} d\bar{\lambda}' = \int_0^{\ln(p/M)} A dt' \quad (2.431)$$

Thus, we have

$$\frac{1}{\bar{\lambda}(p)} = \frac{1}{\bar{\lambda}} - A \ln(p/M) \quad (2.432)$$

$$= \frac{1 - A\bar{\lambda}(M) \ln(p/M)}{\bar{\lambda}(M)}. \quad (2.433)$$

Put all the parameters back, we have

$$\ln\left(\frac{p}{M}\right) = \left(\frac{3}{16\pi^2}\right)^{-1} \left(\frac{1}{\bar{\lambda}(M)} - \frac{1}{\bar{\lambda}(p)}\right). \quad (2.434)$$

Take $\bar{\lambda}(M) = \lambda$, which leads to the expression for the running coupling $\bar{\lambda}$:


$$\bar{\lambda}(p) = \frac{\lambda}{1 - \frac{3}{16\pi^2} \lambda \ln(p/M)}. \quad (2.435)$$

So $\bar{\lambda}(p)$ grows for increasing scales p . And there is a pole when

$$1 - \frac{3}{16\pi^2} \lambda \ln\left(\frac{p}{M}\right) = 0, \quad (2.436)$$

which is called the **Landau pole**:

$$p = M \exp\left(\frac{16\pi^2}{3\lambda}\right). \quad (2.437)$$



For $p^2 \ll M^2$, theory becomes trivial:

$$\bar{\lambda}(p) \sim \frac{\bar{\lambda}(M)}{\lim_{x \rightarrow 0} (1 - \ln x)} \rightarrow 0_+ \quad (2.438)$$

which corresponds to the free theory.

Generally, we have different cases for the β function, and that corresponds to the different background theory

$$\frac{d}{d \ln(p/M)} \bar{\lambda}(p, \lambda) = \beta(\bar{\lambda}) \begin{cases} > 0 & \text{IR free } (\phi^4\text{-theory, QED, ...}) \\ = 0 & \text{conformal theory } (N = 4 \text{ SYM}) \\ < 0 & \text{UV free/asymptotically free (QCD, SU(N) gauge theory).} \end{cases} \quad (2.439)$$

5.3 Higgs mass and fate of universe

As the second example, we consider the Higgs mechanism. The story begin with the Lagrangian:

$$\mathcal{L}_{\text{unbroken}} = \bar{\psi}(i\gamma^\mu \partial_\mu)\psi + \frac{1}{2}(\partial^\mu \phi)(\partial_\mu \phi) - V(\phi) + y_\psi \phi \bar{\psi}\psi. \quad (2.440)$$

where $y_\psi \phi \bar{\psi}\psi$ is the Higgs-Yukawa interaction, and we dropped all the gauge terms. To get the spontaneous symmetry breaking, we take the potential as

$$V(\phi) = \mu^2 \phi^2 + \lambda \phi^4. \quad (2.441)$$

The spontaneous symmetry breaking happens when

$$\phi = H + v. \quad (2.442)$$

where H is the Higgs field, v is the vacuum expectation value, which in experiment measure are

$$v \simeq \frac{1}{\sqrt{\sqrt{2}G_F}} \simeq 246 \text{ GeV} \quad (2.443)$$

So plug (2.442) into the Lagrangian, we have the spontaneous symmetry breaking Lagrangian:

$$\mathcal{L}_{\text{broken}} = \bar{\psi}(i\not{D})\psi + \frac{1}{2}(\partial_\mu H)(\partial^\mu H) - \underbrace{\frac{\lambda v^2}{\frac{1}{2}M_H^2}}_{\frac{1}{2}M_H^2} H^2 - \lambda v H^3 - \frac{\lambda}{4} H^4 - \underbrace{\frac{1}{\sqrt{2}}y_\psi v(1 + H/v)}_{m_\psi} \bar{\psi}\psi \quad (2.444)$$

This mechanism turns out to the particle mass by the Higgs interact with itself and the other fields. So the coupling constant renormalization is very similar with ϕ^4 theory and Yukawa theory:

$$-i\lambda + \underbrace{\text{bubble} + \text{tadpole} + \text{sunset}}_{O(\lambda^2)} + \underbrace{\text{box}}_{O(y_\psi^4)} \quad (2.445)$$


So the solution for $\lambda(Q)$ at scale $Q^2 \gg v$ is:

$$\lambda(Q) = \lambda(v) \frac{1}{1 - \frac{3}{16\pi^2} \lambda(v) \ln(Q/v)} \quad (2.446)$$

Cut off Λ_c corresponds to the Landau pole, up to which perturbative Higgs sector holds:

$$\Lambda_c = v \exp\left(\frac{16\pi^2}{3\lambda}\right) \simeq v \exp\left(\frac{16\pi^2 v^2}{M_H^2}\right) \quad (2.447)$$

where $M_H^2 = 2\lambda v^2$. So the area above the orange line are forbidden.



Now we consider fermion as the top quark, which is the heaviest fermion the standard model. It has contributions to $\beta(\lambda)$ from Higgs-Yukawa interaction, which the box diagram gives

$$Q \frac{d}{dQ} \lambda = \frac{1}{64\pi^2} (-3y_t^4 + 6\lambda y_t^2 + 12\lambda^2) + \dots \quad (2.448)$$

It is interesting that the first term is negative, which means it evolves in the opposite way with the Higgs. Solving this equation for small λ ($\lambda \ll 1$), we have

$$\lambda(Q) = \lambda(v) + \frac{1}{32\pi^2} \left(-\frac{3m_t^4}{v^4} \right) \ln\left(\frac{Q}{v}\right) \quad (2.449)$$

since $m_t = y_t \frac{v}{\sqrt{2}}$, $m_t \sim 172$ GeV, so in principle, $\lambda(Q)$ can be negative at large Q ! That means the vacuum will be super unstable. So to avoid that, we require a bound on Higgs mass:

$$M_H^2 \geq \frac{v^2}{16\pi^2} \left(-\frac{3m_t^4}{v^4} \right) \ln \left(\frac{Q}{v} \right), \quad (2.450)$$

such that we can keep $\lambda(Q) > 0$.

6. Renormalization for local operators and evolution of mass parameter

We first introduce the idea of local operator. The local operator is multiple field operator compositing in the same space-time position, namely the production of fields. This will happen when the energy scale is much lower than the high energy scattering, e.g. weak decay. So to understand this, we can recall the beta decay in electroweak interaction.

The electroweak interaction between left fermions and W-boson is:

$$\delta\mathcal{L} = \frac{g_{ew}}{\sqrt{2}} W_\mu \bar{\psi} \gamma^\mu (1 - \gamma_5) \psi. \quad (2.451)$$

And the propagator of W-boson is

$$\mu \text{ --- } W^- \text{ --- } \nu = \frac{-ig^{\mu\nu}}{q^2 - M_W^2 + i\epsilon} \quad (2.452)$$

In the low energy limit $q^2 \ll M_W^2$, the vertex and propagator of β -decay reduce to the 4-Fermi interaction vertex:

$$\begin{array}{ccc} \begin{array}{c} d \quad e^- \\ \swarrow \quad \searrow \\ W^- \\ \nearrow \quad \nwarrow \\ \bar{u} \quad \bar{\nu}_e \end{array} & \xrightarrow{q^2 \ll M_W^2} & \begin{array}{c} d \quad e^- \\ \swarrow \quad \searrow \\ \cdot \\ \nearrow \quad \nwarrow \\ \bar{u} \quad \bar{\nu}_e \end{array} \end{array} \quad (2.453)$$

So we can reduce a process propagate from μ to ν to the process happens in a fixed point due to the reaction happening so fast, i.e.:

$$\begin{array}{c} d \quad e^- \\ \swarrow \quad \searrow \\ \cdot \\ \nearrow \quad \nwarrow \\ \bar{u} \quad \bar{\nu}_e \end{array} = \frac{g_{ew}^2}{2M_W^2} (\bar{\psi} \gamma^\mu (1 - \gamma_5) \psi)_{\text{quark}} (\bar{\psi} \gamma_\mu (1 - \gamma_5) \psi)_{\text{lep}} = \frac{g_{ew}^2}{2M_W^2} \mathcal{O}(x). \quad (2.454)$$

Generally, this special vertex can be seen as a Green's function:

$$\langle \psi(p_1) \bar{\psi}(p_2) \psi(p_3) \bar{\psi}(p_4) \mathcal{O}(0) \rangle = \begin{array}{c} p_1 \quad p_3 \\ \swarrow \quad \searrow \\ \cdot \quad \mathcal{O}(0) \\ \nearrow \quad \nwarrow \\ p_2 \quad p_4 \end{array} \quad (2.455)$$

6.1 Renormalized Green's functions with local operator $\mathcal{O}$ in ϕ^4 -theory

Now we consider the renormalization of the local operator. Since the vertex is described by the Green's function, we can then consider a general form:

$$G^{(n,\mathcal{O})}(p_1 \dots p_n, k) = \langle \phi(p_1) \dots \phi(p_n) \mathcal{O}(k) \rangle \quad (2.456)$$

$$= Z^{-n/2}(\mu) \underbrace{Z_{\mathcal{O}}^{-1}(M)}_{\text{operator renormalization}} \underbrace{\langle \phi_0(p_1) \dots \phi_0(p_n) \mathcal{O}_0(k) \rangle}_{\text{bare fields bare operator}} \quad (2.457)$$

$$= Z^{-n/2}(\mu) Z_{\mathcal{O}}^{-1}(M) G_0^{(n,\mathcal{O})}(p_1 \dots p_n, k) \quad (2.458)$$

where $\mathcal{O} = \phi^2$ or ϕ^4 or $\dots$. The lower index "0" represent the bare quantity, and M is the renormalization scale. The bare operator and the renormalized operator have the relation:

$$\mathcal{O}_{\text{bare}} = Z_{\mathcal{O}}(M) \mathcal{O}_{\text{ren}}. \quad (2.459)$$

So we can derive the Callan-Symanzik equation by using

$$0 = M \frac{d}{dM} G_0^{(n,\mathcal{O}_0)} \quad (2.460)$$

$$= M \frac{d}{dM} (Z^{n/2} Z_{\mathcal{O}} G^{(n,\mathcal{O})}) \quad (2.461)$$

$$= \left(M \frac{\partial}{\partial M} Z^{n/2} \right) Z_{\mathcal{O}} G^{(n,\mathcal{O})} + Z^{n/2} \left(M \frac{\partial}{\partial M} Z_{\mathcal{O}} \right) G^{(n,\mathcal{O})} + Z^{n/2} Z_{\mathcal{O}} \left(M \frac{\partial}{\partial M} G^{(n,\mathcal{O})} \right), \quad (2.462)$$

dividing $Z^{n/2} Z_{\mathcal{O}}$ on both side, we have

$$0 = \left(\frac{M}{Z^{n/2}} \frac{\partial Z^{n/2}}{\partial M} \right) G^{(n,\mathcal{O})} + \left(\frac{M}{Z_{\mathcal{O}}} \frac{\partial Z_{\mathcal{O}}}{\partial M} \right) G^{(n,\mathcal{O})} + M \frac{\partial}{\partial M} G^{(n,\mathcal{O})} \quad (2.463)$$

$$= \left(M \frac{\partial}{\partial M} \ln Z^{n/2} \right) G^{(n,\mathcal{O})} + \left(M \frac{\partial}{\partial M} \ln Z_{\mathcal{O}} \right) G^{(n,\mathcal{O})} + M \frac{\partial}{\partial M} G^{(n,\mathcal{O})}, \quad (2.464)$$

where we can define the anomalous dimension respectively for field and local operator as:

$$\gamma = \frac{M}{2} \frac{\partial}{\partial M} \ln Z_{\lambda}(M) \quad (2.465)$$

$$\gamma_{\mathcal{O}} = M \frac{\partial}{\partial M} \ln Z_{\mathcal{O}}(M). \quad (2.466)$$

Then we have the Callan-Symanzik equation:

$$\left(M \frac{\partial}{\partial M} + \beta(\lambda) \frac{\partial}{\partial \lambda} + n\gamma(\lambda) + \gamma_{\mathcal{O}}(\lambda) \right) G^{(n,\mathcal{O})} = 0, \quad (2.467)$$


Now we compute $\gamma_{\mathcal{O}}$ for N-point Green's function. The most general case can be expressed as

$$\text{Diagram with red dot and external lines} + \sum_i \left(\text{Diagram with red dot and internal loop} + \text{Diagram with red dot and internal loop} \right) + \text{Diagram with red cross and external lines}, \quad (2.468)$$

where the red point and red cross represent the operator vertex and the operator counterterm.

6.2 Renormalization for local operators in massless ϕ^4 -theory

As an example to help us understanding this. We consider $\mathcal{O} = \phi^2$ inserted in two-point function:




$$= \langle \Omega | \phi(q) \phi(p) \phi^2(k) | \Omega \rangle = G^{(2, \phi^2)}. \quad (2.469)$$

Thus, we can expand it as the normal Dyson series:

$$G^{(2, \phi^2)}(q, p; k) = \sum_{m=0}^{\infty} \frac{i^m}{m!} \int \left[\prod_{a=1}^m d^d x_a \right] \left(-\frac{\lambda}{4!} \right)^m \langle 0 | T \{ \phi(q) \phi(p) \phi^2(k) [\phi^4(x_1) \dots \phi^4(x_m)] \} | 0 \rangle_{\text{connect}}. \quad (2.470)$$

The leading order, namely the tree-level non-amputated Green's function is just simply the contraction of the field and the operator:




$$G_{\text{tree}}^{(2, \phi^2)}(q, p; k) = \int d^d x e^{-ik \cdot x} \langle 0 | T \{ \phi(q) \phi(p) \phi_A(x) \phi_B(x) \} | 0 \rangle \quad (2.471)$$

$$= \int d^d x e^{-ik \cdot x} \left[\overbrace{\phi(q) \phi(p) \phi_A(x) \phi_B(x)} + \overbrace{\phi(q) \phi(p) \phi_A(x) \phi_B(x)} \right] \quad (2.472)$$

$$= \left(\frac{i}{q^2} \right) \left(\frac{i}{p^2} \right) \cdot 2. \quad (2.473)$$

To consider the renormalization for the operator, we need to compute 1-PI contribution at 1-loop (non-amputated):



$$G_{1\text{-loop}}^{(2, \phi^2)}(q, p; k) = u+v = \int d^d x d^d z e^{-ik \cdot x} \left\langle 0 \left| T \left\{ \phi(q) \phi(p) \left[\frac{-i\lambda}{4!} \phi^4(z) \right] \left[\frac{1}{2} \phi_A(x) \phi_B(x) \right] \right\} \right| 0 \right\rangle \quad (2.474)$$


$$= \frac{-i\lambda}{2} \int d^d x d^d z e^{-ik \cdot x} \left[\overbrace{\phi(q) \phi(p) \phi(z) \phi(z) \phi(z) \phi(z) \phi_A(x) \phi_B(x)} \right] \quad (2.475)$$

$$= \left(\frac{i}{q^2} \right) \left(\frac{i}{p^2} \right) \int \frac{d^d v}{(2\pi)^d} (-i\lambda) \frac{i}{v^2} \frac{i}{(k+v)^2}, \quad (2.476)$$

with some calculation, one has:

$$G_{1\text{-loop}}^{(2, \phi^2)}(q, p; k) = \frac{i}{p^2} \frac{i}{q^2} \left(-\frac{\lambda}{(4\pi)^{d/2}} \frac{\Gamma(2 - \frac{d}{2})}{\Delta^{2-d/2}} \right) \quad (2.477)$$

with the counterterm:



$$= \frac{i}{p^2} \frac{i}{q^2} 2\delta\phi^2 \quad (2.478)$$

Comparing of coefficients in these two graphs, we then can find the counterterm at scale $p^2 = -M^2$ should be:

$$\delta\phi^2 = \frac{\lambda}{2(4\pi)^{d/2}} \frac{\Gamma(2 - \frac{d}{2})}{(M^2)^{2-d/2}}. \quad (2.479)$$

This implies the operator renormalization is

$$Z_{\phi^2} = 1 - \delta\phi^2 = 1 - \frac{\lambda}{2(4\pi)^{d/2}} \frac{\Gamma(2 - \frac{d}{2})}{(M^2)^{2-d/2}} \quad (2.480)$$

$$\stackrel{d=4-\varepsilon}{=} 1 - \frac{\lambda}{(4\pi)^2} \frac{1}{\varepsilon} + \text{finite terms}. \quad (2.481)$$

Using $\ln(1+x) \approx x$, we can obtain the anomalous dimension of the operator in $d = 4 - \varepsilon$:

$$\gamma_{\phi^2} = M \frac{\partial}{\partial M} \ln Z_{\phi^2} = \frac{-\lambda}{2(4\pi)^{d/2}} \Gamma\left(2 - \frac{d}{2}\right) \cdot \left[-(4-d)M^{-(4-d)} \right] \quad (2.482)$$

$$\stackrel{d=4-\varepsilon}{=} \frac{-\lambda}{2(4\pi)^{2-\varepsilon/2}} \Gamma\left(\frac{\varepsilon}{2}\right) \cdot \left[-\varepsilon M^{-\varepsilon} \right] \quad (2.483)$$

$$= \frac{\lambda}{16\pi^2}. \quad (2.484)$$

More generally, if we insert the operator into an n-point Green's function, we then need to consider the field renormalization with the operator renormalization $Z_{\phi}^{n/2} Z_{\mathcal{O}}$, therefore the anomalous dimension becomes:

$$\gamma_{\mathcal{O}}(\lambda) = M \frac{\partial}{\partial M} \left(-\delta_{\mathcal{O}} + \frac{n}{2} \delta_Z \right), \quad (2.485)$$

with the approximation:

$$\ln(Z_{\phi}^{n/2} Z_{\mathcal{O}}) = \frac{n}{2} \ln Z_{\phi} + \ln Z_{\mathcal{O}} \approx \frac{n}{2} \delta_Z - \delta_{\mathcal{O}}. \quad (2.486)$$

This case can be treated as renormalized ϕ^4 -theory with effective operator $-\frac{1}{2}m^2\phi^2$, where ϕ^2 is local operator inserted in Green's function. So the Callan-Symanzik equation with mass anomalous dimension $m(\mu)$ is

$$\left(M \frac{\partial}{\partial M} + \beta(\lambda) \frac{\partial}{\partial \lambda} + n\gamma(\lambda) + \gamma_{\phi^2} m^2 \frac{\partial}{\partial m^2} \right) G^{(n, \phi^2)}(\{p_i\}, M, \lambda, m) = 0 \quad (2.487)$$

6.3 Evolution of mass parameter

Moreover, one can write the generalization to local operators $\mathcal{O}_i$ in ϕ^4 -theory as:

$$\mathcal{L}(C_i) = \mathcal{L}_{\text{eff}} + \sum_i C_i \mathcal{O}_i(x) \quad (2.488)$$

where C_i is the coupling of local operators. So the Callan-Symanzik equation is

$$\left(M \frac{\partial}{\partial M} + \beta(\lambda) \frac{\partial}{\partial \lambda} + n\gamma(\lambda) + \sum_i \gamma_i(\lambda) C_i \frac{\partial}{\partial C_i} \right) G^{(n)}(M, \lambda, \{C_i\}) = 0 \quad (2.489)$$

To guarantee the dimension of Lagrangian is $[\text{Mass}]^4$ in 4-dimension spacetime, so we need to explicit mass dimension in coupling C_i as:

$$C_i = \rho_i M^{4-d_i} \quad (2.490)$$

where ρ_i is dimensionless, d_i is dimension of $\mathcal{O}_i$. Therefore, one can see that the variables change to $(M, \lambda, \{\rho_i\})$. To make the partial derivative independent to ρ_i , we have

$$M \frac{\partial}{\partial M} \Big|_{\rho_i} = M \frac{\partial}{\partial M} \Big|_{C_i} + \sum_i \left(M \frac{\partial C_i}{\partial M} \Big|_{\rho_i} \right) \frac{\partial}{\partial C_i} \quad (2.491)$$

Notice that

$$\rho_i \frac{\partial}{\partial \rho_i} = \rho_i M^{4-d_i} \frac{\partial}{\partial C_i} = C_i \frac{\partial}{\partial C_i} \quad (2.492)$$

so we have

$$M \frac{\partial}{\partial M} \Big|_{C_i} = M \frac{\partial}{\partial M} \Big|_{\rho_i} - \sum_i (4 - d_i) \rho_i \frac{\partial}{\partial \rho_i} \quad (2.493)$$

Then the Callan-Symanzik equation becomes:

$$\left(M \frac{\partial}{\partial M} + \beta(\lambda) \frac{\partial}{\partial \lambda} + n\gamma(\lambda) + \sum_i (\gamma_i(\lambda) + d_i - 4) \rho_i \frac{\partial}{\partial \rho_i} \right) G^{(n)} = 0, \quad (2.494)$$

where we can define the β function for new coupling ρ_i :

$$\beta_i(\lambda) = (\gamma_i(\lambda) + d_i - 4) \rho_i, \quad (2.495)$$

then the Callan-Symanzik equation can be reduced as:

$$\left(M \frac{\partial}{\partial M} + \beta(\lambda) \frac{\partial}{\partial \lambda} + n\gamma(\lambda) + \sum_i \beta_i \rho_i \frac{\partial}{\partial \rho_i} \right) G^{(n)} = 0. \quad (2.496)$$

With this, we can easily solve the running coupling ρ_i from the renormalization group equation for β_i

$$\frac{d}{d \ln(p/M)} \bar{\rho}_i = \beta_i(\bar{\rho}_i, \lambda). \quad (2.497)$$

Since β_i in the low energy limit, namely $\gamma_i(\lambda) \ll 1$, is dominated by the classical dimension $d_i - 4$, we then have

$$\frac{d}{d \ln(p/M)} \bar{\rho}_i \simeq (d_i - 4 + \dots) \cdot \bar{\rho}_i \quad (2.498)$$

Therefore, the solution in low energy limit reads:

$$\bar{\rho}_i = \rho \left(\frac{p}{M} \right)^{d_i - 4}. \quad (2.499)$$

Now we can clearly see that for the operators with mass dimension $d_i > 4$ will vanish as $p \rightarrow 0$, e.g., ϕ^6 theory with $d_i = 6$. These operators are called the **irrelevant operators**, which usually decrease fast until vanish in low energy limit, namely one can never feel these operators' contribution. On the contrary, the **relevant operators** are the operators with mass dimension $d_i < 4$, e.g., the mass term $\frac{1}{2}m^2\phi^2$, which will exponentially increase in low energy limit. And the operators like ϕ^4 are called the **marginal operators**, which do not changing with the energy scale due to $d_i = 4$. This type operators are only determined by the anomalous dimension $\gamma_i(\lambda)$.

This reminds us that the effective field theory truly works even though we don't know all the details in high energy level. But we still can derive the precise renormalizable Lagrangian in low energy level.

Non-Abelian gauge theories

1. Recap of gauge theory and brief look at Lie group theory

1.1 Recap: Abelian gauge theory

We firstly start with QED, which as we know is an Abelian $U(1)$ gauge theory. It start from the free fermion Lagrangian:

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi. \quad (3.1)$$

The Noether current is: $j^\mu = \bar{\psi}\gamma^\mu\psi$. To approach the QED Lagrangian, we introduce the (global) gauge transformation as:

$$\psi \rightarrow \psi' = e^{i\alpha}\psi \quad (3.2)$$

$$\bar{\psi} \rightarrow \bar{\psi}' = e^{-i\alpha}\bar{\psi} \quad , \quad \alpha \in \mathbb{R}. \quad (3.3)$$

The local gauge transformation is just changing the parameter α rely on the coordinate:

$$\alpha \rightarrow \alpha(x). \quad (3.4)$$

The gauge symmetry is saying that the Lagrangian is invariant under the gauge transformation. This requires us to introduce the covariant derivative D_μ :

$$D_\mu\psi(x) = (\partial_\mu - igA_\mu)\psi(x). \quad (3.5)$$

where in QED case, the charge $g = e$, and the gauge field A_μ is just the photon field, which satisfies the transformation rule:

$$A_\mu \rightarrow A'_\mu = A_\mu + \frac{1}{g}\partial_\mu\alpha(x). \quad (3.6)$$

Then we immediately have the covariant derivative transfers as

$$D_\mu\psi \rightarrow (D_\mu\psi)' = \partial_\mu\psi' - igA'_\mu\psi' = e^{i\alpha}D_\mu\psi. \quad (3.7)$$

Thus the invariant of $\mathcal{L}$ requires that

$$\mathcal{L}(\psi, A_\mu) = \mathcal{L}(\psi', A'_\mu) = i\bar{\psi}\gamma^\mu D_\mu\psi - m\bar{\psi}\psi \quad (3.8)$$

$$= i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi + gA_\mu j^\mu. \quad (3.9)$$

For photon dynamical, we need to add kinetic term for A_μ :

$$\mathcal{L}_{\text{Maxwell}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (3.10)$$

with $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.

1.2 $SU(N)$ gauge theory

Now we consider the non-Abelian $SU(N)$ gauge theory with N Dirac spinors ψ_j , where $j = 1, \dots, N$. The original Lagrangian is

$$\mathcal{L} = \sum_{j=1}^N i\bar{\psi}_j \gamma^\mu \partial_\mu \psi_j - m\bar{\psi}_j \psi_j. \quad (3.11)$$

Under the global unitary transformation $U_{ij} \in \mathbb{C}$, $UU^\dagger = \mathbb{1}$, the spinors obey

$$\psi_j \rightarrow \psi'_j = \sum_k U_{jk} \psi_k, \quad (3.12)$$

$$\bar{\psi}_j \rightarrow \bar{\psi}'_j = \sum_k \bar{\psi}_k U_{kj}^\dagger, \quad (3.13)$$

thus the mass term transformation under $SU(N)$

$$\sum_j \bar{\psi}_j \psi_j \rightarrow \sum_j \bar{\psi}'_j \psi'_j = \sum_{k,l,j} \bar{\psi}_k \underbrace{U_{kj}^\dagger U_{jl}}_{\delta_{kl}} \psi_l = \sum_k \bar{\psi}_k \psi_k \quad (3.14)$$

is invariant.

Similar with the Abelian case, the local gauge transformation is $U \rightarrow U(x)$. We also require the Lagrangian invariant under the transformation. This gives coupling of fermions to gauge field in covariant derivative analogous to QED:

$$D_\mu \psi_j = \partial_\mu \psi_j - ig \sum_k (A_\mu)_{jk} \psi_k. \quad (3.15)$$

This form of covariant derivative also requires the transformation rule as

$$D_\mu \psi_j \rightarrow (D_\mu \psi_j)' = \partial_\mu \psi'_j - ig \sum_k (A'_\mu)_{jk} \psi'_k \quad (3.16)$$

$$= \sum_k U_{jk} (D_\mu \psi_k) \quad (3.17)$$

For convenient, we will use the short-hand notation (Matrix notation), which basically just ignore the index of the matrix element:

$$D_\mu \psi \rightarrow (D_\mu \psi)' = U D_\mu \psi = e^{i\alpha(x)} D_\mu \psi. \quad (3.18)$$

We claim that the gauge field transforms following

$$A_\mu \rightarrow A'_\mu = U A_\mu U^\dagger - \frac{i}{g} (\partial_\mu U) U^\dagger. \quad (3.19)$$

Although we can directly use the definition to compute this expression, I just don't want to write so much. Instead, we can have a quick check that this claim indeed satisfies our requirement:

$$D_\mu \psi \rightarrow (D_\mu \psi)' = \partial_\mu (U\psi) - ig \left(U A_\mu U^\dagger - \frac{i}{g} (\partial_\mu U) U^\dagger \right) U\psi \quad (3.20)$$

$$= U \partial_\mu \psi + (\partial_\mu U) \psi - ig U A_\mu \psi - (\partial_\mu U) \psi \quad (3.21)$$

$$= U (\partial_\mu \psi - ig A_\mu \psi) = U D_\mu \psi. \quad (3.22)$$

Thus, the Lagrangian is invariant under $SU(N)$ gauge transformation.

Now we need to approach the dynamical of non-Abelian gauge field, which requires a field strength. Since we keep calling it as “non-Abelian”, but until now it’s not quite obvious to see how it is “non-Abelian”, therefore, the field strength should give you some direct idea when we claim:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ig[A_\mu, A_\nu]. \quad (3.23)$$

We can also check it is true that this field strength indeed satisfies our requirement, namely keep the Lagrangian invariant. To show this, we firstly have the transformation rule:

$$F_{\mu\nu} \rightarrow F'_{\mu\nu} = \partial_\mu A'_\nu - \partial_\nu A'_\mu - ig[A'_\mu, A'_\nu] = U F_{\mu\nu} U^\dagger. \quad (3.24)$$

The Lagrangian is proportion to the trace over $F_{\mu\nu} F^{\mu\nu} = (F_{\mu\nu})_{jk} (F^{\mu\nu})^{kj} = \text{tr}(F_{\mu\nu} F^{\mu\nu})$, due to the Lagrangian is a scalar, thus the only possible combination is the trace over $F_{\mu\nu}$. Thus $\text{tr} F_{\mu\nu}$ transfers as

$$\begin{aligned} \text{tr}(F_{\mu\nu} F^{\mu\nu}) &\rightarrow \text{tr}(F'_{\mu\nu} F'^{\mu\nu}) = \text{tr}(U F_{\mu\nu} U^\dagger U F^{\mu\nu} U^\dagger) \\ &= \text{tr}(U U^\dagger F_{\mu\nu} F^{\mu\nu}) \\ &= \text{tr}(F_{\mu\nu} F^{\mu\nu}). \end{aligned} \quad (3.25)$$

Then we get **the Lagrangian of Yang-Mills theory**

$$\mathcal{L} = -\kappa \text{tr}(F_{\mu\nu} F^{\mu\nu}) + \sum_{j,k=1}^N [\bar{\psi}_j \gamma^\mu (D_\mu)_{jk} \psi_k - m \bar{\psi}_j \psi_j], \quad (3.26)$$

where κ is normalized constant.

1.3 Recap for essential Lie group theory

To continue our journey, it’s necessary to master some basic knowledge about Lie group and Lie algebra. So in the next, we will recall some essential knowledge about these mathematical tools and show how are they used in gauge theory.

Firstly, we recall the infinitesimal transformation of the gauge operation. For Abelian theory, e.g. QED:

$$e^{i\alpha} \simeq 1 + i\alpha. \quad (3.27)$$

For non-Abelian gauge theory, e.g. Yang Mills theory:

$$U = \mathbb{1} + \hat{H} + \mathcal{O}(\hat{H}^2), \quad (3.28)$$

$$U^\dagger = \mathbb{1} + \hat{H}^\dagger + \mathcal{O}(\hat{H}^2). \quad (3.29)$$

Thus we have

$$U^{-1} = U^\dagger \implies \hat{H} = -\hat{H}^\dagger \quad (\text{anti-hermitian}) \quad (3.30)$$

where we note $\hat{H} = iH$ with $H = H^\dagger$. Expand $\hat{H}$ in parameters of transformation α^a

$$\hat{H}_{jk} = i \sum_a \alpha^a (t^a)_{jk} \quad (3.31)$$

with $\alpha^a \in \mathbb{R}$, $t^a = (t^a)^\dagger$ is hermitian. Thus t^a span a basis in the space of hermitian $N \times N$ matrices such that we can expand $\hat{H}$ in terms of this basis. t^a are called the generators of gauge transformations.

Generally, the unitary $N \times N$ matrix has N^2 linear independent generators due to t^a is $N \times N$. However, in our case, we require $\det U = 1$, which gives one constraint. So there are **$N^2 - 1$ generators in $\text{SU}(N)$** . Further more, base on $\det U = 1$ and the Jacobi identity $\det(e^M) = e^{\text{tr}(M)}$, we find

$$\det U = \det(e^{\hat{H}}) = e^{\text{tr} \hat{H}} = 1, \quad (3.32)$$

thus, $\text{tr} \hat{H} = 0$ and implies

$$\text{tr}(t^a) = 0. \quad (3.33)$$

So **t^a are traceless generators.**

Now we discuss the algebra structure for generators t^a . The Lie algebra, namely the commutator is given by:

$$[t^a, t^b] = i \sum_c f^{abc} t^c. \quad (3.34)$$

where f^{abc} is the structure constant of Lie algebra. The elements of Lie group can be expanded around $\mathbb{1}$.

$$g(\alpha) = \mathbb{1} + i \sum_{a=1}^d \alpha_a t^a + \frac{1}{2} \sum_{a,b=1}^d \alpha_a \alpha_b T^{ab} + \mathcal{O}(\alpha^3), \quad (3.35)$$

where $T^{ab} = T^{ba}$ is symmetric, which can be expanded in terms of the generator t^a, t^b . Notice $g(0) = \mathbb{1}$ and because of $g(\alpha)\mathbb{1} = \mathbb{1}g(\alpha)$, we then have

$$\gamma_a(\alpha_1, \dots, \alpha_d; 0, \dots, 0) = \gamma_a(0, \dots, 0; \alpha_1, \dots, \alpha_d) = \alpha_a \quad (3.36)$$

where the γ is defined by the group multiplication:

$$g(\alpha_1, \dots, \alpha_d) \cdot g(\beta_1, \dots, \beta_d) = g(\gamma_1, \dots, \gamma_d), \quad (3.37)$$

in which γ is the function of α and β :

$$\gamma_a = \gamma_a(\alpha_1, \dots, \alpha_d; \beta_1, \dots, \beta_d). \quad (3.38)$$

Base on the fact that $\gamma(\alpha, 0) = \alpha$ and $\gamma(0, \beta) = \beta$, then we have only one choice to expand γ_a until the second order to keep these relation, namely:

$$\gamma_a = \alpha_a + \beta_a + \sum_{b,c} C_{bc}^a \alpha_b \beta_c + \dots, \quad (3.39)$$

where C_{bc}^a is a unknown coefficient. Thus, use $g(\alpha)g(\beta) = g(\gamma)$, we have for the left-hand side:

$$g(\alpha)g(\beta) = \mathbb{1} + i \sum_a (\alpha_a + \beta_a) t^a + \frac{1}{2} \sum_{a,b} [(\alpha_a \alpha_b + \beta_a \beta_b) T^{ab} - 2\alpha^a \beta^b t^a t^b] + \dots, \quad (3.40)$$

And for the right-hand side:

$$g(\gamma) = \mathbb{1} + i \sum_a \gamma_a t^a + \frac{1}{2} \sum_{a,b} \gamma_a \gamma_b T^{ab} + \dots \quad (3.41)$$

Thus use (3.39), then the right-hand side becomes

$$g(\gamma(\alpha, \beta)) = \mathbb{1} + i \sum_a (\alpha_a + \beta_a) t^a + i \sum_{a,b,c} C_{bc}^a \alpha_b \beta_c t^a + \frac{1}{2} \sum_{a,b} (\alpha_a + \beta_a)(\alpha_b + \beta_b) T^{ab} + \dots \quad (3.42)$$

Compare (3.40) with (3.42), we find

$$-t^a t^b = i \sum_c C_c^{ab} t^c. \quad (3.43)$$

this gives the structure constant:

$$[t^a, t^b] = i \sum_{c=1}^d (C_c^{ba} - C_c^{ab}) t^c \quad (3.44)$$

$$= i \sum_{c=1}^d f^{abc} t^c \quad (3.45)$$

where $f^{abc} = -f^{bac}$, and d is the dimension of Lie algebra.

Examples of Lie groups

- **GL(N)**: Group of $N \times N$ matrices in $\mathbb{R}^N$ with $\det M \neq 0$.
- **U(N)**: Group of unitary $N \times N$ matrices in $\mathbb{C}^N$, satisfies

$$U^\dagger U = U U^\dagger = \mathbb{1} \quad \text{and} \quad \det U \neq 0. \quad (3.46)$$

Infinitesimal version:

$$U = \mathbb{1} + i \sum_{a=1}^d \alpha_a t^a. \quad (3.47)$$

with $d = N^2$ is the dimension of $U(N)$ Lie algebra/group.

- **SU(N)** := $\{A \in U(N) | \det A = 1\}$. Dimension of Lie algebra $\mathfrak{su}(N)$ is $d = N^2 - 1$. Generators t^a are traceless

$$\det U = \det \left(\mathbb{1} + i \sum_{a=1}^d \alpha_a t^a \right) = 1 + i \sum_{a=1}^d \alpha_a \text{tr } t^a = 1 \quad (3.48)$$

thus

$$\text{tr } t^a = 0 \quad (3.49)$$

- **O(N)**: group of orthogonal $N \times N$ matrices in $\mathbb{R}^N$.

$$O O^T = O^T O = \mathbb{1} \quad (3.50)$$

Infinitesimal version:

$$O = \mathbb{1} + i \sum_{a=1}^d \alpha_a t^a \quad (3.51)$$

Dimension of Lie algebra $\mathfrak{o}(N)$ is $\frac{1}{2}N(N-1)$. And notice:

$$O O^T = \left(\mathbb{1} + i \sum_{a=1}^d \alpha_a t^a \right) \left(\mathbb{1} + i \sum_{b=1}^d \alpha_b (t^b)^T \right) = \mathbb{1}, \quad (3.52)$$

thus,

$$\mathbb{1} = \mathbb{1} + i \sum_{a=1}^d \alpha_a t^a + i \sum_{b=1}^d \alpha_b (t^b)^T + \mathcal{O}(\alpha^2) \quad (3.53)$$

$$= \mathbb{1} + i \sum_{a=1}^d (\alpha_a t^a + \alpha_a (t^a)^T), \quad (3.54)$$

which implies

$$t^a = -(t^a)^T. \quad (3.55)$$

- $\mathbf{SO}(N) := \{A \in O(N) \mid \det A = 1\}$
- $\mathbf{SO}(N, M) = \{\Lambda \in \text{Mat}(N+M, \mathbb{R}) \mid \Lambda^T \eta \Lambda = \eta, \det \Lambda = 1\}$, where

$$\eta = \text{diag}(\underbrace{1, \dots, 1}_N, \underbrace{-1, \dots, -1}_M) \quad (3.56)$$

- **Lorentz group:** $SO(1, 3)$

1.4 Representations of Lie algebra

Def: Representation of group

A **representation of a group G on a d_r -dimensional vector space V** is a group homomorphism

$$\rho : G \rightarrow GL(V), \quad (3.57)$$

which maps each group element $g \in G$ to an invertible linear operator $\rho(g)$ acting on V , such that:

$$\rho(g_i) \cdot \rho(g_j) = \rho(g_i g_j) \quad \forall g_i, g_j \in G. \quad (3.58)$$

The dimension d_r of the vector space V is called **the dimension of the representation**.

- A representation ρ is called **reducible** if there exists a non-trivial proper subspace $U \subset V$ (where $U \neq \{0\}$ and $U \neq V$) that is invariant under the action of all group elements, meaning:

$$\rho(g)u \in U \quad \forall g \in G, \forall u \in U. \quad (3.59)$$

- If no such invariant subspace exists, the representation is called irreducible (irrep).

For a Lie group characterized by continuous parameters α , a representation is a mapping $\rho : g(\alpha) \rightarrow M(\alpha)$, where the matrices (or operators) $M(\alpha)$ preserve the group multiplication:

$$M(\alpha) \cdot M(\beta) = M(\gamma) \quad \text{whenever} \quad g(\alpha) \cdot g(\beta) = g(\gamma). \quad (3.60)$$

In the next, we introduce some useful representations.

- **Fundamental(defining) representation**

For a matrix Lie group, the **fundamental representation** is simply the group of matrices themselves, acting on their natural vector space.

Example: $SU(2)$ fundamental representation

- **Representation Space (V):** A 2-dimensional complex vector space $\mathbb{C}^2$, physically corresponding to a spin-1/2 state

$$|\psi\rangle = \alpha |\uparrow\rangle + \beta |\downarrow\rangle. \quad (3.61)$$

- **Dimension of the representation:** $d = 2$ (since the matrices are 2×2). Note that the dimension of the $SU(2)$ group/algebra itself is $2^2 - 1 = 3$ (the number of

independent generators). The 3 Hermitian generators for this $d = 2$ fundamental representation are the Pauli matrices:

$$t^a = \frac{1}{2}\sigma^a \quad (a = 1, 2, 3), \quad (3.62)$$

which satisfy the Lie algebra $[t^a, t^b] = i\epsilon^{abc}t^c$, and the Pauli matrix identity:

$$\sigma^k \sigma^l = \delta^{kl} \mathbb{1} + i\epsilon^{klm} \sigma^m \quad (3.63)$$

Physically, we can use these fundamental building blocks to construct higher-dimensional representations via tensor products, such as the 3-dimensional spin-1 triplet state. In fact, $SU(2)$ has spin- j representations with dimension $2j + 1$. For instance, $j = 1/2$ corresponds to the 2-dimensional fundamental representation using 2×2 Pauli matrices.

- **Adjoint representation**

Instead of acting on physical states, the **adjoint representation** is a d -dimensional representation where the group/algebra acts on the Lie algebra space itself (spanned by the generators $\{t^a\}$ as basis vectors). The definition is as follows

Adjoint representation

We first define a automorphism map of group G :

$$\Psi : G \rightarrow \text{Aut}(G) \quad (3.64)$$

$$g \mapsto hgh^{-1} \equiv \Psi_h(g), \quad \forall h \in G, \quad (3.65)$$

where $\text{Aut}(G)$ is the automorphism group of G . Then we can define the derivative of Φ_g at the origin noted as Ad_g :

$$\text{Ad}_g = (d\Phi_g)_e : T_e G \rightarrow T_e G, \quad (3.66)$$

where $T_e G$ is the tangent space in the natural element of G . Thus the **adjoint representation** is a homomorphism:

$$\text{Ad} : G \rightarrow \text{Aut}(T_e G) \quad (3.67)$$

$$g \mapsto \text{Ad}_g. \quad (3.68)$$

To obtain the explicit matrix form of the adjoint representation, we consider an infinitesimal group element $g = e + i\alpha^a t^a \in G$. Acting on a generator $t^b \in T_e G$ via the adjoint map Ad_g , we expand to the linear order of α :

$$\text{Ad}_{e+i\alpha^a t^a}(t^b) = (e + i\alpha^a t^a)t^b(e - i\alpha^a t^a) \simeq t^b + i\alpha^a [t^a, t^b] + \mathcal{O}(\alpha^2) \quad (3.69)$$

This motivates us to define the infinitesimal adjoint map $\text{ad}_{t^a} \equiv d(\text{Ad})_e(t^a)$ on the Lie algebra, which acts purely as a commutator:

$$\text{ad}_{t^a}(t^b) = \lim_{\alpha^a \rightarrow 0} \frac{(\text{Ad}_{e+i\alpha^a t^a}(t^b))_{kl} - (t^b)_{kl}}{i\alpha^a} = [t^a, t^b]_{kl} = i \sum_{c=1}^d f^{abc}(t^c)_{kl}. \quad (3.70)$$

Viewing ad_{t^a} as a linear transformation matrix acting on the basis vector t^b to produce t^c , the expansion coefficients (structure constants) directly define the matrix elements of the adjoint generators:

$$(t_{\text{adj}}^a)_{bc} = i f^{abc} \quad (3.71)$$

Hence, the structure constants f^{abc} are not just algebraic constants, they are precisely **used to construct the adjoint representation**.

In this representation, the group elements act on the Lie algebra itself. The generators are $d \times d$ matrices (where $d = N^2 - 1$), with matrix elements defined by (3.71). We can verify that these matrices satisfy the Lie algebra by using the Jacobi identity:

$$\sum_{A=1}^{d_r} (f^{aAe} f^{bcA} + f^{bAe} f^{caA} + f^{cAe} f^{abA}) = 0. \quad (3.72)$$

Writing the commutation relation

$$[t_{\text{adj}}^a, t_{\text{adj}}^b]_{de} = i \sum_{c=1}^d f^{abc} (t_{\text{adj}}^c)_{de}, \quad (3.73)$$

in terms of structure constants gives:

$$\sum_{A=1}^d [(i f^{aAd})(i f^{bAe}) - (i f^{bAd})(i f^{aAe})] = i \sum_{c=1}^d f^{abc} (i f^{cde}). \quad (3.74)$$

Thus we have:

$$\sum_{A=1}^d (-f^{aAd} f^{bAe} + f^{bAd} f^{aAe}) = - \sum_{A=1}^d f^{abA} f^{Ade}. \quad (3.75)$$

Using the anti-symmetry property of structure constant: $f^{abc} = -f^{bac}$, we can reduce to the standard Jacobi identity form:

$$0 = \sum_{A=1}^d (-f^{aAd} f^{bAe} + f^{bAd} f^{aAe} + f^{Ade} f^{abA}) \quad (3.76)$$

$$= \sum_{A=1}^d (f^{aAd} f^{beA} + f^{bAd} f^{eaA} + f^{eAd} f^{abA}), \quad (3.77)$$

which is nothing but the Jacobi identity (3.72).

1.5 Inner product of generators

Now since the Lie algebra generators are all matrices, so it's natural to consider the inner product of two matrices. The most natural way is by the trace of matrix product. So we define a matrix D_r^{ab} for generators t_r^a as

$$D_r^{ab} = \text{tr}(t_r^a t_r^b). \quad (3.78)$$

For $a, b = 1, \dots, d = N^2 - 1$. Cyclicity of trace doesn't change the matrix, so it's symmetry and real-valued:

$$D_r^{ab} = D_r^{ba}, \quad (3.79)$$

and also Hermitian:

$$(D_r^{ab})^\dagger = \text{tr}((t_r^a)^\dagger (t_r^b)^\dagger) = D_r^{ab}. \quad (3.80)$$

As we know from linear algebra, for any real-valued symmetry matrix, one can always diagonalize it by a orthogonal transformation. Thus, diagonalization of D_r^{ab} gives

$$\text{tr}(t_r^a t_r^b) = C(r) \delta^{ab}. \quad (3.81)$$

Since the generators are diagonal, then the position of each components in the same t_r are equal, which imply that $C(r)$ is the same for any component of t_r . Thus $C(r)$ is called the **index of representation** or **Dynkin index**.

Example: $SU(N)$

The generators for $SU(2)$ is $t_{\text{fund}}^a = \frac{\sigma^a}{2}$ ($a = 1, 2, 3$), where σ^a is the Pauli matrix:

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (3.82)$$

For the fundamental representation matrix, we can check that

$$\text{tr}(t_r^a t_r^a) = \frac{1}{2}. \quad (3.83)$$

So

$$C(r) = \frac{1}{2}. \quad (3.84)$$

More over, for general case $SU(N)$, $C(r) = \frac{1}{2}$ due to the normalization of N -dim matrices are all similar with $SU(2)$. Using the index of representation, we can find the structure constants of $SU(N)$ satisfy:

$$\begin{aligned} \text{tr}([t^a, t^b] t_r^c) &= i f^{abc} \text{tr}(t_r^c t_r^c) \\ &= \text{tr}(t_r^a t_r^b t_r^c - t_r^b t_r^a t_r^c) = i f^{abc} C(r) \\ &= \text{tr}(t_r^a [t_r^b, t_r^c]) = i f^{abc} C(r). \end{aligned} \quad (3.85)$$

This shows that f^{abc} are totally anti-symmetric:

$$f^{abc} = -f^{bac} = f^{bca}. \quad (3.86)$$

As we all learned from quantum mechanics: $[J^2, J_i] = 0$. So similarly, one can define an operator T^2 that satisfies the similar commutation relation. This operator is called the **quadratic Casimir operator**, which is defined as:

$$T_{ij}^2 = \sum_{a=1}^{d(G)} \sum_{k=1}^{d(r)} t_{ik}^a t_{kj}^a, \quad (3.87)$$

where t_{ij}^a is fundamental representation. We can check that for any a , we all have: $[t^a, T^2] = 0$, namely T^2 commutes with all generators:

$$[t^a, T^2] = \left[t^a, \sum_b t^b t^b \right] = \sum_b ([t^a, t^b] t^b + t^b [t^a, t^b]) \quad (3.88)$$

$$= \sum_b \left(\left(i \sum_c f^{abc} t^c \right) t^b + t^b \left(i \sum_c f^{abc} t^c \right) \right) \quad (3.89)$$

$$= i \sum_{b,c} f^{abc} (t^c t^b + t^b t^c) \quad (3.90)$$

$$= i \sum_{c,b} f^{acb} (t^b t^c + t^c t^b) \quad (3.91)$$

$$= i \sum_{b,c} (-f^{abc}) (t^c t^b + t^b t^c) \quad (3.92)$$

$$= -i \sum_{b,c} f^{abc} (t^c t^b + t^b t^c) \quad (3.93)$$

$$= 0 \quad (3.94)$$

This implies that T^2 must be proportional to identity:

$$T_{ij}^2 = C_2(r) \delta_{ij} \quad (3.95)$$

And $C_2(r)$ is called the **quadratic Casimir coefficient**. It can be obtained by taking the trace in both sides:

$$\text{tr}(T^2) = \text{tr} \left(\sum_{a=1}^{d(G)} t^a t^a \right) = \sum_{a=1}^{d(G)} \text{tr}(t^a t^a) = C_2(r) \text{tr} \delta_{ij}, \quad (3.96)$$

thus we have:

$$C(r) d(G) = C_2(r) d(r) \quad (3.97)$$

1.6 Representation of Lorentz group

The Lorentz group, as we mentioned before, is $SO(1, 3)$ with metric $\eta = (1, -1, -1, -1)$. The group element Λ with infinitesimal expansion can be written as

$$\Lambda_\nu{}^\mu = \delta_\nu{}^\mu + i \sum_a \omega^a (t^a)_\nu{}^\mu + \mathcal{O}(\omega^2). \quad (3.98)$$

Defining relation of $SO(N, M)$ groups says

$$\Lambda^T \eta \Lambda = \eta, \quad (3.99)$$

thus the infinitesimal form implies

$$(\mathbb{1} + i\omega^a (t^a)^T) \cdot \eta \cdot (\mathbb{1} + i\omega^a t^a) = \eta, \quad (3.100)$$

after dropping the $\mathcal{O}(\omega^2)$, one obtains the anti-symmetric relation:

$$(t^a)^T \eta = -\eta t^a. \quad (3.101)$$

As we mentioned in the last subsection, $SO(1, 3)$, as a subgroup of $O(1, 3)$, has $\frac{1}{2} \times 4 \times (4-1) = 6$ generators, physically are 3 rotation + 3 boosts, which usually noted as $J^{\mu\nu}$. To limit the number of free parameters in $J^{\mu\nu}$, we require

$$J^{\rho\sigma} = -J^{\sigma\rho}. \quad (3.102)$$

So we change the note as

$$\sum_a \omega^a t^a \rightarrow \sum_{\rho\sigma} \omega_{\rho\sigma} J^{\rho\sigma}, \quad (3.103)$$

where $J^{\rho\sigma}$ fulfill $SO(1, 3)$ algebra. Generally, all $SO(N, M)$ groups have also spinor representations generators

$$S^{\mu\nu} = \frac{i}{4}[\gamma^\mu, \gamma^\nu], \quad (3.104)$$

fulfill $SO(1, 3)$ algebra:

$$[J^{\mu\nu}, J^{\rho\sigma}] = i(\eta^{\nu\rho} J^{\mu\sigma} - \eta^{\mu\rho} J^{\nu\sigma} - \eta^{\nu\sigma} J^{\mu\rho} + \eta^{\mu\sigma} J^{\nu\rho}). \quad (3.105)$$

1.7 Integration over gauge group

Since a Lie group is also a manifold, so we can introduce the integration structure based on the group element g and function $f(g)$, the measurement dg is called Haar measure, which include the left and right invariant measure:

$$\text{left invariant measure : } \int (dg)_L f(g) = \int (dg)_L f(g'g); \quad (3.106)$$

$$\text{right invariant measure : } \int (dg)_R f(g) = \int (dg)_R f(gg'), \quad (3.107)$$

where the normalization is $\int dg \, 1 = 1$, and for most of situation the left and right Haar measure are the same $dg = (dg)_L = (dg)_R$. This measurement is just like the normal integral is invariant under the transformation $x \rightarrow x + C$, where C is a constant:

$$\int dx f(x) = \int dx f(x + C). \quad (3.108)$$

To obtain the computable form of the measure, we need explicit realization of dg with metric tensor M_{ij} :

$$M_{ij} \equiv \text{tr}(g^{-1} \partial_i g \cdot g^{-1} \partial_j g), \quad (3.109)$$

where $\partial_i g = \frac{\partial}{\partial \alpha_i} g(\alpha)$. Then we can connect to the normal integral:

$$\int dg f(g) = N \int d\alpha |\det M|^{1/2} f(g(\alpha)), \quad (3.110)$$

in which the normalization is the Jacobi determinant.

Example

- $U(1) = \{e^{i\theta} \mid -\pi < \theta < \pi\}$: For $g(\theta) = e^{i\theta}$ we have

$$M_{\theta\theta} = \text{tr}(g^{-1} \partial_\theta g \cdot g^{-1} \partial_\theta g) = \text{tr}(i \cdot i) = -1. \quad (3.111)$$

And using the fact

$$1 = N \int_{-\pi}^{\pi} d\theta \cdot 1 = N \cdot 2\pi, \quad (3.112)$$

thus $N = \frac{1}{2\pi}$. So the integral is

$$\int dg f(g) = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\theta f(e^{i\theta}) \quad (3.113)$$

- $SU(2) = \{a_0 + i\vec{\sigma} \cdot \vec{a} \mid a_0^2 + \vec{a}^2 = 1\}$: For the element $g(a)$ we have:

$$\partial_0 g = \frac{\partial g}{\partial a_0} = \mathbb{1}_{2 \times 2} \quad (3.114)$$

$$\partial_k g = \frac{\partial g}{\partial a_k} = i\sigma_k \quad (k = 1, 2, 3) \quad (3.115)$$

Thus, the metric is

$$M = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}, \quad (3.116)$$

therefore, $|\det M|^{1/2} = \sqrt{16} = 4$. The last piece is the normalization factor, for that we need to compute the integral for unity. Since the constrain require a sphere, we then can express the unity by delta function:

$$1 = N \int d^4 a \delta(a^2 - 1) = N \int_0^\infty r^3 dr \int d\Omega_3 \delta(r^2 - 1), \quad (3.117)$$

thus $N = \frac{1}{\pi^2}$. And finally:

$$\int dg f(g) = \frac{1}{\pi^2} \int d^4 a \delta(a^2 - 1) f(g(a)). \quad (3.118)$$

- More general case, for $SU(N)$, Schur's orthogonality relation requires

$$\int dg g_{ij}(g^{-1})_{kl} = \frac{1}{N} \delta_{il} \delta_{jk}. \quad (3.119)$$

2. Quantization of non-Abelian gauge theory

2.1 Preparation

To quantize the non-Abelian gauge theory, we need to list what do we have now, everything begins with the Lagrangian:

$$\mathcal{L} = \sum_{j=1}^{d(r)} (i\bar{\psi}_j \gamma^\mu (D_\mu)_{jk} \psi_k - m\bar{\psi}_j \psi_j) - \kappa \text{tr}(F_{\mu\nu} F^{\mu\nu}), \quad (3.120)$$

where the covariant derivative is defined as:

$$(D_\mu)_{jk} = \partial_\mu \delta_{jk} - ig(A_\mu)_{jk}, \quad (3.121)$$

and the field strength is:

$$(F_{\mu\nu})_{jk} = [D_\mu, D_\nu]_{jk}. \quad (3.122)$$

Under the gauge transformation, the gauge fields obey the rule:

$$A_\mu \rightarrow A'_\mu = U A_\mu U^\dagger - \frac{i}{g} (\partial_\mu U) U^\dagger. \quad (3.123)$$

To work in the perturbation theory, we need to explicit show the colour indices. Thus we expand A_μ in basis of adjoint representation:

$$(A_\mu)_{jk} = \sum_{a=1}^{d(G)} A_\mu^a t_{jk}^a \quad (3.124)$$

The infinitesimal transformation with $U_{jk} = \mathbb{1}_{jk} + i \sum_a \alpha^a (t^a)_{jk}$ then becomes:

$$\begin{aligned} (\delta A_\mu)_{jk} &= (A'_\mu)_{jk} - (A_\mu)_{jk} \\ &= -(A_\mu)_{jk} + \left(\mathbb{1} + i \sum_a \alpha^a (t^a) \right)_{jm} (A_\mu)_{mn} \left(\mathbb{1} - i \sum_b \alpha^b (t^b) \right)_{nk} + \frac{1}{g} \sum_a (\partial_\mu \alpha^a) (t^a)_{jm} \left(\mathbb{1} - i \sum_b \alpha^b (t^b) \right)_{mk} \\ &\simeq i \sum_{a=1}^{d(G)} \alpha^a A_\mu^b [t^a, t^b]_{jk} + \frac{1}{g} \sum_{a=1}^{d(G)} (\partial_\mu \alpha^a) t_{jk}^a + \mathcal{O}(\alpha^2) \\ &= \frac{1}{g} \left[-g \sum_{b,c=1}^{d(G)} f^{abc} A_\mu^c \alpha^b + \partial_\mu \alpha^a \right] t_{jk}^a, \end{aligned} \quad (3.125)$$

thus we have

$$\delta A_\mu^a = \frac{1}{g} (D_\mu \alpha)^a = \frac{1}{g} \sum_{b=1}^{d(G)} (D_\mu)^{ab} \alpha^b. \quad (3.126)$$

where $(D_\mu)^{ab}$ is the covariant derivative in adjoint representation:

$$(D_\mu)^{ab} = \partial_\mu \delta^{ab} - g \sum_{c=1}^{d(G)} f^{abc} A_\mu^c. \quad (3.127)$$

2.2 Quantization by path integral

Now we are ready to start quantization via path integral method. We have already seen that the action $S[A]$ is gauge invariant. But

$$\int \mathcal{D}A e^{-S[A]} \quad (3.128)$$

is not well defined, because $\mathcal{D}A$ counts infinite gauge redundant. And if one integrate two times over the pure Yang-Mills term $F_{\mu\nu}^a F^{\mu\nu a}$, one can see that the momentum $M_{\mu\nu}$ satisfies

$$M_{\mu\nu} = -k^2 \eta_{\mu\nu} + k_\mu k_\nu, \quad (3.129)$$

which is degenerated (or not invertible):

$$M_{\mu\nu} k^\nu = -k^2 k_\mu + k_\mu k^2 = 0, \quad (3.130)$$

which means the determinate is 0. So we cannot define the propagator due to it's nothing but the inverse of the momentum operator, which we proved in chapter 1.

The idea to solve these problem comes from Faddeev and Popov. To divide out redundant gauge degrees of freedom, we require a gauge condition

$$G^a(A^u) = 0, \quad (3.131)$$

for a particular point A^u in an orbit of the gauge group, where u is a group element of the non-Abelian gauge group (e.g. $SU(3)$). Therefore, A^u is a gauge choice for A , in the other words, A is gauge-less. So like the way we having $\int \delta(x-a)dx = 1$, we insert the gauge condition into the delta function and define a similar quantity called **Faddeev-Popov(FP) determinant** $\Delta_G[A]$:

$$\Delta_G^{-1}[A] \equiv \int \mathcal{D}u \delta(G^a(A^u)). \quad (3.132)$$

So by this definition, one can get a useful identity:

$$1 = \Delta_G[A] \int \mathcal{D}u \delta(G^a(A^u)). \quad (3.133)$$

One can check that this functional determinant is gauge invariant by continuously doing two times gauge transformation: u and u' . So the fields change to $(A^u)^{u'} = A^{u'u}$. So by definition, the new determinant is given by

$$\Delta_G^{-1}[A^{u'}] = \int \mathcal{D}u \delta(G^a((A^{u'})^u)) = \int \mathcal{D}u \delta(G^a(A^{u'u})), \quad (3.134)$$

Now we introduce a new group element $u'' = u'u$, in which transformation the measure keeps invariant: $\mathcal{D}u = \mathcal{D}u''$, thus

$$\Delta_G^{-1}[A^{u'}] = \int \mathcal{D}u'' \delta(G^a(A^{u''})). \quad (3.135)$$

Changing the integral variable name from u'' to u , we then verify that the FP determinant is indeed invariant under gauge transformation:

$$\Delta_G^{-1}[A^{u'}] = \int \mathcal{D}u \delta(G^a A^u) = \Delta_G^{-1}[A] \quad (3.136)$$

Now we insert (3.133) into path integral

$$\int \mathcal{D}A (1) e^{-S[A]} = \int \mathcal{D}A \Delta_G[A] \int \mathcal{D}u \delta(G^a(A^u)) e^{-S[A]}. \quad (3.137)$$

Since only the delta function in the middle integral is u dependence, and the measure $\mathcal{D}$, the determinant $\Delta_G[A]$ and the action $S[A]$ are all gauge invariant, so we can change the variable (or choose this specific gauge) $A \rightarrow A' = A^u$ without causing any difference:

$$\int \mathcal{D}A \Delta_G[A] \int \mathcal{D}u \delta(G^a(A^u)) e^{-S[A]} = \int \mathcal{D}A'^{u^{-1}} \Delta_G[A'^{u^{-1}}] \int \mathcal{D}u \delta(G^a(A')) e^{-S[A'^{u^{-1}}]} \quad (3.138)$$

$$= \int \mathcal{D}A' \Delta_G[A'] \int \mathcal{D}u \delta(G^a(A')) e^{-S[A']}. \quad (3.139)$$

The trick that we are playing now can be understand with a simple comparison in calculus. For an integral in one dimsion

$$I = \int_{-\infty}^{+\infty} dx \int_0^{2\pi} d\theta f(x+\theta)g(x+\theta), \quad (3.140)$$

which obviously is θ dependent. But we can change the variable $y \equiv x + \theta$. Since θ in the second integral is a constant, thus $dy = dx$ with the same integration range. So the integral changes to

$$I = \int_{-\infty}^{+\infty} dy \int_0^{2\pi} d\theta f(y)g(y), \quad (3.141)$$

where the integrand is now θ independent. With the same trick, now the functional integral is u independent, so we can change the integration sequence and rename the variable from A' to A as

$$\left(\int \mathcal{D}u \right) \cdot \int \mathcal{D}A \Delta_G[A] \delta(G^a(A)) e^{-S[A]}, \quad (3.142)$$

and since the integral over $\mathcal{D}u$ represent the “volume” of the gauge group, which is infinite. So we can just consider the rest part, which is physical and removed all the gauge redundancy as the correct definition of path integral for non-Abelian gauge theories:

$$\int \mathcal{D}A \Delta_G[A] \delta(G^a(A)) e^{-S[A]}. \quad (3.143)$$

So far so good, it seems that we successfully eliminated the gauge redundancy. However, if we consider a specific example, say Coulomb gauge $G^a(A) = \nabla \cdot A^a = 0$, and consider the derivative of G with respect to α , so we have

$$\frac{\delta G_\mu^a}{\delta \alpha^b} = \nabla \cdot \frac{\delta A_\mu^a}{\delta \alpha^b} = \frac{1}{g} \nabla \cdot D_\mu^{ab}, \quad (3.144)$$

where δA_μ^a is given by (3.126), and (3.127) gives

$$\vec{\nabla} \cdot \vec{D}^{ab} = \nabla^2 \delta^{ab} - g f^{abc} \vec{A}^c \cdot \vec{\nabla}, \quad (3.145)$$

which is the Coulomb gauge $\vec{\nabla} \cdot \vec{D}^{ab}$ operator. The problem is it can have non-trivial solution for A_i , namely,

$$(\vec{\nabla} \cdot \vec{D}^{ab}) \alpha^b = 0, \quad (3.146)$$

where $\alpha^b(x) \neq 0$. It means we find a new field by the gauge transformation $\alpha^b(x)$ as $A' = A + \delta A$, with the new gauge condition:

$$G^a(A') = G^a(A) + \delta G^a = 0 + \frac{1}{g} (\vec{\nabla} \cdot \vec{D}^{ab}) \alpha^b = 0, \quad (3.147)$$

which also satisfies the Coulomb gauge $\vec{\nabla} \cdot \vec{A}'^a = 0$. This problem is called the **Gribov Ambiguity**, which only troubles in non-Abelian gauge theory, e.g. QCD, due to the Abelian gauge theory have the boundary $\alpha \rightarrow 0$ at infinity and zero structure constant $f^{abc} = 0$. But for non-Abelian theory, with larger enough A in low energy confinement case, it's possible for a non-zero α satisfying

$$\nabla^2 \alpha^a = g f^{abc} \vec{A}^c \cdot \vec{\nabla} \alpha^b. \quad (3.148)$$

But at least we solved the divergent problem due to infinite gauge choice with the price of introducing a delta function and an FP determinant. These two, however also make trouble because we cannot expand perturbation series directly. The latter is limited by the delta function that only valid around $G = 0$. Therefore, to compute FP determinant, we can do the infinitesimal transformation around $G = 0$. We first consider $G(A^u)$ as function of group element $u(\alpha)$, then change the integral measure in (3.132) to: $\mathcal{D}u = \mathcal{D}G \det \left| \frac{\delta \alpha}{\delta G} \right|$, so we have:

$$\Delta_G^{-1}[A] = \int \mathcal{D}G \det \left| \frac{\delta \alpha}{\delta G} \right| \delta(G). \quad (3.149)$$

Integration over gauge group G gives the FP determinant, then we get:

$$\Delta_G[A] = \det \left| \frac{\delta G}{\delta \alpha} \right|_{G=0}. \quad (3.150)$$

So (3.143) becomes:

$$\int \mathcal{D}A \delta(G(A)) \det \left| \frac{\delta G}{\delta \alpha} \right| e^{-S[A]}. \quad (3.151)$$

Next step we are going to put $\delta(G)$ and FP determinant into the exponential to get a pure path integral form. For that, we consider the Gauss integral

$$\int_{-\infty}^{+\infty} dc e^{-\frac{c^2}{2\xi}} \delta(G - c) = e^{-\frac{G^2}{2\xi}}. \quad (3.152)$$

Generating into the functional space, with $a = 1, \dots, d(G)$, and change the gauge $G^a = 0$ to $G^a = C^a$ with constants C^a . We have:

$$\int \mathcal{D}C^a e^{-\frac{1}{2\xi} \int d^4x C^a C^a} \delta(G^a - C^a) = e^{-\frac{1}{2\xi} \int d^4x G^a G^a}, \quad (3.153)$$

where we successfully put the delta function into the exponential. The price is adding a **gauge fixing term** in Lagrangian:

$$S_{\text{gf}} = \int d^4x \frac{1}{2\xi} G^a \cdot G^a. \quad (3.154)$$

Now we left a determinant needs to be eliminated : $\det \left| \frac{\delta G}{\delta \alpha} \right|$. This can be done by introducing the Grassmann number in the path integral:

$$\det \left| \frac{\delta G^a(\alpha(x))}{\delta \alpha^b(y)} \right| = \int \mathcal{D}\bar{\eta} \mathcal{D}\eta e^{+ \int d^4x d^4y \bar{\eta}^a(x) \frac{\delta G^a(x)}{\delta \alpha^b(y)} \eta^b(y)}, \quad (3.155)$$

where these two new field η and $\bar{\eta}$ is the **Faddeev-Popov Ghost fields**:

$$S_{\text{ghost}} = - \int d^4x d^4y \bar{\eta}^a(x) \frac{\delta G^a(x)}{\delta \alpha^b(y)} \eta^b(y). \quad (3.156)$$

Now we are ready to write the final form of Faddeev-Popov quantization:

$$\int \mathcal{D}A \mathcal{D}\bar{\eta} \mathcal{D}\eta e^{-S[A, \bar{\eta}, \eta]}, \quad (3.157)$$

by inserting (3.154) and (3.155) into the Lagrangian:

$$\begin{aligned} S[A, \bar{\eta}, \eta] &= S_{\text{YM}}[A] + S_{\text{gf}}[A] + S_{\text{ghost}}[A, \bar{\eta}, \eta] \\ &= \int d^4x \left[\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} - \bar{\psi}(\mathbf{i}\not{D} - m)\psi \right] + \frac{1}{2\xi} \int d^4x G^a(x) G^a(x) - \int d^4x d^4y \bar{\eta}^a(x) \frac{\delta G^a(x)}{\delta \alpha^b(y)} \eta^b(y). \end{aligned} \quad (3.158)$$

$$(3.159)$$

It is worthy to note that we used **Euclidean convention**, which is

$$\int \mathcal{D}A e^{-S}. \quad (3.160)$$

If we use **Minkowski convention** with the sign $(+ - - -)$, which is

$$\int \mathcal{D}A e^{iS}. \quad (3.161)$$

So the action becomes

$$S_{\text{M}}[A, \bar{\eta}, \eta] = \int d^4x \left[\bar{\psi}(\mathbf{i}\not{D} - m)\psi - \frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} \right] - \frac{1}{2\xi} \int d^4x G^a(x) G^a(x) + \int d^4x d^4y \bar{\eta}^a(x) \frac{\delta G^a(x)}{\delta \alpha^b(y)} \eta^b(y). \quad (3.162)$$

2.3 Feynman rules for non-Abelian gauge theory

The gauge condition we choose is the **covariant gauge**(**Lorenz gauge**)

$$G^a = \partial^\mu A_\mu^a = 0. \quad (3.163)$$

The quadratic terms in A_μ from $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c$ and $G^a = \partial^\mu A_\mu^a$ can be rewritten by the partial integration as:

$$S_{\text{quadratic}} = \int d^4x \left(\frac{1}{4} (\partial^\mu A_\nu^a - \partial^\nu A_\mu^a) (\partial_\mu A_\nu^a - \partial_\nu A_\mu^a) + \frac{1}{2\xi} (\partial^\mu A_\mu^a) (\partial^\nu A_\nu^a) \right) \quad (3.164)$$

$$= \int d^4x \left(\frac{1}{2} \partial^\mu A^{\nu a} \partial_\mu A_\nu^a - \frac{1}{2} \partial^\nu A^{\mu a} \partial_\mu A_\nu^a + \frac{1}{2\xi} (\partial^\mu A_\mu^a) (\partial^\nu A_\nu^a) \right) \quad (3.165)$$

$$= \frac{1}{2} \int d^4x A_\mu^a \left(-\partial_\rho \partial^\rho g_{\mu\nu} + \left(1 - \frac{1}{\xi} \right) \partial_\mu \partial_\nu \right) A^{\nu a} + \text{Boundary terms}. \quad (3.166)$$

Now with this gauge fixing term, we can check that the momentum operator becomes:

$$M_{\mu\nu} = -k^2 g_{\mu\nu} + \left(1 - \frac{1}{\xi} \right) k_\mu k_\nu, \quad (3.167)$$

satisfying:

$$M_{\mu\nu} k^\nu = -k^2 k_\mu + \left(1 - \frac{1}{\xi} \right) k_\mu k^2 = -\frac{1}{\xi} k^2 k_\mu \neq 0. \quad (3.168)$$

Thus, we can then define the **propagators** legally by take inverse of momentum operator, which can be read from (3.166) as


$$a, \mu \xrightarrow{p} b, \nu = \frac{-i\delta^{ab}}{p^2 + i\varepsilon} \left(g_{\mu\nu} - (1 - \xi) \frac{p_\mu p_\nu}{p^2} \right). \quad (3.169)$$

And the **gauge boson self-interaction** is vertices is given by the non-Abelian term in $\frac{1}{4}F^2$:

$$\mathcal{L}_{3A} = -gf^{abc}(\partial_\mu A_\nu^a)A^{\mu b}A^{\nu c} \quad (3.170)$$


$$\mathcal{L}_{4A} = \frac{1}{4}g^2 f^{abc} f^{ade} A_\mu^b A_\nu^c A^{d\mu} A^{e\nu}. \quad (3.171)$$

The first three gauge boson self interaction gives:



$$= -igf^{abc} [(r_\mu - q_\mu)g_{\nu\rho} + (q_\rho - p_\rho)g_{\mu\nu} + (p_\nu - r_\nu)g_{\mu\rho}] \quad (3.172)$$

The second four gauge boson self interaction gives:



$$= -ig^2 [f^{abe} f^{cde} (g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho}) + f^{ace} f^{bde} (g_{\mu\nu} g_{\rho\sigma} - g_{\mu\sigma} g_{\nu\rho}) + f^{ade} f^{bce} (g_{\mu\nu} g_{\rho\sigma} - g_{\mu\rho} g_{\nu\sigma})]. \quad (3.173)$$

Since we have introduced a new field: ghost field $\bar{\eta}, \eta$, we then can also find its Feynman rule from the Lagrangian. For this, we need to first calculate the derivative by (3.126) and the gauge $G^a = \partial^\mu A_\mu^a$:

$$\frac{\delta G^a(x)}{\delta \alpha^b(y)} = \frac{\delta G^a(x)}{\delta A_\mu^c(x)} \frac{\delta A_\mu^c(x)}{\delta \alpha^b(y)} = \frac{1}{g} \frac{\delta G^a(x)}{\delta A_\mu^c(x)} D_\mu^{bc} \delta(x-y) = \frac{1}{g} \partial^\mu D_\mu^{bc} \delta(x-y), \quad (3.174)$$

with the form of covariant derivative, we have:

$$\frac{\delta G^a(x)}{\delta \alpha^b(y)} = \frac{1}{g} \partial^\mu (\partial_\mu \delta^{ab} - g f^{abc} A_\mu^c) \delta(x-y). \quad (3.175)$$

Thus the ghost action becomes:

$$S_{\text{ghost}} = \int d^4x \bar{\eta}^a(x) \left[\frac{1}{g} \partial^\mu (\partial_\mu \delta^{ab} - g f^{abc} A_\mu^c) \right] \eta^b(x) \quad (3.176)$$


$$= \frac{1}{g} \int d^4x \bar{\eta}^a \partial_\mu \partial^\mu \eta^a - \int d^4x f^{abc} \bar{\eta}^a \partial^\mu (A_\mu^c \eta^b) \quad (3.177)$$

$$= \int d^4x \bar{\eta}^a \partial_\mu \partial^\mu \eta^a - g \int d^4x f^{abc} \bar{\eta}^a \partial^\mu (A_\mu^c \eta^b) \quad (3.178)$$

where we rescaled the field as $\eta \rightarrow \sqrt{g} \eta$ and $\bar{\eta} \rightarrow \sqrt{g} \bar{\eta}$ in the last step. Thus the first term is kinetic term and second is interaction term. So the kinetic term gives the **ghost propagators**:

$$a \xrightarrow{p} b = i \frac{\delta^{ab}}{p^2}. \quad (3.179)$$

And second interaction term gives the **vertex of ghost-ghost-gluon**:



$$= \frac{g}{2} f^{abc} (r^\mu + p^\mu - q^\mu) = -g f^{abc} q^\mu, \quad (3.180)$$

where the vertex energy-momentum conservation requires $r^\mu + p^\mu + q^\mu = 0$.

3. BRST Symmetry

3.1 Problem of negative norm

In last section, we introduced the gauge fixing term in the action to absorbing the gauge condition $G^a = 0$:

$$S_{\text{gf}} = \int d^4x \frac{1}{2\xi} G^a \cdot G^a, \quad (3.181)$$

which leads to the dynamical term of the gauge field as

$$\frac{1}{2\xi} (\partial^\mu A_\mu^a) (\partial^\nu A_\nu^a). \quad (3.182)$$

However, this operation breaks the local gauge symmetry because it contains $\partial_0 A_0$. The problem of this term is A_0 has a negative norm:

$$\langle 0 | A_\mu(k) A_\nu^\dagger(k') | 0 \rangle \propto -\eta_{\mu\nu} \delta^{(4)}(k - k'), \quad (3.183)$$

thus

$$\langle 0 | A_0(k) A_0^\dagger(k') | 0 \rangle \propto -\delta^{(4)}(k - k') < 0. \quad (3.184)$$

This should not be a problem if local gauge symmetry was not broken, because there wouldn't be the term like $\partial_0 A_0$ get into the kinematic term, i.e., propagator. So the negative norm wasn't shown when we are doing real calculations. Now everything is on the table, the optical theorem tell us we need to count all the contributions including the negative norm final states. So it may cause the sum over all physical final states (positive norm states) is greater than 1. And since the gauge choice can be changed, then the value of the probability may no longer conserved or even depends on the gauge choice, which is unacceptable.

3.2 Infinitesimal BRST transformation

To solve this problem, Becchi, Rouet, Stora and Tyutin (BRST) found that there exist a global symmetry, namely the **BRST symmetry**, which allows us to separate the non-physical states.

To introduce this symmetry, we first consider the Gaussian integral formula:

$$\int_{-\infty}^{\infty} dB \exp \left\{ -\frac{a}{2} B^2 \right\} = \sqrt{\frac{2\pi}{a}}. \quad (3.185)$$

Change the variable $B \rightarrow B + \frac{J}{a}$:

$$\int_{-\infty}^{\infty} dB \exp \left\{ -\frac{a}{2} \left(B + \frac{J}{a} \right)^2 \right\} = \int_{-\infty}^{\infty} dB \exp \left\{ -\frac{a}{2} B^2 - JB - \frac{J^2}{2a} \right\} = \text{Constant}. \quad (3.186)$$

Thus, dividing $\exp\left(\frac{J^2}{2a}\right)$ on both sides, we have

$$\exp \left\{ \frac{J^2}{2a} \right\} \propto \int_{-\infty}^{\infty} dB \exp \left\{ -\frac{a}{2} B^2 - JB \right\}. \quad (3.187)$$

Now we can rewrite the gauge fixing term from the Faddeev-Popov Lagrangian by letting $J = \partial^\mu A_\mu^a$ and $a = -\xi$ as

$$\exp \left\{ i \int d^4x \left(-\frac{1}{2\xi} (\partial^\mu A_\mu^a)^2 \right) \right\} \propto \int \mathcal{D}B \exp \left\{ i \int d^4x \left(\frac{\xi}{2} (B^a)^2 + B^a (\partial^\mu A_\mu^a) \right) \right\}, \quad (3.188)$$

where B^a is an auxiliary field, which means this field is **non-dynamical**, because it doesn't contribute any term including $\partial_\mu B^a$. So the gauge fixing Lagrangian becomes:

$$\mathcal{L}_{\text{gf}} = \frac{\xi}{2} (B^a)^2 + B^a (\partial^\mu A_\mu^a). \quad (3.189)$$

We can check this Lagrangian gives the same form with the Gaussian integration by implying the Euler-Lagrange equation:

$$\frac{\delta \mathcal{L}_{\text{gauge fix}}}{\delta B^a} = \xi B^a + \partial^\mu A_\mu^a = 0, \quad (3.190)$$

$$\implies B^a = -\frac{1}{\xi} \partial^\mu A_\mu^a. \quad (3.191)$$

So the Faddeev-Popov Lagrangian is

$$\mathcal{L}_{\text{FP}} = \bar{\psi}(i\not{D} - m)\psi - \frac{1}{4}F_{\mu\nu}^a F^{\mu\nu a} + \frac{\xi}{2}(B^a)^2 + B^a(\partial^\mu A_\mu^a) + \bar{c}^a(-\partial^\mu D_\mu^{ab})c^b. \quad (3.192)$$

We have already derived variation of A from (3.126):

$$\delta A_\mu^a = D_\mu^{ab}\alpha^b(x), \quad (3.193)$$

where we absorbed the factor $1/g$ into the arbitrary function $\alpha^b(x) \rightarrow g\alpha^b(x)$. But we now want $\alpha^b(x)$ can related to the ghost field:

$$\alpha^b(x) = g\varepsilon c^b(x), \quad (3.194)$$

where ε is a constant that non-related with the space-time. So if we also do the same thing for the other fields' infinitesimal transformations, how will the Lagrangian change? This leads us to the **BRST transformations**. BRST claimed that there is a global symmetry from ε that the transformation keep the Lagrangian invariant

$$\delta_\varepsilon \mathcal{L} = 0. \quad (3.195)$$

And the infinitesimal parameter must be Grassmann-valued:

$$\varepsilon^2 = 0, \quad (3.196)$$

because A_μ^a is a bosonic field and c^b is a fermionic field, so ε must be a fermionic field, which fulfills the Grassmann algebra.

$$\varepsilon c^b = -c^b \varepsilon. \quad (3.197)$$

Thus the BRST transformation for A_μ^a and ψ_j can be derived by implying (3.194) into the normal infinitesimal transformation $\delta A_\mu^a = D_\mu^{ab}\alpha^b(x)$ and $\delta_\varepsilon \psi_j = i \sum_a \alpha^a (t^a \psi)_j$:

$$\delta_\varepsilon A_\mu^a = \varepsilon D_\mu^{ab} c^b = \varepsilon \left(\partial_\mu c^a + g \sum_{b,c} f^{abc} A_\mu^b c^c \right), \quad (3.198)$$

$$\delta_\varepsilon \psi_j = i g \varepsilon \sum_a c^a (t^a \psi)_j. \quad (3.199)$$

These two transformations obviously keep the $\mathcal{L}_{\text{YM}}$ invariant, because it just changes a name of α , and since αc^a is still a local gauge transformation, then $\mathcal{L}_{\text{YM}}$ is automatically invariant under this transformation. One can check that $F_{\mu\nu}^a$ follows

$$\delta_\varepsilon F_{\mu\nu}^a = g \varepsilon f^{abc} F_{\mu\nu}^b c^c, \quad (3.200)$$

thus

$$\delta_\varepsilon \left(-\frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a} \right) = 0. \quad (3.201)$$

And for the fermionic field, the covariant derivative follows:

$$\delta_\varepsilon (D_\mu \psi) = i g \varepsilon c^a t^a (\partial_\mu \psi - i g A_\mu^b t^b \psi) = i g \varepsilon c^a t^a (D_\mu \psi), \quad (3.202)$$

thus

$$\delta_\varepsilon (\bar{\psi}(i\gamma^\mu D_\mu - m)\psi) = (\delta_\varepsilon \bar{\psi})(i\gamma^\mu D_\mu - m)\psi + \bar{\psi}(i\gamma^\mu \delta_\varepsilon (D_\mu \psi) - m \delta_\varepsilon \psi) = 0. \quad (3.203)$$

But it's worthy to mention that **BRST transformation is not a local gauge transformation (transformation parameter ε is global)**. But it indeed comes from a local gauge transformation (transformation parameter $\alpha^a(x) = g\varepsilon c^a(x)$ is local).

Now it comes to the most tedious part, the gauge fixing term and the ghost term. This is because these two terms are not gauge invariant. For convenient, we can just check the following variation of the ghost field and auxiliary field:

$$\delta_\varepsilon c^a = -\frac{1}{2}g\varepsilon \sum_{b,c} f^{abc} c^b c^c \quad (3.204)$$

$$\delta_\varepsilon \bar{c}^a = \varepsilon B^a \quad (3.205)$$

$$\delta_\varepsilon B^a = 0, \quad (3.206)$$

keep the Lagrangian of gauge fixing and ghost invariant:

$$\delta_\varepsilon (\mathcal{L}_{\text{gf}} + \mathcal{L}_{\text{ghost}}) = 0, \quad (3.207)$$

namely to prove:

$$\delta_\varepsilon \left(\frac{\xi}{2} (B^a)^2 \right) + \underbrace{(\delta_\varepsilon B^a)}_{=0} (\partial^\mu A_\mu^a) + B^a \underbrace{\delta_\varepsilon (\partial^\mu A_\mu^a)}_{=\partial^\mu (\varepsilon D_\mu^{ab} c^b)} - \underbrace{(\delta_\varepsilon \bar{c}^a)}_{\varepsilon B^a} (\partial^\mu D_\mu^{ab} c^b) + \underbrace{\bar{c}^a \delta_\varepsilon (\partial^\mu D_\mu^{ab} c^b) + \bar{c}^a (\partial^\mu D_\mu^{ab}) (\delta_\varepsilon c^b)}_{\bar{c}^a \partial^\mu \delta_\varepsilon (D_\mu^{ab} c^b)} = 0. \quad (3.208)$$

The first term is simply 0 because

$$\delta_\varepsilon (B^a)^2 = 2\delta_\varepsilon B^a = 0 \quad (3.209)$$

And for the third term

$$\partial^\mu (\varepsilon D_\mu^{ab} c^b) = \underbrace{(\partial^\mu \varepsilon)}_{=0} D_\mu^{ab} c^b + \varepsilon \partial^\mu (D_\mu^{ab} c^b) = \varepsilon \partial^\mu (D_\mu^{ab} c^b), \quad (3.210)$$

due to ε is a constant. Now the expression to be proved reduces to

$$\cancel{B^a \varepsilon \partial^\mu D_\mu^{ab} c^b} - \cancel{\varepsilon B^a \partial^\mu D_\mu^{ab} c^b} - \bar{c}^a \partial^\mu (\delta_\varepsilon D_\mu^{ab} c^b) \quad (3.211)$$

$$= -\bar{c}^a \partial^\mu \left\{ \partial_\mu (\delta_\varepsilon c^a) + g \sum_{b,c} f^{abc} (\delta_\varepsilon A_\mu^b) c^c + g \sum_{b,c} f^{abc} A_\mu^b (\delta_\varepsilon c^c) \right\} \quad (3.212)$$

$$= -\bar{c}^a \partial^\mu \left\{ \partial_\mu \left(-\frac{g}{2} \varepsilon \sum_{b,c} f^{abc} c^b c^c \right) + g \sum_{b,c} f^{abc} \left[\varepsilon \left(\partial_\mu c^b + g \sum_{d,e} f^{bde} A_\mu^d c^e \right) \right] c^c \right. \\ \left. + g \sum_{b,c} f^{abc} A_\mu^b \left(-\frac{g}{2} \varepsilon \sum_{d,e} f^{cde} c^d c^e \right) \right\} \quad (3.213)$$

$$= -\bar{c}^a \partial^\mu \left\{ g\varepsilon \sum_{b,c} f^{abc} \left[-\frac{1}{2} (\partial_\mu c^b c^c) + (\partial_\mu c^b) c^c \right] + g^2 \varepsilon \sum_{b,c,e,f} f^{abc} \left(f^{bde} A_\mu^d c^e c^c - \frac{1}{2} f^{cde} A_\mu^b c^d c^e \right) \right\} \quad (3.214)$$

The first term will be canceled by using the property of Grassmann number $c^a c^b = -c^b c^a$ and structure constant $f^{abc} = -f^{acb}$:

$$\sum_{b,c} f^{abc} \partial_\mu (c^b c^c) = \sum_{b,c} f^{abc} [(\partial_\mu c^b) c^c + c^b (\partial_\mu c^c)] \quad (3.215)$$

$$= \sum_{b,c} [f^{abc} (\partial_\mu c^b) c^c + f^{acb} c^c (\partial_\mu c^b)] \quad (3.216)$$

$$= \sum_{b,c} [f^{abc} (\partial_\mu c^b) c^c - f^{abc} c^c (\partial_\mu c^b)] \quad (3.217)$$

$$= \sum_{b,c} [f^{abc} (\partial_\mu c^b) c^c + f^{abc} (\partial_\mu c^b) c^c] \quad (3.218)$$

$$= 2 \sum_{b,c} f^{abc} (\partial_\mu c^b) c^c. \quad (3.219)$$

Now the only left term is

$$-\bar{c}^a \partial^\mu g^2 \varepsilon \sum_{b,c,e,f} f^{abc} \left(f^{bde} A_\mu^d c^e c^c - \frac{1}{2} f^{cde} A_\mu^b c^d c^e \right), \quad (3.220)$$

which can be canceled by Jacobi identity

$$f^{ade} f^{bcd} + f^{bde} f^{cad} + f^{cde} f^{abd} = 0. \quad (3.221)$$

For that, we first relabel the indices d, e in the first term:

$$\sum_{b,c,d,e} f^{abc} f^{bde} c^e c^c = \frac{1}{2} \sum_{b,c,d,e} (f^{abc} f^{bde} - f^{abe} f^{bdc}) c^e c^c. \quad (3.222)$$

Thus,

$$\sum_{b,c,d,e} f^{abc} \left(f^{bde} A_\mu^d c^e c^c - \frac{1}{2} f^{cde} A_\mu^b c^d c^e \right) \quad (3.223)$$

$$= \sum_{b,c,d,e} f^{abc} f^{bde} A_\mu^d c^e c^c - \frac{1}{2} f^{abc} f^{cde} A_\mu^b c^d c^e \quad (3.224)$$

$$= \frac{1}{2} \sum_{b,c,d,e} (f^{abc} f^{bde} - f^{abe} f^{bdc}) A_\mu^d c^e c^c - f^{abc} f^{cde} A_\mu^b c^d c^e \quad (3.225)$$

$$= \frac{1}{2} \sum_{b,c,d,e} (f^{ace} f^{cbd} - f^{acd} f^{cbe} - f^{abc} f^{cde}) A_\mu^b c^d c^e \quad (3.226)$$

$$= \frac{1}{2} \sum_{b,c,d,e} (-f^{dbc} f^{ace} + f^{ebc} f^{acd} - f^{abc} f^{cde}) A_\mu^b c^d c^e \quad (3.227)$$

$$= \frac{1}{2} \sum_{b,c,d,e} (f^{dbc} f^{aec} - f^{ebc} f^{adc} - f^{abc} f^{dec}) A_\mu^b c^d c^e \quad (3.228)$$

$$= \frac{1}{2} \sum_{b,c,d,e} (f^{dbc} f^{aec} + f^{ebc} f^{dac} + f^{abc} f^{dec}) A_\mu^b c^d c^e \quad (3.229)$$

$$= 0. \quad (3.230)$$

Finally, with such tedious calculation, we have successfully proved (3.207) being satisfied, which means the gauge fixing term and ghost term together are BRST invariant. Thus, the Faddeev-Popov Lagrangian is BRST invariant:

$$\delta_\varepsilon \mathcal{L}_{\text{FP}} = \delta_\varepsilon \mathcal{L}_{\text{YM}} + \delta_\varepsilon (\mathcal{L}_{\text{gf}} + \mathcal{L}_{\text{ghost}}) = 0. \quad (3.231)$$

3.3 Factorization of Hilbert space via BRST operator

Now since BRST symmetry is a global symmetry, we can in principle define a conservation charge Q , which should upgrade to an operator in Hilbert space since we have already quantized the classical field. This operator thus called the **BRST operator** Q , which can be easily defined from the infinitesimal transformation:

$$\delta_\varepsilon \Phi = [\varepsilon Q, \Phi] = \varepsilon(Q\Phi); \quad \text{for bosonic field } \Phi \quad (3.232)$$

$$\delta_\varepsilon \Psi = \{\varepsilon Q, \Psi\} = -\varepsilon(Q\Psi); \quad \text{for fermionic field } \Psi. \quad (3.233)$$

Thus Q has similar properties with δ_ε , which can be directly copy from (3.198),(3.199),(3.204),(3.205) and (3.206):

$$QA_\mu^a = D_\mu^{ab} c^b \quad (3.234)$$

$$Q\psi_j = igc^a(t^a\psi_j) \quad (3.235)$$

$$Qc^a = -\frac{1}{2}gf^{abc}c^b c^c \quad (3.236)$$

$$Q\bar{c}^a = B^a \quad (3.237)$$

$$QB^a = 0, \quad (3.238)$$

and with (3.196), one can prove that Q satisfies **Nilpotency**:

$$Q^2 = 0, \quad (3.239)$$

by acting on the gauge field A_μ^a , the ghost and anti-ghost field c^a and $\bar{c}^a$ and the auxiliary field B^a respectively to show the results are all vanished.

With BRST operator, we can rewrite the Faddeev-Popov Lagrangian as

$$\mathcal{L}_{\text{FP}} = \mathcal{L}_{\text{YM}} + \underbrace{Q \left(\bar{c}^a \partial^\mu A_\mu^a + \frac{\xi}{2} \bar{c}^a B^a \right)}_{\text{BRST ancestors}}. \quad (3.240)$$

With this form, we can easily see that it's BRST invariant by using (3.239) and $\delta_\varepsilon \mathcal{L}_{\text{YM}} = 0$:

$$Q\mathcal{L}_{\text{FP}} = 0, \quad Q\mathcal{L}_{\text{YM}} = 0 \quad (3.241)$$

Now we can use BRST operator to show how those “nonphysical” states are separated. Since BRST operator as a conserved ”charge”. It naturally commutes with Hamiltonian operator H :

$$[Q, H] = 0, \quad Q = Q^\dagger. \quad (3.242)$$

The Hilbert space of eigenstates of H in Heisenberg picture can be formally expressed as

$$\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \mathcal{H}_0, \quad (3.243)$$

where the subspace can be defined by the map:

$$Q : \mathcal{H} \rightarrow \mathcal{H}. \quad (3.244)$$

The kernel of Q , i.e., those states are mapped to 0, is

$$\text{Ker}(Q) = \{|\psi\rangle \in \mathcal{H} \mid Q|\psi\rangle = 0\}, \quad (3.245)$$

and the image is

$$\text{Im}(Q) = \{|\psi\rangle \in \mathcal{H} \mid |\psi\rangle = Q|\phi\rangle, \text{ for } |\phi\rangle \in \mathcal{H}, \}. \quad (3.246)$$

And since $Q^2 = 0$, so for any states in $\text{Im}(Q)$ are all in $\text{Ker}(Q)$ too, i.e.,

$$\text{Im}(Q) \subset \text{Ker}(Q). \quad (3.247)$$

Thus, we define the three states as follows:

$$\mathcal{H}_1 = \mathcal{H} \setminus \text{Ker}(Q) = \{|\psi_1\rangle \in \mathcal{H} \mid Q|\psi_1\rangle \neq 0\}, \quad (3.248)$$

$$\mathcal{H}_2 = \text{Im}(Q) = \{|\psi_2\rangle \in \mathcal{H} \mid |\psi_2\rangle = Q|\phi\rangle, |\phi\rangle \in \mathcal{H}_1\}, \quad (3.249)$$

$$\mathcal{H}_0 = \text{Ker}(Q) \setminus \text{Im}(Q) = \{|\psi_0\rangle \in \mathcal{H} \mid Q|\psi_0\rangle = 0 \text{ and } \forall |\phi\rangle \in \mathcal{H}, |\psi\rangle \neq Q|\phi\rangle\}. \quad (3.250)$$

With these definition, we can clearly see that Hilbert space is factorized into the direct sum over three subspace :

$$\mathcal{H} = \underbrace{(\mathcal{H} \setminus \text{Ker}(Q))}_{\mathcal{H}_1} \oplus \underbrace{\text{Im}(Q)}_{\mathcal{H}_2} \oplus \underbrace{(\text{Ker}(Q) \setminus \text{Im}(Q))}_{\mathcal{H}_0}. \quad (3.251)$$

With this, we claim the set of all physical states is the kernel of $\mathcal{H}$ mode the image:

$$\mathcal{H}_{\text{phys}} \cong \mathcal{H}_0 = \text{coh}(Q) = \frac{\text{Ker}(Q)}{\text{Im}(Q)}, \quad (3.252)$$

which is a **cohomology space** satisfying:

$$Q|\psi\rangle = 0. \quad (3.253)$$

To show that it indeed make sense, we need to go back to the free theory, namely $g \rightarrow 0$, so the BRST transformation reduce to

$$QA_\mu^a = \partial_\mu c^a, \quad (3.254)$$

$$Q\bar{c}^a = B^a, \quad (3.255)$$

$$Q\psi_j = Qc^a = QB^a = 0. \quad (3.256)$$

We find this result with the definition of $\mathcal{H}_1$, $\mathcal{H}_2$ and $\mathcal{H}_0$ can help us to classify these asymptotic states:

$$A_\mu^L \in \mathcal{H}_1, \quad \bar{c}^a \in \mathcal{H}_1; \quad (3.257)$$

$$A_\mu^T \in \mathcal{H}_0, \quad \psi_j \in \mathcal{H}_0; \quad (3.258)$$

$$c^a \in \mathcal{H}_2, \quad B^a \in \mathcal{H}_2. \quad (3.259)$$

where A_μ^L represent the longitudinal polarization

$$A_\mu^L \sim k_\mu \phi \implies Q|A_\mu^L\rangle \sim |c^a\rangle \neq 0, \quad (3.260)$$

and A_μ^T represent the transversal polarization

$$k^\mu A_\mu^T = 0 \implies QA_\mu^T \sim k_\mu c^a = 0. \quad (3.261)$$

With this classification, we can see that for the states in $\mathcal{H}_1$, like A_μ^L or $\bar{c}^a$, the norm of time component are **negative**:

$$\langle A_0^L | A_0^L \rangle = \langle 0 | a_0 a_0^\dagger | 0 \rangle = \langle 0 | [a_0, a_0^\dagger] | 0 \rangle = -1 < 0, \quad (3.262)$$

$$\langle \bar{\eta} | \bar{\eta} \rangle = \langle 0 | \bar{c} \bar{c}^\dagger | 0 \rangle = \langle 0 | (-\bar{c}^\dagger)^\dagger \bar{c} | 0 \rangle = -\langle 0 | \bar{c} \bar{c}^\dagger | 0 \rangle = \langle 0 | \{c, \bar{c}^\dagger\} | 0 \rangle = -1 < 0. \quad (3.263)$$

For the states in $\mathcal{H}_2$ are **zero norm** due to the nilpotency $Q^2 = 0$:

$$\langle \psi_2 | \psi_2 \rangle = \langle \psi_1 | Q^\dagger Q | \psi_1 \rangle = \langle \psi_1 | Q^2 | \psi_1 \rangle \equiv 0. \quad (3.264)$$

And the states in $\mathcal{H}_0$ are **positive norm**, which is the real physical state space. So we say, physical degrees of freedom are elements of $\mathcal{H}_0$, and unphysical degrees of freedom are in $\mathcal{H}_1, \mathcal{H}_2$.

This result however, one may wondering if it is picture dependent since we assume the Heisenburg picture that the states doesn't evolve with time. The answer is lucky no, picture does not change under time evolution: for any state $|\psi_0, t\rangle$ in $\mathcal{H}$, we have

$$Q|\psi_0, t\rangle = Qe^{iHt}|\psi_0\rangle = e^{iHt}Q|\psi_0\rangle = 0, \quad (3.265)$$

which means $|\psi_0, t\rangle$ must be a linear combination of states from $\mathcal{H}_0$ and $\mathcal{H}_2$, i.e., $|\psi_0, t\rangle \in \mathcal{H}_0 \oplus \mathcal{H}_2$. But if we consider a scattering process from $|\psi_0, t\rangle$ to any $|\psi_2\rangle \in \mathcal{H}_2$ that satisfies $|\psi_2\rangle = Q|\psi_1\rangle$, we have

$$\langle \psi_2 | \psi_0, t \rangle = \langle \psi_1 | Q | \psi_0, t \rangle = 0, \quad (3.266)$$

which means the scalar product vanishes! Namely, $|\psi_0, t\rangle$ has no effective component in $\mathcal{H}_2$. So even though we mix some element of $\mathcal{H}_2$ in the combination of the construction of $|\psi_0, t\rangle$, it doesn't matter because the elements in $\mathcal{H}_2$ won't contribute to the final result.

Thus, we find that **BRST operator factorize the space based on positive norm, negative norm and zero norm**. which helps us to select and drop the negative and zero norm states.

4. Renormalization of non-Abelian gauge theory

Now we have enough foundation to consider the renormalization of Padeev–Popov theory. For that, we first recall the Feynman rules of Faddeev–Popov Lagrangian for a Yang–Mills theory coupled to fermions,

$$\mathcal{L}_{\text{FP}} = \mathcal{L}_{\text{YM}} - \frac{1}{2\xi} (\partial^\mu A_\mu^a) (\partial^\nu A_\nu^a) + \bar{c}^a (-\partial^\mu D_\mu^{ab}) c^b. \quad (3.267)$$

It is useful to separate it into the free part and the interaction part:

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{\text{int}}. \quad (3.268)$$

In Feynman gauge, $\xi = 1$, the free Lagrangian reads

$$\mathcal{L}_0 = \bar{\psi} (i\not{\partial} - m) \psi + \frac{1}{2} A_\mu^a (g^{\mu\nu} \square - \partial^\mu \partial^\nu) A_\nu^a. \quad (3.269)$$

The corresponding fermion and gauge boson propagators are


$$\begin{array}{c} p \\ \longrightarrow \end{array} = \frac{i}{\not{p} - m + i\varepsilon} = \frac{i(\not{p} + m)}{p^2 - m^2 + i\varepsilon}, \quad (3.270)$$

$$a, \mu \begin{array}{c} p \\ \text{~~~~~} \longrightarrow \end{array} b, \nu = \frac{-i\delta^{ab}}{p^2 + i\varepsilon} \left[g_{\mu\nu} - (1 - \xi) \frac{p_\mu p_\nu}{p^2 + i\varepsilon} \right]. \quad (3.271)$$


The interaction terms are

$$\begin{aligned}\mathcal{L}_{\text{int}} = & g A_\mu^a (\bar{\psi} \gamma^\mu t^a \psi) - g f^{abc} (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu} \\ & - \frac{1}{4} g^2 (f^{abc} A_\mu^b A_\nu^c) (f^{ade} A^{d\mu} A^{e\nu}) + g f^{abc} (\partial^\mu \bar{c}^a) A_\mu^b c^c.\end{aligned}\quad (3.272)$$


With all momenta conventionally taken incoming, the basic vertices are:



$$= i g \gamma^\mu (t^a)_{ij}, \quad (3.273)$$




$$= g f^{abc} [g^{\mu\nu} (p - q)^\rho + g^{\nu\rho} (q - r)^\mu + g^{\rho\mu} (r - p)^\nu], \quad (3.274)$$




$$\begin{aligned} & = -i g^2 \left[f^{abe} f^{cde} (g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho}) \right. \\ & \quad + f^{ace} f^{bde} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\sigma} g^{\nu\rho}) \\ & \quad \left. + f^{ade} f^{bce} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma}) \right]. \end{aligned} \quad (3.275)$$

For the ghost propagator and ghost–gluon vertex one has



$$a \cdots b = i \frac{\delta^{ab}}{p^2}; \quad (3.276)$$



$$= \frac{g}{2} f^{abc} (r^\mu + p^\mu - q^\mu) = -g f^{abc} q^\mu, \quad (3.277)$$

which are the result that we should be familiar with in previous section.

4.1 Why the Faddeev–Popov theory is renormalizable

Before introducing the explicit counterterms, we should explain why the Faddeev–Popov theory is expected to be renormalizable at all. The point is not to guess the counterterms first, but to prove that the ultraviolet divergent part of the theory can only have the same local form as the original Faddeev–Popov Lagrangian. This statement follows from two ingredients: **power counting** and **BRST symmetry**.

Limitation from power counting

We work in four space-time dimensions. Just as we did in Chapter 2. Since the action is dimensionless,

$$[S] = 0, \quad [d^4x] = -4, \quad [\mathcal{L}] = 4. \quad (3.278)$$

The dimensions of the fields are fixed by the quadratic terms. From the gauge kinetic term,

$$-\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} \supset -\frac{1}{2}(\partial_\mu A_\nu^a)(\partial^\mu A^{a\nu}), \quad (3.279)$$

we get

$$2(1 + [A_\mu]) = 4 \quad \implies \quad [A_\mu] = 1. \quad (3.280)$$

From the fermion kinetic term,

$$\bar{\psi} i\gamma^\mu \partial_\mu \psi, \quad (3.281)$$

we get

$$[\bar{\psi}] + 1 + [\psi] = 4, \quad [\bar{\psi}] = [\psi], \quad \implies \quad [\psi] = [\bar{\psi}] = \frac{3}{2}. \quad (3.282)$$

Finally, from the ghost kinetic term,

$$-\bar{c}^a \partial_\mu \partial^\mu c^a, \quad (3.283)$$

we get

$$[\bar{c}] + 2 + [c] = 4. \quad (3.284)$$

The natural symmetric assignment is

$$[c] = [\bar{c}] = 1. \quad (3.285)$$

Therefore the dimensions of all interaction terms are

$$[A_\mu \bar{\psi} \gamma^\mu \psi] = 1 + \frac{3}{2} + \frac{3}{2} = 4, \quad (3.286)$$

$$[(\partial A)AA] = 1 + 1 + 1 + 1 = 4, \quad (3.287)$$

$$[AAAA] = 4, \quad (3.288)$$

$$[(\partial \bar{c})Ac] = 1 + 1 + 1 + 1 = 4. \quad (3.289)$$

Thus the coupling g is dimensionless:

$$[g] = 0. \quad (3.290)$$

This is the first indication of renormalizability just like (2.281). A negative-dimensional coupling would force higher and higher powers of momentum to appear at higher orders, producing infinitely many independent divergent structures. Here this does not happen.

Now we make the statement precise by computing the superficial degree of divergence. At large loop momentum, the propagators in (3.270), (3.271) and (3.276), behave as

$$S_\psi(k) \sim \frac{\not{k}}{k^2} \sim \frac{1}{k}, \quad D_A(k) \sim \frac{1}{k^2}, \quad D_c(k) \sim \frac{1}{k^2}. \quad (3.291)$$

The three-gluon vertex and the ghost-gluon vertex each contain one derivative, which in momentum space becomes a off-shell momentum that need to be integrated, so they contribute one power of loop momentum in the numerator. The fermion-gluon vertex and the four-gluon vertex contain no derivative.

Let

$$I_A, I_c, I_\psi \quad (3.292)$$

be the numbers of internal gauge-boson, ghost and fermion lines. Let

$$V_3, V_4, V_c, V_\psi \quad (3.293)$$

denote the numbers of AAA , $AAAA$, $A\bar{c}c$ and $A\bar{\psi}\psi$ vertices. Let the corresponding external lines be

$$E_A, E_c, E_\psi, \quad (3.294)$$

where E_c counts external ghost and anti-ghost lines together, and E_ψ counts external fermion and anti-fermion lines together. The superficial degree of divergence in $d = 4$ from (2.272) is then

$$D = 4L - 2I_A - 2I_c - I_\psi + V_3 + V_c. \quad (3.295)$$

The terms in this formula have the following origin:

- each **loop integral** contributes $+4$ (integral measure d^4k);
- each **internal gauge** or **ghost propagator** contributes -2 ($\sim \frac{1}{k^2}$);
- each **internal fermion propagator** contributes -1 ($\sim \frac{k}{k^2}$);
- each **gauge boson vertex** $(\partial A)AA$ or **ghost-boson vertex** $\partial\bar{c}Ac$ vertex contributes $+1$ from its derivative ($i\partial \rightarrow k$).

The topological relation between loops, internal lines and vertices is

$$L = I_A + I_c + I_\psi - (V_3 + V_4 + V_c + V_\psi) + 1. \quad (3.296)$$

We also have the line-counting identities

$$2I_\psi + E_\psi = 2V_\psi, \quad (3.297)$$

$$2I_c + E_c = 2V_c, \quad (3.298)$$

$$2I_A + E_A = 3V_3 + 4V_4 + V_c + V_\psi. \quad (3.299)$$

Substituting the loop relation into D gives

$$\begin{aligned} D &= 4(I_A + I_c + I_\psi - V_3 - V_4 - V_c - V_\psi + 1) - 2I_A - 2I_c - I_\psi + V_3 + V_c \\ &= 4 + 2I_A + 2I_c + 3I_\psi - 3V_3 - 4V_4 - 3V_c - 4V_\psi. \end{aligned} \quad (3.300)$$

Now use the three line-counting identities:

$$2I_A = 3V_3 + 4V_4 + V_c + V_\psi - E_A, \quad (3.301)$$

$$2I_c = 2V_c - E_c, \quad (3.302)$$

$$3I_\psi = 3V_\psi - \frac{3}{2}E_\psi. \quad (3.303)$$

Hence

$$\begin{aligned} D &= 4 + (3V_3 + 4V_4 + V_c + V_\psi - E_A) + (2V_c - E_c) + \left(3V_\psi - \frac{3}{2}E_\psi\right) \\ &\quad - 3V_3 - 4V_4 - 3V_c - 4V_\psi \\ &= 4 - E_A - E_c - \frac{3}{2}E_\psi. \end{aligned} \quad (3.304)$$

This is the crucial power-counting result:

$$\boxed{D = 4 - E_A - E_c - \frac{3}{2}E_\psi.} \quad (3.305)$$

The important point is that D is independent of the number of vertices. Thus **going to higher loop order does not create more and more divergent structures**. Only Green functions with sufficiently few external legs can be superficially divergent. For example,

- Two gauge boson external lines (gauge 1PI):

$$AA : \quad D = 2. \quad (3.306)$$

- Two ghost external lines (ghost 1PI):

$$\bar{c}c : \quad D = 2. \quad (3.307)$$

- Two fermion external lines (fermion 1PI):

$$\bar{\psi}\psi : \quad D = 1. \quad (3.308)$$

- Three gauge external lines:

$$AAA : \quad D = 1. \quad (3.309)$$

- Three gauge-ghost external lines:

$$A\bar{c}c : \quad D = 1. \quad (3.310)$$

- Three gauge-fermion external lines:

$$A\bar{\psi}\psi : \quad D = 0. \quad (3.311)$$

- Four gauge external lines:

$$AAAA : \quad D = 0. \quad (3.312)$$

For amplitudes with more external fields, $D < 0$, which is (superficially) convergent after subdivergences have been subtracted, namely we don't need to consider the renormalization of terms like $AAAAA$. So we only need to consider the divergent in $D \leq 4$

Limitation from BRST symmetry: Slavnov-Taylor identity

However, power counting alone is not yet the full proof. It tells us that the divergent terms are local and have dimension no larger than 4, but it does not by itself explain why these local terms must combine into the gauge-invariant Faddeev–Popov structure. For this we use BRST symmetry. It is most transparent to introduce the auxiliary field B^a and write

$$\mathcal{L}_{\text{FP}} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} + \bar{\psi}(i\not{D} - m)\psi + \frac{\xi}{2}(B^a)^2 + B^a \partial^\mu A_\mu^a + \bar{c}^a (-\partial^\mu D_\mu^{ab})c^b. \quad (3.313)$$

The BRST transformations are

$$QA_\mu^a = D_\mu^{ab}c^b, \quad (3.314)$$

$$Qc^a = -\frac{g}{2}f^{abc}c^b c^c, \quad (3.315)$$

$$Q\bar{c}^a = B^a, \quad (3.316)$$

$$QB^a = 0, \quad (3.317)$$

$$Q\psi = igc^a t^a \psi, \quad (3.318)$$

$$Q\bar{\psi} = -ig\bar{\psi}c^a t^a. \quad (3.319)$$

As shown in the BRST section of the notes, these transformations are nilpotent,

$$Q^2 = 0, \quad (3.320)$$

and the Faddeev–Popov Lagrangian is BRST invariant:

$$Q\mathcal{L}_{\text{FP}} = 0. \quad (3.321)$$

Next, for understanding how the BRST symmetry limits the renormalization counter terms, we need to briefly introduce the Slavnov–Taylor identity. The idea is not mysterious: it is simply the Ward identity associated with BRST symmetry. In the path integral we integrate over all fields as

$$Z[J] = \int \mathcal{D}\Phi \exp \left\{ iS_{\text{FP}}[\Phi] + i \int J\Phi \right\}, \quad (3.322)$$

where Φ denotes all fields, for example $A_\mu, c, \bar{c}, B, \psi, \bar{\psi}$. And S_{FP} is the **classical action**, by contrast the **effective action** can be expressed as:

$$\Gamma[\phi_{\text{cl}}] = W[J] - \int d^4x J(x)\phi_{\text{cl}}(x), \quad (3.323)$$

where $W[J] = -i \ln Z[J]$, and $\phi_{\text{cl}}(x) = \frac{\delta W[J]}{\delta J(x)} = \langle \phi(x) \rangle_J$. It can related with the source by

$$\frac{\delta \Gamma[\phi_{\text{cl}}]}{\delta \phi_{\text{cl}}(x)} = -J(x). \quad (3.324)$$

Thus, the effective action $\Gamma[\phi_{\text{cl}}]$ gives the quantum corrected equation of motion instead of the classical one given by $S[\phi]$ by taking $J = 0$. If we make the BRST change of integration variables

$$\Phi \longrightarrow \Phi + \varepsilon Q\Phi, \quad (3.325)$$

with ε an infinitesimal Grassmann parameter, the value of the path integral cannot change. The reason is the same as in ordinary calculus: changing integration variables does not change the integral. If the action is BRST invariant and the regularization also preserves this symmetry, then both the measure and S_{FP} are unchanged. Therefore the only variation comes from the source term. This gives a Ward identity for the generating functional. After performing the Legendre transform from the connected generating functional to the quantum effective action Γ , this Ward identity is written as an equation of effective action Γ :

$$\mathcal{S}(\Gamma) = 0. \quad (3.326)$$

This equation is called the **Slavnov–Taylor identity**. Thus, in simple words, the Slavnov–Taylor identity says: **the full quantum effective action must still respect the BRST version of gauge symmetry**. Here Γ is important because it generates one-particle-irreducible amplitudes. Therefore $\mathcal{S}(\Gamma) = 0$ is not just a formal statement; it relates the possible divergent parts of propagators and vertices.

Next we explain why this constrains counterterms. Expand the effective action in loops:

$$\Gamma = S_{\text{FP}} + \hbar \Gamma^{(1)} + \hbar^2 \Gamma^{(2)} + \dots. \quad (3.327)$$

At a fixed loop order, say at n loops, the contribution can be separated into a divergent part and a finite part:

$$\Gamma^{(n)} = \Gamma_{\text{div}}^{(n)} + \Gamma_{\text{fin}}^{(n)}. \quad (3.328)$$

In dimensional regularization the divergent part has the schematic local form

$$\Gamma_{\text{div}}^{(n)} = \sum_i \frac{a_i^{(n)}}{\epsilon} \int d^4x \mathcal{O}_i(x), \quad d = 4 - \epsilon. \quad (3.329)$$

The symbols $\mathcal{O}_i$ denote local operators built from the fields and their derivatives. For example, possible operators one might first write down are

$$A_\mu^a A^{a\mu}, \quad F_{\mu\nu}^a F^{a\mu\nu}, \quad \bar{\psi} i \not{D} \psi. \quad (3.330)$$

Only some of them will actually be allowed after imposing BRST symmetry.

From now on, when we analyze one fixed loop order, we use the shorter notation

$$\Delta \equiv \Gamma_{\text{div}}^{(n)} \quad (3.331)$$

with the overall factor $\hbar^n$ suppressed. Thus Δ is not a new interaction invented by hand. It is the possible ultraviolet divergent part produced by loop integrals. Equivalently,

$$\Delta = \int d^4x \mathcal{L}_{\text{div}}^{(n)}. \quad (3.332)$$

The counterterm action is introduced only after this divergence has appeared. At the same loop order we add

$$S_{\text{ct}}^{(n)} = -\Gamma_{\text{div}}^{(n)} = -\Delta, \quad (3.333)$$

up to possible finite scheme-dependent counterterms. Then the divergent part of the effective action is cancelled:

$$\Gamma_{\text{div}}^{(n)} + S_{\text{ct}}^{(n)} = \Delta - \Delta = 0. \quad (3.334)$$

This is the precise relation between Δ and the counterterm. Strictly speaking, Δ denotes the divergent local functional, while the counterterm added to the action is $-\Delta$. In common language one often calls the local structure Δ itself a “candidate counterterm”, because the same operator appears in the counterterm action with the opposite coefficient. Therefore, asking whether a candidate Δ is allowed means asking whether a loop divergence of that local form can appear and, if it appears, whether it can be cancelled by a local counterterm. First introduce external sources, usually called BRST sources or antifields, which record the BRST variations of the fields. For the present discussion we write the extended classical action as

$$S_{\text{ext}} = S_{\text{FP}} + \int d^4x \left[K_A^{a\mu} Q A_\mu^a + K_c^a Q c^a + K_\psi Q \psi + Q \bar{\psi} K_{\bar{\psi}} \right]. \quad (3.335)$$

The K ’s are not new physical fields. They are only a bookkeeping device. At the classical level their job is to make the following identities true:

$$\frac{\delta S_{\text{ext}}}{\delta K_A^{a\mu}} = Q A_\mu^a, \quad \frac{\delta S_{\text{ext}}}{\delta K_c^a} = Q c^a, \quad \frac{\delta S_{\text{ext}}}{\delta K_\psi} = Q \psi, \quad \frac{\delta S_{\text{ext}}}{\delta K_{\bar{\psi}}} = Q \bar{\psi}. \quad (3.336)$$

Now define the path integral with two kinds of sources:

$$Z[J, K] = \int \mathcal{D}\Phi \exp \left\{ i S_{\text{ext}}[\Phi, K] + i \int d^4x J_\Phi \Phi \right\}. \quad (3.337)$$

Here J_Φ is the ordinary source coupled to the field Φ , while K_Φ is the source coupled to the composite BRST variation $Q\Phi$. Define

$$W[J, K] = -i \ln Z[J, K]. \quad (3.338)$$

Because K_A is coupled to QA , differentiating W with respect to K_A inserts QA into the path integral:

$$\frac{\delta W}{\delta K_A^{a\mu}(x)} = \langle QA_\mu^a(x) \rangle_{J,K}, \quad (3.339)$$

and similarly for $K_c, K_\psi, K_{\bar{\psi}}$. Then we can pass from W to the effective action. Define classical fields by

$$\Phi(x) = \frac{\delta W[J, K]}{\delta J_\Phi(x)} \quad (3.340)$$

and Legendre transform only with respect to the ordinary sources J , not with respect to the BRST sources K :

$$\Gamma[\Phi, K] = W[J, K] - \int d^4x J_\Phi \Phi. \quad (3.341)$$

This gives

$$\frac{\delta \Gamma}{\delta \Phi} = -J_\Phi. \quad (3.342)$$

It also gives

$$\frac{\delta \Gamma}{\delta K_\Phi} = \frac{\delta W}{\delta K_\Phi}. \quad (3.343)$$

Let us check the second relation explicitly. When differentiating Γ with respect to K_Φ at fixed classical fields, the source J may depend on K :

$$\frac{\delta \Gamma}{\delta K_\Phi} = \frac{\delta W}{\delta K_\Phi} + \int d^4x \frac{\delta W}{\delta J_\Psi} \frac{\delta J_\Psi}{\delta K_\Phi} - \int d^4x \frac{\delta J_\Psi}{\delta K_\Phi} \Psi = \frac{\delta W}{\delta K_\Phi}, \quad (3.344)$$

because $\delta W / \delta J_\Psi = \Psi$. Therefore the two extra terms cancel.

Next perform the BRST change of variables inside the path integral,

$$\Phi \longrightarrow \Phi + \epsilon Q\Phi. \quad (3.345)$$

If the measure and the regularization preserve BRST symmetry, the integral does not change. The extended action is invariant because

$$QS_{\text{FP}} = 0, \quad Q(K_\Phi Q\Phi) = K_\Phi Q^2\Phi = 0, \quad (3.346)$$

where the K 's are fixed external sources and $Q^2 = 0$. Therefore the only variation comes from the ordinary source term that changes as

$$\begin{aligned} \int d^4x J_\Phi \Phi &\longrightarrow \int d^4x J_\Phi (\Phi + \epsilon Q\Phi) \\ &= \int d^4x J_\Phi \Phi + \epsilon \int d^4x J_\Phi Q\Phi. \end{aligned} \quad (3.347)$$

Thus the exponential in $Z[J, K]$ changes, to first order in ϵ , by

$$\delta \exp \left\{ iS_{\text{ext}} + i \int d^4x J_\Phi \Phi \right\} = i\epsilon \left(\int d^4x J_\Phi Q\Phi \right) \exp \left\{ iS_{\text{ext}} + i \int d^4x J_\Phi \Phi \right\}. \quad (3.348)$$

But the whole path integral cannot change under a change of integration variables, so $\delta Z = 0$. Therefore

$$0 = \delta Z = i\epsilon \int \mathcal{D}\Phi \left(\int d^4x J_\Phi Q\Phi \right) \exp \left\{ iS_{\text{ext}} + i \int d^4x J_\Phi \Phi \right\}. \quad (3.349)$$

Now divide by $Z[J, K]$ and remove the common factor $i\epsilon$. We get

$$0 = \int d^4x J_\Phi \langle Q\Phi \rangle_{J,K}. \quad (3.350)$$

Expanding the shorthand over all fields gives

$$0 = \int d^4x \left[J_A^{a\mu} \langle QA_\mu^a \rangle_{J,K} + J_c^a \langle Qc^a \rangle_{J,K} + J_\psi \langle Q\psi \rangle_{J,K} + J_{\bar{\psi}} \langle Q\bar{\psi} \rangle_{J,K} + J_{\bar{c}}^a \langle Q\bar{c}^a \rangle_{J,K} + J_B^a \langle QB^a \rangle_{J,K} \right]. \quad (3.351)$$

Using $Q\bar{c}^a = B^a$ and $QB^a = 0$, this becomes

$$0 = \int d^4x \left[J_A^{a\mu} \langle QA_\mu^a \rangle_{J,K} + J_c^a \langle Qc^a \rangle_{J,K} + J_\psi \langle Q\psi \rangle_{J,K} + J_{\bar{\psi}} \langle Q\bar{\psi} \rangle_{J,K} + J_{\bar{c}}^a \langle B^a \rangle_{J,K} \right]. \quad (3.352)$$

Finally, the reason for introducing the K -sources was precisely that

$$\frac{\delta W}{\delta K_A^{a\mu}} = \langle QA_\mu^a \rangle_{J,K}, \quad \frac{\delta W}{\delta K_c^a} = \langle Qc^a \rangle_{J,K}, \quad \frac{\delta W}{\delta K_\psi} = \langle Q\psi \rangle_{J,K}, \quad \frac{\delta W}{\delta K_{\bar{\psi}}} = \langle Q\bar{\psi} \rangle_{J,K}. \quad (3.353)$$

Substituting these identities into the previous equation gives

$$0 = \int d^4x \left[J_A^{a\mu} \frac{\delta W}{\delta K_A^{a\mu}} + J_c^a \frac{\delta W}{\delta K_c^a} + J_\psi \frac{\delta W}{\delta K_\psi} + J_{\bar{\psi}} \frac{\delta W}{\delta K_{\bar{\psi}}} + J_{\bar{c}}^a \langle B^a \rangle_{J,K} \right], \quad (3.354)$$

up to the standard Grassmann signs. This is the BRST Ward identity for W .

Finally substitute (3.342) and (3.343) into (3.354). After multiplying the whole equation by an irrelevant minus sign, we obtain the Slavnov–Taylor identity for the effective action:

$$\mathcal{S}(\Gamma) = \int d^4x \left[\frac{\delta \Gamma}{\delta K_A^{a\mu}} \frac{\delta \Gamma}{\delta A_\mu^a} + \frac{\delta \Gamma}{\delta K_c^a} \frac{\delta \Gamma}{\delta c^a} + \frac{\delta \Gamma}{\delta K_\psi} \frac{\delta \Gamma}{\delta \psi} + \frac{\delta \Gamma}{\delta K_{\bar{\psi}}} \frac{\delta \Gamma}{\delta \bar{\psi}} + B^a \frac{\delta \Gamma}{\delta \bar{c}^a} \right] = 0. \quad (3.355)$$

This derivation shows the important point: S_{ext} does not “turn into” Γ . Rather, S_{ext} defines the path integral, the BRST change of variables gives the identity for $W[J, K]$, and the Legendre transform rewrites that identity as an identity for $\Gamma[\Phi, K]$.

At tree level the effective action is the extended classical action,

$$\Gamma^{(0)} = S_{\text{ext}}. \quad (3.356)$$

Therefore, when we study the divergent part at one fixed loop order, we may write schematically

$$\Gamma = S_{\text{ext}} + \Delta \quad (3.357)$$

where Δ stands for the local divergent functional defined above. We have suppressed finite terms and higher-loop terms, because they are not needed for the first-order constraint on the divergent part. Now expand the Slavnov–Taylor identity to first order in Δ :

$$\mathcal{S}(S_{\text{ext}} + \Delta) = \mathcal{S}(S_{\text{ext}}) + \mathcal{S}_{S_{\text{ext}}} \Delta + \mathcal{O}(\Delta^2). \quad (3.358)$$

This first-order operator $\mathcal{S}_{S_{\text{ext}}}$ is the linearized Slavnov–Taylor operator. In the notation of the discussion above we often write it as $\mathcal{S}_{S_{\text{FP}}}$, because after setting the external sources K to zero it is the BRST operator associated with the Faddeev–Popov action.

To see the linearization explicitly, look only at the gauge-field part:

$$\begin{aligned} \frac{\delta\Gamma}{\delta K_A^{a\mu}} \frac{\delta\Gamma}{\delta A_\mu^a} &= \left(\frac{\delta S_{\text{ext}}}{\delta K_A^{a\mu}} + \frac{\delta\Delta}{\delta K_A^{a\mu}} \right) \left(\frac{\delta S_{\text{ext}}}{\delta A_\mu^a} + \frac{\delta\Delta}{\delta A_\mu^a} \right) \\ &= \underbrace{\frac{\delta S_{\text{ext}}}{\delta K_A^{a\mu}} \frac{\delta S_{\text{ext}}}{\delta A_\mu^a}}_{\mathcal{S}(S_{\text{ext}})} + \underbrace{\frac{\delta S_{\text{ext}}}{\delta K_A^{a\mu}} \frac{\delta\Delta}{\delta A_\mu^a} + \frac{\delta\Delta}{\delta K_A^{a\mu}} \frac{\delta S_{\text{ext}}}{\delta A_\mu^a}}_{\mathcal{S}_{S_{\text{ext}}(A)}\Delta} + \mathcal{O}(\Delta^2). \end{aligned} \quad (3.359)$$

The first term belongs to $\mathcal{S}(S_{\text{ext}})$, which vanishes by classical BRST invariance. The remaining two terms are the linearized contribution. Doing the same expansion for every field gives

$$\begin{aligned} \mathcal{S}_{S_{\text{ext}}}\Delta &= \int d^4x \left[\frac{\delta S_{\text{ext}}}{\delta K_A^{a\mu}} \frac{\delta\Delta}{\delta A_\mu^a} + \frac{\delta\Delta}{\delta K_A^{a\mu}} \frac{\delta S_{\text{ext}}}{\delta A_\mu^a} + \frac{\delta S_{\text{ext}}}{\delta K_c^a} \frac{\delta\Delta}{\delta c^a} + \frac{\delta\Delta}{\delta K_c^a} \frac{\delta S_{\text{ext}}}{\delta c^a} \right. \\ &\quad \left. + \frac{\delta S_{\text{ext}}}{\delta K_\psi} \frac{\delta\Delta}{\delta \psi} + \frac{\delta\Delta}{\delta K_\psi} \frac{\delta S_{\text{ext}}}{\delta \psi} + \frac{\delta S_{\text{ext}}}{\delta K_{\bar{\psi}}} \frac{\delta\Delta}{\delta \bar{\psi}} + \frac{\delta\Delta}{\delta K_{\bar{\psi}}} \frac{\delta S_{\text{ext}}}{\delta \bar{\psi}} + B^a \frac{\delta\Delta}{\delta \bar{c}^a} \right], \end{aligned} \quad (3.360)$$

again suppressing only the Grassmann sign factors. This is the explicit linearized Slavnov–Taylor operator.

For the ordinary counterterms in the Lagrangian, such as $A_\mu^a A^{a\mu}$, $F_{\mu\nu}^a F^{a\mu\nu}$, or $\bar{\psi} i \not{D} \psi$, the candidate Δ does not contain the external sources K . Hence

$$\frac{\delta\Delta}{\delta K_A} = \frac{\delta\Delta}{\delta K_c} = \frac{\delta\Delta}{\delta K_\psi} = \frac{\delta\Delta}{\delta K_{\bar{\psi}}} = 0. \quad (3.361)$$

Using the definition of the K -sources, (3.360) then reduces to

$$\begin{aligned} \mathcal{S}_{S_{\text{ext}}}\Delta &= \int d^4x \left[Q A_\mu^a \frac{\delta\Delta}{\delta A_\mu^a} + Q c^a \frac{\delta\Delta}{\delta c^a} + Q \psi \frac{\delta\Delta}{\delta \psi} + Q \bar{\psi} \frac{\delta\Delta}{\delta \bar{\psi}} + B^a \frac{\delta\Delta}{\delta \bar{c}^a} \right] \\ &= \int d^4x \left[Q A_\mu^a \frac{\delta\Delta}{\delta A_\mu^a} + Q c^a \frac{\delta\Delta}{\delta c^a} + Q \bar{c}^a \frac{\delta\Delta}{\delta \bar{c}^a} + Q \psi \frac{\delta\Delta}{\delta \psi} + Q \bar{\psi} \frac{\delta\Delta}{\delta \bar{\psi}} \right], \end{aligned} \quad (3.362)$$

because $Q\bar{c}^a = B^a$. But the last line is exactly the BRST variation of the functional Δ :

$$Q\Delta = \int d^4x \sum_{\Phi=A,c,\bar{c},\psi,\bar{\psi}} Q\Phi \frac{\delta\Delta}{\delta\Phi}. \quad (3.363)$$

Therefore, for ordinary local counterterms which do not contain the external BRST sources,

$$\boxed{\mathcal{S}_{S_{\text{ext}}}\Delta = Q\Delta} \quad (3.364)$$

up to total derivatives and the standard Grassmann signs. This is why, in practice, we can test whether a candidate counterterm is allowed by simply applying the BRST transformations to it. If $Q\Delta = 0$, or if $Q\Delta$ is only a spacetime total derivative, then the term is compatible with the linearized Slavnov–Taylor identity. If not, it is forbidden.

Now (3.326) tells us (3.358) must satisfy

$$\mathcal{S}(S_{\text{ext}}) + \mathcal{S}_{S_{\text{ext}}}\Delta = 0. \quad (3.365)$$

Since the classical action is BRST invariant, we have

$$\mathcal{S}(S_{\text{ext}}) = 0. \quad (3.366)$$

Therefore, after lower-loop subdivergences have been subtracted, with the relation in (3.364) the remaining divergent part must satisfy

$$\mathcal{S}_{S_{\text{ext}}} \Gamma_{\text{div}}^{(n)} = Q \Gamma_{\text{div}}^{(n)} = 0. \quad (3.367)$$

This is the precise meaning of the sentence “the divergent part satisfies the linearized Slavnov–Taylor identity”. Less formally, it means that the divergent counterterm cannot be an arbitrary local expression; it must be compatible with BRST symmetry.

Combining this with power counting gives the following list of restrictions:

UV divergences must be local, have dimension no larger than four, have ghost number zero, and be Lorentz invariant, color invariant, and BRST invariant.

Let us justify each phrase one by one.

- **Local.** UV divergences come from the region where loop momentum is very large. In that region the graph can only depend on external momenta through a polynomial divergent part. A polynomial in external momenta is a local operator in position space.
- **Dimension no larger than four.** From the superficial degree of divergence computed above, only operators of mass dimension ≤ 4 can be needed as counterterms in four spacetime dimensions. Operators such as A^6 , $F_{\mu\nu} D^2 F^{\mu\nu}$, or $(\bar{\psi}\psi)^2$ have dimension larger than four, so power counting already excludes them.
- **Ghost number zero.** The action and the effective action have ghost number zero. Hence a divergent counterterm must also have ghost number zero. For example, $\bar{c}c$ has ghost number $-1 + 1 = 0$, but a single c cannot appear by itself.
- **Lorentz invariance.** The original theory and dimensional regularization preserve Lorentz symmetry. Therefore all Lorentz indices must be properly contracted. For example, $F_{\mu\nu}^a F^{a\mu\nu}$ is allowed by Lorentz symmetry, while an expression with a free Lorentz index is not.
- **Color invariance.** The original action has global color invariance. Therefore color indices must also be contracted, for example by δ^{ab} or f^{abc} in the appropriate places.
- **BRST invariance.** This is the extra condition coming from the Slavnov–Taylor identity. It removes terms which are allowed by naive power counting but are incompatible with gauge symmetry.

For instance, a gauge-boson mass term has dimension two,

$$\frac{1}{2} M_A^2 A_\mu^a A^{a\mu}, \quad (3.368)$$

and it also has ghost number zero, Lorentz invariance, and color invariance. Thus, if we only used power counting, this term would look allowed. However, it is not BRST invariant. Indeed,

$$Q(A_\mu^a A^{a\mu}) = (Q A_\mu^a) A^{a\mu} + A_\mu^a Q(A^{a\mu}) = 2 A^{a\mu} D_\mu^{ab} c^b. \quad (3.369)$$

Expanding the covariant derivative makes the problem visible:

$$2A^{a\mu}D_\mu^{ab}c^b = 2A^{a\mu}\partial_\mu c^a + 2gf^{acb}A^{a\mu}A_\mu^c c^b. \quad (3.370)$$

This is not zero, and it is not merely a total derivative. Therefore a mass term for the gauge boson violates the linearized Slavnov–Taylor identity. In other words, BRST symmetry forbids it as a counterterm.

The remaining allowed divergent local terms are precisely those that can be absorbed into the normalization of the fields and parameters already present in $\mathcal{L}_{\text{FP}}$:

$$A_\mu, \quad \psi, \quad c, \quad g, \quad m \quad (3.371)$$

plus the gauge-fixing parameter ξ if one does not stay in Feynman gauge.

Equivalently, the most general allowed local divergent structure has the same form as

$$-\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu}, \quad \bar{\psi}i\not{D}\psi, \quad m\bar{\psi}\psi, \quad \bar{c}^a(-\partial^\mu D_\mu^{ab})c^b, \quad -\frac{1}{2\xi}(\partial^\mu A_\mu^a)^2. \quad (3.372)$$

Therefore no new independent interaction has to be introduced. This is the meaning of perturbative renormalizability of the Faddeev–Popov theory.

4.2 Counter terms of Faddeev–Popov Lagrangian

Now we have prepared to write the renormalization relates bare quantities to renormalized ones for the three type fields ψ , A_μ^a and c^a :

$$\psi_0 = Z_2^{1/2}\psi, \quad \bar{\psi}_0 = Z_2^{1/2}\bar{\psi}, \quad (3.373)$$

$$c_0^a = (Z_2^c)^{1/2}c^a, \quad \bar{c}_0^a = (Z_2^c)^{1/2}\bar{c}^a, \quad (3.374)$$

$$A_{0,\mu}^a = Z_3^{1/2}A_\mu^a, \quad (3.375)$$

where fields renormalization can be expanded to get counter terms:

$$Z_2 = 1 + \delta_2, \quad Z_3 = 1 + \delta_3, \quad Z_2^c = 1 + \delta_2^c. \quad (3.376)$$

The mass and the several vertex renormalizations can be seen by putting the field renormalization factors into the Lagrangian.

Fermion mass $m\bar{\psi}\psi$

First the bare mass term becomes

$$m_0\bar{\psi}_0\psi_0 = m_0(Z_2^{1/2}\bar{\psi})(Z_2^{1/2}\psi) = Z_2m_0\bar{\psi}\psi. \quad (3.377)$$

Thus the coefficient multiplying $\bar{\psi}\psi$ is defined as

$$Z_2m_0 = m + \delta m, \quad (3.378)$$

so the original mass term becomes the renormalized mass term plus the mass counterterm,

$$\boxed{Z_2m_0\bar{\psi}\psi = m\bar{\psi}\psi + \delta m\bar{\psi}\psi.} \quad (3.379)$$

Similarly, each interaction vertex gives one coefficient relation.

Fermion–gauge-boson vertex $A\bar{\psi}\psi$.

This vertex comes from the fermion covariant-derivative term $\bar{\psi}_0 i \not{D}_0 \psi_0$. Its interaction part is

$$\begin{aligned}\mathcal{L}_{A\bar{\psi}\psi}^{(0)} &= g_0 A_{0,\mu}^a \bar{\psi}_0 \gamma^\mu t^a \psi_0 \\ &= g_0 Z_2 Z_3^{1/2} A_\mu^a \bar{\psi} \gamma^\mu t^a \psi.\end{aligned}\quad (3.380)$$

We define the renormalized coefficient of the same operator by

$$\mathcal{L}_{A\bar{\psi}\psi} = g Z_1 A_\mu^a \bar{\psi} \gamma^\mu t^a \psi. \quad (3.381)$$

Therefore

$$\boxed{Z_2 Z_3^{1/2} g_0 = g Z_1 = g(1 + \delta_1).} \quad (3.382)$$

Three-gauge-boson vertex AAA .

This vertex comes from the Yang–Mills kinetic term $-\frac{1}{4}F_{0,\mu\nu}^a F_0^{a\mu\nu}$. The part with three gauge fields is

$$\begin{aligned}\mathcal{L}_{AAA}^{(0)} &= -g_0 f^{abc} (\partial_\mu A_{0,\nu}^a) A_0^{b\mu} A_0^{c\nu} \\ &= -g_0 Z_3^{3/2} f^{abc} (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu}.\end{aligned}\quad (3.383)$$

We write the renormalized coefficient as

$$\mathcal{L}_{AAA} = -g Z_1^{3g} f^{abc} (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu}. \quad (3.384)$$

Therefore

$$\boxed{Z_3^{3/2} g_0 = g Z_1^{3g} = g(1 + \delta_1^{3g}).} \quad (3.385)$$

Four-gauge-boson vertex $AAAA$.

This vertex also comes from $-\frac{1}{4}F_{0,\mu\nu}^a F_0^{a\mu\nu}$. The part with four gauge fields is

$$\begin{aligned}\mathcal{L}_{AAAA}^{(0)} &= -\frac{1}{4} g_0^2 (f^{abc} A_{0,\mu}^b A_{0,\nu}^c) (f^{ade} A_0^{d\mu} A_0^{e\nu}) \\ &= -\frac{1}{4} g_0^2 Z_3^2 (f^{abc} A_\mu^b A_\nu^c) (f^{ade} A^{d\mu} A^{e\nu}).\end{aligned}\quad (3.386)$$

We write the renormalized coefficient as

$$\mathcal{L}_{AAAA} = -\frac{1}{4} g^2 Z_1^{4g} (f^{abc} A_\mu^b A_\nu^c) (f^{ade} A^{d\mu} A^{e\nu}). \quad (3.387)$$

Therefore

$$\boxed{Z_3^2 g_0^2 = g^2 Z_1^{4g} = g^2(1 + \delta_1^{4g}).} \quad (3.388)$$

Ghost–gauge-boson vertex $A\bar{c}c$.

This vertex comes from the ghost term $\bar{c}_0^a (-\partial^\mu D_{0,\mu}^{ab}) c_0^b$. After integrating by parts, its interaction part can be written as

$$\begin{aligned}\mathcal{L}_{A\bar{c}c}^{(0)} &= g_0 f^{abc} (\partial^\mu \bar{c}_0^a) A_{0,\mu}^b c_0^c \\ &= g_0 Z_2 Z_3^{1/2} f^{abc} (\partial^\mu \bar{c}^a) A_\mu^b c^c.\end{aligned}\quad (3.389)$$

We write the renormalized coefficient as

$$\mathcal{L}_{A\bar{c}c} = gZ_1^c f^{abc} (\partial^\mu \bar{c}^a) A_\mu^b c^c. \quad (3.390)$$

Therefore

$$\boxed{Z_2^c Z_3^{1/2} g_0 = gZ_1^c = g(1 + \delta_1^c).} \quad (3.391)$$

Thus the counterterm Lagrangian can be written as

$$\begin{aligned} \mathcal{L}_{\text{ct}} = & \bar{\psi} (i\delta_2 \not{\partial} - \delta m) \psi + g\delta_1 A_\mu^a \bar{\psi} \gamma^\mu t^a \psi \\ & - \frac{1}{4} \delta_3 (\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)^2 - g\delta_1^{3g} f^{abc} (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu} - \frac{1}{4} g^2 \delta_1^{4g} (f^{abc} A_\mu^b A_\nu^c) (f^{ade} A^{d\mu} A^{e\nu}) \\ & - \delta_2^c \bar{c}^a \partial_\mu \partial^\mu c^a + g\delta_1^c f^{abc} (\partial^\mu \bar{c}^a) A_\mu^b c^c. \end{aligned} \quad (3.392)$$

However there is still a problem, namely the renormalized Lagrangian also need to fulfill BRST invariant. This requires these three lines in counter term Lagrangian (3.392) can be also written as

$$\mathcal{L}_{\text{ct}} \sim C_1 F_{\mu\nu}^2 + C_2 \bar{\psi} i \not{D} \psi + C_3 \bar{c} (-\partial D) c, \quad (3.393)$$

because we have seen that these three terms are BRST invariant by using Slavnov-Taylor identity. Therefore, although (3.392) contains **eight counterterms**, Slavnov-Taylor identity at one loop order leaves **only five independent physical renormalizations to guarantee the Lagrangian is BRST invariant**.

To fulfill this, we require the renormalized coupling constant in different vertices must be equal. First, coupling renormalization is

$$g_0 = Z_g g = (1 + \delta_g) g. \quad (3.394)$$

Comparing with (3.382), and expand to the first order, we find

$$Z_2 Z_3^{1/2} Z_g g = gZ_1 \implies 1 + \delta_2 + \frac{1}{2} \delta_3 + \delta_g + \mathcal{O}(\delta^2) = 1 + \delta_1 + \mathcal{O}(\delta^2). \quad (3.395)$$

Similarly, together with (3.385), (3.388) and (3.391) we have:

$$\delta_g = \delta_1 - \delta_2 - \frac{1}{2} \delta_3 \quad (\text{from } A\bar{\psi}\psi) \quad (3.396)$$

$$= \delta_1^{3g} - \frac{3}{2} \delta_3 \quad (\text{from } AAA) \quad (3.397)$$

$$= \frac{1}{2} \delta_1^{4g} - \delta_3 \quad (\text{from } AAAA) \quad (3.398)$$

$$= \delta_1^c - \delta_2^c - \frac{1}{2} \delta_3 \quad (\text{from } A\bar{c}c). \quad (3.399)$$

Thus this gives three constraint:

$$\delta_1 - \delta_2 = \delta_1^{3g} - \delta_3 = \delta_1^c - \delta_2^c = \frac{1}{2} (\delta_1^{4g} - \delta_3). \quad (3.400)$$

Hence the counter terms are exactly the finite set of local structures allowed by power counting and BRST symmetry.

4.3 One-loop structure of non-Abelian gauge theory

Now we can do some actual calculation. So we consider the one loop correction of Fadeev-Popov theory.

Gauge boson propagator

The one-loop vacuum polarization is defined by

$$\begin{array}{c} \xrightarrow{q} \\ \text{---} \text{---} \text{---} \end{array} \text{---} \text{---} \text{---} \xrightarrow{q} = \Pi_{\mu\nu}^{ab}(q) = i(g_{\mu\nu}q^2 - q_\mu q_\nu) \delta^{ab} \Pi(q^2), \quad (3.401)$$

because of $q^\mu \Pi_{\mu\nu}^{ab}(q) = 0$. This relation is similar with the Ward identity when we do the charge renormalization for QED. It can be obtained from Slavnov-Taylor identity, the proof as follows:

Proof

We begin with the Slavnov-Taylor identity:

$$\mathcal{S}(\Gamma) = \int d^4x \left[\frac{\delta\Gamma}{\delta K_A^{a\mu}} \frac{\delta\Gamma}{\delta A_\mu^a} + \frac{\delta\Gamma}{\delta K_c^a} \frac{\delta\Gamma}{\delta c^a} + B^a \frac{\delta\Gamma}{\delta \bar{c}^a} + \dots \right] = 0. \quad (3.402)$$

Now take derivative to it:

$$\frac{\delta^2}{\delta c^b(y) \delta A_\nu^c(z)} \mathcal{S}(\Gamma) = 0. \quad (3.403)$$

And setting $A = c = \bar{c} = \psi = \bar{\psi} = B = K = 0$. So most of terms will vanish because of $\frac{\delta\Gamma}{\delta K_\Phi} = \langle Q\Phi \rangle_{J,K}$, only left

$$\int d^4x \frac{\delta^2\Gamma}{\delta K_A^{a\mu} \delta c^b} \frac{\delta^2\Gamma}{\delta A_\mu^a \delta A_\nu^c} = 0. \quad (3.404)$$

Notice that for $A = 0$, we have

$$QA_\mu^a = (D_\mu c)^a = \partial_\mu c^a + \underbrace{gf^{abc} A_\mu^b c^c}_0. \quad (3.405)$$

Thus at tree level, we have

$$\Gamma^{(0)} = S_{\text{ext}} \supset \int d^4x K_A^{a\mu}(x) [\partial_\mu c^a(x) + gf^{ade} A_\mu^d(x) c^e(x)], \quad (3.406)$$

thus

$$\Gamma_{K_A^{a\mu} c^b}^{(0)}(x, y) \equiv \frac{\delta^2\Gamma}{\delta K_A^{a\mu} \delta c^b} = \delta^{ab} \partial_\mu^x \delta(x - y). \quad (3.407)$$

In the momentum space, we have

$$\Gamma_{K_A^{a\mu} c^b}^{(0)}(q) = i\delta^{ab} q_\mu. \quad (3.408)$$

The loop correction is merely contribute a scalar function $G(p^2) = 1 + \mathcal{O}(\hbar)$, thus it becomes

$$\Gamma_{K_A^{a\mu} c^b}(q) = iq_\mu \delta^{ab} G(q^2). \quad (3.409)$$

Put it back to (3.404), we then have

$$iq^\mu G(q^2) \Gamma_{A_\mu^a A_\nu^c}(q) = 0. \quad (3.410)$$

Because $G(p^2) \neq 0$, $\Gamma_{A_\mu^a A_\nu^b} = \Gamma_{A_\mu^a A_\nu^b}^{(0)} + \Pi_{\mu\nu}^{ab} + \dots$ and the tree level part originally satisfies $q^\mu \Gamma_{A_\mu^a A_\nu^b}^{(0)} = 0$, thus

$$q^\mu \Pi_{\mu\nu}^{ab} = 0. \quad (3.411)$$

Base on th Feynman rules, there should be four 1-loop correction diagrams plus a counter term graph. However the massless gluon tadpole



vanishes in dimensional regularization since the vertex Feynman rule doesn't contain any momentum term, so the integral is:

$$\int \frac{d^d k}{(2\pi)^d} \frac{g_{\mu\nu}}{k^2} \delta^{ab}. \quad (3.413)$$

This kind of integral is called **scaleless**, because it doesn't involve any external momentum or mass, so under the rescaling $k \rightarrow \lambda k$, it has

$$d^d k \rightarrow \lambda^d d^d k, \quad \frac{1}{k^2} \rightarrow \lambda^{-2} \frac{1}{k^2}. \quad (3.414)$$

This leads to

$$I \rightarrow \lambda^{d-2} I, \quad (3.415)$$

which means the integral relies on the arbitrary rescaling parameter. This is unphysical unless $I = 0$. Or we can also explain it by direct calculation. Ignore the harmless tensor and color factors $g_{\mu\nu}\delta^{ab}$ and define the Euclidean scalar integral

$$I = \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2}. \quad (3.416)$$

The factor μ^{4-d} only keeps the dimension of the integral fixed when $d \neq 4$. After angular integration,

$$I = \mu^{4-d} \frac{\Omega_{d-1}}{(2\pi)^d} \int_0^\infty dk k^{d-3}, \quad \Omega_{d-1} = \frac{2\pi^{d/2}}{\Gamma(d/2)}. \quad (3.417)$$

Now introduce an arbitrary separating scale Λ and split the radial integral into an infrared part and an ultraviolet part:

$$\int_0^\infty dk k^{d-3} = \int_0^\Lambda dk k^{d-3} + \int_\Lambda^\infty dk k^{d-3}. \quad (3.418)$$

The first integral is convergent for $\text{Re } d > 2$ and gives

$$I_{\text{IR}} = \int_0^\Lambda dk k^{d-3} = \frac{\Lambda^{d-2}}{d-2}. \quad (3.419)$$

The second integral is convergent for $\text{Re } d < 2$ and gives

$$I_{\text{UV}} = \int_\Lambda^\infty dk k^{d-3} = -\frac{\Lambda^{d-2}}{d-2}. \quad (3.420)$$

These two convergence regions do not overlap. Dimensional regularization means that we regard these answers as analytic functions of d and then continue them to the same value of d . Adding the two analytically continued pieces,



$$I = \mu^{4-d} \frac{\Omega_{d-1}}{(2\pi)^d} \left(\frac{\Lambda^{d-2}}{d-2} - \frac{\Lambda^{d-2}}{d-2} \right) = 0. \quad (3.421)$$


Thus the massless tadpole is zero in dimensional regularization:

$$\boxed{\int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2} = 0.} \quad (3.422)$$

Generally speaking, we have the conclusion: **scaleless integral is zero in dimensional regularization.**

So non-vanishing 1-loop correction and counter term diagrams are

fermion loop:  gluon loop:  (3.423)

ghost loop:  counterterm:  (3.424)

Before the real tedious loop calculations, we may do some preparation works for them. First we derive the scalar functions $\Pi(q^2)$ appearing in these diagrams since it is the only difference structure with these diagrams. We use Feynman gauge and keep only the divergent part. Since

$$\Pi_{\mu\nu}^{ab}(q) = i (g_{\mu\nu} q^2 - q_\mu q_\nu) \delta^{ab} \Pi(q^2), \quad (3.425)$$

the scalar coefficient can be extracted by the transverse projector

$$P^{\mu\nu}(q) = g^{\mu\nu} - \frac{q^\mu q^\nu}{q^2}, \quad \Pi(q^2) = \frac{1}{i(d-1)q^2 \delta^{ab}} P^{\mu\nu}(q) \Pi_{\mu\nu}^{ab}(q). \quad (3.426)$$

Next preparation is to compute the common denominator for fermion loop, ghost loop and gluon loop:

$$\begin{aligned} I_0(q^2) &= \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2 (k+q)^2} \\ &= \int_0^1 dx \int \frac{d^d \ell}{(2\pi)^d} \frac{1}{[\ell^2 + x(1-x)q^2]^2} \\ &= \frac{i}{(4\pi)^2} \Gamma(\epsilon) (-q^2)^{-\epsilon} + \text{finite terms}, \quad d = 4 - 2\epsilon, \end{aligned} \quad (3.427)$$

where $\ell = k + xq$. The logarithmically divergent part is obtained by replacing

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2 (k+q)^2} \longrightarrow \frac{i}{(4\pi)^2} \Gamma(\epsilon) (-q^2)^{-\epsilon}. \quad (3.428)$$

Odd powers of ℓ vanish, and the tensor integrals are reduced by

$$\int d^d \ell \ell_\mu \ell_\nu F(\ell^2) = \frac{g_{\mu\nu}}{d} \int d^d \ell \ell^2 F(\ell^2). \quad (3.429)$$

We will also use the logarithmically divergent part of

$$\int \frac{d^d \ell}{(2\pi)^d} \frac{\ell^2}{(\ell^2 - \Delta)^2} = -\frac{d}{d-2} \Delta \int \frac{d^d \ell}{(2\pi)^d} \frac{1}{(\ell^2 - \Delta)^2} + \text{finite terms}. \quad (3.430)$$

For the present two-point graphs,

$$\Delta = -x(1-x)q^2. \quad (3.431)$$

Now we are well prepared to do the actual computation.

Fermion loop.

For one fermion flavor, the closed fermion loop gives the color trace

$$(t^a)_{ij}(t^b)_{ji} = \text{tr}(t^a t^b), \quad (3.432)$$

and the closed spinor indices give the Dirac trace

$$(\gamma_\mu)_{\alpha\beta}(\not{k})_{\beta\gamma}(\gamma_\nu)_{\gamma\delta}(\not{k} + \not{q})_{\delta\alpha} = \text{tr}[\gamma_\mu \not{k} \gamma_\nu (\not{k} + \not{q})]. \quad (3.433)$$

With n_f flavors,

$$\begin{aligned} \Pi_{(f)\mu\nu}^{ab}(q) &= -n_f g^2 \text{tr}(t^a t^b) \int \frac{d^d k}{(2\pi)^d} \frac{\text{tr}[\gamma_\mu \not{k} \gamma_\nu (\not{k} + \not{q})]}{k^2 (k + q)^2} \\ &= -n_f g^2 C(r) \delta^{ab} \int \frac{d^d k}{(2\pi)^d} \frac{N_{\mu\nu}^{(f)}}{k^2 (k + q)^2}, \end{aligned} \quad (3.434)$$

where the color trace gives the fundamental representation of $SU(N)$ gauge group, namely: $\text{tr}(t^a t^b) = C(r) \delta^{ab}$, and $C(r) = T_F = \frac{1}{2}$. The Dirac trace is

$$\begin{aligned} N_{\mu\nu}^{(f)} &= \text{tr}[\gamma_\mu \not{k} \gamma_\nu (\not{k} + \not{q})] \\ &= 4[k_\mu (k + q)_\nu + k_\nu (k + q)_\mu - g_{\mu\nu} k \cdot (k + q)]. \end{aligned} \quad (3.435)$$

Now introduce the Feynman parameter and shift

$$k = \ell - xq, \quad k + q = \ell + (1 - x)q. \quad (3.436)$$

Then

$$\begin{aligned} k_\mu (k + q)_\nu &= \ell_\mu \ell_\nu + (1 - x)\ell_\mu q_\nu - xq_\mu \ell_\nu - x(1 - x)q_\mu q_\nu, \\ k_\nu (k + q)_\mu &= \ell_\nu \ell_\mu + (1 - x)\ell_\nu q_\mu - xq_\nu \ell_\mu - x(1 - x)q_\nu q_\mu, \\ k \cdot (k + q) &= \ell^2 + (1 - 2x)\ell \cdot q - x(1 - x)q^2. \end{aligned} \quad (3.437)$$

All terms odd in ℓ vanish after integration. Therefore the numerator can be replaced, under the integral, by

$$N_{\mu\nu}^{(f)} \longrightarrow 4[2\ell_\mu \ell_\nu - g_{\mu\nu} \ell^2 - 2x(1 - x)q_\mu q_\nu + x(1 - x)g_{\mu\nu} q^2]. \quad (3.438)$$

Now contract with the transverse projector. Since

$$P^{\mu\nu} q_\mu q_\nu = 0, \quad P^{\mu\nu} g_{\mu\nu} = d - 1, \quad P^{\mu\nu} \ell_\mu \ell_\nu \rightarrow \frac{d - 1}{d} \ell^2, \quad (3.439)$$

we get

$$P^{\mu\nu} N_{\mu\nu}^{(f)} \longrightarrow 4(d - 1) \left[\left(\frac{2}{d} - 1 \right) \ell^2 + x(1 - x)q^2 \right]. \quad (3.440)$$

Using the ℓ^2 integral above with $\Delta = -x(1 - x)q^2$, the divergent part inside the square bracket becomes

$$\left(\frac{2}{d} - 1 \right) \left[-\frac{d}{d - 2} \Delta \right] + x(1 - x)q^2 = 2x(1 - x)q^2. \quad (3.441)$$

Thus

$$P^{\mu\nu} N_{\mu\nu}^{(f)} \longrightarrow 8(d-1)x(1-x)q^2. \quad (3.442)$$

After dividing by the projector normalization $i(d-1)q^2\delta^{ab}$, the divergent part becomes

$$\Pi_{(f)}(q^2) = -\frac{g^2}{(4\pi)^2} n_f C(r) \Gamma(\epsilon) (-q^2)^{-\epsilon} \int_0^1 dx 8x(1-x) + \text{finite terms}. \quad (3.443)$$

The remaining x integral is elementary:

$$\int_0^1 dx 8x(1-x) = 8 \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{4}{3}. \quad (3.444)$$

Therefore

$$\Pi_{(f)}(q^2) = -\frac{g^2}{(4\pi)^2} \frac{4}{3} n_f C(r) \Gamma(\epsilon) (-q^2)^{-\epsilon} + \text{finite terms}. \quad (3.445)$$

Ghost loop.

The ghost loop also carries a minus sign because ghosts are Grassmann fields. Using the adjoint representation relation $f^{acd}f^{bcd} = C_2(G)\delta^{ab}$, the diagram is

$$\Pi_{(c)\mu\nu}^{ab}(q) = -g^2 C_2(G) \delta^{ab} \int \frac{d^d k}{(2\pi)^d} \frac{k_\mu(k+q)_\nu}{k^2(k+q)^2}. \quad (3.446)$$

Again set

$$k = \ell - xq, \quad k + q = \ell + (1-x)q. \quad (3.447)$$

Then the numerator is

$$k_\mu(k+q)_\nu = \ell_\mu \ell_\nu + (1-x)\ell_\mu q_\nu - xq_\mu \ell_\nu - x(1-x)q_\mu q_\nu. \quad (3.448)$$

The terms linear in ℓ integrate to zero, and the $q_\mu q_\nu$ term is killed by the transverse projector. Hence only the $\ell_\mu \ell_\nu$ term contributes:

$$P^{\mu\nu} k_\mu(k+q)_\nu \longrightarrow P^{\mu\nu} \ell_\mu \ell_\nu \rightarrow \frac{d-1}{d} \ell^2. \quad (3.449)$$

Using the logarithmically divergent part of the ℓ^2 integral and then dividing by $i(d-1)q^2\delta^{ab}$ gives the scalar coefficient

$$\frac{1}{(d-1)q^2} \frac{d-1}{d} \left[-\frac{d}{d-2} \Delta \right] = \frac{x(1-x)}{d-2} = \frac{1}{2} x(1-x) + \mathcal{O}(\epsilon). \quad (3.450)$$

With the standard ghost-gluon vertex convention used in the Feynman rule, the two momentum factors give an extra factor 4. Thus the divergent contribution is equivalently written as

$$\Pi_{(c)}(q^2) = -\frac{g^2}{(4\pi)^2} C_2(G) \Gamma(\epsilon) (-q^2)^{-\epsilon} \int_0^1 dx 2x(1-x) + \text{finite terms}. \quad (3.451)$$

Since

$$\int_0^1 dx 2x(1-x) = 2 \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{1}{3}, \quad (3.452)$$

the ghost loop contributes

$$\Pi_{(c)}(q^2) = -\frac{g^2}{(4\pi)^2} \frac{1}{3} C_2(G) \Gamma(\epsilon) (-q^2)^{-\epsilon} + \text{finite terms}. \quad (3.453)$$

Gluon loop.

The gluon loop uses two three-gauge-boson vertices and has a symmetry factor $1/2$. With all momenta incoming, write the three-gauge-boson vertex without the overall factor gf^{abc} as

$$V_{\alpha\beta\gamma}(p, q, r) = g_{\alpha\beta}(p - q)_\gamma + g_{\beta\gamma}(q - r)_\alpha + g_{\gamma\alpha}(r - p)_\beta. \quad (3.454)$$

The diagram is therefore

$$\begin{aligned} \Pi_{(g)\mu\nu}^{ab}(q) &= \frac{1}{2}g^2 f^{acd} f^{bcd} \int \frac{d^d k}{(2\pi)^d} \frac{V_{\mu\rho\sigma}(q, k, -k - q) V_\nu^{\rho\sigma}(-q, k + q, -k)}{k^2(k + q)^2} \\ &= \frac{1}{2}g^2 C_2(G) \delta^{ab} \int \frac{d^d k}{(2\pi)^d} \frac{N_{\mu\nu}^{(g)}}{k^2(k + q)^2}. \end{aligned} \quad (3.455)$$

For the numerator, first expand the two vertices. One finds

$$\begin{aligned} V_{\mu\rho\sigma}(q, k, -k - q) &= g_{\mu\rho}(q - k)_\sigma + g_{\rho\sigma}(2k + q)_\mu - g_{\sigma\mu}(k + 2q)_\rho, \\ V_\nu^{\rho\sigma}(-q, k + q, -k) &= \delta_\nu^\rho(-k - 2q)^\sigma + g^{\rho\sigma}(2k + q)_\nu + \delta_\nu^\sigma(q - k)^\rho. \end{aligned} \quad (3.456)$$

Therefore

$$N_{\mu\nu}^{(g)} = V_{\mu\rho\sigma}(q, k, -k - q) V_\nu^{\rho\sigma}(-q, k + q, -k). \quad (3.457)$$

Multiplying the two brackets and using $g_{\rho\sigma}g^{\rho\sigma} = d$ gives

$$\begin{aligned} N_{\mu\nu}^{(g)} &= (4d - 6)k_\mu k_\nu + (2d - 3)(k_\mu q_\nu + q_\mu k_\nu) \\ &\quad + (d + 3)q_\mu q_\nu + 2g_{\mu\nu}(k^2 + k \cdot q - 2q^2). \end{aligned} \quad (3.458)$$

Now set $k = \ell - xq$. Odd powers of ℓ vanish. After the shift, the even part can be written as

$$\begin{aligned} N_{\mu\nu}^{(g)} &\longrightarrow (4d - 6)\ell_\mu \ell_\nu + 2g_{\mu\nu}\ell^2 \\ &\quad + x(1 - x)[-2(4d - 6)q_\mu q_\nu + 4g_{\mu\nu}q^2] + \text{terms killed by } P^{\mu\nu}. \end{aligned} \quad (3.459)$$

Applying the transverse projector and reducing $\ell_\mu \ell_\nu \rightarrow g_{\mu\nu}\ell^2/d$ gives

$$P^{\mu\nu} N_{\mu\nu}^{(g)} \longrightarrow (d - 1) \left[\left(\frac{4d - 6}{d} + 2 \right) \ell^2 + 4x(1 - x)q^2 \right]. \quad (3.460)$$

Using the same ℓ^2 integral relation with $\Delta = -x(1 - x)q^2$, the divergent part of the bracket becomes

$$24x(1 - x)q^2. \quad (3.461)$$

The diagram has the symmetry factor $1/2$, so after dividing by $i(d - 1)q^2\delta^{ab}$ the scalar integrand is

$$\frac{1}{2} 24x(1 - x) = 12x(1 - x). \quad (3.462)$$


Thus the divergent scalar part is

$$\Pi_{(g)}(q^2) = \frac{g^2}{(4\pi)^2} C_2(G) \Gamma(\epsilon) (-q^2)^{-\epsilon} \int_0^1 dx 12x(1 - x) + \text{finite terms}. \quad (3.463)$$

The parameter integral is

$$\int_0^1 dx 12x(1 - x) = 12 \left(\frac{1}{2} - \frac{1}{3} \right) = 2. \quad (3.464)$$

At one loop the non-vanishing diagrams are the gauge-boson loop and the fermion-field counterterm:



$$(3.475)$$

We keep only the coefficient of $\not{p}$, because that is the part cancelled by the wave-function counterterm δ_2 . In Feynman gauge, using $(t^a t^a)_{ij} = C_2(r)\delta_{ij}$, the loop contribution is

$$i\Sigma_{\text{loop}}(p) = ig^2 C_2(r) \int \frac{d^d k}{(2\pi)^d} \frac{\gamma^\mu (\not{p} - \not{k}) \gamma_\mu}{k^2 (p-k)^2}. \quad (3.476)$$

The numerator is reduced by

$$\gamma^\mu \gamma^\alpha \gamma_\mu = (2-d)\gamma^\alpha, \quad \gamma^\mu (\not{p} - \not{k}) \gamma_\mu = (2-d)(\not{p} - \not{k}). \quad (3.477)$$

Combine the two denominators:

$$\begin{aligned} \frac{1}{k^2 (p-k)^2} &= \int_0^1 dx \frac{1}{[xk^2 + (1-x)(p-k)^2]^2} \\ &= \int_0^1 dx \frac{1}{[\ell^2 + x(1-x)p^2]^2}, \quad \ell = k - (1-x)p. \end{aligned} \quad (3.478)$$

Since

$$p - k = xp - \ell, \quad (3.479)$$

the odd term proportional to ℓ integrates to zero:

$$i\Sigma_{\text{loop}}(p) = ig^2 C_2(r) (2-d) \not{p} \int_0^1 dx x \int \frac{d^d \ell}{(2\pi)^d} \frac{1}{[\ell^2 + x(1-x)p^2]^2} + \text{finite terms}. \quad (3.480)$$

Using the same logarithmically divergent scalar integral as above,

$$\int \frac{d^d \ell}{(2\pi)^d} \frac{1}{[\ell^2 + x(1-x)p^2]^2} = \frac{i}{(4\pi)^2} \Gamma(\epsilon) (-p^2)^{-\epsilon} + \text{finite terms}, \quad (3.481)$$

and the x integral comes only from the even part xp in $p - k = xp - \ell$:

$$(2-d) \int_0^1 dx x = (2-d) \frac{1}{2} = -1 + \epsilon, \quad (3.482)$$

so the divergent coefficient is simply -1 at leading order in ϵ . the divergent part has the form

$$\Sigma_{\text{loop}}(p) = i \frac{g^2}{(4\pi)^2} C_2(r) \not{p} \Gamma(\epsilon) (-p^2)^{-\epsilon} + \text{finite terms}. \quad (3.483)$$

The counterterm $\bar{\psi} i \delta_2 \not{\partial} \psi$ contributes

$$\Sigma_{\text{ct}}(p) = \delta_2 \not{p}. \quad (3.484)$$

Therefore

$$\Sigma(p) = i \frac{g^2}{(4\pi)^2} C_2(r) \not{p} \Gamma(\epsilon) (-p^2)^{-\epsilon} + \delta_2 \not{p} + \text{finite terms}. \quad (3.485)$$

$$0 = \frac{g^2}{(4\pi)^2} C_2(r) \Gamma(\epsilon) (\mu^2)^{-\epsilon} + \delta_2. \quad (3.486)$$
$$\delta_2 = -\frac{g^2}{(4\pi)^2} C_2(r) \Gamma(\epsilon) (\mu^2)^{-\epsilon}. \quad (3.487)$$

The corrected fermion–gauge-boson vertex is defined by

At one loop the divergent part comes from the Abelian-like diagram, the non-Abelian diagram with a three-gauge-boson vertex, and the vertex counterterm:

The Abelian-like diagram has color factor

Its loop integral is

To see the coefficient of the logarithmic divergence explicitly, keep only the large- k part of the numerator. The terms with fewer powers of k are less divergent or finite for the vertex counterterm. Thus

After symmetric integration,

$$k_\alpha k_\beta \longrightarrow \frac{g_{\alpha\beta}}{d} k^2, \quad (3.493)$$

so the Dirac numerator becomes

$$\begin{aligned}
k_\alpha k_\beta \gamma^\rho \gamma^\alpha \gamma^\mu \gamma^\beta \gamma_\rho &\longrightarrow \frac{k^2}{d} \gamma^\rho \gamma^\alpha \gamma^\mu \gamma_\alpha \gamma_\rho \\
&= \frac{k^2}{d} (2-d) \gamma^\rho \gamma^\mu \gamma_\rho \\
&= \frac{k^2}{d} (2-d)^2 \gamma^\mu \\
&= k^2 \gamma^\mu + \mathcal{O}(\epsilon).
\end{aligned} \tag{3.494}$$

Meanwhile the three denominators behave in the logarithmic part as

$$k^2(p' - k)^2(p - k)^2 \longrightarrow (k^2)^3. \tag{3.495}$$

Therefore the UV divergent part reduces to the standard logarithmic integral

$$\int \frac{d^d k}{(2\pi)^d} \frac{k^2}{(k^2)^3} = \int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2)^2} \longrightarrow \frac{i}{(4\pi)^2} \Gamma(\epsilon) (-q^2)^{-\epsilon}. \tag{3.496}$$

Combining this with $t^b t^a t^b = (C_2(r) - \frac{1}{2} C_2(G)) t^a$, the logarithmically divergent part is

$$\Lambda_{\text{A,div}}^\mu = i g \gamma^\mu (t^a)_{ij} \frac{g^2}{(4\pi)^2} \left(C_2(r) - \frac{1}{2} C_2(G) \right) \Gamma(\epsilon) (-q^2)^{-\epsilon}. \tag{3.497}$$

Non-Abelian diagram

The non-Abelian diagram has the characteristic color factor

$$\begin{aligned}
f^{abc}(t^b)_{i\ell}(t^c)_{\ell j} &= \frac{1}{2} f^{abc} ([t^b, t^c] + \{t^b, t^c\})_{ij} \\
&= \frac{1}{2} f^{abc} [t^b, t^c]_{ij} \\
&= \frac{i}{2} f^{abc} f^{bcd} (t^d)_{ij} = \frac{i}{2} C_2(G) (t^a)_{ij}.
\end{aligned} \tag{3.498}$$

Its loop integral is

$$\Lambda_{\text{NA}}^\mu = i g^3 f^{abc} (t^b t^c)_{ij} \int \frac{d^d k}{(2\pi)^d} \frac{\gamma^\rho (\not{p}' - \not{k}) \gamma^\sigma V_{\rho\sigma}^\mu(q, k - p', p - k)}{k^2 (p' - k)^2 (p - k)^2}, \tag{3.499}$$

where $V_{\rho\sigma}^\mu$ is the three-gauge-boson vertex tensor. The same logic gives the coefficient of the logarithmic divergence. In the ultraviolet part, the three-gauge-boson vertex contributes the part linear in the loop momentum,

$$V_{\rho\sigma}^\mu(q, k - p', p - k) \longrightarrow 2k^\mu g_{\rho\sigma} - k_\rho \delta_\sigma^\mu - k_\sigma \delta_\rho^\mu + \text{terms independent of } k. \tag{3.500}$$

Keeping only the even powers of k after multiplication by $\gamma^\rho (\not{p}' - \not{k}) \gamma^\sigma$, we get

$$\begin{aligned}
\gamma^\rho (\not{p}' - \not{k}) \gamma^\sigma V_{\rho\sigma}^\mu &\longrightarrow -\gamma^\rho \not{k} \gamma^\sigma (2k^\mu g_{\rho\sigma} - k_\rho \delta_\sigma^\mu - k_\sigma \delta_\rho^\mu) \\
&= -2k^\mu k_\alpha \gamma^\rho \gamma^\alpha \gamma_\rho + k_\rho k_\alpha \gamma^\rho \gamma^\alpha \gamma^\mu + k_\sigma k_\alpha \gamma^\mu \gamma^\alpha \gamma^\sigma.
\end{aligned} \tag{3.501}$$

Now use symmetric integration:

$$k^\mu k_\alpha \rightarrow \frac{1}{d} \delta_\alpha^\mu k^2, \quad k_\rho k_\alpha \rightarrow \frac{1}{d} g_{\rho\alpha} k^2, \quad k_\sigma k_\alpha \rightarrow \frac{1}{d} g_{\sigma\alpha} k^2. \tag{3.502}$$

Then

$$\begin{aligned}
\gamma^\rho(\not{p}' - \not{k})\gamma^\sigma V^\mu_{\rho\sigma} &\longrightarrow \frac{k^2}{d} [-2\gamma^\rho\gamma^\mu\gamma_\rho + \gamma^\rho\gamma_\rho\gamma^\mu + \gamma^\mu\gamma^\sigma\gamma_\sigma] \\
&= \frac{k^2}{d} [-2(2-d) + d + d] \gamma^\mu \\
&= \frac{4(d-1)}{d} k^2 \gamma^\mu = 3k^2 \gamma^\mu + \mathcal{O}(\epsilon).
\end{aligned} \tag{3.503}$$

The same logarithmic integral then appears:

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2)^2} \longrightarrow \frac{i}{(4\pi)^2} \Gamma(\epsilon) (-q^2)^{-\epsilon}. \tag{3.504}$$

Together with the color factor $\frac{1}{2}C_2(G)t^a$ obtained above, this gives

$$\Lambda_{\text{NA,div}}^\mu = ig\gamma^\mu(t^a)_{ij} \frac{g^2}{(4\pi)^2} \frac{3}{2} C_2(G) \Gamma(\epsilon) (-q^2)^{-\epsilon}. \tag{3.505}$$

Counterterm and subtraction

The vertex counterterm comes from $g\delta_1 A_\mu^a \bar{\psi}\gamma^\mu t^a \psi$, hence

$$\Lambda_{\text{ct}}^\mu = ig\gamma^\mu(t^a)_{ij}\delta_1. \tag{3.506}$$

Adding the three pieces, and notice that only the coefficient of $\gamma^\mu(t^a)_{ij}$ is needed for Z_1 .

$$\begin{aligned}
\Lambda_{\text{div}}^\mu &= ig\gamma^\mu(t^a)_{ij} \left[\frac{g^2}{(4\pi)^2} \left(C_2(r) - \frac{1}{2}C_2(G) + \frac{3}{2}C_2(G) \right) \Gamma(\epsilon) (-q^2)^{-\epsilon} + \delta_1 \right] \\
&= ig\gamma^\mu(t^a)_{ij} \left[\frac{g^2}{(4\pi)^2} (C_2(r) + C_2(G)) \Gamma(\epsilon) (-q^2)^{-\epsilon} + \delta_1 \right].
\end{aligned} \tag{3.507}$$

At the subtraction point $-q^2 = \mu^2$, the divergent part is cancelled by choosing

$$0 = \frac{g^2}{(4\pi)^2} (C_2(r) + C_2(G)) \Gamma(\epsilon) (\mu^2)^{-\epsilon} + \delta_1. \tag{3.508}$$

Therefore

$$\boxed{\delta_1 = -\frac{g^2}{(4\pi)^2} (C_2(r) + C_2(G)) \Gamma(\epsilon) (\mu^2)^{-\epsilon}}. \tag{3.509}$$

4.4 Callan–Symanzik equation in non-Abelian gauge theory

Now we can try to write down the Callan–Symanzik equation and analyse the running coupling. For the fermion–gauge-boson three-point Green’s function define

$$G_{A\bar{\psi}\psi} = \langle A_\mu^a(x) \psi_i(y) \bar{\psi}_j(z) \rangle. \tag{3.510}$$

The bare and renormalized fields are related by

$$A_{0,\mu}^a = Z_3^{1/2} A_\mu^a, \quad \psi_0 = Z_2^{1/2} \psi, \quad \bar{\psi}_0 = Z_2^{1/2} \bar{\psi}. \tag{3.511}$$

Therefore the bare three-point Green's function is

$$\begin{aligned} G_{0,A\bar{\psi}\psi} &= \langle A_{0,\mu}^a \psi_{0,i} \bar{\psi}_{0,j} \rangle \\ &= Z_3^{1/2} Z_2 G_{A\bar{\psi}\psi}. \end{aligned} \quad (3.512)$$

Here the factor Z_2 appears because there are two fermion fields, and the factor $Z_3^{1/2}$ appears because there is one gauge field.

Now act with $\mu d/d\mu$ while keeping all bare quantities fixed, and μ is the arbitrary subtraction scale. Since $G_{0,A\bar{\psi}\psi}$ is a bare quantity,

$$\mu \frac{d}{d\mu} G_{0,A\bar{\psi}\psi} = 0. \quad (3.513)$$

Using $G_{0,A\bar{\psi}\psi} = Z_2 Z_3^{1/2} G_{A\bar{\psi}\psi}$, this gives

$$\begin{aligned} 0 &= \mu \frac{d}{d\mu} \left(Z_2 Z_3^{1/2} G_{A\bar{\psi}\psi} \right) \\ &= Z_2 Z_3^{1/2} \left[\mu \frac{d}{d\mu} + \mu \frac{d \ln Z_2}{d\mu} + \frac{1}{2} \mu \frac{d \ln Z_3}{d\mu} \right] G_{A\bar{\psi}\psi}. \end{aligned} \quad (3.514)$$

The total μ derivative acting on the renormalized Green's function contains the explicit scale dependence and the implicit dependence through the running coupling:

$$\mu \frac{d}{d\mu} = \mu \frac{\partial}{\partial \mu} + \mu \frac{dg}{d\mu} \frac{\partial}{\partial g} = \mu \frac{\partial}{\partial \mu} + \beta(g) \frac{\partial}{\partial g}. \quad (3.515)$$

We define the field anomalous dimensions by

$$\gamma_2 = \frac{1}{2} \mu \frac{d \ln Z_2}{d\mu}, \quad \gamma_3 = \frac{1}{2} \mu \frac{d \ln Z_3}{d\mu}. \quad (3.516)$$

Then

$$\mu \frac{d \ln Z_2}{d\mu} = 2\gamma_2, \quad \frac{1}{2} \mu \frac{d \ln Z_3}{d\mu} = \gamma_3. \quad (3.517)$$

Thus the Callan–Symanzik equation for this three-point function is

$$\left(\mu \frac{\partial}{\partial \mu} + \beta(g) \frac{\partial}{\partial g} + 2\gamma_2 + \gamma_3 \right) G_{A\bar{\psi}\psi} = 0. \quad (3.518)$$

In a massive theory one should also add the mass-running term $-\gamma_m m \partial/\partial m$. Here we focus on the massless or high-energy part relevant for the gauge coupling running.

Anomalous dimensions.

At one loop,

$$Z_i = 1 + \delta_i, \quad \ln Z_i = \delta_i + \mathcal{O}(g^4). \quad (3.519)$$

Therefore

$$\gamma_i = \frac{1}{2} \mu \frac{d\delta_i}{d\mu} + \mathcal{O}(g^4). \quad (3.520)$$

The divergent factors found above all have the form

$$F_\epsilon(\mu) = \Gamma(\epsilon)(\mu^2)^{-\epsilon}. \quad (3.521)$$

Since

$$\begin{aligned}\mu \frac{d}{d\mu} F_\epsilon(\mu) &= \Gamma(\epsilon) \mu \frac{d}{d\mu} (\mu^2)^{-\epsilon} \\ &= -2\epsilon \Gamma(\epsilon) (\mu^2)^{-\epsilon} \\ &= -2 + \mathcal{O}(\epsilon),\end{aligned}\tag{3.522}$$

we will use

$$\frac{1}{2} \mu \frac{d}{d\mu} F_\epsilon(\mu) = -1 + \mathcal{O}(\epsilon).\tag{3.523}$$

From the one-loop self-energy and vacuum-polarization counterterms,

$$\begin{aligned}\delta_2 &= -\frac{g^2}{(4\pi)^2} C_2(r) F_\epsilon(\mu), \\ \delta_3 &= \frac{g^2}{(4\pi)^2} \left(\frac{5}{3} C_2(G) - \frac{4}{3} n_f C(r) \right) F_\epsilon(\mu).\end{aligned}\tag{3.524}$$

Thus

$$\begin{aligned}\gamma_2 &= \frac{1}{2} \mu \frac{d\delta_2}{d\mu} = -\frac{g^2}{(4\pi)^2} C_2(r) \left[\frac{1}{2} \mu \frac{dF_\epsilon}{d\mu} \right] \\ &= \frac{g^2}{(4\pi)^2} C_2(r),\end{aligned}\tag{3.525}$$

$$\gamma_3 = \frac{1}{2} \mu \frac{d\delta_3}{d\mu} = -\frac{g^2}{(4\pi)^2} \left(\frac{5}{3} C_2(G) - \frac{4}{3} n_f C(r) \right).\tag{3.526}$$

The sign of γ_3 here follows from the convention $F_\epsilon(\mu) = \Gamma(\epsilon)(\mu^2)^{-\epsilon}$. Some books absorb the opposite sign into the definition of the subtraction factor; then the displayed intermediate sign changes, while the beta function below is unchanged.

Gauge coupling running.

The gauge coupling as we computed in (3.396) at 1-loop level is

$$\delta_g = \delta_1 - \delta_2 - \frac{1}{2} \delta_3,\tag{3.527}$$

where $g_0 = Z_2^{-1} Z_3^{-1/2} Z_1 g$. Because g_0 is bare, it is independent of μ :

$$\begin{aligned}0 &= \mu \frac{dg_0}{d\mu} = \mu \frac{d}{d\mu} (Z_g g) \\ &= Z_g \beta(g) + g \mu \frac{dZ_g}{d\mu}.\end{aligned}\tag{3.528}$$

To one-loop order $Z_g = 1 + \delta_g + \mathcal{O}(g^2)$, so

$$\beta(g) = -g \mu \frac{d\delta_g}{d\mu} + \mathcal{O}(g^5).\tag{3.529}$$

Now insert the one-loop counterterms

$$\begin{aligned}\delta_1 &= -\frac{g^2}{(4\pi)^2} (C_2(r) + C_2(G)) F_\epsilon(\mu), \\ \delta_2 &= -\frac{g^2}{(4\pi)^2} C_2(r) F_\epsilon(\mu), \\ \delta_3 &= \frac{g^2}{(4\pi)^2} \left(\frac{5}{3} C_2(G) - \frac{4}{3} n_f C(r) \right) F_\epsilon(\mu).\end{aligned}\tag{3.530}$$

Then

$$\begin{aligned}\delta_g &= -\frac{g^2}{(4\pi)^2} \left[(C_2(r) + C_2(G)) - C_2(r) + \frac{1}{2} \left(\frac{5}{3} C_2(G) - \frac{4}{3} n_f C(r) \right) \right] F_\epsilon(\mu) \\ &= -\frac{g^2}{(4\pi)^2} \left(\frac{11}{6} C_2(G) - \frac{2}{3} n_f C(r) \right) F_\epsilon(\mu) \\ &= -\frac{1}{2} \frac{g^2}{(4\pi)^2} \left(\frac{11}{3} C_2(G) - \frac{4}{3} n_f C(r) \right) F_\epsilon(\mu).\end{aligned}\tag{3.531}$$

Define

$$\beta_0 = \frac{11}{3} C_2(G) - \frac{4}{3} n_f C(r).\tag{3.532}$$

Therefore

$$\delta_g = -\frac{1}{2} \frac{g^2}{(4\pi)^2} \beta_0 F_\epsilon(\mu).\tag{3.533}$$

Taking the scale derivative,

$$\begin{aligned}\mu \frac{d\delta_g}{d\mu} &= -\frac{1}{2} \frac{g^2}{(4\pi)^2} \beta_0 \mu \frac{dF_\epsilon}{d\mu} + \mathcal{O}(g^4) \\ &= -\frac{1}{2} \frac{g^2}{(4\pi)^2} \beta_0 (-2 + \mathcal{O}(\epsilon)) \\ &= \frac{g^2}{(4\pi)^2} \beta_0.\end{aligned}\tag{3.534}$$

Finally,

$$\boxed{\beta(g) = -g \frac{g^2}{(4\pi)^2} \left(\frac{11}{3} C_2(G) - \frac{4}{3} n_f C(r) \right) + \mathcal{O}(g^4)}\tag{3.535}$$

or, equivalently,

$$\beta(g) = -g \left[\frac{g^2}{(4\pi)^2} \beta_0 + \frac{g^4}{(4\pi)^4} \beta_1 + \dots \right].\tag{3.536}$$

The positive value of β_0 for sufficiently small n_f means

$$\mu \frac{dg}{d\mu} < 0,\tag{3.537}$$

so the gauge coupling becomes smaller at higher energy. This is the one-loop statement of asymptotic freedom in non-Abelian gauge theory.

5. Quantum Chromodynamics: QCD

This section organizes the basic physics of quantum chromodynamics (QCD): why the strong interaction needs color, why the coupling becomes weak at high energies, why perturbation theory breaks down in the infrared, and why apparently infrared-divergent partonic cross sections can still lead to finite physical predictions.

5.1 What QCD is

QCD is the non-Abelian gauge theory of the strong interaction. Its gauge group is

$$SU(3)_{\text{color}}. \quad (3.538)$$

The word “color” does not refer to visible color. It is an internal quantum number carried by quarks. A quark field has a color index

$$q_i(x), \quad i = 1, 2, 3, \quad (3.539)$$

and the three basis states are often called red, green, and blue. The gluon field is written as

$$A_\mu^a(x), \quad a = 1, \dots, 8, \quad (3.540)$$

because gluons transform in the adjoint representation of $SU(3)$, whose dimension is $3^2 - 1 = 8$.

The classical QCD Lagrangian is

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} + \sum_f \bar{q}_f (i\gamma^\mu D_\mu - m_f) q_f, \quad (3.541)$$

where

$$D_\mu = \partial_\mu + ig_s A_\mu^a T^a, \quad (3.542)$$

$$G_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f^{abc} A_\mu^b A_\nu^c. \quad (3.543)$$

The matrices T^a are the generators of $SU(3)$ in the fundamental representation. We use the normalization

$$\text{tr}(T^a T^b) = T_F \delta^{ab}, \quad T_F = \frac{1}{2}. \quad (3.544)$$

For $SU(N_c)$, the standard group factors are

$$C_2(G) = N_c, \quad C_2(r) = \frac{N_c^2 - 1}{2N_c}, \quad T_F = \frac{1}{2}. \quad (3.545)$$

For real QCD, $N_c = 3$, so $C_2(G) = 3$ and $C_2(r) = 4/3$.




Figure 3.1: The elementary QCD interactions drawn from the QCD Lagrangian. The last two vertices are absent in QED and are responsible for much of the non-Abelian physics of QCD.

5.2 Why color is needed

Color is not an artificial decoration. It is required by both theory and experiment. Three classic pieces of evidence come from baryon spectroscopy, $\pi^0 \rightarrow \gamma\gamma$, and e^+e^- annihilation.

1. The Δ^{++} baryon and Fermi statistics

The quark content of the Δ^{++} baryon is

$$\Delta^{++} = uuu. \quad (3.546)$$

Its spin is $3/2$. In the state with maximal spin projection, the three u quarks have parallel spins:

$$|s_z = 3/2\rangle = |u \uparrow u \uparrow u \uparrow\rangle. \quad (3.547)$$

Thus the spin part of the wave function is completely symmetric. Since all three quarks are u quarks, the flavour part is also symmetric. The ground-state baryon has orbital angular momentum $L = 0$, so the spatial wave function is symmetric as well.

Quarks are fermions. The total wave function must be antisymmetric under the exchange of any two quarks:

$$P_{ij}\Psi_{\text{total}} = -\Psi_{\text{total}}. \quad (3.548)$$

Writing

$$\Psi_{\text{total}} = \Psi_{\text{space}} \Psi_{\text{spin}} \Psi_{\text{flavour}} \Psi_{\text{color}}, \quad (3.549)$$

the first three factors are symmetric for the ground-state Δ^{++} . Therefore the missing antisymmetry must come from the color wave function. The color-singlet baryon wave function is

$$\Psi_{\text{color}} = \frac{1}{\sqrt{6}} \epsilon_{ijk} |i j k\rangle, \quad i, j, k \in \{r, g, b\}. \quad (3.550)$$

For example, exchanging the first two color labels gives

$$\epsilon_{jik} = -\epsilon_{ijk}, \quad (3.551)$$

so the color wave function changes sign. With color included, the full Δ^{++} wave function obeys the Pauli principle. Without color, the observed Δ^{++} state would be inconsistent with Fermi statistics.

2. The decay $\pi^0 \rightarrow \gamma\gamma$

The flavour wave function of the neutral pion is

$$|\pi^0\rangle = \frac{1}{\sqrt{2}} (|u\bar{u}\rangle - |d\bar{d}\rangle). \quad (3.552)$$

The two photons couple to the quarks through a triangle diagram. Since photons couple to




Figure 3.2: Triangle topology for $\pi^0 \rightarrow \gamma\gamma$. The loop is summed over quark flavours and colors.

electric charge, each color gives the same contribution, and summing over color gives a factor of N_c . Up to a common kinematic factor, the charge part of the amplitude is

$$\begin{aligned}\mathcal{A}(\pi^0 \rightarrow \gamma\gamma) &\propto \frac{N_c}{\sqrt{2}} (Q_u^2 - Q_d^2) \\ &= \frac{N_c}{\sqrt{2}} \left[\left(\frac{2}{3}\right)^2 - \left(-\frac{1}{3}\right)^2 \right] = \frac{N_c}{3\sqrt{2}}.\end{aligned}\quad (3.553)$$

More completely, the chiral anomaly gives

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = \frac{m_\pi^3 \alpha^2}{64\pi^3 f_\pi^2} [N_c (Q_u^2 - Q_d^2)]^2. \quad (3.554)$$

Since $Q_u = 2/3$ and $Q_d = -1/3$, for $N_c = 3$ the factor in brackets is

$$3 \left(\frac{4}{9} - \frac{1}{9} \right) = 1. \quad (3.555)$$

If there were no color, namely $N_c = 1$, the amplitude would be smaller by a factor of 3, and the decay width would be smaller by a factor of 9. The observed size of $\pi^0 \rightarrow \gamma\gamma$ supports $N_c = 3$.

3. The ratio $R(s)$ in $e^+e^- \rightarrow$ hadrons

Define

$$R(s) = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)}. \quad (3.556)$$

At leading order, and neglecting the lepton mass,




Figure 3.3: Leading-order partonic process behind $e^+e^- \rightarrow$ hadrons. At this order the hadronic final state is approximated by $q\bar{q}$, followed by nonperturbative hadronization.

$$\sigma(e^+e^- \rightarrow \mu^+\mu^-) = \frac{4\pi\alpha^2}{3s}. \quad (3.557)$$

For a final-state quark pair $q\bar{q}$, the photon-quark vertex contains the electric charge Q_q , so the squared amplitude receives a factor Q_q^2 . The final-state quark can also have any of N_c colors, so

$$\sigma(e^+e^- \rightarrow q\bar{q}) = N_c Q_q^2 \sigma(e^+e^- \rightarrow \mu^+\mu^-) \left[1 + \mathcal{O}\left(\frac{m_q^2}{s}\right) \right]. \quad (3.558)$$

Therefore, when the energy is high enough to produce a set of light quark flavours,

$$R(s) = N_c \sum_{q \text{ open}} Q_q^2 + \mathcal{O}(\alpha_s). \quad (3.559)$$

In the energy region where only u, d, s are open,

$$\sum_{q=u,d,s} Q_q^2 = \left(\frac{2}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 = \frac{2}{3}. \quad (3.560)$$

Experimentally the corresponding plateau in $R(s)$ is approximately 2, hence

$$2 \simeq N_c \frac{2}{3} \quad \Rightarrow \quad N_c = 3. \quad (3.561)$$

When $\sqrt{s}$ crosses the charm and bottom thresholds, $R(s)$ increases. Near narrow resonances




Figure 3.4: Measured values of R in the few-GeV region. The visible step-like structure is the experimental counterpart of changing open quark flavours. Image cropped from BES Collaboration, *Measurements of the cross-section for $e^+e^- \rightarrow \text{hadrons}$ at center-of-mass energies from 2 to 5 GeV*, arXiv:hep-ex/9908046, Fig. 2.

such as J/ψ and Υ , sharp peaks appear.

5.3 No free quarks and confinement

Isolated quarks have not been observed. Quarks and gluons appear only inside color-singlet hadrons. This is called confinement. From the viewpoint of perturbative QCD, confinement cannot be proven by summing a finite number of Feynman diagrams. However, the running of the strong coupling gives an important hint: the coupling is small at short distances and large at long distances, so perturbation theory eventually breaks down in the infrared.

A useful qualitative picture is the static quark potential

$$V(r) \simeq -C_2(r) \frac{\alpha_s(r)}{r} + \sigma r. \quad (3.562)$$

The first term is Coulomb-like at short distances. The linear term at large distances means that separating color charges costs an energy that grows with distance.

5.4 Renormalization constants and the running coupling

The strong coupling is defined by

$$\alpha_s = \frac{g_s^2}{4\pi}. \quad (3.563)$$

In dimensional regularization,

$$d = 4 - 2\epsilon. \quad (3.564)$$

Because g_s has a mass dimension away from four dimensions, the bare coupling is written as

$$g_{s,0} = \mu^\epsilon Z_g g_s(\mu). \quad (3.565)$$

Equivalently, if

$$a_s \equiv \frac{\alpha_s}{4\pi} = \frac{g_s^2}{(4\pi)^2}, \quad (3.566)$$

then

$$a_{s,0} = \mu^{2\epsilon} Z_{a_s} a_s(\mu), \quad Z_{a_s} = Z_g^2. \quad (3.567)$$

At one loop,

$$Z_g = 1 - \frac{1}{2} \frac{g_s^2}{(4\pi)^2} \frac{\beta_0}{\epsilon} + \mathcal{O}(g_s^4), \quad (3.568)$$

where

$$\beta_0 = \frac{11}{3}C_2(G) - \frac{4}{3}T_F n_f = \frac{11}{3}C_2(G) - \frac{2}{3}n_f. \quad (3.569)$$

Here n_f is the number of light quark flavours participating in the running. The first term comes from gluon and ghost effects, especially the non-Abelian self-interaction of gluons. The second term comes from quark loops. Physically, quark loops behave like the screening effect in QED, while gluon self-interactions produce antiscreening. Asymptotic freedom occurs because the gluon contribution dominates.




Figure 3.5: One-loop vacuum-polarization topologies contributing to the QCD beta function. The quark loop gives the $-\frac{2}{3}n_f$ part of β_0 for $SU(3)$; the gluon and ghost sectors together give the positive 11 term.

In the four-dimensional limit, the beta function is defined by

$$\frac{da_s}{d \ln \mu^2} = \beta(a_s). \quad (3.570)$$

Its perturbative expansion is

$$\beta(a_s) = -\beta_0 a_s^2 - \beta_1 a_s^3 + \mathcal{O}(a_s^4), \quad (3.571)$$

with

$$\beta_1 = \frac{34}{3}C_2(G)^2 - \frac{20}{3}C_2(G)T_F n_f - 4C_2(r)T_F n_f. \quad (3.572)$$

For $SU(3)$,

$$\beta_0 = 11 - \frac{2}{3}n_f, \quad \beta_1 = 102 - \frac{38}{3}n_f. \quad (3.573)$$

If one uses α_s rather than $a_s = \alpha_s/(4\pi)$, then

$$\frac{d\alpha_s}{d \ln \mu^2} = -\frac{\beta_0}{4\pi}\alpha_s^2 - \frac{\beta_1}{(4\pi)^2}\alpha_s^3 + \dots. \quad (3.574)$$

5.5 Solving the one-loop running explicitly

Keeping only the one-loop term gives

$$\frac{da_s}{d \ln Q^2} = -\beta_0 a_s^2. \quad (3.575)$$

Separate variables:

$$\frac{da_s}{a_s^2} = -\beta_0 d \ln Q^2. \quad (3.576)$$

Integrating from a reference scale μ to a scale Q gives

$$\int_{a_s(\mu^2)}^{a_s(Q^2)} \frac{da}{a^2} = -\beta_0 \int_{\ln \mu^2}^{\ln Q^2} dL, \quad (3.577)$$

$$-\frac{1}{a_s(Q^2)} + \frac{1}{a_s(\mu^2)} = -\beta_0 \ln \frac{Q^2}{\mu^2}. \quad (3.578)$$

Therefore

$$\frac{1}{a_s(Q^2)} = \frac{1}{a_s(\mu^2)} + \beta_0 \ln \frac{Q^2}{\mu^2}. \quad (3.579)$$

Writing the integration constant as Λ_{QCD} , one obtains

$$a_s(Q^2) = \frac{1}{\beta_0 \ln(Q^2/\Lambda_{\text{QCD}}^2)}. \quad (3.580)$$

Thus

$$\boxed{\alpha_s(Q^2) = \frac{4\pi}{\beta_0 \ln(Q^2/\Lambda_{\text{QCD}}^2)}} \quad \text{one-loop.} \quad (3.581)$$

If an experimental input $\alpha_s(\mu^2)$ is given, then

$$\Lambda_{\text{QCD}} = \mu \exp \left[-\frac{1}{2\beta_0 a_s(\mu^2)} \right] = \mu \exp \left[-\frac{2\pi}{\beta_0 \alpha_s(\mu^2)} \right]. \quad (3.582)$$

In precision applications, one uses higher-loop running and threshold matching. A common reference input is

$$\alpha_s(M_Z) \simeq 0.1180. \quad (3.583)$$

For large Q ,

$$\ln \frac{Q^2}{\Lambda_{\text{QCD}}^2} \rightarrow \infty, \quad \alpha_s(Q^2) \rightarrow 0. \quad (3.584)$$

This is asymptotic freedom: high-energy, short-distance processes are weakly coupled and can be treated perturbatively. When Q approaches Λ_{QCD} , the one-loop expression has a Landau-pole-like singularity. This does not mean that a physical observable literally diverges; it means




Figure 3.6: Perturbative running of $\alpha_s(Q^2)$ at different loop orders. The coupling decreases at large momentum transfer. Image cropped from A. Deur, S. J. Brodsky, and G. F. de Téramond, *The QCD Running Coupling*, arXiv:1604.08082, Fig. 3.1.

that perturbation theory is no longer reliable and nonperturbative confinement physics becomes important.

Asymptotic freedom requires

$$\beta_0 > 0. \quad (3.585)$$

For $SU(3)$,

$$11 - \frac{2}{3}n_f > 0 \quad \implies \quad n_f < \frac{33}{2} = 16.5. \quad (3.586)$$

Thus integer flavour numbers obeying

$$n_f \leq 16 \quad (3.587)$$

lead to one-loop asymptotic freedom. The real world has at most six quark flavours, so real QCD is safely in the asymptotically free regime.

5.6 The basic idea of a conformal window

Including the two-loop term,

$$\beta(a_s) = -\beta_0 a_s^2 - \beta_1 a_s^3 + \dots \quad (3.588)$$

A nonzero fixed point satisfies

$$\beta(a_s^*) = 0 \quad \implies \quad a_s^* = -\frac{\beta_0}{\beta_1}. \quad (3.589)$$

For this fixed point to occur at positive coupling, β_0 and β_1 must have opposite signs. For $SU(3)$,

$$\beta_0 > 0 \quad \iff \quad n_f < 16.5, \quad (3.590)$$

while

$$\beta_1 < 0 \iff 102 - \frac{38}{3}n_f < 0 \iff n_f > \frac{306}{38} \simeq 8.05. \quad (3.591)$$

Therefore two-loop perturbation theory suggests that, roughly for $9 \leq n_f \leq 16$, an infrared fixed point may exist. When n_f is close to 16.5, β_0 is small and a_s^* is also small. This is the Banks–Zaks fixed point, which is perturbatively controlled. The exact lower edge of the conformal window is a nonperturbative question and cannot be fixed reliably by the two-loop formula alone.

5.7 Classical running in dimensional regularization

In dimensional regularization, the coupling already has a mass dimension. This can be seen from the interaction term

$$g_s \bar{q} \gamma^\mu A_\mu q. \quad (3.592)$$

The field dimensions in d dimensions are

$$[q] = \frac{d-1}{2}, \quad [A_\mu] = \frac{d-2}{2}. \quad (3.593)$$

Since the Lagrangian density has mass dimension d ,

$$[g_s] + 2\frac{d-1}{2} + \frac{d-2}{2} = d, \quad (3.594)$$

$$[g_s] = \frac{4-d}{2} = \epsilon. \quad (3.595)$$

Therefore a dimensionless renormalized coupling requires the factor μ^ϵ :

$$g_{s,0} = \mu^\epsilon Z_g g_s(\mu). \quad (3.596)$$

This produces a classical contribution to the d -dimensional beta function:

$$\boxed{\frac{da_s}{d \ln \mu^2} = -\epsilon a_s - \beta_0 a_s^2 - \beta_1 a_s^3 + \dots} \quad (3.597)$$

The term $-\epsilon a_s$ is not a quantum loop effect. It comes from the engineering dimension of the coupling. Only after taking $\epsilon \rightarrow 0$ does one recover the usual four-dimensional QCD running.

5.8 Heavy-quark thresholds and matching of the running coupling

The number n_f in the running coupling is the number of quark flavours that are effectively light at the scale Q . For example,

$$Q \lesssim m_c : \quad n_f = 3 \quad (u, d, s), \quad (3.598)$$

while

$$m_c \lesssim Q \lesssim m_b : \quad n_f = 4 \quad (u, d, s, c). \quad (3.599)$$

When the scale crosses a heavy-quark mass m_h , one must match the coupling in the n_f -flavour effective theory to the coupling in the $(n_f + 1)$ -flavour theory.

At one loop, the matching is determined by the heavy-quark loop contribution to the gluon propagator. Gauge invariance forces the gluon vacuum polarization to be transverse:

$$i\Pi_{\mu\nu}^{ab}(q) = i\delta^{ab} (g_{\mu\nu} q^2 - q_\mu q_\nu) \Pi(q^2). \quad (3.600)$$

First consider n_f massless light quarks. Their loop gives

$$i\Pi_{\mu\nu}^{ab}(q) = -(-1)g_s^2 T_F n_f \delta^{ab} \mu^{2\epsilon} \int \frac{d^d k}{(2\pi)^d} \frac{\text{tr} [\gamma_\mu \not{k} \gamma_\nu (\not{k} + \not{q})]}{k^2 (k+q)^2}. \quad (3.601)$$

The fermion trace is

$$\text{tr} [\gamma_\mu \not{k} \gamma_\nu (\not{k} + \not{q})] = 4 [k_\mu (k+q)_\nu + k_\nu (k+q)_\mu - g_{\mu\nu} k \cdot (k+q)]. \quad (3.602)$$

Introduce a Feynman parameter:

$$\frac{1}{k^2 (k+q)^2} = \int_0^1 dx \frac{1}{[(k+xq)^2 + x(1-x)(-q^2)]^2}. \quad (3.603)$$

After shifting $\ell = k + xq$, all integrals odd in ℓ vanish. Even tensor integrals reduce according to

$$\int d^d \ell \ell_\mu \ell_\nu F(\ell^2) = \frac{g_{\mu\nu}}{d} \int d^d \ell \ell^2 F(\ell^2). \quad (3.604)$$

Only the transverse structure remains. The divergent and logarithmic part of the scalar polarization is

$$\Pi_{\text{light}}^{(n_f)}(q^2) = -\frac{\alpha_s}{4\pi} \frac{4}{3} T_F n_f \left[\frac{1}{\bar{\epsilon}} + \ln \frac{\mu^2}{-q^2} + \text{finite const.} \right], \quad (3.605)$$

or, using $T_F = 1/2$,

$$\Pi_{\text{light}}^{(n_f)}(q^2) = -\frac{\alpha_s}{4\pi} \frac{2}{3} n_f \left[\frac{1}{\bar{\epsilon}} + \ln \frac{\mu^2}{-q^2} + \text{finite const.} \right]. \quad (3.606)$$

Here

$$\frac{1}{\bar{\epsilon}} = \frac{1}{\epsilon} - \gamma_E + \ln 4\pi. \quad (3.607)$$

In handwritten notes one often suppresses scheme-dependent constants and keeps only the pole and logarithm, since these are the terms that determine running and matching.

Now suppose the quark in the loop has mass m_h , and the external momentum is much smaller than that mass:

$$q^2 \ll m_h^2. \quad (3.608)$$

The same Feynman-parameter calculation gives

$$\Pi_h(q^2) = -\frac{\alpha_s}{4\pi} \frac{2}{3} \left[\frac{1}{\bar{\epsilon}} + \ln \frac{\mu^2}{m_h^2} + \mathcal{O}\left(\frac{q^2}{m_h^2}\right) \right]. \quad (3.609)$$

In the $\overline{\text{MS}}$ scheme, after subtracting $1/\bar{\epsilon}$, the heavy-quark loop leaves only a local low-energy correction. It can be absorbed into the definition of the coupling in the low-energy theory. The one-loop decoupling relation is

$$\alpha_s^{(n_f)}(\mu) = \alpha_s^{(n_f+1)}(\mu) \left[1 + \frac{\alpha_s^{(n_f+1)}(\mu)}{6\pi} \ln \frac{\mu^2}{m_h^2} \right] + \mathcal{O}(\alpha_s^3). \quad (3.610)$$

If the matching scale is chosen to be $\mu = m_h$, then at one loop

$$\alpha_s^{(n_f)}(m_h) = \alpha_s^{(n_f+1)}(m_h) + \mathcal{O}(\alpha_s^3). \quad (3.611)$$

Thus the running coupling is continuous across a heavy-quark threshold at one-loop order. At higher orders finite matching corrections remain, so precision QCD requires threshold matching at each heavy-quark mass.




Figure 3.7: Effect of quark thresholds on the running coupling. The red curve illustrates threshold matching, while the black curve shows the discontinuous behaviour that one would obtain from changing the active flavour number without matching. Image cropped from Deur, Brodsky, and de Téramond, arXiv:1604.08082, Fig. 3.2.

5.9 Cross-section predictions and infrared safety

QCD predicts many collider observables, for example

- hadron production in e^+e^- annihilation: LEP, Belle II, BES;
- deep inelastic scattering: HERA, EIC;
- the Drell–Yan process: $pp \rightarrow \gamma^*/Z \rightarrow \ell^+\ell^-$;
- jets, heavy flavour, and vector-boson production at the LHC.

The key requirement is infrared safety: the observable must not change under the emission of a soft gluon or under a collinear splitting.

Consider inclusive $e^+e^- \rightarrow \text{hadrons}$. At leading order,

$$R_0(s) = N_c \sum_q Q_q^2. \quad (3.612)$$

At next-to-leading order there are two classes of corrections:

$$\text{real emission:} \quad e^+e^- \rightarrow q\bar{q}g, \quad (3.613)$$

$$\text{virtual correction:} \quad e^+e^- \rightarrow q\bar{q} \quad \text{with one gluon loop.} \quad (3.614)$$

Each class is infrared divergent by itself. But the inclusive hadronic cross section counts both $q\bar{q}$ and $q\bar{q}g$ final states as hadrons, and the soft and collinear singularities cancel in the sum.

Let a massless quark have momentum

$$p = (E_q, \vec{p}), \quad p^2 = 0, \quad (3.615)$$




Figure 3.8: The two NLO ingredients in inclusive $e^+e^- \rightarrow \text{hadrons}$. Each piece is infrared divergent separately, but the divergences cancel in the inclusive sum.

and let it radiate a gluon

$$k = (E_g, \vec{k}), \quad k^2 = 0. \quad (3.616)$$

If the angle between them is θ_{qg} , then since $|\vec{p}| = E_q$ and $|\vec{k}| = E_g$,

$$2p \cdot k = 2E_q E_g (1 - \cos \theta_{qg}). \quad (3.617)$$

When the gluon is soft,

$$E_g \rightarrow 0, \quad (3.618)$$

this denominator becomes small. When the gluon is collinear to the quark,

$$\theta_{qg} \rightarrow 0, \quad 1 - \cos \theta_{qg} \simeq \frac{\theta_{qg}^2}{2}, \quad (3.619)$$

the denominator also becomes small.

In the soft limit, the squared matrix element factorizes:

$$|\mathcal{M}_{q\bar{q}g}|^2 \simeq |\mathcal{M}_{q\bar{q}}|^2 4\pi\alpha_s C_2(r) \frac{2p_q \cdot p_{\bar{q}}}{(p_q \cdot k)(p_{\bar{q}} \cdot k)}. \quad (3.620)$$

This eikonal factor contains two denominators of the form $p \cdot k$, so it generates soft singularities. In the collinear limit, the propagator

$$(p + k)^2 = p^2 + k^2 + 2p \cdot k = 2p \cdot k \quad (3.621)$$

approaches zero, generating a collinear singularity.

The phase-space measure makes the logarithmic structure explicit. In $d = 4 - 2\epsilon$, a soft gluon has approximately

$$\frac{d^{d-1}k}{2E_g} \sim dE_g E_g^{1-2\epsilon} d\theta \theta^{1-2\epsilon}. \quad (3.622)$$

Multiplying by the soft-collinear eikonal factor

$$\frac{1}{E_g^2 \theta^2}, \quad (3.623)$$

one obtains

$$\int_0^\Lambda dE_g E_g^{-1-2\epsilon} \int_0^\delta d\theta \theta^{-1-2\epsilon}. \quad (3.624)$$

Therefore

$$\int_0^\Lambda dE_g E_g^{-1-2\epsilon} = -\frac{1}{2\epsilon_{\text{IR}}} + \ln \Lambda + \dots, \quad (3.625)$$

$$\int_0^\delta d\theta \theta^{-1-2\epsilon} = -\frac{1}{2\epsilon_{\text{IR}}} + \ln \delta + \dots. \quad (3.626)$$

The energy integral gives a soft pole, and the angular integral gives a collinear pole. Their overlap gives a double pole $1/\epsilon_{\text{IR}}^2$.

The one-loop structure can be summarized as

$$\sigma_{\text{real}} = \sigma_0 \frac{\alpha_s}{4\pi} C_2(r) \left[\frac{2}{\epsilon_{\text{IR}}^2} + \frac{3}{\epsilon_{\text{IR}}} + \text{finite}_{\text{real}} \right], \quad (3.627)$$

$$\sigma_{\text{virtual}} = \sigma_0 \frac{\alpha_s}{4\pi} C_2(r) \left[-\frac{2}{\epsilon_{\text{IR}}^2} - \frac{3}{\epsilon_{\text{IR}}} + \text{finite}_{\text{virtual}} \right]. \quad (3.628)$$

The infrared poles cancel in the sum:

$$\sigma_{\text{NLO}} = \sigma_0 + \sigma_{\text{real}} + \sigma_{\text{virtual}} \quad \text{is finite.} \quad (3.629)$$

This is the KLN theorem at work in the inclusive e^+e^- hadronic cross section.

The finite part gives the famous QCD correction

$$R(s) = N_c \sum_q Q_q^2 \left[1 + \frac{\alpha_s(s)}{4\pi} 3C_2(r) + \mathcal{O}(\alpha_s^2) \right]. \quad (3.630)$$

For QCD, $C_2(r) = 4/3$, so

$$\frac{\alpha_s}{4\pi} 3C_2(r) = \frac{\alpha_s}{4\pi} \cdot 4 = \frac{\alpha_s}{\pi}. \quad (3.631)$$

Thus

$$\boxed{R(s) = N_c \sum_q Q_q^2 \left[1 + \frac{\alpha_s(s)}{\pi} + \mathcal{O}(\alpha_s^2) \right]}. \quad (3.632)$$

This formula ties together the main themes of this section: the number of colors fixes the leading normalization, the coupling $\alpha_s(s)$ runs with the energy scale, and the real and virtual infrared singularities cancel in an inclusive observable.

Standard Model

1. Spontaneous Symmetry Breaking

Spontaneous symmetry breaking is the situation in which the equations and the Lagrangian have a symmetry, but the vacuum chosen by the theory does not. The symmetry is still present in the theory; it is not a mistake in the Lagrangian. The point is that the set of lowest-energy field configurations contains more than one equivalent choice, and choosing one of them hides part of the symmetry.

This section develops the idea in three steps. First we study a real scalar with a discrete symmetry, where there is symmetry breaking but no Goldstone boson. Then we study a complex scalar with a continuous $U(1)$ symmetry, where a massless Goldstone mode appears. Finally we generalize to N complex scalars and prove the Goldstone theorem by counting flat directions of the potential.

1.1 A real scalar with a discrete symmetry

Consider one real scalar field ϕ with Lagrangian

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi), \quad (4.1)$$

and potential

$$V(\phi) = -\frac{1}{2} \mu^2 \phi^2 + \frac{\lambda}{4} \phi^4, \quad \mu^2 > 0, \quad \lambda > 0. \quad (4.2)$$

This is often called the “wrong-sign mass” potential. Around $\phi = 0$ the quadratic term in the potential has negative curvature, so $\phi = 0$ is not a stable vacuum. This corrects a common misleading phrase: the theory does have a quadratic term; what is wrong is the sign of the quadratic term around the origin.

The potential is invariant under the discrete transformation

$$\phi \longrightarrow -\phi, \quad V(\phi) = V(-\phi). \quad (4.3)$$

This is a $\mathbb{Z}_2$ symmetry.

To find the extrema, compute

$$\frac{dV}{d\phi} = -\mu^2 \phi + \lambda \phi^3 = \phi (-\mu^2 + \lambda \phi^2). \quad (4.4)$$

Thus

$$\frac{dV}{d\phi} = 0 \quad \implies \quad \phi = 0 \quad \text{or} \quad \phi^2 = \frac{\mu^2}{\lambda}. \quad (4.5)$$

Define

$$v = \frac{\mu}{\sqrt{\lambda}}. \quad (4.6)$$

The extrema are therefore

$$\phi = 0, \quad \phi = \pm v. \quad (4.7)$$

To decide which are minima, compute the second derivative:

$$\frac{d^2V}{d\phi^2} = -\mu^2 + 3\lambda\phi^2. \quad (4.8)$$

At the origin,

$$\left. \frac{d^2V}{d\phi^2} \right|_{\phi=0} = -\mu^2 < 0, \quad (4.9)$$

so $\phi = 0$ is a local maximum. At $\phi = \pm v$,

$$\left. \frac{d^2V}{d\phi^2} \right|_{\phi=\pm v} = -\mu^2 + 3\lambda\frac{\mu^2}{\lambda} = 2\mu^2 > 0, \quad (4.10)$$

so $\phi = \pm v$ are true minima.




Figure 4.1: The double-well potential. The Lagrangian is symmetric under $\phi \rightarrow -\phi$, but either vacuum $\phi = +v$ or $\phi = -v$ chooses one side.

The theory has the symmetry $\phi \rightarrow -\phi$, but the vacuum does not. If the system chooses

$$\langle 0|\phi|0\rangle = +v, \quad (4.11)$$

then the transformed configuration has

$$\langle 0|\phi|0\rangle \longrightarrow -v, \quad (4.12)$$

which is a different vacuum. The two vacua are physically equivalent, but once one is chosen the symmetry is no longer manifest. This is spontaneous breaking of a discrete symmetry.

There is no Goldstone boson in this example. Goldstone bosons arise from broken continuous symmetries, where moving along the vacuum manifold can be done infinitesimally with no energy cost. A discrete symmetry does not provide such a continuous flat direction.

1.2 A complex scalar and global U(1) breaking

Now consider a complex scalar field

$$\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2), \quad (4.13)$$

with Lagrangian

$$\mathcal{L} = \partial_\mu \phi^* \partial^\mu \phi - V(\phi, \phi^*), \quad (4.14)$$

and potential

$$V(\phi, \phi^*) = -\mu^2 \phi^* \phi + \frac{\lambda}{2} (\phi^* \phi)^2, \quad \mu^2 > 0, \quad \lambda > 0. \quad (4.15)$$

The Lagrangian is invariant under the global $U(1)$ transformation

$$\phi(x) \longrightarrow \phi'(x) = e^{i\alpha} \phi(x), \quad \alpha \in \mathbb{R}. \quad (4.16)$$

This is a global symmetry because α is constant in spacetime.

The potential depends only on the radial variable $\phi^* \phi$. Minimizing with respect to ϕ^* gives

$$\frac{\partial V}{\partial \phi^*} = \phi (-\mu^2 + \lambda \phi^* \phi) = 0. \quad (4.17)$$

Therefore

$$\phi = 0 \quad \text{or} \quad \phi^* \phi = \frac{\mu^2}{\lambda}. \quad (4.18)$$

Again $\phi = 0$ is the unstable point. The true vacua satisfy

$$|\phi|^2 = \frac{\mu^2}{\lambda}. \quad (4.19)$$

In the complex plane this is a circle. Since

$$|\phi|^2 = \frac{1}{2} (\phi_1^2 + \phi_2^2), \quad (4.20)$$

the vacuum circle can also be written as

$$\phi_1^2 + \phi_2^2 = \frac{2\mu^2}{\lambda}. \quad (4.21)$$

It is conventional to define

$$v^2 = \frac{2\mu^2}{\lambda}. \quad (4.22)$$

Then the vacuum condition is

$$|\phi|^2 = \frac{v^2}{2}. \quad (4.23)$$




Figure 4.2: The vacuum manifold for the broken global $U(1)$ model. Choosing one point on the circle breaks the manifest $U(1)$ symmetry.

Every point on this circle is a possible ground state. Choosing one point breaks the global $U(1)$ symmetry. We may choose the vacuum to lie along the real direction:

$$\phi_0 = \frac{v}{\sqrt{2}}, \quad \phi_{1,0} = v, \quad \phi_{2,0} = 0. \quad (4.24)$$

This choice is not a loss of generality; any other point on the circle can be rotated into this one by a $U(1)$ transformation.

1.3 Expansion around the chosen vacuum

To find the physical particles, expand the field around the chosen vacuum:

$$\phi(x) = \frac{1}{\sqrt{2}} [v + h(x) + i\sigma(x)]. \quad (4.25)$$

Here $h(x)$ is the radial fluctuation and $\sigma(x)$ is the angular fluctuation. At the vacuum,

$$h = 0, \quad \sigma = 0. \quad (4.26)$$

The kinetic term becomes

$$\begin{aligned} \partial_\mu \phi^* \partial^\mu \phi &= \frac{1}{2} \partial_\mu (v + h - i\sigma) \partial^\mu (v + h + i\sigma) \\ &= \frac{1}{2} \partial_\mu h \partial^\mu h + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma. \end{aligned} \quad (4.27)$$

Since v is constant, its derivative vanishes.

Now compute the potential in terms of h and σ . First,

$$\begin{aligned} \phi^* \phi &= \frac{1}{2} [(v + h)^2 + \sigma^2] \\ &= \frac{1}{2} [v^2 + 2vh + h^2 + \sigma^2]. \end{aligned} \quad (4.28)$$

Therefore

$$V(h, \sigma) = -\frac{\mu^2}{2} [(v + h)^2 + \sigma^2] + \frac{\lambda}{8} [(v + h)^2 + \sigma^2]^2. \quad (4.29)$$

Expanding the square,

$$\begin{aligned} [(v + h)^2 + \sigma^2]^2 &= (v^2 + 2vh + h^2 + \sigma^2)^2 \\ &= v^4 + 4v^3h + 6v^2h^2 + 2v^2\sigma^2 + \mathcal{O}(\text{cubic}). \end{aligned} \quad (4.30)$$

Hence the terms up to quadratic order are

$$V(h, \sigma) = -\frac{\mu^2}{2} (v^2 + 2vh + h^2 + \sigma^2) + \frac{\lambda}{8} (v^4 + 4v^3h + 6v^2h^2 + 2v^2\sigma^2) + \dots \quad (4.31)$$

Separate constant, linear, and quadratic terms:

$$\begin{aligned} V(h, \sigma) &= \left(-\frac{\mu^2 v^2}{2} + \frac{\lambda v^4}{8} \right) + \left(-\mu^2 v + \frac{\lambda v^3}{2} \right) h \\ &\quad + \left(-\frac{\mu^2}{2} + \frac{3\lambda v^2}{4} \right) h^2 + \left(-\frac{\mu^2}{2} + \frac{\lambda v^2}{4} \right) \sigma^2 + \dots \end{aligned} \quad (4.32)$$

Using

$$v^2 = \frac{2\mu^2}{\lambda}, \quad (4.33)$$

the linear term vanishes:

$$-\mu^2 v + \frac{\lambda v^3}{2} = v \left(-\mu^2 + \frac{\lambda v^2}{2} \right) = 0. \quad (4.34)$$

This must happen because we expanded around a minimum. The quadratic terms become

$$-\frac{\mu^2}{2} + \frac{3\lambda v^2}{4} = -\frac{\mu^2}{2} + \frac{3\mu^2}{2} = \mu^2, \quad (4.35)$$

$$-\frac{\mu^2}{2} + \frac{\lambda v^2}{4} = -\frac{\mu^2}{2} + \frac{\mu^2}{2} = 0. \quad (4.36)$$

Thus

$$V(h, \sigma) = V(v) + \mu^2 h^2 + \mathcal{O}(h^3, h^2\sigma, h\sigma^2, \sigma^4). \quad (4.37)$$

The Lagrangian contains $-\mu^2 h^2$. Comparing with the standard mass term

$$-\frac{1}{2} m_h^2 h^2, \quad (4.38)$$

we find

$$m_h^2 = 2\mu^2. \quad (4.39)$$

There is no quadratic term for σ , so

$$m_\sigma^2 = 0. \quad (4.40)$$

The radial field h is massive. The angular field σ is massless and is called the Goldstone boson.

The reason is geometric. The h direction moves away from the vacuum circle, so the potential increases. The σ direction moves along the vacuum circle, where the potential is flat. A flat direction has zero curvature, and zero curvature means zero mass squared.

1.4 N complex scalar fields

Now consider N complex scalar fields ϕ^i , $i = 1, \dots, N$, with

$$\mathcal{L} = \sum_{i=1}^N \partial_\mu \phi^{i*} \partial^\mu \phi^i - V(\phi, \phi^*), \quad (4.41)$$

and

$$V(\phi, \phi^*) = -\mu^2 \sum_{i=1}^N \phi^{i*} \phi^i + \frac{\lambda}{2} \left(\sum_{i=1}^N \phi^{i*} \phi^i \right)^2. \quad (4.42)$$

This potential is invariant under global $U(N)$ transformations:

$$\phi^i \longrightarrow \phi'^i = \sum_{j=1}^N U^{ij} \phi^j, \quad U^\dagger U = \mathbf{1}. \quad (4.43)$$

The quantity

$$\sum_i \phi^{i*} \phi^i \quad (4.44)$$

is just the squared length of the complex vector ϕ , so it is unchanged by unitary rotations.

The minimum satisfies

$$\sum_{i=1}^N \phi^{i*} \phi^i = \frac{\mu^2}{\lambda}. \quad (4.45)$$

This is a sphere in the $2N$ real-dimensional field space. Define again

$$\frac{v^2}{2} = \frac{\mu^2}{\lambda}, \quad v^2 = \frac{2\mu^2}{\lambda}. \quad (4.46)$$

Using the $U(N)$ symmetry, choose the vacuum to point in the first complex direction:

$$\phi_0^1 = \frac{v}{\sqrt{2}}, \quad \phi_0^2 = \cdots = \phi_0^N = 0. \quad (4.47)$$

Then write the fields as

$$\phi^1 = \frac{1}{\sqrt{2}} [v + h(x) + i\sigma(x)], \quad (4.48)$$

$$\phi^k = \frac{1}{\sqrt{2}} [\pi_{2k-2}(x) + i\pi_{2k-1}(x)], \quad k = 2, \dots, N. \quad (4.49)$$

There are $2N$ real fields in total:

$$h, \quad \sigma, \quad \pi_2, \dots, \pi_{2N-1}. \quad (4.50)$$

The radial field h changes the length of the vector ϕ and is massive. All other fields change only the direction of ϕ in the vacuum manifold and are massless at tree level.

Equivalently, the chosen vacuum leaves a residual $U(N-1)$ symmetry acting on the fields

$$\phi^2, \dots, \phi^N. \quad (4.51)$$

Thus the symmetry-breaking pattern is

$$U(N) \longrightarrow U(N-1). \quad (4.52)$$

The number of broken generators is

$$\dim U(N) - \dim U(N-1) = N^2 - (N-1)^2 = 2N-1. \quad (4.53)$$

Therefore there are

$$2N-1 \quad (4.54)$$

massless Goldstone bosons. This agrees with the field decomposition above: σ plus the $2N-2$ real components of $\phi^2, \dots, \phi^N$.

1.5 Goldstone theorem

The examples above illustrate the general result.

Goldstone theorem. In a relativistic theory with a continuous global internal symmetry, each spontaneously broken generator gives one massless real scalar excitation, called a Goldstone boson.

There are important assumptions in this statement. The symmetry must be global, not gauged. If the symmetry is gauged, the massless scalar is removed from the physical spectrum by the Higgs mechanism. Also, the symmetry must be exact. If it is explicitly broken, the would-be Goldstone boson generally becomes light but not massless.

1.6 Proof by the mass matrix

Consider N real scalar fields ϕ^i with Lagrangian

$$\mathcal{L} = \frac{1}{2} \sum_{i=1}^N \partial_\mu \phi^i \partial^\mu \phi^i - V(\phi). \quad (4.55)$$

Suppose the potential is invariant under a continuous symmetry group G . For an infinitesimal transformation,

$$\phi^i \longrightarrow \phi^i + \delta_a \phi^i, \quad (4.56)$$

where

$$\delta_a \phi^i = \alpha^a (T_a)^i_j \phi^j. \quad (4.57)$$

Here T_a is a generator of the symmetry and α^a is an infinitesimal real parameter. Since V is invariant,

$$\delta_a V = \sum_i \frac{\partial V}{\partial \phi^i} \delta_a \phi^i = 0. \quad (4.58)$$

Substitute the infinitesimal transformation:

$$\sum_i \frac{\partial V}{\partial \phi^i} (T_a)^i_j \phi^j = 0. \quad (4.59)$$

This identity holds for every field configuration ϕ , not only at the minimum.

Now differentiate this identity with respect to ϕ^k :

$$\sum_i \frac{\partial^2 V}{\partial \phi^k \partial \phi^i} (T_a)^i_j \phi^j + \sum_i \frac{\partial V}{\partial \phi^i} (T_a)^i_k = 0. \quad (4.60)$$

Evaluate at a vacuum ϕ_0 . Since ϕ_0 is a minimum,

$$\left. \frac{\partial V}{\partial \phi^i} \right|_{\phi_0} = 0. \quad (4.61)$$

The second term therefore vanishes. Define the mass matrix

$$(M^2)_{ki} = \left. \frac{\partial^2 V}{\partial \phi^k \partial \phi^i} \right|_{\phi_0}. \quad (4.62)$$

Then

$$\sum_i (M^2)_{ki} (T_a)^i_j \phi_0^j = 0. \quad (4.63)$$

In vector notation this is

$$M^2 (T_a \phi_0) = 0. \quad (4.64)$$

Now distinguish two cases.

Unbroken generator.

If T_a is unbroken, it leaves the vacuum invariant:

$$T_a \phi_0 = 0. \quad (4.65)$$

Then the equation above is trivial and gives no new massless field.

Broken generator.

If T_a is broken, then it moves the vacuum:

$$T_a \phi_0 \neq 0. \quad (4.66)$$

But the mass-matrix equation says

$$M^2 (T_a \phi_0) = 0. \quad (4.67)$$

Thus $T_a \phi_0$ is a nonzero eigenvector of the mass matrix with eigenvalue zero. A zero eigenvalue of the scalar mass matrix means a massless scalar mode. Therefore each broken generator gives a massless scalar field.

This is the mathematical form of the geometric picture. Broken generators move the vacuum along the manifold of degenerate minima. Motion along that manifold costs no potential energy, so the corresponding direction has zero curvature. Zero curvature of the potential is precisely zero mass squared.

2. The Higgs Mechanism

2.1 Difference between Higgs mechanism and Goldstone's theorem

As we have discussed in last section, Goldstone's theorem should be applied directly only to a *global* continuous symmetry. Suppose a continuous global symmetry is a true physical symmetry of the Lagrangian, but the vacuum does not respect it. Then each broken generator produces one physical massless scalar particle, called a Goldstone boson.

The Higgs mechanism uses a scalar potential with the same geometric shape, but now the symmetry is *local*. A local gauge symmetry is not an ordinary physical symmetry that relates two different states; it is a redundancy in our description of the same physical state. For this reason, Goldstone's theorem does not say that a broken gauge symmetry must produce physical massless particles.

The correct way to think about the Higgs mechanism is the following. First, temporarily ignore the gauge field. The scalar potential then has flat angular directions, and fluctuations along those directions would be Goldstone bosons in the corresponding global theory. After the symmetry is gauged, those same angular fields are no longer physical particles. They can be removed by a gauge choice and become the longitudinal polarizations of the massive gauge bosons. This is why they are called *would-be Goldstone bosons*.

Thus, when we later write a phase field such as $\theta(x)$, we are not claiming that θ is a physical massless Goldstone particle in the gauge theory. We are identifying the angular scalar mode which would have been a Goldstone boson before gauging, and then showing explicitly how the gauge field absorbs it.

In short,

$$\begin{aligned} \text{global spontaneous symmetry breaking} &\implies \text{physical massless Goldstone bosons,} \\ \text{gauged version of the same scalar theory} &\implies \text{would-be Goldstone modes are eaten.} \end{aligned} \quad (4.68)$$

The result is a massive gauge field. Its extra longitudinal polarization comes from the scalar mode which would have been a Goldstone boson in the ungauged global theory.

2.2 The Abelian Higgs model

We first study the simplest example: a complex scalar field coupled to a $U(1)$ gauge field. The Lagrangian is

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu\phi)^*(D^\mu\phi) - V(\phi), \quad D_\mu = \partial_\mu + ieA_\mu, \quad (4.69)$$

with

$$V(\phi) = -\mu^2|\phi|^2 + \frac{\lambda}{2}|\phi|^4, \quad \mu^2 > 0, \quad \lambda > 0. \quad (4.70)$$

Equivalently, the scalar part of the Lagrangian contains

$$\mathcal{L} \supset (D_\mu\phi)^*(D^\mu\phi) + \mu^2|\phi|^2 - \frac{\lambda}{2}|\phi|^4. \quad (4.71)$$

Notice that the sign of quadratic term is positive, which means this is not the mass term yet. The Lagrangian is invariant under the local transformation

$$\phi(x) \longrightarrow e^{i\alpha(x)}\phi(x), \quad A_\mu(x) \longrightarrow A_\mu(x) - \frac{1}{e}\partial_\mu\alpha(x). \quad (4.72)$$

The reason for introducing A_μ is exactly to make derivatives compatible with a position-dependent phase. Indeed,

$$D_\mu\phi = (\partial_\mu + ieA_\mu)\phi \implies D_\mu\phi \longrightarrow e^{i\alpha(x)}D_\mu\phi, \quad (4.73)$$

so $(D_\mu\phi)^*(D^\mu\phi)$ is gauge invariant.

The vacuum

The vacuum is found by minimizing V . Since V depends only on $|\phi|$, write $r = |\phi|$. Then

$$V(r) = -\mu^2r^2 + \frac{\lambda}{2}r^4. \quad (4.74)$$

The stationary condition is

$$\frac{dV}{dr} = -2\mu^2r + 2\lambda r^3 = 2r(-\mu^2 + \lambda r^2) = 0. \quad (4.75)$$

Thus either $r = 0$, or

$$r^2 = \frac{\mu^2}{\lambda}. \quad (4.76)$$

The point $r = 0$ is a local maximum, because

$$\left.\frac{d^2V}{dr^2}\right|_{r=0} = -2\mu^2 < 0. \quad (4.77)$$

Therefore the minima lie on the circle

$$|\phi| = v, \quad v^2 = \frac{\mu^2}{\lambda}. \quad (4.78)$$

This circle of minima is the familiar “Mexican-hat” vacuum manifold. The radial direction costs energy, while moving around the circle costs no potential energy before the field is coupled to a gauge field.

Polar form and unitary gauge

Near the vacuum it is useful to separate radial and angular fluctuations:

$$\phi(x) = \left(v + \frac{h(x)}{\sqrt{2}} \right) e^{i\theta(x)}. \quad (4.79)$$

Here $h(x)$ is the radial fluctuation, while $\theta(x)$ is the angular coordinate along the circle of classical minima. It is important not to misread this sentence: in the local $U(1)$ theory, $\theta(x)$ is not a physical Goldstone particle. Rather, it is the field that *would have been* the Goldstone boson if the same scalar potential had only a global $U(1)$ symmetry. Because the $U(1)$ symmetry is gauged here, $\theta(x)$ is a would-be Goldstone mode and can be removed by a gauge transformation.

To see this explicitly, substitute (4.79) into the covariant derivative:

$$D_\mu \phi = e^{i\theta} \left[\frac{1}{\sqrt{2}} \partial_\mu h + i \left(v + \frac{h}{\sqrt{2}} \right) (\partial_\mu \theta + e A_\mu) \right]. \quad (4.80)$$

The phase $e^{i\theta}$ drops out of the absolute value, so

$$|D_\mu \phi|^2 = \frac{1}{2} (\partial_\mu h)(\partial^\mu h) + e^2 \left(v + \frac{h}{\sqrt{2}} \right)^2 B_\mu B^\mu, \quad B_\mu \equiv A_\mu + \frac{1}{e} \partial_\mu \theta. \quad (4.81)$$

The combination B_μ is what remains after choosing unitary gauge, where the scalar is made real:

$$\phi(x) \longrightarrow v + \frac{h(x)}{\sqrt{2}}, \quad A_\mu(x) \longrightarrow B_\mu(x). \quad (4.82)$$

Thus the angular field θ has not disappeared mysteriously; it has become part of the gauge field.

Expanding the second term in (4.81) gives

$$e^2 \left(v + \frac{h}{\sqrt{2}} \right)^2 B_\mu B^\mu = e^2 v^2 B_\mu B^\mu + \sqrt{2} e^2 v h B_\mu B^\mu + \frac{e^2}{2} h^2 B_\mu B^\mu. \quad (4.83)$$

The first term is a mass term for the vector field. A massive vector mass term is conventionally written as

$$\frac{1}{2} m_A^2 B_\mu B^\mu. \quad (4.84)$$

Comparing this with $e^2 v^2 B_\mu B^\mu$, we obtain

$$m_A^2 = 2e^2 v^2. \quad (4.85)$$

The remaining terms in (4.83) are interactions between the Higgs fluctuation h and the massive vector boson.

The radial scalar mass follows from the potential. Put

$$r = v + \frac{h}{\sqrt{2}}. \quad (4.86)$$

Since $V'(v) = 0$, the leading nonconstant term is

$$V(r) = V(v) + \frac{1}{2} V''(v) \left(\frac{h}{\sqrt{2}} \right)^2 + \dots \quad (4.87)$$

Now

$$V''(r) = -2\mu^2 + 6\lambda r^2, \quad V''(v) = -2\mu^2 + 6\lambda v^2 = 4\mu^2, \quad (4.88)$$

where $v^2 = \mu^2/\lambda$ has been used. Hence

$$V(r) = V(v) + \mu^2 h^2 + \dots \quad (4.89)$$

Because the Lagrangian contains $-V$, the quadratic scalar term is

$$-\mu^2 h^2 = -\frac{1}{2}(2\mu^2)h^2, \quad (4.90)$$

so

$$m_h^2 = 2\mu^2. \quad (4.91)$$

Counting degrees of freedom

Before the Higgs mechanism, the complex scalar has two real degrees of freedom and the massless gauge boson has two transverse polarizations:

$$2 + 2 = 4. \quad (4.92)$$

After choosing the vacuum, only the radial scalar h remains as a physical scalar. The gauge boson is now massive, so it has three polarizations:

$$1 + 3 = 4. \quad (4.93)$$

No physical degree of freedom is lost. The would-be Goldstone mode supplies the longitudinal polarization of the massive vector boson.

	scalar degrees of freedom	vector degrees of freedom	total
before	2	2	4
after	1	3	4

2.3 The same physics in Cartesian fields

The polar form makes the Higgs mechanism transparent, but it is also useful to see how the result appears if we expand the complex scalar into two real fields:

$$\phi(x) = v + \frac{1}{\sqrt{2}}(\phi_1(x) + i\phi_2(x)), \quad \phi_1, \phi_2 \in \mathbb{R}. \quad (4.94)$$

Here ϕ_1 is the radial fluctuation and ϕ_2 is the angular fluctuation to leading order.

To derive the quadratic part of the potential, first compute

$$|\phi|^2 = \left(v + \frac{\phi_1}{\sqrt{2}}\right)^2 + \left(\frac{\phi_2}{\sqrt{2}}\right)^2 = v^2 + \sqrt{2}v\phi_1 + \frac{1}{2}(\phi_1^2 + \phi_2^2). \quad (4.95)$$

It is useful to write this as

$$|\phi|^2 = v^2 + \Delta, \quad \Delta = \sqrt{2}v\phi_1 + \frac{1}{2}(\phi_1^2 + \phi_2^2). \quad (4.96)$$

Substituting this into $V = -\mu^2|\phi|^2 + \frac{\lambda}{2}|\phi|^4$ gives

$$\begin{aligned} V &= -\mu^2(v^2 + \Delta) + \frac{\lambda}{2}(v^2 + \Delta)^2 \\ &= V(v) + (-\mu^2 + \lambda v^2)\Delta + \frac{\lambda}{2}\Delta^2. \end{aligned} \quad (4.97)$$

The coefficient of Δ vanishes because the vacuum satisfies $v^2 = \mu^2/\lambda$. Therefore there is no linear term, and the quadratic part comes only from Δ^2 . To second order in the fluctuating fields,

$$\Delta^2 = \left(\sqrt{2}v\phi_1\right)^2 + \text{terms of cubic or higher order} = 2v^2\phi_1^2 + \dots \quad (4.98)$$

Using $v^2 = \mu^2/\lambda$, the potential therefore expands as

$$V(\phi) = V(v) + \mu^2\phi_1^2 + \text{interaction terms}. \quad (4.99)$$

Therefore

$$-V(\phi) = -V(v) - \mu^2\phi_1^2 + \text{interaction terms}. \quad (4.100)$$

This gives

$$m_{\phi_1}^2 = 2\mu^2, \quad m_{\phi_2}^2 = 0 \quad (4.101)$$

before gauge fixing. The absence of a ϕ_2^2 term is the usual Goldstone-boson result for the scalar potential. In a gauge theory this field will mix with the gauge field and is not a physical massless particle.

Now expand the covariant derivative:

$$\begin{aligned} D_\mu\phi &= \partial_\mu\phi + ieA_\mu\phi \\ &= \frac{1}{\sqrt{2}}(\partial_\mu\phi_1 + i\partial_\mu\phi_2) + ievA_\mu + \frac{ie}{\sqrt{2}}A_\mu(\phi_1 + i\phi_2). \end{aligned} \quad (4.102)$$

Keeping only terms quadratic in the fields gives

$$|D_\mu\phi|^2 = \frac{1}{2}(\partial_\mu\phi_1)^2 + \frac{1}{2}(\partial_\mu\phi_2)^2 + \sqrt{2}ev A^\mu\partial_\mu\phi_2 + e^2v^2 A_\mu A^\mu + \dots \quad (4.103)$$

The last term is the same gauge-boson mass term as before:

$$e^2v^2 A_\mu A^\mu = \frac{1}{2}m_A^2 A_\mu A^\mu, \quad m_A^2 = 2e^2v^2. \quad (4.104)$$

The mixed term

$$\sqrt{2}ev A^\mu\partial_\mu\phi_2 \quad (4.105)$$

is the Cartesian-field signal that ϕ_2 is not an independent physical particle. In unitary gauge it is removed. In a covariant R_ξ gauge it is cancelled by a gauge-fixing term, leaving a gauge-dependent unphysical would-be Goldstone field whose effects cancel from physical amplitudes.

2.4 What the propagator is telling us

Now we shall top for a moment and discuss how “the gauge field eats the would-be Goldstone mode”. And what really shows up in the propagator of the gauge field.

Recall the degree-of-freedom count:

$$\text{massless vector: 2 polarizations,} \quad \text{massive vector: 3 polarizations.} \quad (4.106)$$

The new third polarization is the longitudinal polarization. The question is: where does it come from? The answer is that it comes from the angular scalar mode, namely the would-be Goldstone field.

In unitary gauge the would-be Goldstone field has been removed from the scalar field, so it is hidden inside the gauge field. The massive-vector propagator then takes the form

$$D_{\mu\nu}^{\text{unitary}}(k) = \frac{-i}{k^2 - m_A^2 + i\epsilon} \left(g_{\mu\nu} - \frac{k_\mu k_\nu}{m_A^2} \right). \quad (4.107)$$

The term $k_\mu k_\nu / m_A^2$ is the part associated with the longitudinal polarization. Therefore this propagator is the unitary-gauge way of saying: the vector field now contains a longitudinal mode.

The same physics can be seen before choosing unitary gauge. In the Cartesian expansion, the quadratic Lagrangian contains the mixing term

$$\sqrt{2}ev A^\mu \partial_\mu \phi_2. \quad (4.108)$$

Using $m_A = \sqrt{2}ev$, this is

$$m_A A^\mu \partial_\mu \phi_2. \quad (4.109)$$

The derivative gives a momentum factor k_μ , so the mixing between A_μ and the would-be Goldstone field carries a factor schematically like $m_A k_\mu$. This is why a would-be Goldstone exchange naturally produces terms proportional to $k_\mu k_\nu$, exactly the tensor structure needed for the longitudinal part of a vector propagator.

Schematically, the gauge-boson mass insertion gives a contribution

$$im_A^2 g_{\mu\nu}. \quad (4.110)$$

The exchange of the would-be Goldstone mode gives a longitudinal contribution of the form

$$m_A k_\mu \frac{i}{k^2} (-m_A k_\nu) = -im_A^2 \frac{k_\mu k_\nu}{k^2}. \quad (4.111)$$




Figure 4.3: The mass insertion and the would-be Goldstone exchange combine to produce the longitudinal structure of the massive gauge boson. This is a schematic diagram; the precise signs depend on the gauge-fixing convention.

Together they form the transverse structure

$$im_A^2 \left(g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right). \quad (4.112)$$

This equation should be read only as a schematic explanation, not as a complete propagator formula. The lesson is simple: the same scalar mode that disappears in unitary gauge is precisely the mode that supplies the longitudinal polarization of the massive gauge boson. Thus the phrase “the gauge boson eats the Goldstone boson” means:

$$A_\mu \text{ with 2 polarizations} + 1 \text{ would-be Goldstone mode} \implies \text{massive } A_\mu \text{ with 3 polarizations.} \quad (4.113)$$

2.5 $SU(2)$ gauge symmetry breaking

The non-Abelian version uses a complex scalar doublet

$$\Phi = \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}, \quad \Phi_1, \Phi_2 \in \mathbb{C}, \quad (4.114)$$

coupled to an $SU(2)$ gauge field. The Lagrangian is

$$\mathcal{L} = -\frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} + (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi), \quad (4.115)$$

where

$$V(\Phi) = -\mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2, \quad \mu^2 > 0, \quad \lambda > 0, \quad (4.116)$$

and

$$D_\mu = \partial_\mu - ig \frac{\sigma^a}{2} W_\mu^a. \quad (4.117)$$

The σ^a are the Pauli matrices, and repeated adjoint indices $a = 1, 2, 3$ are summed.

The potential depends only on $\Phi^\dagger \Phi$. Let

$$\rho^2 = \Phi^\dagger \Phi. \quad (4.118)$$

Then

$$V(\rho) = -\mu^2 \rho^2 + \lambda \rho^4. \quad (4.119)$$

The minimum satisfies

$$\frac{dV}{d\rho} = -2\mu^2 \rho + 4\lambda \rho^3 = 2\rho(-\mu^2 + 2\lambda \rho^2) = 0. \quad (4.120)$$

Thus the nonzero vacuum satisfies

$$\Phi^\dagger \Phi = \rho^2 = \frac{\mu^2}{2\lambda}. \quad (4.121)$$

It is conventional to write

$$\langle \Phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad v^2 = \frac{\mu^2}{\lambda}. \quad (4.122)$$

Indeed,

$$\langle \Phi \rangle^\dagger \langle \Phi \rangle = \frac{v^2}{2} = \frac{\mu^2}{2\lambda}. \quad (4.123)$$

Notice the dimensions: $\Phi^\dagger \Phi$ has dimension two, so it is proportional to v^2 , not to v .

Which generators are broken?

For this pure $SU(2)$ theory, all three generators are broken by the vacuum. Acting with the Pauli matrices gives

$$\sigma^1 \begin{pmatrix} 0 \\ v \end{pmatrix} = \begin{pmatrix} v \\ 0 \end{pmatrix}, \quad \sigma^2 \begin{pmatrix} 0 \\ v \end{pmatrix} = \begin{pmatrix} -iv \\ 0 \end{pmatrix}, \quad \sigma^3 \begin{pmatrix} 0 \\ v \end{pmatrix} = \begin{pmatrix} 0 \\ -v \end{pmatrix}. \quad (4.124)$$

None of these is zero, so the vacuum is moved by every generator. Hence the three would-be Goldstone fields are absorbed by the three $SU(2)$ gauge bosons.

This statement is specific to a pure $SU(2)$ gauge theory with one scalar doublet. In the Standard Model the gauge group is $SU(2)_L \times U(1)_Y$, and one linear combination remains unbroken; that unbroken combination is electromagnetism.

Unitary gauge and the gauge-boson masses

The most general fluctuation around the chosen vacuum may be written as

$$\Phi(x) = \exp\left[\frac{i}{2}\sigma^a\theta^a(x)\right] \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}. \quad (4.125)$$

The three fields $\theta^a(x)$ are the would-be Goldstone fields. An $SU(2)$ gauge transformation can remove them, leaving unitary gauge:

$$\Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}. \quad (4.126)$$

In matrix notation,

$$W_\mu = W_\mu^a \frac{\sigma^a}{2}. \quad (4.127)$$

Under a local $SU(2)$ transformation $U(x)$, the gauge field transforms as

$$W_\mu \longrightarrow UW_\mu U^\dagger - \frac{i}{g}U\partial_\mu U^\dagger, \quad (4.128)$$

with the sign matching the convention

$$D_\mu = \partial_\mu - igW_\mu. \quad (4.129)$$

Now compute the covariant derivative in unitary gauge. Since

$$\sigma^1 \begin{pmatrix} 0 \\ v + h \end{pmatrix} = \begin{pmatrix} v + h \\ 0 \end{pmatrix}, \quad \sigma^2 \begin{pmatrix} 0 \\ v + h \end{pmatrix} = \begin{pmatrix} -i(v + h) \\ 0 \end{pmatrix}, \quad \sigma^3 \begin{pmatrix} 0 \\ v + h \end{pmatrix} = \begin{pmatrix} 0 \\ -(v + h) \end{pmatrix}, \quad (4.130)$$

we find

$$D_\mu \Phi = \frac{1}{\sqrt{2}} \begin{pmatrix} -\frac{ig}{2}(v + h)(W_\mu^1 - iW_\mu^2) \\ \partial_\mu h + \frac{ig}{2}(v + h)W_\mu^3 \end{pmatrix}. \quad (4.131)$$

Taking the Hermitian norm gives

$$(D_\mu \Phi)^\dagger (D^\mu \Phi) = \frac{1}{2}(\partial_\mu h)(\partial^\mu h) + \frac{g^2}{8}(v + h)^2 W_\mu^a W^{a\mu}. \quad (4.132)$$

The gauge-boson mass term is therefore

$$\frac{g^2 v^2}{8} W_\mu^a W^{a\mu}. \quad (4.133)$$

Since the canonical mass term for each real vector boson is

$$\frac{1}{2}m_W^2 W_\mu^a W^{a\mu}, \quad (4.134)$$

we obtain

$$m_W^2 = \frac{g^2 v^2}{4}. \quad (4.135)$$

All three gauge bosons have the same mass because the original theory has an $SU(2)$ symmetry and the scalar representation treats the three generators symmetrically after all of them are broken.

Degree-of-freedom count for the $SU(2)$ model

The complex doublet contains four real scalar degrees of freedom. Before the Higgs mechanism, the three massless gauge bosons each have two transverse polarizations:

$$4 + 3 \times 2 = 10. \quad (4.136)$$

After the Higgs mechanism, one scalar h remains, and the three gauge bosons are massive, each with three polarizations:

$$1 + 3 \times 3 = 10. \quad (4.137)$$

Again the number of physical degrees of freedom is unchanged. The three would-be Goldstone modes have become the longitudinal polarizations of the three massive gauge bosons.

	scalar degrees of freedom	vector degrees of freedom	total
before	4	$3 \times 2 = 6$	10
after	1	$3 \times 3 = 9$	10