# Plant Stress Detection Based on Spectral Imaging and Deep Learning

Xiaosheng Luo

Imperial College London

Supervisor: Oliver Windram, Department of Life Sciences (Silwood Park), Imperial College London

### Introduction

- Organic growers face a number of challenges they must overcome to deliver high quality food to their
- 2 customers. The lack of reactive chemical control measures in organic farming requires additional levels
- 3 of insight in order to optimise plant and hervest schedule. Traditional manual detection can only wait
- 4 until late crop stress morphological changes, and is generally time-consuming and subjective. However,
- 5 using spectral imaging technology, the differences in crop physiological structure characteristics will
- 6 lead to differences in light reflection, absorption and transmission. Studying these differences in
- <sup>7</sup> spectral imaging can help identify the crop growth status, nutrient and disease.
- 8 Previous studies have made extensive fundamental research on crop detection using spectral technol-
- 9 ogy, mainly based on spectral index features and image features. For instance, the olive Verticillium
- wilt can be diagnosed by images collected from UAV-mounted multi-spectral camera and thermal
- infrared camera, and it was found that early Verticillium wilt was related to green light band, and
- chlorophyll fluorescence index decreased with the disease aggravating (?). Moreover, applied hyper-
- 13 spectral imaging techniques to identify the angular leaf spot of cucumber by detecting the contents of
- 14 chlorophyll and carotenoids showed the feasible for visualizing the pigment distribution in cucumber
- leaves in response to angular leaf spot (?). Besides, progress has also been made in the research of
- crop identification (??), as well as crop growth status monitoring (?).
- Although the identification of different plant stresses has reached a considerable level of accuracy, the
- pratical plant stress process may be more complicated and the understanding of the spectral images
- of plant stress could be deeper. In order to explore the feasibility of using spectral imaging to identify
- 20 complex plant stress and better apply spectroscopy technology to pratical production, we propose the
- 21 application and research and as follow:
- 1) How can we build a robust classifier to detecte the quality of organically grown broccoli on
- conveyor belts using spetral imaging?
- 24 2) Can we build a robust classifier based on limited data set, using the spectral images collected
- under different combinations of stress (temperature, light, drought)? How can we extend to the
- 26 field?

27

3) Deep into a metabolic levels, can we find the evidence to support the classification?

keywords: Organic farming, Deep learning, Plant stress, Spectral imaging, Detection, Metabolomics

### Methods

- 28 First, use multi-spectral imaging technology combined with deep learning algorithms to detecte the
- <sup>29</sup> quality of organically grown broccoli on conveyor belts.
- <sup>30</sup> Second, build the same detection model using the data collected by the drone.
- Third, under the environment controlled by the laboratory, collect spectral image data under different
- combinations of stress (temperature, light, drought) and explore the possibility of establishing a robust
- 33 classifier in a limited data set.
- Fourth, sample and explore molecular differences by GC-MS, to provide classification basis in metabolic
- 35 levels.

# **Expected Outcomes**


Robust spectral images classifiers and matabolomics analysis.

# **Budget**

\$270 £500 funded by school for computing hours, printing costs and for travel related to the project.

# **Project Feasibility**

Feasible equipment for data collecting and timeline as bellow



### References

- R Calderón, Juan Antonio Navas-Cortés, C Lucena, and Pablo J Zarco-Tejada. High-resolution airborne hyperspectral and thermal imagery for early detection of verticillium wilt of olive using fluorescence, temperature and narrow-band spectral indices. *Remote Sensing of Environment*, 139:231–245, 2013.
- Yan-Ru Zhao, Xiaoli Li, Ke-Qiang Yu, Fan Cheng, and Yong He. Hyperspectral imaging for determining pigment contents in cucumber leaves in response to angular leaf spot disease. Scientific reports, 6:27790, 2016.
- J Torres-Sánchez, JM Peña-Barragán, D Gómez-Candón, AI De Castro, and F López-Granados. Imagery from unmanned aerial vehicles for early site specific weed management. In *Precision agriculture13*, pages 193–199. Springer, 2013.
- Andrea S Laliberte and Albert Rango. Texture and scale in object-based analysis of subdecimeter resolution unmanned aerial vehicle (uav) imagery. *IEEE Transactions on Geoscience and Remote Sensing*, 47(3):761–770, 2009.
- Francisco Agüera Vega, Fernando Carvajal Ramírez, Mónica Pérez Saiz, and Francisco Orgaz Rosúa. Multi-temporal imaging using an unmanned aerial vehicle for monitoring a sunflower crop. *Biosystems Engineering*, 132:19–27, 2015.