



Analytical Geometry and Linear Algebra II, Lab 1

The geometry of linear equation

Gaussian elimination

Matrix notation and matrix multiplication

Task 1

Warm up



Rewrite the following system in the matrix form:

$$\begin{cases} 3x + 4y - 2z = 1 \\ 3y - 2z + x = -2 \\ 5x - 7z - 2y = 3 \end{cases}$$

Task 2



Describe geometrically (line, plane, or whole space) all linear combinations of:

1. $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$

2. $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}$

3. $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$

Task 2



Describe geometrically (line, plane, or whole space) all linear combinations of:

1. $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$ *Ans: line*

2. $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}$

3. $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$

Task 2



Describe geometrically (line, plane, or whole space) all linear combinations of:

1. $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$ *Ans: line*

2. $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}$ *Ans: plane*

3. $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$

Task 2



Describe geometrically (line, plane, or whole space) all linear combinations of:

1. $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$ *Ans: line*

2. $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}$ *Ans: plane*

3. $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$ *Ans: whole space*

Task 3



Draw $v = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$ and $w = \begin{bmatrix} -2 \\ 2 \end{bmatrix}$ and $v + w, v - w$ in a single xy plane

Task 4



Explain, why the system is singular?

$$\begin{cases} u + v + w = 2 \\ u + 2v + 3w = 1 \\ v + 2w = 0 \end{cases}$$

What value should replace that last zero on the right side, to allow the equations to have solutions, and what is one of the solutions?

What does it mean, singular?

Lec 1, page 11



Example 2. Singular (incurable)

$$\begin{cases} u + v + w = \\ 2u + 2v + 5w = \\ 4u + 4v + 8w = \end{cases} \Rightarrow \begin{cases} u + v + w = \\ 3w = \\ 4w = \end{cases}$$

There is no exchange of equations that can avoid zero in the second pivot position. The equations themselves may be solvable or unsolvable. If the last two equations are $3w=6$ and $4w=7$, there is no solution. If those two equations happen to be consistent as in $3w=6$ and $4w=8$ then this singular case has an infinity of solutions. We know that $w=2$, but the first equation cannot decide both u and v .



What does it mean, pivot point?

Definition

The pivot or pivot element is the element of a matrix, or an array, which is selected first on a particular step by an algorithm (e.g. Gaussian elimination, simplex algorithm, etc.), to do certain calculations.

Task 4



Explain, why the system is singular?

$$\begin{cases} u + v + w = 2 \\ u + 2v + 3w = 1 \\ v + 2w = 0 \end{cases}$$

What value should replace that last zero on the right side, to allow the equations to have solutions, and what is one of the solutions?

Task 5



1. Choose a coefficient b that makes this system singular.
2. Then choose a right-hand side g that makes it solvable.
3. Find two solutions in that singular case.

$$\begin{cases} 2x + by = 16 \\ 4x + 8y = g \end{cases}$$



Task 6

Give 3x3 examples (not just the zero matrix):

1. a diagonal matrix: $a_{ij} = 0$, if $i \neq j$;
2. a symmetric matrix: $a_{ij} = a_{ji}$ for all i and j ;
3. an upper triangular matrix: $a_{ij} = 0$, if $i > j$;
4. a skew-symmetric matrix: $a_{ij} = -a_{ji}$ for all i and j .

Task 7



Obtain a Reduced Row Echelon Form (rref) of

$$\begin{cases} 2u + 3v + 0w = 0 \\ 4u + 5v + w = 3 \\ 2u - 1v - 3w = 5 \end{cases}$$

Reference material



- Lectures 1 – 3