



All these steps use the λ 's and the x 's. This section solves the constant coefficient problems that turn into linear algebra. It clarifies these simplest but most important differential equations—whose solution is completely based on growth factors $e^{\lambda t}$.

Second Order Equations

The most important equation in mechanics is $my'' + by' + ky = 0$. The first term is the mass m times the acceleration $a = y''$. This term ma balances the force F (that is *Newton's Law*). The force includes the damping $-by'$ and the elastic force $-ky$, proportional to distance moved. This is a second-order equation because it contains the second derivative $y'' = d^2y/dt^2$. It is still linear with constant coefficients m, b, k .

In a differential equations course, the method of solution is to substitute $y = e^{\lambda t}$. Each derivative of y brings down a factor λ . We want $y = e^{\lambda t}$ to solve the equation:

$$m \frac{d^2y}{dt^2} + b \frac{dy}{dt} + ky = 0 \quad \text{becomes} \quad (m\lambda^2 + b\lambda + k)e^{\lambda t} = 0. \quad (8)$$

Everything depends on $m\lambda^2 + b\lambda + k = 0$. This equation for λ has two roots λ_1 and λ_2 . Then the equation for y has two pure solutions $y_1 = e^{\lambda_1 t}$ and $y_2 = e^{\lambda_2 t}$. Their combinations $c_1y_1 + c_2y_2$ give the complete solution unless $\lambda_1 = \lambda_2$.

In a linear algebra course we expect matrices and eigenvalues. Therefore we turn the scalar equation (with y'') into a *vector equation for y and y'* : first derivative only. Suppose the mass is $m = 1$. Two equations for $u = (y, y')$ give $du/dt = Au$:

$$\begin{aligned} dy/dt &= y' \\ dy'/dt &= -ky - by' \end{aligned} \quad \text{converts to} \quad \frac{d}{dt} \begin{bmatrix} y \\ y' \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -k & -b \end{bmatrix} \begin{bmatrix} y \\ y' \end{bmatrix} = Au. \quad (9)$$

The first equation $dy/dt = y'$ is trivial (but true). The second is equation (8) connecting y'' to y' and y . Together they connect u' to u . So we solve $u' = Au$ by eigenvalues of A :

$$A - \lambda I = \begin{bmatrix} -\lambda & 1 \\ -k & -b - \lambda \end{bmatrix} \quad \text{has determinant} \quad \lambda^2 + b\lambda + k = 0.$$

The equation for the λ 's is the same as (8)! It is still $\lambda^2 + b\lambda + k = 0$, since $m = 1$. The roots λ_1 and λ_2 are now *eigenvalues of A* . The eigenvectors and the solution are

$$x_1 = \begin{bmatrix} 1 \\ \lambda_1 \end{bmatrix} \quad x_2 = \begin{bmatrix} 1 \\ \lambda_2 \end{bmatrix} \quad u(t) = c_1 e^{\lambda_1 t} \begin{bmatrix} 1 \\ \lambda_1 \end{bmatrix} + c_2 e^{\lambda_2 t} \begin{bmatrix} 1 \\ \lambda_2 \end{bmatrix}.$$

The first component of $u(t)$ has $y = c_1 e^{\lambda_1 t} + c_2 e^{\lambda_2 t}$ —the same solution as before. It can't be anything else. In the second component of $u(t)$ you see the velocity dy/dt . The vector problem is completely consistent with the scalar problem. The 2 by 2 matrix A is called a *companion matrix*—a companion to the second order equation with y'' .