

Supplementary Table 3. Pneumococcal shedding isolates antibiogram among HIV uninfected adults

Characteristic	Aerosol shedding vs Nasopharyngeal carriage			Mechanical shedding vs Nasopharyngeal carriage		
	Aerosol shedding, N = 7 <sup>1</sup>	Nasopharyngeal carriage, N = 7 <sup>1</sup>	p-value <sup>2</sup>	Mechanical shedding, N = 8 <sup>1</sup>	Nasopharyngeal carriage, N = 8 <sup>1</sup>	p-value <sup>2</sup>
<b>Tetracycline</b>	6 (86%)	5 (71%)	>0.9	6 (75%)	6 (75%)	>0.9
<b>Erythromycin</b>	6 (86%)	4 (57%)	0.6	6 (75%)	6 (75%)	>0.9
<b>Cotrimoxazole</b>	6 (86%)	5 (71%)	>0.9	8 (100%)	7 (88%)	>0.9
<b>Benzylpenicillin</b> <sup>3</sup>	4 (57%)	5 (71%)	>0.9	3 (38%)	3 (38%)	>0.9
<b>MDR</b>	6 (86%)	5 (71%)	>0.9	6 (75%)	6 (75%)	>0.9

<sup>1</sup> n (%)

<sup>2</sup> Fisher's exact test

<sup>3</sup> i.e. Minimum inhibitory concentration