


Question ID 145337bc

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: 145337bc



Note: Figure not drawn to scale.

In the triangle shown, what is the value of x ?

- A. 24
- B. 45
- C. 48
- D. 69

ID: 145337bc Answer

Correct Answer:
A

Rationale

Choice A is correct. Since the two acute angles have the same measure and the third angle is a right angle, the triangle shown is an isosceles right triangle. In an isosceles right triangle, the two legs have the same length. The figure shows that the length of one leg of the triangle is **24** and the length of the other leg of the triangle is x . It follows that the value of x is **24**.

Choice B is incorrect. This is the measure, in degrees, of one of the angles shown.

Choice C is incorrect and may result from conceptual errors.


Choice D is incorrect and may result from conceptual errors.

Question Difficulty:
Easy

Question ID c9f8d1e9

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: c9f8d1e9



Note: Figure not drawn to scale.

For the right triangle shown, $a = 4$ and $b = 5$. Which expression represents the value of c ?

- A. $4 + 5$
- B. $\sqrt{(4)(5)}$
- C. $\sqrt{4 + 5}$
- D. $\sqrt{4^2 + 5^2}$

ID: c9f8d1e9 Answer

Correct Answer:
D

Rationale

Choice D is correct. By the Pythagorean theorem, if a right triangle has a hypotenuse with length c and legs with lengths a and b , then $c^2 = a^2 + b^2$. In the right triangle shown, the hypotenuse has length c and the legs have lengths a and b . It's given that $a = 4$ and $b = 5$. Substituting 4 for a and 5 for b in the Pythagorean theorem yields $c^2 = 4^2 + 5^2$. Taking the square root of both sides of this equation yields $c = \pm\sqrt{4^2 + 5^2}$. Since the length of a side of a triangle must be positive, the value of c is $\sqrt{4^2 + 5^2}$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.


Choice C is incorrect and may result from conceptual or calculation errors.

Question Difficulty:
Easy

Question ID 36661021

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: 36661021



Note: Figure not drawn to scale.

In the right triangle shown, which of the following is closest to the value of c ?

- A. 7.5
- B. 10.8
- C. 15
- D. 58.5

ID: 36661021 Answer

Correct Answer:
B

Rationale

Choice B is correct. By the Pythagorean theorem, if a right triangle has a hypotenuse with length t and legs with lengths r and s , then $r^2 + s^2 = t^2$. It's given in the right triangle shown that the legs have lengths of 9 and 6 and the hypotenuse has a length of c . Substituting 9 for r , 6 for s , and c for t in $r^2 + s^2 = t^2$ yields $9^2 + 6^2 = c^2$, or $117 = c^2$. Taking the square root of both sides of this equation yields $\pm\sqrt{117} = c$. Since the length of a side of a triangle must be positive, the value of c is $\sqrt{117}$, which is approximately equal to 10.8167. Of the choices, 10.8 is the closest to the value of c .

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:
Easy

Question ID 379ffefb

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: 379ffefb

A right triangle has legs with lengths of **11** centimeters and **9** centimeters. What is the length of this triangle's hypotenuse, in centimeters?

- A. $\sqrt{40}$

B. $\sqrt{202}$

C. 20

D. 202

ID: 379ffefb Answer

Correct Answer:
B

Rationale

Choice B is correct. The Pythagorean theorem states that for a right triangle, $c^2 = a^2 + b^2$, where c represents the length of the hypotenuse and a and b represent the lengths of the legs. It's given that a right triangle has legs with lengths of **11** centimeters and **9** centimeters. Substituting **11** for a and **9** for b in the formula $c^2 = a^2 + b^2$ yields $c^2 = 11^2 + 9^2$, which is equivalent to $c^2 = 121 + 81$, or $c^2 = 202$. Taking the square root of each side of this equation yields $c = \pm\sqrt{202}$. Since c represents a length, c must be positive. Therefore, the length of the triangle's hypotenuse, in centimeters, is $\sqrt{202}$.

Choice A is incorrect. This is the result of solving the equation $c^2 = 11(2) + 9(2)$, not $c^2 = 11^2 + 9^2$.

Choice C is incorrect. This is the result of solving the equation $c(2) = 11(2) + 9(2)$, not $c^2 = 11^2 + 9^2$.


Choice D is incorrect. This is the result of solving the equation $c = 11^2 + 9^2$, not $c^2 = 11^2 + 9^2$.

Question Difficulty:
Easy

Question ID a6097ec2

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: a6097ec2



Note: Figure not drawn to scale.

In the right triangle shown, what is the value of $\sin x^\circ$?

- A. $\frac{1}{73}$
- B. $\frac{10}{73}$
- C. $\frac{63}{73}$
- D. $\frac{136}{73}$

ID: a6097ec2 Answer

Correct Answer:

C

Rationale

Choice C is correct. The sine of an acute angle in a right triangle is the ratio of the length of the side opposite that angle to the length of the hypotenuse. In the right triangle shown, it's given that the length of the side opposite the angle with measure x° is **63** units and the length of the hypotenuse is **73** units. Therefore, the value of $\sin x^\circ$ is $\frac{63}{73}$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Easy

Question ID bbaac300

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: bbaac300

Triangle ABC is similar to triangle DEF , where angle A corresponds to angle D , and angles C and F are right angles. If $\cos B = \frac{1}{22}$, what is the value of $\cos E$?

- A. $\frac{1}{22}$
- B. $\frac{1}{23}$
- C. $\frac{21}{22}$
- D. $\frac{22}{23}$

ID: bbaac300 Answer

Correct Answer:

A

Rationale

Choice A is correct. The cosine of an acute angle in a right triangle is defined as the ratio of the length of the leg adjacent to that angle to the length of the hypotenuse. It's given that angle C is a right angle in triangle ABC and that angle F is a right angle in triangle DEF . Therefore, $\cos B$ is equal to the ratio of the length of side BC to the length of side AB , and $\cos E$ is equal to the ratio of the length of side EF to the length of side DE . It's also given that triangle ABC is similar to triangle DEF , where angle A corresponds to angle D . Since similar triangles have proportional side lengths, $\frac{BC}{AB} = \frac{EF}{DE}$. Therefore, the value of $\cos B$ is equal to the value of $\cos E$. Since $\cos B = \frac{1}{22}$, the value of $\cos E$ is $\frac{1}{22}$.

Choice B is incorrect and may result from conceptual errors.

Choice C is incorrect and may result from conceptual errors.

Choice D is incorrect and may result from conceptual errors.


Question Difficulty:

Easy

Question ID 64c1f044

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: 64c1f044



Note: Figure not drawn to scale.

In the triangle shown, what is the value of $\tan x^\circ$?

- A. $\frac{1}{26}$
- B. $\frac{19}{26}$
- C. $\frac{26}{7}$
- D. $\frac{33}{7}$

ID: 64c1f044 Answer

Correct Answer:

C

Rationale

Choice C is correct. The tangent of an acute angle in a right triangle is defined as the ratio of the length of the side opposite the angle to the length of the shorter side adjacent to the angle. In the triangle shown, the length of the side opposite the angle with measure x° is **26** units and the length of the side adjacent to the angle with measure x° is **7** units. Therefore, the value of $\tan x^\circ$ is $\frac{26}{7}$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.


Question Difficulty:

Easy

Question ID b0c5ece5

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: b0c5ece5



Note: Figure not drawn to scale.

Which equation shows the relationship between the side lengths of the given triangle?

- A. $4b = 19$
- B. $4 + b = 19$
- C. $4^2 + b^2 = 19^2$
- D. $4^2 - b^2 = 19^2$

ID: b0c5ece5 Answer

Correct Answer:
C

Rationale

Choice C is correct. The Pythagorean theorem states that in a right triangle, the sum of the squares of the lengths of the two legs is equal to the square of the length of the hypotenuse. Therefore, $a^2 + b^2 = c^2$, where a and b are the lengths of the legs and c is the length of the hypotenuse. For the given right triangle, the lengths of the legs are 4 and b , and the length of the hypotenuse is 19 . Substituting 4 for a and 19 for c in the equation $a^2 + b^2 = c^2$ yields $4^2 + b^2 = 19^2$. Thus, the relationship between the side lengths of the given triangle is $4^2 + b^2 = 19^2$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.


Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:
Easy

Question ID e6f2ace7

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: e6f2ace7



Note: Figure not drawn to scale.

The lengths of the legs of a right triangle are shown. Which of the following is closest to the length of the triangle's hypotenuse?

- A. 3.2
- B. 5
- C. 7.6
- D. 20

ID: e6f2ace7 Answer

Correct Answer:
C

Rationale

Choice C is correct. The Pythagorean theorem states that for a right triangle, $a^2 + b^2 = c^2$, where a and b represent the lengths of the legs of the triangle and c represents the length of its hypotenuse. In the triangle shown, the legs have lengths of 3 and 7. Substituting 3 for a and 7 for b in the equation $a^2 + b^2 = c^2$ yields $3^2 + 7^2 = c^2$, which is equivalent to $9 + 49 = c^2$, or $58 = c^2$. Taking the positive square root of both sides of this equation yields $\sqrt{58} = c$. Thus, the value of c is approximately 7.6. Therefore, of the given choices, 7.6 is the closest to the length of the triangle's hypotenuse.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.


Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:
Easy

Question ID 87a9a2d4

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Right triangles and trigonometry	

ID: 87a9a2d4



Note: Figure not drawn to scale.

In the right triangle shown, what is the value of $\sin A$?

- A. $\frac{1}{171}$
- B. $\frac{35}{171}$
- C. $\frac{171}{35}$
- D. 171

ID: 87a9a2d4 Answer

Correct Answer:

B

Rationale

Choice B is correct. The sine of an acute angle in a right triangle is the ratio of the length of the side opposite that angle to the length of the hypotenuse. The hypotenuse of a right triangle is the side opposite the right angle. In right triangle ABC , side BC is the side opposite angle A and side AB is the hypotenuse. It's given that the length of side BC is 35 units and the length of side AB is 171 units. Therefore, the value of $\sin A$ is $\frac{35}{171}$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect. This is the ratio of the length of the hypotenuse to the length of the side opposite angle A rather than the ratio of the length of the side opposite angle A to the length of the hypotenuse.

Choice D is incorrect. This is the length of the hypotenuse rather than $\sin A$.

Question Difficulty:

Easy