

Question ID 628300a9

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 628300a9

A science teacher is preparing the 5 stations of a science laboratory. Each station will have either Experiment A materials or Experiment B materials, but not both. Experiment A requires 6 teaspoons of salt, and Experiment B requires 4 teaspoons of salt. If x is the number of stations that will be set up for Experiment A and the remaining stations will be set up for Experiment B, which of the following expressions represents the total number of teaspoons of salt required?

- A. $5x$
- B. $10x$
- C. $2x + 20$
- D. $10x + 20$

ID: 628300a9 Answer

Correct Answer:
C

Rationale

Choice C is correct. It is given that x represents the number of stations that will be set up for Experiment A and that there will be 5 stations total, so it follows that $5 - x$ is the number of stations that will be set up for Experiment B. It is also given that Experiment A requires 6 teaspoons of salt and that Experiment B requires 4 teaspoons of salt, so the total number of teaspoons of salt required is $6x + 4(5 - x)$, which simplifies to $2x + 20$.

Choices A, B, and D are incorrect and may be the result of not understanding the description of the context.

Question Difficulty:
Hard

Question ID f5ff91b2

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: f5ff91b2

If $\frac{x-5}{7} = \frac{x-5}{9}$, the value of $x - 5$ is between which of the following pairs of values?

- A. -9 and -7
- B. -3 and 3
- C. 4.5 and 5.5
- D. 6.75 and 9.25

ID: f5ff91b2 Answer

Correct Answer:

B

Rationale

Choice B is correct. Multiplying both sides of the given equation by $(7)(9)$, or **63**, yields $(63)\left(\frac{x-5}{7}\right) = (63)\left(\frac{x-5}{9}\right)$, or $9(x - 5) = 7(x - 5)$. Subtracting $7(x - 5)$ from both sides of this equation yields $2(x - 5) = 0$. Dividing both sides of this equation by **2** yields $x - 5 = 0$. Therefore, if $\frac{x-5}{7} = \frac{x-5}{9}$, then the value of $x - 5$ is **0**. It follows that of the given choices, the value of $x - 5$ is between -3 and **3**.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID 429fb7c0

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 429fb7c0

What value of t is the solution to the equation $0.8t - 0.46 = 8(t - 0.001) + 1.9$?

ID: 429fb7c0 Answer

Correct Answer:
-.3266, -.3267, -49/150

Rationale

The correct answer is **-.3267**. Applying the distributive property to the right-hand side of the given equation yields $0.8t - 0.46 = 8t - 0.008 + 1.9$, or $0.8t - 0.46 = 8t + 1.892$. Subtracting $0.8t$ from both sides of this equation yields $-0.46 = 7.2t + 1.892$. Subtracting 1.892 from both sides of this equation yields $-2.352 = 7.2t$. Dividing both sides of this equation by 7.2 yields $\frac{-2.352}{7.2} = t$. Therefore, the value of t is approximately **-0.32667**. Note that -.3267, -.3266, -0.326, and -0.327 are examples of ways to enter a correct answer.

Question Difficulty:
Hard

Question ID 3f8a701b

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 3f8a701b

The equation $9x + 5 = a(x + b)$, where a and b are constants, has no solutions. Which of the following must be true?

- I. $a = 9$
 - II. $b = 5$
 - III. $b \neq \frac{5}{9}$
- A. None
- B. I only
- C. I and II only
- D. I and III only

ID: 3f8a701b Answer

Correct Answer:
D

Rationale

Choice D is correct. For a linear equation in a form $ax + b = cx + d$ to have no solutions, the x-terms must have equal coefficients and the remaining terms must not be equal. Expanding the right-hand side of the given equation yields $9x + 5 = ax + ab$.

Inspecting the x-terms, 9 must equal a, so statement I must be true. Inspecting the remaining terms, 5 can't equal $9b$. Dividing

both of these quantities by 9 yields that b can't equal $\frac{5}{9}$. Therefore, statement III must be true. Since b can have any value other than $\frac{5}{9}$, statement II may or may not be true.

Choice A is incorrect. For the given equation to have no solution, both $a = 9$ and $b \neq \frac{5}{9}$ must be true. Choice B is incorrect because it must also be true that $b \neq \frac{5}{9}$. Choice C is incorrect because when $a = 9$, there are many values of b that lead to an equation having no solution. That is, b might be 5, but b isn't required to be 5.

Question Difficulty:
Hard

Question ID 4771a64a

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 4771a64a

If $5 - 7(2 - 4x) = 16 - 8(2 - 4x)$, what is the value of $2 - 4x$?

ID: 4771a64a Answer

Correct Answer:
11

Rationale

The correct answer is **11**. Subtracting **5** from each side of the given equation yields $-7(2 - 4x) = 11 - 8(2 - 4x)$. Adding $8(2 - 4x)$ to each side of this equation yields $2 - 4x = 11$. Therefore, the value of $2 - 4x$ is **11**.

Question Difficulty:
Hard

Question ID 0393ba15

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 0393ba15

$-49x = -98x$

How many solutions does the given equation have?

- A. Zero
- B. Exactly one
- C. Exactly two
- D. Infinitely many

ID: 0393ba15 Answer

Correct Answer:

B

Rationale

Choice B is correct. Adding $98x$ to each side of the given equation yields $49x = 0$. Dividing each side of this equation by 49 yields $x = 0$. This means that 0 is the only solution to the given equation. Therefore, the given equation has exactly one solution.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID aee9fd2d

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: aee9fd2d

If $\frac{x+6}{3} = \frac{x+6}{13}$, the value of $x + 6$ is between which of the following pairs of values?

- A. -7 and -3
- B. -2 and 2
- C. 2 and 7
- D. 8 and 13

ID: aee9fd2d Answer

Correct Answer:

B

Rationale

Choice B is correct. Multiplying both sides of the given equation by $(3)(13)$, or 39 , yields $(39)\left(\frac{x+6}{3}\right) = (39)\left(\frac{x+6}{13}\right)$, or $13(x + 6) = 3(x + 6)$. Subtracting $3(x + 6)$ from both sides of this equation yields $10(x + 6) = 0$. Dividing both sides of this equation by 10 yields $x + 6 = 0$. Therefore, if $\frac{x+6}{3} = \frac{x+6}{13}$, then the value of $x + 6$ is 0 . It follows that of the given choices, the value of $x + 6$ is between -2 and 2 .

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID 153ee763

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 153ee763

$$-3x + 21px = 84$$

In the given equation, p is a constant. The equation has no solution. What is the value of p ?

- A. 0
- B. $\frac{1}{7}$
- C. $\frac{4}{3}$
- D. 4

ID: 153ee763 Answer

Correct Answer:

B

Rationale

Choice B is correct. A linear equation in one variable has no solution if and only if the equation is false; that is, when there is no value of x that produces a true statement. It's given that in the equation $-3x + 21px = 84$, p is a constant and the equation has no solution for x . Therefore, the value of the constant p is one that results in a false equation. Factoring out the common factor of $-3x$ on the left-hand side of the given equation yields $-3x(1 - 7p) = 84$. Dividing both sides of this equation by -3 yields $x(1 - 7p) = -28$. Dividing both sides of this equation by $(1 - 7p)$ yields $x = \frac{-28}{1-7p}$. This equation is false if and only if $1 - 7p = 0$. Adding $7p$ to both sides of $1 - 7p = 0$ yields $1 = 7p$. Dividing both sides of this equation by 7 yields $\frac{1}{7} = p$. It follows that the equation $x = \frac{-28}{1-7p}$ is false if and only if $p = \frac{1}{7}$. Therefore, the given equation has no solution if and only if the value of p is $\frac{1}{7}$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID 0cb57740

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 0cb57740

Each side of a **30**-sided polygon has one of three lengths. The number of sides with length **8 centimeters (cm)** is **5** times the number of sides n with length **3 cm**. There are **6** sides with length **4 cm**. Which equation must be true for the value of n ?

- A. $5n + 6 = 30$
- B. $6n + 6 = 30$
- C. $8n + 3n + 4n = 30$
- D. $8(5n) + 3n + 4(6) = 30$

ID: 0cb57740 Answer

Correct Answer:

B

Rationale

Choice B is correct. It's given that each side of a **30**-sided polygon has one of three lengths. It's also given that the number of sides with length **8 centimeters (cm)** is **5** times the number of sides n with length **3 cm**. Therefore, there are $5 \times n$, or $5n$, sides with length **8 cm**. It's also given that there are **6** sides with length **4 cm**. Therefore, the number of **3 cm**, **4 cm**, and **8 cm** sides are n , **6**, and $5n$, respectively. Since there are a total of **30** sides, the equation $n + 6 + 5n = 30$ represents this situation. Combining like terms on the left-hand side of this equation yields $6n + 6 = 30$. Therefore, the equation that must be true for the value of n is $6n + 6 = 30$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID 69f609b2

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 69f609b2

How many solutions does the equation $12(x - 3) = -3(x + 12)$ have?

- A. Exactly one
- B. Exactly two
- C. Infinitely many
- D. Zero

ID: 69f609b2 Answer

Correct Answer:

A

Rationale

Choice A is correct. Distributing **12** on the left-hand side and **−3** on the right-hand side of the given equation yields $12x - 36 = -3x - 36$. Adding **3x** to each side of this equation yields $15x - 36 = -36$. Adding **36** to each side of this equation yields $15x = 0$. Dividing each side of this equation by **15** yields $x = 0$. This means that **0** is the only solution to the given equation. Therefore, the given equation has exactly one solution.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID 771bd0ca

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 771bd0ca

$5(t + 3) - 7(t + 3) = 38$

What value of t is the solution to the given equation?

ID: 771bd0ca Answer

Correct Answer:

-22

Rationale

The correct answer is -22 . The given equation can be rewritten as $-2(t + 3) = 38$. Dividing both sides of this equation by -2 yields $t + 3 = -19$. Subtracting 3 from both sides of this equation yields $t = -22$. Therefore, -22 is the value of t that is the solution to the given equation.

Question Difficulty:

Hard

Question ID 7d5d1b32

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 7d5d1b32

$$2(kx - n) = -\frac{28}{15}x - \frac{36}{19}$$

In the given equation, k and n are constants and $n > 1$. The equation has no solution. What is the value of k ?

ID: 7d5d1b32 Answer

Correct Answer:

-.9333, -14/15

Rationale

The correct answer is $-\frac{14}{15}$. A linear equation in the form $ax + b = cx + d$ has no solution only when the coefficients of x on each side of the equation are equal and the constant terms are not equal. Dividing both sides of the given equation by 2 yields $kx - n = -\frac{28}{30}x - \frac{36}{38}$, or $kx - n = -\frac{14}{15}x - \frac{18}{19}$. Since it's given that the equation has no solution, the coefficient of x on both sides of this equation must be equal, and the constant terms on both sides of this equation must not be equal. Since $\frac{18}{19} < 1$, and it's given that $n > 1$, the second condition is true. Thus, k must be equal to $-\frac{14}{15}$. Note that -14/15, -.9333, and -0.933 are examples of ways to enter a correct answer.

Question Difficulty:

Hard

Question ID e6cb2402

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: e6cb2402

$$3(kx + 13) = \frac{48}{17}x + 36$$

In the given equation, k is a constant. The equation has no solution. What is the value of k ?

ID: e6cb2402 Answer

Correct Answer:
.9411, .9412, 16/17

Rationale

The correct answer is $\frac{16}{17}$. It's given that the equation $3(kx + 13) = \frac{48}{17}x + 36$ has no solution. A linear equation in the form $ax + b = cx + d$, where a, b, c , and d are constants, has no solution only when the coefficients of x on each side of the equation are equal and the constant terms aren't equal. Dividing both sides of the given equation by 3 yields $kx + 13 = \frac{48}{51}x + \frac{36}{3}$, or $kx + 13 = \frac{16}{17}x + 12$. Since the coefficients of x on each side of the equation must be equal, it follows that the value of k is $\frac{16}{17}$. Note that 16/17, .9411, .9412, and 0.941 are examples of ways to enter a correct answer.

Question Difficulty:
Hard

Question ID 25e1cfed

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 25e1cfed

How many solutions does the equation $10(15x - 9) = -15(6 - 10x)$ have?

- A. Exactly one
- B. Exactly two
- C. Infinitely many
- D. Zero

ID: 25e1cfed Answer

Correct Answer:

C

Rationale

Choice C is correct. Applying the distributive property to each side of the given equation yields $150x - 90 = -90 + 150x$. Applying the commutative property of addition to the right-hand side of this equation yields $150x - 90 = 150x - 90$. Since the two sides of the equation are equivalent, this equation is true for any value of x . Therefore, the given equation has infinitely many solutions.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty:

Hard

Question ID b7e6394d

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: b7e6394d

Alan drives an average of 100 miles each week. His car can travel an average of 25 miles per gallon of gasoline. Alan would like to reduce his weekly expenditure on gasoline by \$5. Assuming gasoline costs \$4 per gallon, which equation can Alan use to determine how many fewer average miles, m , he should drive each week?

- A. $\frac{25}{4}m = 95$
- B. $\frac{25}{4}m = 5$
- C. $\frac{4}{25}m = 95$
- D. $\frac{4}{25}m = 5$

ID: b7e6394d Answer

Correct Answer:
D

Rationale

Choice D is correct. Since gasoline costs \$4 per gallon, and since Alan’s car travels an average of 25 miles per gallon, the expression $\frac{4}{25}$ gives the cost, in dollars per mile, to drive the car. Multiplying $\frac{4}{25}$ by m gives the cost for Alan to drive m miles in his car. Alan wants to reduce his weekly spending by \$5, so setting $\frac{4}{25}m$ equal to 5 gives the number of miles, m , by which he must reduce his driving.

Choices A, B, and C are incorrect. Choices A and B transpose the numerator and the denominator in the fraction. The fraction $\frac{25}{4}$ would result in the unit miles per dollar, but the question requires a unit of dollars per mile. Choices A and C set the expression equal to 95 instead of 5, a mistake that may result from a misconception that Alan wants to reduce his driving by 5 miles each week; instead, the question says he wants to reduce his weekly expenditure by \$5.

Question Difficulty:
Hard

Question ID 38bf4e04

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 38bf4e04

A factory makes 9-inch, 7-inch, and 4-inch concrete screws. During a certain day, the number of 9-inch concrete screws that the factory makes is 5 times the number n of 7-inch concrete screws, and the number of 4-inch concrete screws is 22. During this day, the factory makes 100 concrete screws total. Which equation represents this situation?

- A. $9(5n) + 7n + 4(22) = 100$
- B. $9n + 7n + 4n = 100$
- C. $5n + 22 = 100$
- D. $6n + 22 = 100$

ID: 38bf4e04 Answer

Correct Answer:
D

Rationale

Choice D is correct. It's given that during a certain day at a factory, the number of 7-inch concrete screws the factory makes is n and the number of 4-inch concrete screws the factory makes is 22. It's also given that during this day the number of 9-inch concrete screws the factory makes is 5 times the number of 7-inch concrete screws, or $5n$. Therefore, the total number of 7-inch, 9-inch, and 4-inch concrete screws is $n + 5n + 22$, or $6n + 22$. It's given that during this day, the factory makes 100 concrete screws total. Thus, the equation $6n + 22 = 100$ represents this situation.

Choice A is incorrect. This equation represents a situation where the total length, in inches, of all the concrete screws, rather than the total number of concrete screws, is 100.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect. This equation represents a situation where the total number of 9-inch concrete screws and 4-inch concrete screws, not including the 7-inch concrete screws, is 100.

Question Difficulty:
Hard

Question ID f14484a5

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: f14484a5

A manufacturing plant makes 10-inch, 9-inch, and 7-inch frying pans. During a certain day, the number of 10-inch frying pans that the manufacturing plant makes is 4 times the number n of 9-inch frying pans it makes, and the number of 7-inch frying pans it makes is 10. During this day, the manufacturing plant makes 100 frying pans total. Which equation represents this situation?

- A. $10(4n) + 9n + 7(10) = 100$
- B. $10n + 9n + 7n = 100$
- C. $4n + 10 = 100$
- D. $5n + 10 = 100$

ID: f14484a5 Answer

Correct Answer:
D

Rationale

Choice D is correct. It's given that during a certain day, the number of 9-inch frying pans the manufacturing plant makes is n and the number of 7-inch frying pans it makes is 10. It's also given that during this day the number of 10-inch frying pans that the manufacturing plant makes is 4 times the number of 9-inch frying pans, or $4n$. Therefore, the total number of 7-inch, 9-inch, and 10-inch frying pans the manufacturing plant makes is $n + 10 + 4n$, or $5n + 10$. It's given that during this day the manufacturing plant makes 100 frying pans total. Thus, the equation $5n + 10 = 100$ represents this situation.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Question Difficulty:
Hard

Question ID 2937ef4f

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: 2937ef4f

Hector used a tool called an auger to remove corn from a storage bin at a constant rate. The bin contained 24,000 bushels of corn when Hector began to use the auger. After 5 hours of using the auger, 19,350 bushels of corn remained in the bin. If the auger continues to remove corn at this rate, what is the total number of hours Hector will have been using the auger when 12,840 bushels of corn remain in the bin?

- A. 3
- B. 7
- C. 8
- D. 12

ID: 2937ef4f Answer

Correct Answer:

D

Rationale

Choice D is correct. After using the auger for 5 hours, Hector had removed $24,000 - 19,350 = 4,650$ bushels of corn from the storage bin. During the 5-hour period, the auger removed corn from the bin at a constant rate of $\frac{4,650}{5} = 930$ bushels per hour. Assuming the auger continues to remove corn at this rate, after x hours it will have removed $930x$ bushels of corn. Because the bin contained 24,000 bushels of corn when Hector started using the auger, the equation $24,000 - 930x = 12,840$ can be used to find the number of hours, x , Hector will have been using the auger when 12,840 bushels of corn remain in the bin. Subtracting 12,840 from both sides of this equation and adding $930x$ to both sides of the equation yields $11,160 = 930x$. Dividing both sides of this equation by 930 yields $x = 12$. Therefore, Hector will have been using the auger for 12 hours when 12,840 bushels of corn remain in the storage bin.

Choice A is incorrect. Three hours after Hector began using the auger, $24,000 - 3(930) = 21,210$ bushels of corn remained, not 12,840. Choice B is incorrect. Seven hours after Hector began using the auger, $24,000 - 7(930) = 17,490$ bushels of corn will remain, not 12,840. Choice C is incorrect. Eight hours after Hector began using the auger, $24,000 - 8(930) = 16,560$ bushels of corn will remain, not 12,840.

Question Difficulty:

Hard

Question ID ac472881

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in one variable	

ID: ac472881

$$\frac{12x+28}{4} - \frac{s}{13} = r(x - 8)$$

In the given equation, s and r are constants, and $s > 0$. If the equation has infinitely many solutions, what is the value of s ?

ID: ac472881 Answer

Correct Answer:

403

Rationale

The correct answer is **403**. For a linear equation in one variable to have infinitely many solutions, the coefficients of the variable must be equal on both sides of the equation and the constant terms must also be equal on both sides of the equation. The given equation can be rewritten as $\frac{4(3x+7)}{4} - \frac{s}{13} = r(x - 8)$, or $3x + 7 - \frac{s}{13} = r(x - 8)$. Applying the distributive property to the right-hand side of this equation yields $3x + 7 - \frac{s}{13} = rx - 8r$. For this equation to have infinitely many solutions, the coefficients of x must be equal, so it follows that $3 = r$. Additionally, the constant terms must be equal, which means $7 - \frac{s}{13} = -8r$. Substituting **3** for r in this equation yields $7 - \frac{s}{13} = -8(3)$, or $7 - \frac{s}{13} = -24$. Adding $\frac{s}{13}$ to both sides of this equation yields $7 = -24 + \frac{s}{13}$. Adding **24** to both sides of this equation yields $31 = \frac{s}{13}$. Multiplying both sides of this equation by **13** yields **403** = s . Therefore, if the equation has infinitely many solutions, the value of s is **403**.

Question Difficulty:

Hard