A decorative graphic on the left side of the page, consisting of a network of blue lines and circles. The lines are of varying thickness and connect to circles of different sizes, creating a circuit-like or orbital pattern that extends from the top to the bottom of the page.

SPACE ENGINEERING 3
Assignment 1
24th March 2016

**ORBIT SIMULATION AND
DETERMINATION**

Lydia Drabsch
311217591
ldra3557@uni.sydney.edu.au

**STUDENT PLAGIARISM: COURSE WORK – POLICY AND PROCEDURE****MTRX 2700 COMPLIANCE STATEMENT****INDIVIDUAL / COLLABORATIVE WORK**

I/We certify that:

1. I/We have read and understood the *University of Sydney Student Plagiarism: Coursework Policy and Procedure*;
2. I/We understand that failure to comply with the *University of Sydney Student Plagiarism: Coursework Policy and Procedure* can lead to the University commencing proceedings against me/us for potential student misconduct under Chapter 8 of the *University of Sydney By-Law 1999* (as amended);
3. The Work undertaken in this course is substantially my/our own, and to the extent that any part of this Work is not my/our own I/we have indicated that it is not my/our own by Acknowledging the Source of that part or those parts of the Work.

| Name | Signature | SID | Date |
|-------------------|-----------|-----------|------------|
| Melissa Mitrevski | | 440207636 | 05/06/2015 |
| Meg Flannery | | 440291196 | 05/06/2015 |
| Lydia Drabsch | | 311217591 | 05/06/2015 |

CONTENTS

| | | |
|----------|--|----------|
| 1 | Simulation of Orbits with Classical Elements | 1 |
| 1.1 | Introduction | 1 |
| 1.2 | Methodology | 1 |
| 1.2.1 | Orbital Period | 2 |
| 1.3 | Results/Discussion | 2 |
| 1.3.1 | Van Allen Probes | 2 |
| 1.3.2 | Solar Radiation and Climate Experiment (SORCE) | 2 |
| 1.3.3 | Orbital Properties | 2 |
| 2 | Simulating Perturbations | 3 |
| 2.1 | Introduction | 3 |
| 2.1.1 | Earth oblateness | 3 |
| 2.1.2 | Van Allen Probes | 4 |
| 2.2 | Methodology | 4 |
| 2.3 | Results/Discussion | 4 |
| 3 | Orbital Determination | 5 |
| 3.1 | Introduction | 5 |
| 3.2 | Methodology | 5 |
| 3.3 | Results/Discussion | 5 |
| 4 | Conclusions | 5 |
| 5 | Appendix | 7 |

LIST OF FIGURES

| | | |
|-----|---|---|
| 1.1 | | 3 |
| 2.1 | Classical Elements over 12 days | 4 |
| 2.2 | Equinoctial Elements over 12 days | 5 |

LIST OF TABLES

| | | |
|-----|---|---|
| 1.1 | Orbital properties of simulated satellites. (* indicates data taken directly from TLEs) | 3 |
|-----|---|---|

INTRODUCTION

Each mainQN.m file has a section called 'User Input' where the animations and state plots can be turned on/off. Also the timestep and the number of days to simulate are defined. The default settings are $dt = 100$ seconds and $days = 1$.

1. SIMULATION OF ORBITS WITH CLASSICAL ELEMENTS

1.1 Introduction

- keplers three laws
- perifocal frame
- The true anomaly θ is the angle taken at the focus of the perifocal frame to the satellite from the perigee. The eccentric anomaly E is the angle taken at the centre of perifocal frame to the satellite from the perigee.
- The mean anomaly M_t is the mean number of orbits per day.
- LEO,MEO
- TLE's

1.2 Methodology

From Kepler's second law, the mean anomaly at time t is calculated using the mean motion n from an epoch time described by $M_0(t_0)$.

$$M_t = M_0 + n(t - t_0) \quad (1)$$

To solve for the eccentric anomaly, newtons method was used

$$E_{i+1} = E_i - \frac{f(E_i)}{f'(E_i)} \quad (2)$$

$$E_{i+1} = E_i - \frac{E - e \sin(E_i) - M_t}{1 - e \cos(E_i)} \quad (3)$$

The true anomaly was calculated using

$$\theta = 2 \tan^{-1} \left(\sqrt{\frac{1+e}{1-e}} \tan \left(\frac{E}{2} \right) \right) \quad (4)$$

The general equation for the radius of an ellipse

$$r = \frac{p}{1 + e \cos \theta} \quad (5)$$

The two parameters, θ and r can completely define the orbit in the perifocal frame as polar coordinates. The conversion to cartesian coordinates in the perifocal frame are as follows:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix}_{perifocal} = \begin{bmatrix} r \cos \theta \\ r \sin \theta \\ 0 \end{bmatrix} \quad (6)$$

$$\begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix}_{perifocal} = \begin{bmatrix} -\sqrt{\frac{\mu}{p}} \sin \theta \\ \sqrt{\frac{\mu}{p}} (e + \cos \theta) \\ 0 \end{bmatrix} \quad (7)$$

$$(8)$$

The perifocal parameters were transformed to the ECI frame to animate the 3D model of the satellite orbit around the Earth using the function *orbit2ECI*. From the ECI coordinates, the ECEF and LLHGD coordinates were calculated for the ground trace.

1.2.1 Orbital Period

The period was calculated analytically and computationally. The period is inversely proportional to the mean motion. From the simulation, the ECI coordinates of the satellite were analysed by auto correlation based on the Wiener-Khinchin Theorem. The data was fast fourier transformed using the built-in matlab function, multiplied with the resultant complex conjugate then transformed back to the time domain by inverse fourier transform.

```
%% L Drabsch 24/3/16
% Find period of orbit with autocorrelation based on Wiener-Khinchin Theorem
% Inputs: X = signal to analyse, eg X_ECI
%         dt = timestep (s)
% Output: P = period (hours)

function P = Period_AC(X,dt)
    Fr = fft(X);
    S = Fr.*conj(Fr);
    R = ifft(S);
    [~,loc] = findpeaks(R);
    P = loc(1)*dt/(60*60);
end
```

1.3 Results/Discussion

1.3.1 Van Allen Probes

The satellites Van Allen Probes, previously known as the Radiation Belt Storm Probes (RBSP A and B), are in a highly eccentric orbit. RBSP-A was modelled based on TLEs from spacetrack.com, see Table 1.1. It was launched on 30th August 2012

The Van Allen Belts are regions of plasma that surround the Earth that are contained by the Earth's magnetic field. The highly energetic particles become trapped in the Earth's magnetic field by travelling along the field lines to a pole and being reflected back and travelling to the opposite pole. This phenomenon is called a magnetic mirror, and is the reason the Van Allen belts have a torus shape about the Earth's magnetic axis, see Figure 1.1. Therefore, the radiation is at the highest intensity about the magnetic equatorial plane, currently at 11.5° inclination. The satellite was put in an orbit close to this inclination at about 10° , with the mission objective to be in an orbit of no greater than 18° . The inclination has fluctuated between $9-11^\circ$ over the course of it's mission.

The orbit was highly eccentric in order to cover the entire radiation belt region. The perigee altitude is 500 km, which is below the inner radiation belt and the apogee altitude is 30600 km, which is above the outer edge of the outer radiation belt.

1.3.2 Solar Radiation and Climate Experiment (SORCE)

- long term measurements of total solar irradiance in UV and VNIR - how and why variability occurs at the sun and how it affects our atmosphere and climate - launched in 2003

1.3.3 Orbital Properties




Figure 1.1

Table 1.1: Orbital properties of simulated satellites. (* indicates data taken directly from TLEs)

| Orbital Properties | Variable Name | Units | Van Allen Probe A | SORCE |
|---------------------------------|---------------|----------------|-------------------|------------------|
| NORAD ID | | | | 27651 |
| *Julian year/day data was taken | t0 | year/day | 2016 68.37507029 | 2016 82.69823837 |
| Semimajor axis | a | m | | |
| *Eccentricity | e | - | 0.6813430 | 0.0024311 |
| *Inclination angle | inc | degrees | 10.1687 | 39.9961 |
| *Right ascension of the node | Rasc | degrees | 46.5607 | 317.0592 |
| *Argument of Perigee | omega | degrees | 77.2770 | 301.6655 |
| *Mean Anomaly | Mt | degrees | 346.0491 | 58.1763 |
| *Mean Motion | MM | orbits per day | 2.681033090 | 14.88275952 |
| Analytical Period | | | | |
| Computational Period | | | | |
| Altitude at Perigee | | | | |
| Altitude at Apogee | | | | |

2. SIMULATING PERTURBATIONS

2.1 Introduction

The basic keperian model only accounted for an ideal two body system with no other effects when in reality, other forces act of the satellite. Compared to the gravitational motion between the Earth and the satellite however, these other effects are small. Therefore, they can be mathematically modelled as perturbations. While it is still an approximation, the error is reduced. Types of perturbations include;

- gravitational effects: such as the oblate Earth, third-body interactions such as the Moon, Sun or other planets
- drag forces: from the Earth's atmosphere and from solar radiation pressure
- unexpected thrusting: from malfunctioning thrusters

2.1.1 Earth oblateness

For the standard simulation model in Question 1, it was assumed that the Earth was spherical. A perfectly spherical mass has an inverse square relation of the gravitational field to the force applied

on a body. However, the Earth is slightly oblate, as it is flatter at the poles and wider at the equator than a sphere.

Unlike using the keplerin model of Question 1, the classical orbital parameters change through time.

2.1.2 Van Allen Probes

Due to the highly elliptical orbit of the Van Allen Probes, most of the time the satellite is in MEO away from the Earth's atmosphere. As the inclination of the orbit keeps the satellite close to the equator, the perturbation due to the non-spherical earth

Newtown's second law and his law of gravitation results in

$$\ddot{\mathbf{r}} + \frac{\mu \mathbf{r}}{r^3} = \mathbf{a}_p \quad (9)$$

2.2 Methodology

Equinoctial elements were used to remove the possible singularities that affect classical elements. The

2.3 Results/Discussion

Figure 2.1: Classical Elements over 12 days





Figure 2.2: Equinoctial Elements over 12 days



3. ORBITAL DETERMINATION

3.1 Introduction

3.2 Methodology

The perturbation model from Question 2 was used

3.3 Results/Discussion

<http://vanallenprobes.jhuapl.edu/mission/conversation/overview/index.php>

Three networks are used to communicate with the Van Allen Probes. The main communications ground station is at the Applied Physics Laboratory in Washington, US.

<http://www.scf.jhuapl.edu/>

The LLH coordinates are 39.10N 76.53W 140 m above mean seal level.

For backup, the satellite communicates with the United Space Network using the Near Earth Network with ground stations in Hawaii and Australia. The data rates that can be achieved with this network is high enough to gather scientific data from the probe. Also as a backup, the NASA program Space Network is used for monitoring data, which relays through geosynchronous Tracking and Data Relay Satellites (TDRS). The satellite TDRS-12 was modelled with the basic Keplerian model in Question 1.

The TLEs are collected at approximately the same point in the orbit, at a mean anomaly of 345° .

4. CONCLUSIONS

REFERENCES

5. APPENDIX